Risk-Monotonicity in Statistical Learning

Zakaria Mhammedi[*]
Massachusetts Institute of Technology
mhammedi@mit.edu

Abstract

Acquisition of data is a difficult task in many applications of machine learning, and it is only natural that one hopes and expects the population risk to decrease (better performance) *monotonically* with increasing data points. It turns out, somewhat surprisingly, that this is not the case even for the most standard algorithms that minimize the empirical risk. Non-monotonic behavior of the risk and instability in training have manifested and appeared in the popular deep learning paradigm under the description of *double descent*. These problems highlight the current lack of understanding of learning algorithms and generalization. It is, therefore, crucial to pursue this concern and provide a characterization of such behavior. In this paper, we derive the first consistent and risk-monotonic algorithms for a general statistical learning setting under weak assumptions, consequently resolving an open problem [53] on how to avoid non-monotonic behavior of risk curves. We further show that risk monotonicity need not necessarily come at the price of worse excess risk rates. To achieve this, we derive new empirical Bernstein-like concentration inequalities of independent interest that hold for certain non-i.i.d. processes such as Martingale Difference Sequences.

1 Introduction

Guarantees on the performance of machine learning algorithms are desirable, especially given the widespread deployment. A traditional performance guarantee often takes the form of a generalization bound, where the *expected* risk associated with hypotheses returned by an algorithm is bounded in terms of the corresponding empirical risk plus an additive error which typically converges to zero as the sample size increases. However, interpreting such bounds is not always straight forward and can be somewhat ambiguous. In particular, given that the error term in these bounds goes to zero, it is tempting to conclude that more data would monotonically decrease the expected risk of an algorithm such as the Empirical Risk Minimizer (ERM). However, this is not always the case; for example, [33] showed that increasing the sample size by one, can sometimes make the test performance worse in expectation for commonly used algorithms such as ERM in popular settings including linear regression. This type of non-monotonic behavior is still poorly understood and indeed not a desirable feature of an algorithm since it is expensive to acquire more data in many applications.

Non-monotonic behavior of *risk curves* [47]—the curve of the expected risk as a function of the sample size—has been observed in many previous works [18, 43, 48, 20] (see also [33, 52] for nice accounts of the literature). At least two phenomena have been identified as being the cause behind such behavior. The first one, coined *peaking* [30, 19], or *double descent* according to more recent literature [6, 49, 7, 17, 15, 37, 40, 41, 16, 12, 11, 14, 42], is the phenomenon where the risk curve peaks at a certain sample size n. This sample size typically represents the cross-over point from an over-parameterized to under-parameterized model. For example, when the number of data points is less than the number of parameters of a model (over-parameterized model), such as Neural

---

[*]Work done while at the Australian National University.

35th Conference on Neural Information Processing Systems (NeurIPS 2021), Sydney, Australia.

Networks, the expected risk can typically increase until the number of data points exceeds the number of parameters (under-parameterized model). The second phenomenon is known as *dipping* [32, 31], where the risk curve reaches a minimum at a certain sample size n and increases after that—never reaching the minimum again even for very large n. This phenomenon typically happens when the algorithm is trained on a surrogate loss that differs from the one used to evaluate the risk [8].

It is becoming more apparent that the two phenomena just mentioned (double descent and dipping) do not fully characterize when non-monotonic risk behavior occurs [34]. [33] showed that non-monotonic risk behavior could happen outside these settings and formally prove that the risk curve of ERM is non-monotonic in linear regression with prevalent losses. The most striking aspect of their findings is that the risk curves in some of the cases they study can display a perpetual "oscillating" behavior; there is no sample size beyond which the risk curve becomes monotone—see Figure 1. In such cases, the risk's non-monotonicity cannot be attributed to the peaking/double descent phenomenon. Moreover, they rule out the dipping phenomenon by studying the ERM on the actual loss (not a surrogate loss).

Figure 1: Expected risk of ERM on a 1d linear regression problem with absolute loss and two instances $z_1 = (x_1, y_1) = (1, 1)$ and $z_2 = (x_2, y_2) = (1/10, 1)$ such that $\mathbf{P}[Z = z_1] = 0.1$ and $\mathbf{P}[Z = z_2] = 0.9$. The set of hypotheses is the real line, i.e. $\mathcal{H} = \mathbb{R}$. The ERM solution $\hat{h}_n$ admits a closed form in this case—see [33] for details.

The findings of [33] stress our current lack of understanding of generalization. This was echoed more particularly by [53], who posed the following question as part of a COLT open problem:

How can we provably avoid non-monotonic behavior?

While excess risk bounds are typically monotonic, this does not guarantee the monotonicity of the actual risk. In this work, we study under which assumptions on the learning problem there exist consistent and risk monotonic algorithms. We also aim to quantify the price to pay, in terms of corresponding excess risk rates, for achieving risk monotonicity.

Contributions.   In this work, we answer the question posed by [53] by presenting an algorithm that is both consistent and risk-monotonic under weak assumptions on the learning problem. Our algorithm is technically a "wrapper" that takes as input any base learning algorithm B and makes up a new algorithm A that is risk monotonic and enjoys essentially the same excess risk rate as B. Crucially, our results show that risk monotonicity need not come at the expense of worse excess risk rates. In fact, we show that fast rates are achievable under a Bernstein condition (Definition 3).

Our results hold under the general statistical learning setting with a bounded loss. We even go beyond the standard i.i.d. assumption on the loss process. Our relaxed technical condition on the loss process, which is formalized in Assumption 1 below, is reminiscent of the condition characterizing Martingale Difference Sequences (MDS). In a nutshell, we will assume a setting where the instance random variables $Z_1, Z_2, \ldots$ and the loss ℓ satisfy, for all hypotheses h, $\mathbf{E}[\ell(h, Z_t) \mid Z_1, \ldots, Z_{t-1}] = L(h)$, for some risk function L. This is trivially satisfied in the i.i.d. case, where L corresponds to the standard risk function. In general, this condition may be satisfied even if $Z_1, Z_2, \ldots$ are dependent or have different marginal distributions. We argue that our relaxed assumption on the loss process is the weakest assumption under which studying risk-monotonicity still makes sense.

To achieve risk monotonicity under our loss process assumption, we derive a new concentration inequality/generalization bound of PAC-Bayesian flavor for MDS (see Proposition 5). This concentration inequality may be thought of as an empirical Freedman's inequality [23] or as an extension of the empirical Bernstein inequality [35] to MDS. Our concentration inequalities also have the advantage of being time-uniform with the optimal dependence on the number of samples. Here, time-uniform means that the inequalities hold for all sample sizes simultaneously. While standard concentration inequalities can be turned into time-uniform ones using a union bound over the number of samples

n, the resulting bounds will have a sub-optimal $\ln n$ factor instead of the optimal[2] $\ln \ln n$ that we are able to get. Finally, our concentration bounds are easily derived using the guarantee of a recent parameter-free online learning algorithm—FREEGRAD[39]. Our approach opens up the door for obtaining new concentration inequalities through the design of online learning algorithms.

Approach Overview.   Our approach to deriving the new concentration inequalities is based on the guarantee of the recent FREEGRAD algorithm. The algorithm operates in rounds, where at each round t, FREEGRAD outputs $\widehat{w}_t$ in some convex set $\mathcal{W}$, say $\mathbb{R}^d$, then observes a vector $g_t \in \mathbb{R}^d$, typically the sub-gradient of a loss function at the iterate $\widehat{w}_t$. The algorithm guarantees a regret bound of the form $\sum_{t=1}^T g_t^\top (\widehat{w}_t - w) \leq \widetilde{O}(\|w\|\sqrt{Q_T})$, for all $w \in \mathcal{W}$, where $Q_T := \sum_{t=1}^T \|g_t\|^2$. What is more, FREEGRAD's outputs $(\widehat{w}_t)$ ensure the following (see [39, Theorem 5]):

$$\widehat{w}_t^\top g_t + \Phi(S_t, Q_t) \leq \Phi(S_{t-1}, Q_{t-1}), \quad \forall t \geq 1, \tag{1}$$

where $S_t := \|\sum_{i=1}^t g_i\|$, $Q_t := \sum_{i=1}^t \|g_i\|^2$, and $\Phi(S, V) := \exp(\frac{S^2/2}{\gamma^2+V+|S|} - \frac{1}{2}\ln\frac{\gamma^2}{\gamma^2+V})$, for any $\gamma > 0$. Instantiating this guarantee in 1d with (g_t) set to an MDS (X_t) and taking (conditional) expectation in (1) shows that $\Phi_t := \Phi(\sum_{i=1}^t X_i, \sum_{i=1}^t X_i^2)$ is a non-negative supermartingale, from which concentration results can be obtained via Ville's inequality (a generalization of Markov's inequality—see Lemma 18). Our proof technique is similar to the one introduced in [28],with the difference that we use the specific shape of FREEGRAD's potential function to build our supermartingale, which leads to a desirable empirical variance term in the final concentration bound.

On the side of risk monotonicity, given n samples, the key idea behind our approach is to iteratively generate a sequence of distributions $P_1, P_2, \ldots$ leading up to P_n over hypotheses, where we only allow consecutive distributions, say P_{k-1} and P_k to differ if we can guarantee (with high enough confidence) that the risk associated with P_k is lower than that of P_{k-1}. To test for this, we compare the average empirical losses of hypotheses sampled from P_{k-1} versus ones sampled from P_k, taking into account the potential gap between empirical and population expectations. Applying our new concentration bounds to quantify this gap not only allows us to achieve risk monotonicity under a non-i.i.d. loss process but also enables us to achieve fast excess risk rates under the Bernstein condition. For the latter, it was crucial to have an empirical loss variance term in the concentration inequality.

Related Works.   Much work has already been done in efforts to mitigate the non-monotonic behavior of risk curves [54, 42, 33]. For example, in the supervised learning setting with the zero-one loss, [9] introduced the "memorize" algorithm that predicts the majority label on any test instance x that was observed during training; otherwise, a default label is predicted. [9] showed that this algorithm is risk-monotonic. However, it is unclear how their result could generalize beyond the particular setting they considered. Risk-monotonic algorithms are also known for the case where the model is correctly specified (see [33] for an overview); in this paper, we do not make such an assumption.

Closer to our work is that of [54] who, like us, also used the idea of only updating the current predictor for sample size n if it has a lower risk than the predictor for sample size $n-1$. They determine whether this is the case by performing statistical tests on a validation set (or through cross-validation). They introduce algorithm wrappers that ensure that the risk curves of the final algorithms are monotonic with high probability. However, their results are specialized to the 0-1 loss, and they do not answer the question by [53] on the existence of learners that guarantee a monotonic risk in *expectation*.

On the side of concentration bounds for non-i.i.d. processes, our results are somewhat similar to those found in e.g. [27, 26]. However, our technique for deriving them, which relies on the guarantee of a parameter-free online learning algorithm, is entirely different. The theoretical link between online regret and concentration inequalities was previously drawn—see e.g. [45, 22]. However, our approach is slightly different as we use the monotonicity of an online algorithm's potential function to get our concentration results. Thus, new concentration inequalities may be derived similarly by modifying the explicit potential function directly. Our approach is more similar to that of [28] who also derived concentration inequalities using guarantees of online betting algorithms that bet fractions smaller than one of their wealth at each round.

---

[2]One can not improve on the $\ln \ln n$ factor by the law of iterated logarithm [13].

We allow for a parameter $h \in \mathcal{H}$ in our concentration bounds to make the results useful in the statistical learning setting. These bounds are of PAC-Bayesian type and are somewhat reminiscent of those in [50, 38]. We refer the reader to [25] for an overview of existing PAC-Bayesian bounds.

Outline. In Section 2, we introduce the setting, notation, and relevant definitions. In Section 3, we present our new concentration inequalities for Martingale Difference Sequences and loss processes we are interested in (those that satisfy Assumption 1 below). In Section 4, we present our risk-monotonic algorithm wrapper and show that it answers the question posed by [53]. We conclude with a discussion in Section 5. The proofs of the new concentration inequalities and risk monotonicity are differed to Appendices B and C, respectively. Appendix A presents existing and new technical results needed in some of our proofs.

2 Preliminaries

In this section, we present the setting, notation, and relevant definitions for the rest of the paper.

Setting and Notation. Throughout, we will assume an underlying probability space $(\Omega, \mathcal{F}, \mathbf{P})$. Let $\mathcal{Z}$ [resp. $\mathcal{H}$] be an arbitrary feature [resp. hypothesis] space, and let $\ell : \mathcal{H} \times \mathcal{Z} \to [0,1]$ be a bounded loss function. We denote by $\triangle(\mathcal{H})$ the set of probability measures on $\mathcal{H}$. Data is represented by random variables $Z_1, Z_2, \dots \in \mathcal{Z}$ that we assume are accessible to a learning algorithm in a sequential fashion. We use the concise notation $Z_{1:t}$ for the tuple $(Z_1, \dots, Z_t)$ and denote by $\mathcal{G}_t$ the σ-algebra generated by the random variables $Z_1, \dots, Z_t$, with the convention that $\mathcal{G}_0 = \varnothing$. We will write $\mathbf{E}_{t-1}[\cdot] := \mathbf{E}[\cdot \mid \mathcal{G}_{t-1}]$, for $t \geq 1$.

We will not assume that the random variables (Z_t) are independent and identically distributed. Instead, we will make the following weaker assumption on the loss process $(\ell(h, Z_t))$:

Assumption 1 (Process Assumption). *There exists a risk function $L : \mathcal{H} \to [0,1]$ such that the sequence of random variables $Z_1, Z_2, \dots$ satisfy $\mathbf{E}_{t-1}[\ell(h, Z_t)] = L(h)$, for all $h \in \mathcal{H}$ and $t \geq 1$.*

When the random variables $Z_1, Z_2, \dots$ are i.i.d. and $Z_i \sim P_Z$, $i \in \mathbb{N}$, then Assumption 1 trivially holds with the standard risk function $L(h) = \mathbf{E}_{P_Z(z)}[\ell(h, z)]$. In Assumption 1, the conditional distribution of Z_t given $\mathcal{G}_{t-1}$ may be arbitrary as long as the corresponding conditional expectation $\mathbf{E}_{t-1}$ of the loss of a given hypothesis h is the same (equal to $L(h)$) for all $t \geq 1$. Arguably, Assumption 1 represents the weakest condition under which studying risk monotonicity in the statistical learning setting still makes sense. We touch more on this point after defining risk monotonicity below. To simplify notation for the rest of this paper, we let

$$L(Q) := \mathbf{E}_{Q(h)}[L(h)], \quad \text{for all } Q \in \triangle(\mathcal{H}),$$

where L is as in Assumption 1.

A learning algorithm A is a map from $\bigcup_{i=1}^{\infty} \mathcal{Z}^i$ to $\triangle(\mathcal{H})$; given data $Z_{1:t}$, the output $\mathsf{A}(Z_{1:t})$ of the algorithm is a distribution over hypotheses in $\mathcal{H}$. This definition includes deterministic algorithms for which the distribution $\mathsf{A}(Z_{1:t})$ is a Dirac at some $h = h(Z_{1:t})$. We will use the notation $\mathsf{A}(\cdot \mid Z_{1:t}) := \mathsf{A}(Z_{1:t})(\cdot)$.

Throughout, we will make use of a fixed "prior" distribution P_0 over hypotheses in $\mathcal{H}$. In Section 4, we will present an algorithm wrapper that takes any base algorithm B as input and makes a risk-monotonic algorithm out of it with essentially the same excess risk rate. The results of this paper are useful for base algorithms that output distributions that are absolutely continuous w.r.t. our choice of prior P_0; that is $\mathsf{B}(Z_{1:t}) \ll P_0$, for all $t \geq 1$[3]. In practice, if $\mathcal{H} = \mathbb{R}^d$, P_0 may be a multivariate Gaussian around the origin and $\mathsf{B}(Z_{1:t})$ may also be a multivariate Gaussian around the ERM $\hat{h}_n \in \arg\inf_{h \in \mathcal{H}} \sum_{i=1}^{n} \ell(h, Z_i)$, in which case $\mathsf{B}(Z_{1:t}) \ll P_0$ holds for all $t \geq 1$. We now define the notion of risk monotonicity we will work with:

Definition 1 (Risk Monotonicity). *For $\delta \in (0, 1)$ and $N \geq 1$, we say that a learning algorithm $\mathsf{A} : \bigcup_{i=1}^{\infty} \mathcal{Z}^i \to \triangle(\mathcal{H})$ is (δ, N)-risk-monotonic if, with probability at least $1 - \delta$,*

$$\forall t \geq N, \quad \mathbf{E}_{A(h|Z_{1:t})}\left[\mathbf{E}_t[\ell(h, Z_{t+1})]\right] \leq \mathbf{E}_{A(h|Z_{1:t-1})}\left[\mathbf{E}_{t-1}[\ell(h, Z_t)]\right]. \tag{2}$$

---

[3]We inherit this restriction from the PAC-Bayesian approach that we use to quantify generalization. In Appendix D, we show how this restriction can be removed in the i.i.d. setting (see also Remark 7).

Note that since the loss ℓ is positive, Fubini's theorem implies that $\mathbf{E}_{\mathsf{A}(h|Z_{1:t-1})}\left[\mathbf{E}_{t-1}[\ell(h, Z_t)]\right] = \mathbf{E}_{t-1}\left[\mathbf{E}_{\mathsf{A}(h|Z_{1:t-1})}[\ell(h, Z_t)]\right]$, for all $t \geq 1$. Thus, the condition in (2) requires the expected loss of algorithm A on the next sample, conditioned on the past data, to decrease with the size of the data. We note that if Assumption 1 does not hold and $\mathbf{E}_t[\ell(h, Z_{t+1})]$ depends on t in an arbitrary fashion, then requiring (2) would be too strong. Thus, we will restrict our attention to processes that satisfy Assumption 1. We stress that the condition in this assumption is weaker than i.i.d., and allows the random variables $Z_1, Z_2, \ldots$ to have different (conditional) distributions as long as the moment constraint $\mathbf{E}_{t-1}[\ell(h, Z_t)] = L(h)$ is satisfied for all $h \in \mathcal{H}$ and $t \geq 1$.

The notion of monotonicity presented in [33, 53] concerned only i.i.d. random variables, and requires the risk to be monotonic in expectation as opposed to in high probability. In particular, the strongest notation of monotonicity in [33], which they refer to as global $\mathcal{Z}$-monotonicity, can be expressed as

$$\forall t \geq 1, \quad \mathbf{E}\left[\mathbf{E}_{\mathsf{A}(h|Z_{1:t})}[L(h)]\right] \leq \mathbf{E}\left[\mathbf{E}_{\mathsf{A}(h|Z_{1:t-1})}[L(h)]\right], \tag{3}$$

where $L(h) \coloneqq \mathbf{E}_{P_Z(z)}[\ell(h, z)]$ and $Z_1, Z_2 \ldots \overset{\text{i.i.d.}}{\sim} P_Z$. We will show that achieving risk-monotonicity in expectation is as easy as achieving (δ, N)-risk-monotonicity—at least for bounded losses. In fact, we will show how our risk-monotonic (in the sense of Definition 1) algorithm can easily be turned into one that satisfies (3).

Monotonicity alone is rather easy to achieve; it suffices to output a fixed hypothesis $h \in \mathcal{H}$ regardless of the training dataset. In this case, the risk would be constant, and so risk monotonicity is achieved by definition. In practice, it is important to generate hypotheses with low risk, and so a fixed hypothesis that does not dependent on data is likely to be useless. Formally, we want algorithms that are risk monotonic and *consistent*:

Definition 2 (Consistency). *Under Assumption 1, we say that algorithm* A *is* consistent *if for any* $\epsilon > 0$, $\lim_{n \to \infty} \mathbf{P}\left[|\mathbf{E}_{\mathsf{A}(h|Z_{1:n})}[L(h)] - \inf_{h \in \mathcal{H}} L(h)| > \epsilon\right] = 0$.

We will go beyond the notation of consistency and study the rate of convergence of the risk of our algorithm to the optimal risk. We do this under the assumption that the loss process $\ell(h, Z_t)$ satisfies the Bernstein condition for $h \in \mathcal{H}$:

Definition 3 (Bernstein Condition). *For $\beta \in [0, 1]$ and $B > 0$, the (β, B)-Bernstein condition holds if the random variables $Z_1, Z_2, \ldots$ and the loss ℓ satisfy, for all $t \geq 1$ and all $h \in \mathcal{H}$,*

$$\mathbf{E}_{t-1}\left[(\ell(h, Z_t) - \ell(h_\star, Z_t))^2\right] \leq B\mathbf{E}_{t-1}[\ell(h, Z_t) - \ell(h_\star, Z_t)]^\beta,$$

for $h_\star \in \arg\inf_{h \in \mathcal{H}} \mathbf{E}_{t-1}[\ell(h, Z_t)]$.

The Bernstein condition [3, 4, 5, 21, 29] essentially characterizes the easiness of the learning problem. In particular, it implies that the conditional variance of the excess-loss random variable $\ell(h, Z_t) - \ell(h_\star, Z_t)$ vanishes when the risk associated with the hypothesis $h \in \mathcal{H}$ gets closer to the $\mathcal{H}$-optimal risk $L(h_\star)$. For bounded loss functions, the Bernstein condition with $\beta = 0$ always holds, and so the results of this paper are always true for $\beta = 0$. The Bernstein condition with $\beta = 1$ corresponds to the easiest learning setting. The case where $\beta \in (0, 1)$ interpolates naturally between these two extremes, where intermediate excess-risk rates are achievable. We refer the reader to [29, Section 3] for examples of learning settings where a Bernstein condition holds.

Additional useful definitions. For $\rho > 1$ and $\delta \in (0, 1)$, we define

$$c \coloneqq \sum_{k \geq 1} \frac{1}{k \ln^2(k+1)} \approx 3.2; \quad \phi_\rho(n) \coloneqq c\sqrt{\rho + 1}(\ln_\rho(n) + 1) \ln^2(\ln_\rho(n) + 2); \tag{4}$$

$$\text{and} \quad n_\delta \coloneqq \sup\left\{n \in \mathbb{N} \,:\, 8\ln(\phi_\rho(n)/\delta) > n\right\}. \tag{5}$$

Note that n_δ is not too large as a function of $1/\delta$. In fact, the definitions of ϕ_ρ and n_δ imply that $n_\delta \leq O(\ln(1/\delta))$. In the next section, we present some new concentration inequalities of independent interest that will be useful in achieving risk-monotonicity under Assumption 1 while maintaining good excess risk rates.

3   New (PAC-Bayesian) Concentration Inequalities

In this section, we present some new concentration inequalities of PAC-Bayesian flavor that will be crucial to deriving our risk monotonic algorithm wrapper under Assumption 1. These concentration

inequalities hold for non-i.i.d. data (which we require to accommodate Assumption 1), and are so-called *time-uniform*; the inequalities hold for all sample sizes simultaneously given a fixed confidence level. We explain below the advantage that this has in our setting.

We start by a new concentration inequality for Martingale Difference Sequences, from which we derive the bound we need under Assumption 1. First, we give the formal definition of an MDS:

Definition 4. *Let $(\mathcal{F}_t)_{t\in\mathbb{N}}$ be a filtration w.r.t. the underlying probability space $(\Omega, \mathcal{F}, \mathbf{P})$, i.e. $(\mathcal{F}_t)_{t\in\mathbb{N}}$ is a sequence of non-decreasing sub-σ-algebras of $\mathcal{F}$. A sequence of random variables (X_t) is an MDS w.r.t. $(\mathcal{F}_t)_{t\in\mathbb{N}}$, if for all $t \geq 1$, X_t is $\mathcal{F}_t$-measurable; $\mathbf{E}[|X_t|] < \infty$; and $\mathbf{E}[X_t \mid \mathcal{F}_{t-1}] = 0$ a.s.*

With this in hand, we present our first concentration inequality:

Proposition 5 (PAC-Bayes for MDS)**.** *Let $\rho > 1$ and ϕ_ρ be as in (4). Further, let $\{X_t^h\}$ be a family of random variables taking values in $[-1, 1]$ and $(\mathcal{F}_t)_{t\in\mathbb{N}}$ be a filtration such that (X_t^h) is an MDS w.r.t to $(\mathcal{F}_t)_{t\in\mathbb{N}}$, for all $h \in \mathcal{H}$. Then, for any distribution P_0 on $\mathcal{H}$ and all $\delta \in (0, 1)$, we have*

$$\mathbf{P}\left[\forall n \geq 1, \forall P, \quad \frac{\mathbf{E}_{P(h)}[|S_n^h|]^2}{2(\rho+1)\mathbf{E}_{P(h)}[V_n^h] + 2\mathbf{E}_{P(h)}[|S_n^h|]} \leq \mathrm{KL}(P\|P_0) + \ln\frac{\phi_\rho(n)}{\delta}\right] \geq 1-\delta, \quad (6)$$

where $S_n^h \coloneqq \sum_{t=1}^n X_t^h$ and $V_n^h \coloneqq 1 + \sum_{t=1}^n (X_t^h)^2$.

The proof of the theorem is in Appendix B. The bound in Proposition 5 does not look like the typical PAC-Bayesian bound. However, simple algebra reveals that for any $C > 0$,

$$\frac{\mathbf{E}_{P(h)}[|S_n^h|]^2}{2(\rho+1)\mathbf{E}_{P(h)}[V_n^h] + 2\mathbf{E}_{P(h)}[|S_n^h|]} \leq C \implies \mathbf{E}_{P(h)}[|S_n^h|] \leq 2C + \sqrt{2(\rho+1)\mathbf{E}_{P(h)}[V_n^h] \cdot C}.$$

Combining this with the fact that $|\mathbf{E}_{P(h)}[S_n^h]| \leq \mathbf{E}_{P(h)}[|S_n^h|]$ (by Jensen's inequality), and (6), we obtain, under the same conditions as Proposition 5 that

$$\mathbf{P}\left[\forall n \geq 1, \forall P \in \triangle(\mathcal{H}), \;\; \left|\mathbf{E}_{P(h)}[S_n^h]\right| \leq 2C_n(P) + \sqrt{2(\rho+1)\mathbf{E}_{P(h)}[V_n^h] \cdot C_n(P)}\right] \geq 1-\delta, \quad (7)$$

where $C_n(P) \coloneqq \mathrm{KL}(P\|P_0) + \ln(\phi_\rho(n)/\delta)$. When $\mathcal{H}$ is a singleton, the concentration inequality in (7) can be viewed as an empirical version of Freedman's inequality [23] for MDS. The inequality is also reminiscent of the PAC-Bayesian empirical Bernstein inequality due to [50]. In addition to it holding for MDS, another advantage of (7) is that it is time-uniform—it holds for all sample sizes n, simultaneously. While standard concentration inequalities that hold for a fixed sample size can be turned into time-uniform ones by applying a union bound over sample sizes, the resulting inequalities will have sub-optimal $\ln n$ factors under the main square-root error term. In contrast, the concentration inequality in (7) has a $\ln(\phi_\rho(n)) = O(\ln\ln n)$ term, which matches the optimal dependence in n according to the law of iterated logarithm [13]. MDS is a central concept in probability theory [44] and machine learning [10], and so our result in Proposition 5 is of independent interest.

Interestingly, the concentration inequality in Proposition 5 is derived using the guarantee of a parameter-free online algorithm—FREEGRAD. This opens the door for new ways of deriving such concentration inequalities through online learning algorithms, adding to existing results due to [45, 22, 28].

Using the result of Proposition 5, we are now going to derive a time-uniform "Empirical Bernstein" concentration inequality that holds under Assumption 1:

Theorem 6. *Let $\delta \in (0, 1)$, $\rho > 1$, and ϕ_ρ be as in (4). Further, for $P_0 \in \triangle(\mathcal{H})$ define $\epsilon_n(P) \coloneqq \frac{2(\rho+1)}{n}(\mathrm{KL}(P\|P_0) + \ln\frac{\phi_\rho(n)}{\delta})$ and $\mathcal{P}_n \coloneqq \{P : 1 - \epsilon_n(P) > 0\}$. Under Assumption 1 we have*

$$\mathbf{P}\left[\forall n \geq n_\delta, \forall P \in \mathcal{P}_n, \;\; \left|\frac{1}{n}\sum_{t=1}^n \mathbf{E}_{P(h)}\left[\ell(h, Z_t) - L(h)\right]\right| \leq \frac{\sqrt{\epsilon_n(P) \cdot \widehat{V}_n(P)} + \frac{\epsilon_n(P)}{\rho+1}}{1 - \epsilon_n(P)}\right] \geq 1-\delta,$$

where $\widehat{V}_n(P) \coloneqq \frac{1}{n}\sum_{t=1}^n \mathbf{E}_{P(h)}\left[\left(\ell(h, Z_t) - \frac{1}{n}\sum_{i=1}^n \mathbf{E}_{P(\theta)}[\ell(\theta, Z_i)]\right)^2\right] + \frac{1}{n}$ and n_δ is as in (5).

The restriction that $P \in \mathcal{P}_n$ in the theorem merely ensures that the denominator in the concentration bound remains positive. The set P_n is guaranteed to be non-empty for all $n \geq n_\delta$. In this case, P is in

$\mathcal{P}_n$ whenever $8(\rho + 1)\,\mathrm{KL}(P\|P_0) \le n$, which is a fairly weak condition on the distribution P; even for large models such as Neural Networks the KL-divergence $\mathrm{KL}(P\|P_0)$ does not typically grow superlinearly with the size n of the sample used to generate the posterior P [55]. Nevertheless, we note that if one only cares about the i.i.d. setting then other concentration inequalities may be used to achieve risk monotonicity without restrictions on the posterior P (see Appendix D).

The concentration inequality in Theorem 6 can be viewed as an extension of the empirical Bernstein inequality in [35, 50] that holds under the non-i.i.d. condition described in Assumption 1. Our new bound is also time-uniform with the optimal dependence in the sample size n. As mentioned before, simply applying a union bound to a standard (non-time-uniform) concentration inequalities to obtain its time-uniform version will lead to a sub-optimal dependence in n. When $\mathcal{H}$ is a singleton (i.e. no learning), the concentration inequality becomes reminiscent of an existing one due to [26]. However, the latter has a term that looks like, but is different than, the empirical variance, and so it is not directly comparable to ours. The proof of Theorem 6 is postponed to Appendix B.

Finally, we note that for any sample size n the value of ρ in Theorem 6 that minimizes the bound would typically fall within the interval $(1, 2)$. One can tune ρ as a function of the data by treating ρ as an extra "hypothesis" parameter. In the case of a finite grid $\mathcal{G} \subset (1, +\infty)$ of ρ's, the result of Theorem 6 would hold for all $\rho \in \mathcal{G}$ inside the probability event as long as any $\mathrm{KL}(P\|P_0)$ instance is replaced by $\mathrm{KL}(P\|P_0) + \ln|\mathcal{G}|$.

We now move on to describing our risk monotonic procedure that makes use of our new concentration inequality.

4 Risk Monotonicity in Statistical Learning

In this section, we combine the concentration inequality from Theorem 6 with a novel "greedy" procedure for selecting distributions over hypotheses to derive a risk monotonic algorithm in the statistical learning setting. With the right choice of gap sequence (ξ_n), the procedure we present in Algorithm 1 takes as input a base learning algorithm $\mathsf{B} : \bigcup_{i=1}^{\infty} \mathcal{Z}^i \to \triangle(\mathcal{H})$ together with samples $Z_{1:n}$ whose generating process satisfies Assumption 1, and returns a distribution over $\mathcal{H}$ that has a monotonic risk as a function of n with high probability. By leveraging our new concentration inequality in Theorem 6 to specify the gap sequence (ξ_n), we further show that achieving risk monotonicity need not deteriorate rates of convergence to the optimal risk $\inf_{h \in \mathcal{H}} L(h)$. In fact, we show that it is possible to attain fast rates under the Bernstein condition (Definition 3). To arrive at this result, it was crucial for our concentration inequality to have an empirical variance term.

Algorithm 1 A Risk Monotonic Algorithm Wrapper

Require:
        A base learning algorithm $\mathsf{B} : \bigcup_{i=1}^{\infty} \mathcal{Z}^i \to \triangle(\mathcal{H})$.
        Samples $Z_1, \ldots, Z_n$.
        A sequence of gap functions (ξ_k), where $\xi_k = \xi_k(Z_{1:k}, \mathsf{B})$ for all k. $\triangleright$ (ξ_k) will be set according to (9) [resp. (11)] for risk-monotonicity in probability [resp. expectation].

1: **for** $k = 1, \ldots, n$ **do**
2:      **if** $\dfrac{1}{k}\sum_{i=1}^{k} \mathbf{E}_{\mathsf{B}(h|Z_{1:k})}[\ell(h, Z_i)] - \dfrac{1}{k}\sum_{i=1}^{k} \mathbf{E}_{P_{k-1}(h)}[\ell(h, Z_i)] \le -\xi_k$ **then**
3:          Set $P_k = \mathsf{B}(Z_{1:k})$.
4:      **else**
5:          Set $P_k = P_{k-1}$.
6: Return P_n.

To simplify our analysis, we will focus our attention on base algorithms that are restricted in the following way:

Assumption 2 (Base Algorithm Restriction). *We assume access to a base algorithm* $\mathsf{B} : \bigcup_{i=1}^{\infty} \mathcal{Z}^i \to \triangle(\mathcal{H})$ *such that for* $\rho > 1$, *prior distribution* P_0 *on* $\mathcal{H}$, *and all* $k \ge 1$,

$$\mathsf{B}(Z_{1:k}) \in \mathcal{Q}_k := \{Q \in \triangle(\mathcal{H}) : 16(\rho + 1)\,\mathrm{KL}(Q\|P_0) \le k\}. \tag{8}$$

In other words, we are restricting our attention to algorithms whose posteriors given samples of size k do not have a KL divergence to the prior P_0 that grow too quickly with k. We make this assumption to satisfy the technical conditions needed for the concentration bound in Theorem 6 to be non-vacuous. Assumption 2 is reasonable even for large Neural Network models [55][4].

In practice, if $\mathcal{H}$ is $\mathbb{R}^d$, an option for B is the algorithm that outputs a multivariate Gaussian distribution around a regularized ERM. The KL-divergence between the outputs of B and P_0 can be controlled by tuning the regularisation parameter(s) and/or the co-variance matrix of the multivariate Gaussian.

Remark 7. *If one only cares about i.i.d. loss processes, Assumption 2 can be removed by using generalization bounds based on the standard Bernstein concentration inequality, e.g. [50, 36], which would also enable our risk decomposition in Theorem 9 below (and thus fast rates). We make Assumption 2 only to bound the denominator in our (non-i.i.d) concentration in Theorem 6 away from zero, which is not needed for other concentration bounds in the i.i.d. setting such as those in [50, 36] (see Appendix D for more detail).*

To specify the sequence of gaps (ξ_n) that our wrapper Algorithm 1 requires, we will use our concentration bound in Theorem 6. We recall that using this concentration bound instead of other existing ones allows us to I) achieve risk monotonicity under a weaker condition than i.i.d. on the loss process (Assumption 1 in this case); and II) to achieve potentially fast excess risk rates under the Bernstein condition. The latter is made possible by the fact that the concentration bound in Theorem 6 has an empirical loss variance term that allows a particularly useful decomposition of the excess risk under the Bernstein condition (see Theorem 9 below).

To give a concise expression of the gaps (ξ_k), we let $Q_k \coloneqq \mathsf{B}(Z_{1:k}) \times P_{k-1}$, where (P_k) are the intermediate distributions generated internally by Algorithm 1. With this, and the convention that $1/0 = +\infty$, we define

$$\xi_k \coloneqq \frac{\sqrt{\epsilon_k \cdot \widehat{V}_k} + \frac{2\epsilon_k}{\rho+1}}{|1 - \epsilon_k|}, \quad \text{where} \quad \epsilon_k \coloneqq \frac{2\,\mathrm{KL}(Q_k \| P_0 \times P_0) + 2\ln\frac{\phi_\rho(k)}{\delta}}{k \cdot (\rho+1)^{-1}}; \quad \text{and} \tag{9}$$

$$\widehat{V}_k \coloneqq \frac{1}{k}\sum_{t=1}^{k}\mathbf{E}_{Q_k(h,h')}\left[(\ell(h, Z_t) - \ell(h', Z_t))^2\right] - \left(\frac{1}{k}\sum_{t=1}^{k}\mathbf{E}_{Q_k(h,h')}\left[\ell(h, Z_t) - \ell(h', Z_t)\right]\right)^2.$$

We note that our Assumption 2 ensures that $\epsilon_n \leq 1/2$ (which in turn ensures that the gaps (ξ_n) are not too large), for all $n \geq n_\delta$, where n_δ is as in (5). This follows from the fact that $\mathrm{KL}(Q_k \| P_0 \times P_0) = \mathrm{KL}(\mathsf{B}(Z_{1:k}) \| P_0) + \mathrm{KL}(P_{k-1} \| P_0)$, and that there exists $m < k$ such that $P_{k-1} = \mathsf{B}(Z_{1:m}) \in \mathcal{Q}_m \subseteq \mathcal{Q}_k$ (by definition of (P_k) in Algorithm 1). Furthermore, $\mathcal{Q}_n \neq \varnothing$ for all $n \geq n_\delta$.

Before presenting our results for this section, we also note that Algorithm 1 assumes we can evaluate expectations over the output distributions of B. Such expectations can be approximated via Monte Carlo sampling, and any approximation errors need be added to the gaps (ξ_n) to maintain the guarantees we present. Alternatively, one can avoid estimating expectations by applying recent derandomization techniques, see e.g. [46, 51], or by using other, non PAC-Bayesian generalization bounds (see Appendix D).

We now state the guarantees of Algorithm 1. We start by the statement of risk-monotonicity (the proof is postponed to Appendix C):

Theorem 8 (Risk Monotonicity). *Let $\delta \in (0, 1)$ and n_δ be as in (5). Under Assumptions 1 and 2, Algorithm 1 with (ξ_k) as in (9) is (δ, n_δ)-risk-monotonic according to Definition 1.*

We now show that risk monotonicity need not come at a worse excess risk rate. Under the Bernstein condition, we have the following excess risk decomposition for the output of Algorithm 1:

Theorem 9 (Risk Decomposition). *Let $B > 1$, $\beta \in [0,1]$, and suppose that the (β, B)-Bernstein condition holds. Further, for $\rho > 1$ and $\delta \in (0,1)$, let n_δ and ϵ_n be as in (5) and (9), respectively. Then, under Assumptions 1 and 2, the outputs (P_k) of Alg. 1 satisfy, with probability at least $1 - 2\delta$,*

$$\forall n \geq n_\delta, \quad L(P_n) - L(h_\star) \leq 3(L(\mathsf{B}(Z_{1:n})) - L(h_\star)) + O(\epsilon_n)^{\frac{1}{2-\beta}}, \tag{10}$$

where $h_\star \in \arg\inf_{h \in \mathcal{H}} L(h)$.

---

[4]Besides, when the KL divergence is very large, it is often very hard to infer non-vacuous generalization bounds (which one may view as a prerequisite to achieving risk monotonicity) using a PAC-Bayesian approach.

We stress that this risk decomposition was only made possible by the fact that our concentration bound in Theorem 6 has an empirical loss variance term.

Theorem 9 shows that the excess risk of Algorithm 1 is at most a constant times the excess risk of the base algorithm B, plus a potentially lower-order term $O(\epsilon_n^{\frac{1}{2-\beta}})$. To appreciate what this additional term is doing, consider the case of a finite hypothesis class $\mathcal{H}$. In this case, the definition of ϵ_n in (9) implies that $\epsilon_n \leq O(\frac{1}{n} \ln \frac{|\mathcal{H}| \ln n}{\delta})$. Thus, $\epsilon_n^{\frac{1}{2-\beta}}$ interpolates between the fast $\frac{1}{n} \ln \frac{|\mathcal{H}| \ln n}{\delta}$ rate under the best Bernstein condition with $\beta = 1$ and the standard (up to log-log-factors) rate $\sqrt{\frac{1}{n} \ln \frac{|\mathcal{H}| \ln n}{\delta}}$ under the Bernstein condition with $\beta = 0$, which we recall always holds for bounded losses. What is more, if $\mathcal{H}$ is finite and algorithm B is the ERM, i.e. if $\mathsf{B}(Z_{1:k})$ is a Dirac at $\hat{h}_n \in \arg\inf_{h \in \mathcal{H}} \sum_{i=1}^{n} \ell(h, Z_i)$, then we have the following explicit excess risk rate for Algorithm 1:

Proposition 10. *Under the setting of Theorem 9, if $\mathcal{H}$ is finite, P_0 is set to the uniform prior over $\mathcal{H}$, and $\mathsf{B}(Z_{1:n})$ is a point mass around the ERM $\hat{h}_n \in \arg\min_{h \in \mathcal{H}} \sum_{i=1}^{n} \ell(h, Z_i)$, $\forall n \geq 1$, then Alg. 1 is risk monotonic according to Def. 1 and its outputs (P_k) satisfy, with probability at least $1 - 2\delta$,*

$$\forall n \geq n_\delta \vee (16(\rho+1) \ln |\mathcal{H}|), \quad L(P_n) - L(h_\star) \leq O\left(\frac{\ln \frac{|\mathcal{H}| \ln n}{\delta}}{n} \right)^{\frac{1}{2-\beta}}.$$

The story is not much different for a continuous set $\mathcal{H}$. The standard excess risk rate one would expect from algorithm B is $\sqrt{(\mathrm{KL}(\mathsf{B}(Z_{1:k}) \| P_0) + \ln(1/\delta))/k}$, which can dominate the right-most term in our risk decomposition (10) since[5]

$$(\epsilon_n)^{\frac{1}{2-\beta}} \leq O\left(\frac{\max_{k \leq n} \mathrm{KL}(\mathsf{B}(Z_{1:k}) \| P_0) + \ln \frac{\ln n}{\delta}}{n} \right)^{\frac{1}{2-\beta}}.$$

Together, the above inequality and Theorem 9 show that fast rates for algorithm 1 are achievable whenever the Bernstein condition holds with $\beta > 0$ and the base algorithm B itself achieves a fast rate (when $\mathcal{H}$ is finite and B is the ERM, Proposition 10 shows that it is sufficient that $\beta > 0$). Thus, risk monotonicity need not come at the price of a worse rate of convergence to the optimal risk. Finally, we note that the factor 3 in our risk decomposition (10) is just an artifact of our analysis. In fact, by slightly modifying our proof of Theorem 9, one can show that any factor in the interval $(1, 3]$ is achievable at the cost of a larger lower-order term in (10).

Risk monotonicity in expectation. The original open problem due to [53] was around risk monotonicity in expectation—as in (3). Our algorithm wrapper also allows us to achieve risk monotonicity in expectation with a slight modification of the sequence (ξ_k) in (9). In fact, we have the following result, which essentially answers open question by [53] on the existence of risk monotonic algorithms.

Theorem 11. *Let $\rho > 1$ and ϕ_ρ be as in (4). Under Assumptions 1 and 2, Algorithm 1 with the sequence of gaps (ξ_k) given by*

$$\xi_k = \xi_k' := \frac{1}{|1 - \epsilon_k'|} \left(\sqrt{\epsilon_k' \cdot \widehat{V}_k} + \frac{2\epsilon_k'}{\rho+1} \right) + \frac{1}{k}, \tag{11}$$

with $\epsilon_k' := \frac{2(\rho+1)}{k} \left(\mathrm{KL}(\mathsf{B}(Z_{1:k}) \times P_{k-1} \| P_0 \times P_0) + 2\ln(k\phi_\rho(k)) \right)$ and (P_k) being as in Algorithm 1, is risk-monotonic in expectation; that is, for all $t \geq N := \sup \{n : 8\ln(n\phi_\rho(n)) > n\}$,

$$\mathbf{E}\left[\mathbf{E}_{P_t(h)}[L(h)] \right] \leq \mathbf{E}\left[\mathbf{E}_{P_{t-1}(h)}[L(h)] \right]. \tag{12}$$

We note that N need not be too large since $\phi_\rho(n) \leq O(\ln n)$. The notion of risk-monotonicity in theorem 11 corresponds to the notion of weak $\mathcal{Z}$-monotonicity in [33]—one of their strongest notions of risk monotonicity since it holds for all data generating distributions (granted Assumption 1 holds, or the data is i.i.d.). We can modify Algorithm 1 to make it globally $\mathcal{Z}$-monotonic, where (12) would hold for all $t \geq 1$ (instead of $t \geq N$), by forcing the outputs P_k of the algorithm to be P_0 for all $k < N$.

With the same choice of gap sequence (ξ_k) in Theorem 11 it is easy to show (following almost the same proof steps as in our previous results) that the excess risk rates in Theorem 9 and Proposition 10 will hold with probability $1/n$ (instead of δ) for every sample size n.

---

[5]We note that it is typical that $\max_{k \leq n} \mathrm{KL}(\mathsf{B}(Z_{1:k}) \| P_0) \leq O(\mathrm{KL}(\mathsf{B}(Z_{1:n}) \| P_0))$, for $n \geq 1$.

5 Discussion and Future Work

The primary goal of this paper was to answer the fundamental question around the existence of a consistent, risk-monotonic algorithm in the general statistical learning setting. We answer this in the affirmative and further show that there is virtually no cost for achieving this when it comes to excess-risk rates. We believe this is an important milestone in the search for risk-monotonic algorithms.

From a computational perspective, the main setback of Algorithm 1 is that returning the distribution $P_n = P_n(Z_{1:n})$ requires n-calls to the base algorithm B. This implies that 1 may run n times slower than B in the worst case. However, when the base algorithm outputs distributions centered at ERMs, it may be possible to efficiently generate the sequence of distributions $(B(Z_{1:k}))_k$ by leveraging the fact that ERM solutions for sample sizes k and $k + 1$ can be close to each other.

When the loss ℓ is convex in the first argument, there is no need for a randomized algorithm[6] (i.e. we do not need $B(Z_{1:k}) \ll P_0$), and it is possible to efficiently generate a sequence of predictors $(\hat{h}_n)$ with monotonic risk using an online convex optimization algorithm as the base algorithm B. However, in general, it is unclear whether risk-monotonicity can be achieved without the (greedy) for-loop procedure of Algorithm 1. We note also that if one only wants a decreasing risk after some sample size $s \in \mathbb{N}$, then computing the distributions $(P_k)_{k<s}$ is unnecessary. In this case, the for-loop in Algorithm 1 need only start at $k = s$; the resulting hypotheses would satisfy the monotonicity condition in Definition 1 for all $n \geq s \vee n_\delta$.

Through Theorem 9, we showed that the excess risk of Algorithm 1 is at most three times that of the base algorithm plus a lower-order term. It remains to evaluate the empirical performance of the algorithm to identify for which applications the additional cost in the excess risk is worth it to achieve risk monotonicity.

Some important questions remain open along the axes of assumptions. In particular, can we remove the boundedness condition on the loss while retaining risk-monotonicity? It might be possible to achieve this for unbounded convex losses using the non-exponential weighted aggregation techniques recently suggested by [2]. Lifting the boundedness assumption may be key in resolving another COLT open problem [24] regarding achievable risk rates of log-loss Bayesian predictors. Our results build foundations for these avenues, which are promising subjects for future work.

Finally, it would be interesting to explore what other concentration inequalities for non-i.i.d. processes can be derived using other parameter-free online learning algorithms. An obvious starting point is to look at MATRIX-FREEGRAD [39].

---

[6]Randomization is also not needed if one is only interested in i.i.d. processes (see Appendix D).

Acknowledgments and Disclosure of Funding

We would like to thank Hisham Husain for instrumental discussions during the early phases of the project. We also thank anonymous reviewers for their valuable feedback. This work was supported by the Australian Research Council and Data61.

References

[1] Amir Ahmadi-Javid. Entropic value-at-risk: A new coherent risk measure. *Journal of Optimization Theory and Applications*, 155(3):1105–1123, 2012.

[2] Pierre Alquier. Non-exponentially weighted aggregation: regret bounds for unbounded loss functions. *arXiv preprint arXiv:2009.03017*, 2020.

[3] Jean-Yves Audibert. PAC-Bayesian statistical learning theory. *These de doctorat de l'Université Paris*, 6:29, 2004.

[4] Peter L. Bartlett, Michael I. Jordan, and Jon D. McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006.

[5] Peter L. Bartlett and Shahar Mendelson. Empirical minimization. *Probability Theory and Related Fields*, 135(3):311–334, 2006.

[6] Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine learning and the bias-variance trade-off. *arXiv preprint arXiv:1812.11118*, 2018.

[7] Mikhail Belkin, Daniel Hsu, and Ji Xu. Two models of double descent for weak features. *arXiv preprint arXiv:1903.07571*, 2019.

[8] Shai Ben-David, David Loker, Nathan Srebro, and Karthik Sridharan. Minimizing the misclassification error rate using a surrogate convex loss. In *Proceedings of the 29th International Coference on International Conference on Machine Learning*, pages 83–90, 2012.

[9] Shai Ben-David, Nathan Srebro, and Ruth Urner. Universal learning vs. no free lunch results. In *Philosophy and Machine Learning Workshop NIPS*, 2011.

[10] Nicolo Cesa-Bianchi and Gábor Lugosi. *Prediction, learning, and games*. Cambridge university press, 2006.

[11] Prasad Cheema and Mahito Sugiyama. A geometric look at double descent risk: Volumes, singularities, and distinguishabilities. *arXiv preprint arXiv:2006.04366*, 2020.

[12] Lin Chen, Yifei Min, Mikhail Belkin, and Amin Karbasi. Multiple descent: Design your own generalization curve. *arXiv preprint arXiv:2008.01036*, 2020.

[13] DA Darling and Herbert Robbins. Iterated logarithm inequalities. *Proceedings of the National Academy of Sciences of the United States of America*, 57(5):1188, 1967.

[14] Stéphane d'Ascoli, Levent Sagun, and Giulio Biroli. Triple descent and the two kinds of overfitting: Where & why do they appear? *arXiv preprint arXiv:2006.03509*, 2020.

[15] Zeyu Deng, Abla Kammoun, and Christos Thrampoulidis. A model of double descent for high-dimensional binary linear classification. *arXiv preprint arXiv:1911.05822*, 2019.

[16] Michal Derezinski, Feynman T Liang, and Michael W Mahoney. Exact expressions for double descent and implicit regularization via surrogate random design. *Advances in Neural Information Processing Systems*, 33, 2020.

[17] Michał Dereziński, Feynman Liang, and Michael W. Mahoney. Exact expressions for double descent and implicit regularization via surrogate random design, 2019.

[18] Robert P.W. Duin. Small sample size generalization. In *Proceedings of the Scandinavian Conference on Image Analysis*, volume 2, pages 957–964, 1995.

[19] Robert P.W. Duin. Classifiers in almost empty spaces. In *Proceedings 15th International Conference on Pattern Recognition. ICPR-2000*, volume 2, pages 1–7. IEEE, 2000.

[20] Robert P.W. Duin. Learning to generalize. *Frontiers of Life*, 3(part 2):763–775, 2001.

[21] Tim Van Erven, Nishant A. Mehta, Mark D. Reid, and Robert C. Williamson. Fast rates in statistical and online learning. *Journal of Machine Learning Research*, 16:1793–1861, 2015.

[22] Dylan J. Foster, Alexander Rakhlin, and Karthik Sridharan. Online learning: Sufficient statistics and the Burkholder method. In *Conference On Learning Theory, COLT 2018, Stockholm, Sweden, 6-9 July 2018.*, pages 3028–3064, 2018.

[23] David A Freedman. On tail probabilities for martingales. *the Annals of Probability*, pages 100–118, 1975.

[24] Peter D Grünwald and Wojciech Kotłowski. Bounds on individual risk for log-loss predictors. In *Proceedings of the 24th Annual Conference on Learning Theory*, pages 813–816, 2011.

[25] Benjamin Guedj. A primer on pac-bayesian learning. *arXiv preprint arXiv:1901.05353*, 2019.

[26] Steven R Howard, Aaditya Ramdas, Jon McAuliffe, and Jasjeet Sekhon. Time-uniform, nonparametric, nonasymptotic confidence sequences. *The Annals of Statistics*, 49(2):1055–1080, 2021.

[27] Steven R Howard, Aaditya Ramdas, Jon McAuliffe, Jasjeet Sekhon, et al. Time-uniform chernoff bounds via nonnegative supermartingales. *Probability Surveys*, 17:257–317, 2020.

[28] Kwang-Sung Jun and Francesco Orabona. Parameter-free online convex optimization with sub-exponential noise. In *Conference on Learning Theory*, pages 1802–1823, 2019.

[29] Wouter M. Koolen, Peter D. Grünwald, and Tim Van Erven. Combining adversarial guarantees and stochastic fast rates in online learning. In *Advances in Neural Information Processing Systems*, pages 4457–4465, 2016.

[30] Nicole Krämer. On the peaking phenomenon of the lasso in model selection. *arXiv preprint arXiv:0904.4416*, 2009.

[31] Marco Loog. Contrastive pessimistic likelihood estimation for semi-supervised classification. *IEEE transactions on pattern analysis and machine intelligence*, 38(3):462–475, 2015.

[32] Marco Loog and Robert P.W. Duin. The dipping phenomenon. In *Joint IAPR International Workshops on Statistical Techniques in Pattern Recognition (SPR) and Structural and Syntactic Pattern Recognition (SSPR)*, pages 310–317. Springer, 2012.

[33] Marco Loog, Tom Viering, and Alexander Mey. Minimizers of the empirical risk and risk monotonicity. In *Advances in Neural Information Processing Systems*, pages 7476–7485, 2019.

[34] Marco Loog, Tom Viering, Alexander Mey, Jesse H. Krijthe, and David M. J. Tax. A brief prehistory of double descent. *Proceedings of the National Academy of Sciences*, 117(20):10625–10626, 2020.

[35] Andreas Maurer and Massimiliano Pontil. Empirical bernstein bounds and sample variance penalization. *arXiv preprint arXiv:0907.3740*, 2009.

[36] Andreas Maurer and Massimiliano Pontil. Empirical bernstein bounds and sample-variance penalization. In *COLT 2009 - The 22nd Conference on Learning Theory, Montreal, Quebec, Canada, June 18-21, 2009*, 2009.

[37] Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and double descent curve. *arXiv preprint arXiv:1908.05355*, 2019.

[38] Zakaria Mhammedi, Peter Grünwald, and Benjamin Guedj. Pac-bayes un-expected bernstein inequality. In *Advances in Neural Information Processing Systems*, pages 12202–12213, 2019.

[39] Zakaria Mhammedi and Wouter M. Koolen. Lipschitz and comparator-norm adaptivity in online learning. In *Conference on Learning Theory, COLT 2020, 9-12 July 2020, Virtual Event [Graz, Austria]*, volume 125 of *Proceedings of Machine Learning Research*, pages 2858–2887. PMLR, 2020.

[40] Preetum Nakkiran. More data can hurt for linear regression: Sample-wise double descent. *arXiv preprint arXiv:1912.07242*, 2019.

[41] Preetum Nakkiran, Gal Kaplun, Yamini Bansal, Tristan Yang, Boaz Barak, and Ilya Sutskever. Deep double descent: Where bigger models and more data hurt. In *International Conference on Learning Representations*, 2020.

[42] Preetum Nakkiran, Prayaag Venkat, Sham Kakade, and Tengyu Ma. Optimal regularization can mitigate double descent. *arXiv preprint arXiv:2003.01897*, 2020.

[43] Manfred Opper and Wolfgang Kinzel. Statistical mechanics of generalization. In *Models of neural networks III*, pages 151–209. Springer, 1996.

[44] V.H. Peña, T.L. Lai, and Q.M. Shao. *Self-Normalized Processes: Limit Theory and Statistical Applications*. Probability and Its Applications. Springer Berlin Heidelberg, 2008.

[45] Alexander Rakhlin and Karthik Sridharan. On equivalence of martingale tail bounds and deterministic regret inequalities. In *Conference on Learning Theory*, pages 1704–1722. PMLR, 2017.

[46] Omar Rivasplata, Ilja Kuzborskij, Csaba Szepesvári, and John Shawe-Taylor. Pac-bayes analysis beyond the usual bounds. In *NeurIPS*, 2020.

[47] Shai Shalev-Shwartz and Shai Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.

[48] Alexander J. Smola, Peter J. Bartlett, Dale Schuurmans, and Bernhard Schölkopf. *Advances in large margin classifiers*. MIT press, 2000.

[49] Stefano Spigler, Mario Geiger, Stéphane d'Ascoli, Levent Sagun, Giulio Biroli, and Matthieu Wyart. A jamming transition from under-to over-parametrization affects loss landscape and generalization. *arXiv preprint arXiv:1810.09665*, 2018.

[50] Ilya O. Tolstikhin and Yevgeny Seldin. Pac-bayes-empirical-bernstein inequality. In *Advances in Neural Information Processing Systems 26: 27th Annual Conference on Neural Information Processing Systems 2013. Proceedings of a meeting held December 5-8, 2013, Lake Tahoe, Nevada, United States*, pages 109–117, 2013.

[51] Paul Viallard, Pascal Germain, Amaury Habrard, and Emilie Morvant. A general framework for the disintegration of pac-bayesian bounds, 2021.

[52] Tom Viering and Marco Loog. The shape of learning curves: a review. *arXiv preprint arXiv:2103.10948*, 2021.

[53] Tom Viering, Alexander Mey, and Marco Loog. Open problem: Monotonicity of learning. In *Conference on Learning Theory*, pages 3198–3201, 2019.

[54] Tom J. Viering, Alexander Mey, and Marco Loog. Making learners (more) monotone, 2019.

[55] Wenda Zhou, Victor Veitch, Morgane Austern, Ryan P. Adams, and Peter Orbanz. Non-vacuous generalization bounds at the imagenet scale: a pac-bayesian compression approach. In *7th International Conference on Learning Representations, ICLR 2019, New Orleans, LA, USA, May 6-9, 2019*, 2019.