A Universal Law of Robustness via Isoperimetry

Abstract

Classically, data interpolation with a parametrized model class is possible as long as the number of parameters is larger than the number of equations to be satisfied. A puzzling phenomenon in deep learning is that models are trained with many more parameters than what this classical theory would suggest. We propose a theoretical explanation for this phenomenon. We prove that for a broad class of data distributions and model classes, overparametrization is *necessary* if one wants to interpolate the data *smoothly*. Namely we show that *smooth* interpolation requires d times more parameters than mere interpolation, where d is the ambient data dimension. We prove this universal law of robustness for any smoothly parametrized function class with polynomial size weights, and any covariate distribution verifying isoperimetry (or a mixture thereof). In the case of two-layer neural networks and Gaussian covariates, this law was conjectured in prior work by Bubeck, Li and Nagaraj.

1 Introduction

Solving n equations generically requires only n unknowns[1]. However, the revolutionary deep learning methodology revolves around highly overparametrized models, with many more than n parameters to learn from n training data points. We propose an explanation for this enigmatic phenomenon, showing in great generality that finding a *smooth* function to fit d-dimensional data requires at least nd parameters. In other words, overparametrization by a factor of d is *necessary* for *smooth* interpolation, suggesting that perhaps the large size of the models used in deep learning is a *necessity* rather than a weakness of the framework. Another way to phrase the result is as a *tradeoff* between the size of a model (as measured by the number of parameters) and its "robustness" (as measured by its Lipschitz constant): either one has a small model (with n parameters) which must then be non-robust, or one has a robust model (constant Lipschitz) but then it must be very large (with nd parameters). Such a tradeoff was conjectured for the specific case of two-layer neural networks and Gaussian data in [BLN21]. Our result shows that in fact it is a *universal* phenomenon, which applies to essentially any parametrized function class (including in particular deep neural networks) as well as a much broader class of data distributions. As in [BLN21] we obtain an entire tradeoff curve between size and robustness: our universal law of robustness states that, for any function class smoothly parametrized by p parameters, and for any d-dimensional dataset satisfying mild regularity conditions, any function in this class that fits the data *below the noise level* must have its (Euclidean) Lipschitz constant larger than $\sqrt{\frac{nd}{p}}$.

Theorem 1 (Informal version of Theorem 3). *Let $\mathcal{F}$ be a class of functions from $\mathbb{R}^d \to \mathbb{R}$ and let $(x_i, y_i)_{i=1}^n$ be i.i.d. input-output pairs in $\mathbb{R}^d \times [-1, 1]$. Assume that:*

> *1. $\mathcal{F}$ admits a Lipschitz parametrization by p real parameters, each of size at most $poly(n, d)$.*

---

[1]As in, for instance, the inverse function theorem in analysis or Bézout's theorem in algebraic geometry.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

    2. The distribution μ of the covariates x_i satisfies isoperimetry (or is a mixture theoreof).

    3. The expected conditional variance of the output (i.e., the "noise level") is strictly positive, denoted $\sigma^2 := \mathbb{E}^{\mu}[Var[y|x]] > 0$.

Then, with high probability over the sampling of the data, one has simultaneously for all $f \in \mathcal{F}$:

$$\frac{1}{n}\sum_{i=1}^{n}(f(x_i) - y_i)^2 \le \sigma^2 - \epsilon \;\Rightarrow\; \mathrm{Lip}(f) \ge \widetilde{\Omega}\left(\sqrt{\frac{nd}{p}}\right).$$

Remark 1.1. For the distributions μ we have in mind, for instance uniform on the unit sphere, there exists with high probability some $O(1)$-Lipschitz function $f : \mathbb{R}^d \to \mathbb{R}$ satisfying $f(x_i) = y_i$ for all i. Indeed, with probability $1 - e^{-\Omega(d)}$ we have $||x_i - x_j|| \ge 1$ for all $1 \le i \ne j \le n$ so long as $n \le poly(d)$. In this case we may apply the Kirszbraun extension theorem to find a suitable f regardless of the labels y_i.

1.1 Speculative implication for real data

To put Theorem 1 in context, we compare to the empirical results presented in [MMS$^+$18]. In the latter work, they consider the MNIST dataset which consists of $n = 6 \times 10^4$ images in dimension $28^2 = 784$. They trained robustly different architectures, and reported in Figure 4 the size of the architecture versus the obtained robust test accuracy (third plot from the left). One can see a sharp transition from roughly 10% accuracy to roughly 90% accuracy at around 2×10^5 parameters (capacity scale 4 in their notation). Moreover the robust accuracy keeps climbing up with more parameters, to roughly 95% accuracy at roughly 3×10^6 parameters.

How can we compare these numbers to the law of robustness? There are a number of difficulties that we discuss below, and we emphasize that this discussion is highly speculative in nature, though we find that, with a few leaps of faith, our universal law of robustness sheds light on the potential parameter regimes of interest for robust deep learning.

The first difficulty is to evaluate the "correct" dimension of the problem. Certainly the number of pixels per image gives an upper bound, however one expects that the data lies on something like a lower dimensional sub-manifold. Optimistically, we hope that Theorem 1 will continue to apply for an appropriate *effective dimension* which may be rather smaller than the literal number of pixels. This hope is partially justified by the fact that isoperimetry holds in many less-than-picturesque situations, some of which are stated in the next subsection.

The next difficulty is to estimate/interpret the noise value σ^2. From a theoretical point of view, this noise assumption is necessary for otherwise there could exist a smooth classifier with perfect accuracy in $\mathcal{F}$, defeating the point of any lower bound on the size of $\mathcal{F}$. We tentatively would like to think of σ^2 as capturing the contribution of the "difficult" part of the learning problem, that is σ^2 could be thought of as the non-robust generalization error of reasonably good models, so a couple of % of error in the case of MNIST. With that interpretation, one gets "below the noise level" in MNIST with a training error of a couple of %.

Finally another subtlety to interpret the empirical results of [MMS$^+$18] is that there is a mismatch between what they measure and our quantities of interest. Namely the law of robustness talks about two things: the training error, and the worst-case robustness (i.e., the Lipschitz constant). On the other hand [MMS$^+$18] measures the *robust generalization error*. Understanding the interplay between those three quantities is a fantastic open problem. Here we take the perspective that a small robust generalization error should imply a small training error and a small Lipschitz constant. Another important mismatch is that we stated our universal law of robustness for Lipschitzness in ℓ_2, while the experiments in [MMS$^+$18] are for robustness in ℓ_∞. We believe that a variant of the law of robustness remains true for ℓ_∞, a belief again partially justified by how broad isoperimetry is (see next subsection).

With all the caveats described above, we can now look at the numbers as follows: in the [MMS$^+$18] experiments, smooth models with accuracy below the noise level are attained with a number of

parameters somewhere in the range $2 \times 10^5 - 3 \times 10^6$ parameters (possibly even larger depending on the interpretation of the noise level), while the law of robustness would predict any such model must have at least nd parameters, and this latter quantity should be somewhere in the range $10^6 - 10^7$ (corresponding to an effective dimension between 15 and 150). While far from perfect, the law of robustness prediction is far more accurate than the classical rule of thumb # parameters $\simeq$ # equations (which here would predict a number of parameters of the order 10^4).

Perhaps more interestingly, one could apply a similar reasoning to the ImageNet dataset, which consists of 1.4×10^7 images of size roughly 2×10^5. Estimating that the effective dimension is a couple of order of magnitudes smaller than this size, the law of robustness predicts that to obtain good robust models on ImageNet one would need at least $10^{10} - 10^{11}$ parameters. This number is larger than the size of current neural networks trained robustly for this task, which sports between $10^8 - 10^9$ parameters. Thus, we arrive at the tantalizing possibility that robust models for ImageNet do not exist yet simply because we are a couple orders of magnitude off in the current scale of neural networks trained for this task.

1.2 Related work

Theorem 1 is a direct follow-up to the conjectured law of robustness in [BLN21] for (arbitrarily weighted) two-layer neural networks with Gaussian data. Our result does not actually prove their conjecture, because we assume here polynomially bounded weights. While this assumption is reasonable from a practical perspective, it remains mathematically interesting to prove the full conjecture for the two-layer case. We prove however in Section 4 that the polynomial weights assumption is necessary as soon as one considers three-layer neural networks.

In addition to [MMS+18], several recent works have experimentally studied the relationship between a neural network scale and its achieved robustness, see e.g., [XY20, GQU+21]. It has been consistently reported that larger networks help tremendously for robustness, beyond what is typically seen for classical non-robust accuracy. We view our universal law of robustness as putting this empirical observation on a more solid footing: scale is actually *necessary* to achieve robustness.

The law of robustness setting is closely related to the interpolation setting: in the former case one considers models optimizing "beyond the noise level", while in the latter case one studies models with perfect fit on the training data. The study of generalization in this interpolation regime has been a central focus of learning theory in the last few years (see e.g., [BHMM19, MM19, BLLT20, NKB+20]), as it seemingly contradicts classical theory about regularization. More broadly though, generalization remains a mysterious phenomon in deep learning, and the exact interplay between the law of robustness' setting (interpolation regime/worst-case robustness) and (robust) generalization error is a fantastic open problem. Interestingly, we note that one could potentially avoid the conclusion of the law of robustness (that is, that large models are necessary for robustness), with early stopping methods that could stop the optimization once the noise level is reached. In fact, this theoretically motivated suggestion has already been empirically tested and confirmed in the recent work [RWK20], showing again a close tie between the conclusions one can draw from the law of robustness and actual practical settings.

Classical lower bounds on the gradient of a function include Poincaré type inequalities, but they are of a qualitately different nature compared to the law of robustness lower bound. We recall that a measure μ on $\mathbb{R}^d$ satisfies a Poincaré inequality if for any function f, one has $\mathbb{E}^\mu[\|\nabla f\|^2] \geq C \cdot \mathrm{Var}(f)$ (for some constant $C > 0$). In our context, such a lower bound has essentially no consequence since the variance of a function interpolating a finite dataset could be exponentially small. In fact this is tight, as one easily use similar constructions to those in [BLN21] to show that one can interpolate with an exponentially small expected norm squared of the gradient (in particular it is crucial in the law of robustness to consider the Lipschitz constant, i.e., the supremum of the norm of the gradient). On the other hand, our isoperimetry assumption is related to a certain strenghtening of the Poincaré inequality known as log-Sobolov inequality (see e.g., [Led01]). If the covariate measure satisfies only a Poincaré inequality, then we could prove a weaker law of robustness of the form $\mathrm{Lip} \gtrsim \frac{n\sqrt{d}}{p}$ (using for example the concentration result obtained in [BL97]).

1.3 Isoperimetry

Concentration of measure and isoperimetry are perhaps the most ubiquitous features of high-dimensional geometry. In short, they assert in many cases that Lipschitz functions on high-dimensional space concentrate tightly around their mean. Our result assumes that the distribution μ of the covariates x_i satisfies such an inequality in the following sense.

Definition 1.1. *A probability measure μ on $\mathbb{R}^d$ satisfies c-isoperimetry if for any bounded L-Lipschitz $f : \mathbb{R}^d \to \mathbb{R}$,*

$$\mathbb{P}[|f(x) - \mathbb{E}[f]| \geq t] \leq 2e^{-\frac{dt^2}{2cL^2}}. \tag{1.1}$$

In general, if a scalar random variable X satisfies $\mathbb{P}[|X| \geq t] \leq 2e^{-t^2/C}$ then X is said to be C-subgaussian. Hence isoperimetry states that the output of any Lipschitz function is $O(1)$-subgaussian under suitable rescaling. Distributions satisfying $O(1)$-isoperimetry include high dimensional Gaussians $\mu = \mathcal{N}\left(0, \frac{I_d}{d}\right)$ and uniform distributions on spheres and hypercubes (normalized to have diameter 1). Isoperimetry also holds for mild perturbations of these idealized scenarios, including[2]:

- The sum of a Gaussian and an independent random vector of small norm [CCNW21].

- Strongly log-concave measures in any normed space [BL00, Proposition 3.1].

- Manifolds with positive Ricci curvature [Gro86, Theorem 2.2].

Due to the last condition above, we believe our results are realistic even under the *manifold hypothesis* that high-dimensional data tends to lie on a lower-dimensional submanifold. This viewpoint on learning has been studied for decades, see e.g. [HS89, KL93, RS00, TDSL00, NM10, FMN16]. We also note that our formal theorem (Theorem 3) actually applies to distributions that can be written as a mixture of distributions satisfying isoperimetry.

Our proofs will repeatedly use the following simple fact:

Proposition 1.2. *[Ver18, Proposition 2.6.1],[vH14, Exercise 3.1] If $X_1, \ldots, X_n$ are independent, C-subgaussian, with mean 0, then $\frac{1}{\sqrt{n}} \sum_{i=1}^{n} X_i$ is $18C$-subgaussian.*

2 A finite approach to the law of robustness

For the function class of two-layer neural networks, [BLN21] investigated several approaches to prove the law of robustness. At a high level, the proof strategies there relied on various ways to measure how "large" the set of two-layer neural networks can be (specifically, they tried a geometric approach based on relating to multi-index models, a statistical approach based on the Rademacher complexity, and an algebraic approach for the case of polynomial activations).

In this work we take here a different route: we shift the focus from the function class $\mathcal{F}$ to an *individual* function $f \in \mathcal{F}$. Namely, our proof starts by asking the following question: for a fixed function f, what is the probability that it would give a good approximate fit on the (random) data? For simplicity, consider for a moment the case where we require f to actually interpolate the data (i.e., perfect fit), and say that y_i are random ± 1 labels. The key insight is that isoperimetry implies that *either* the 0-level set of f *or* the 1-level set of f must have probability smaller than $\exp\left(-\frac{d}{\text{Lip}(f)^2}\right)$. Thus, the probability that f fits all the n points is at most $\exp\left(-\frac{nd}{\text{Lip}(f)^2}\right)$ so long as both labels $y_i \in \{-1, 1\}$ actually appear a constant fraction of the time. In particular, using an union bound[3], for a finite function class $\mathcal{F}$ of size N with L-Lipschitz functions, the probability that there exists a

---

[2]The first two examples satisfy a logarithmic Sobolev inequality, which implies isoperimetry [Led99, Proposition 2.3].

[3]In this informal argument we ignore the possibility that the labels y_i are not well-balanced. Note that the probability of this rare event is not amplified by a union bound over $f \in \mathcal{F}$.

function $f \in \mathcal{F}$ fitting the data is at most

$$N \exp\left(-\frac{nd}{L^2}\right) = \exp\left(\log(N) - \frac{nd}{L^2}\right).$$

Thus we see that, if $L \ll \sqrt{\frac{nd}{\log(N)}}$, then the probability of finding a fitting function in $\mathcal{F}$ is very small. This basically concludes the proof, since via a standard discretization argument, for a smoothly parametrized family with p (bounded) parameters one expects $\log(N) = \tilde{O}(p)$.

We now give the formal proof, which applies in particular to approximate fit rather than exact fit in the argument above. The only difference is that we will identify a well-chosen subgaussian random variable in the problem. We start with the finite function class case:

Theorem 2. *Let (x_i, y_i) be i.i.d. input-output pairs in $\mathbb{R}^d \times [-1, 1]$ such that:*

 1. *The distribution μ of the covariates x_i can be written as $\mu = \sum_{\ell=1}^{k} \alpha_\ell \mu_\ell$, where each μ_ℓ satisfies c-isoperimetry and $\alpha_\ell \geq 0, \sum_{\ell=1}^{k} \alpha_\ell = 1$.*

 2. *The expected conditional variance of the output is strictly positive, denoted $\sigma^2 := \mathbb{E}^\mu[Var[y|x]] > 0$.*

Then one has:

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} (y_i - f(x_i))^2 \leq \sigma^2 - \epsilon \text{ and } \mathrm{Lip}(f) \leq L\right)$$

$$\leq 4k \exp\left(-\frac{n\epsilon^2}{8^3 k}\right) + 2\exp\left(\log(|\mathcal{F}|) - \frac{\epsilon^2 nd}{8^4 c L^2}\right).$$

We start with a lemma showing that, to optimize beyond the noise level one must necessarily correlate with the noise part of the labels. In what follows we denote $g(x) = \mathbb{E}[y|x]$ for the target function, and $z_i = y_i - g(x_i)$ for the noise part of the observed labels (namely y_i is the sum of the target function $g(x_i)$ and the noise term z_i).

Lemma 2.1. *One has*

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} (y_i - f(x_i))^2 \leq \sigma^2 - \epsilon\right) \leq 2\exp\left(-\frac{n\epsilon^2}{8^3}\right) + \mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} f(x_i) z_i \geq \frac{\epsilon}{3}\right).$$

Proof. The sequence (z_i^2) is i.i.d., with mean σ^2, and such that $|z_i|^2 \leq 4$. Thus Hoeffding's inequality yields:

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^{n} z_i^2 \leq \sigma^2 - \frac{\epsilon}{6}\right) \leq \exp\left(-\frac{n\epsilon^2}{8^3}\right). \tag{2.1}$$

On the other hand the sequence $(z_i g(x_i))$ is i.i.d., with mean 0 (since $\mathbb{E}[z_i|x_i] = 0$), and such that $|z_i g(x_i)| \leq 2$, Thus Hoeffding's inequality yields:

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^{n} z_i g(x_i) \leq -\frac{\epsilon}{6}\right) \leq \exp\left(-\frac{n\epsilon^2}{8^3}\right). \tag{2.2}$$

Let us write $Z = \frac{1}{\sqrt{n}}(z_1, \ldots, z_n), G = \frac{1}{\sqrt{n}}(g(x_1,) \ldots, g(x_n))$, and $F = \frac{1}{\sqrt{n}}(f(x_1), \ldots, f(x_n))$. We claim that if $\|Z\|^2 \geq \sigma^2 - \frac{\epsilon}{6}$ and $\langle Z, G\rangle \geq -\frac{\epsilon}{6}$, then for any $f \in \mathcal{F}$ one has

$$\|G + Z - F\|^2 \leq \sigma^2 - \epsilon \Rightarrow \langle F, Z\rangle \geq \frac{\epsilon}{3}.$$

This claim together with (2.1) and (2.2) conclude the proof. On the other hand the claim itself directly follows from:

$$\sigma^2 - \epsilon \geq \|G + Z - F\|^2 = \|Z + G - F\|^2 = \|Z\|^2 + 2\langle Z, G - F\rangle + \|G - F\|^2 \geq \sigma^2 - \frac{\epsilon}{3} - 2\langle Z, F\rangle.$$

$\square$

211 We can now proceed to the proof of Theorem 2:

212 *Proof.* First note that without loss of generality we can assume that the range of any function in $\mathcal{F}$
213 is included in $[-1,1]$ (indeed clipping the values improves both the fit to any $y \in [-1,1]$ and the
214 Lipschitz constant). We also assume wlog that all functions in $\mathcal{F}$ are L-Lipschitz.

215 For clarity let us start with the case $k = 1$. By the isoperimetry assumption we have that
216 $\sqrt{\frac{d}{c}} \frac{f(x_i) - \mathbb{E}[f]}{L}$ is 1-subgaussian. Since $|z_i| \leq 2$, we also have that $\sqrt{\frac{d}{c}} \frac{(f(x_i) - \mathbb{E}[f]) z_i}{L}$ is 4-
217 subgaussian. Moreover, the latter random variable has zero-mean since $\mathbb{E}[z|x] = 0$. Thus by
218 Proposition 1.2 we have:

$$\mathbb{P}\left(\sqrt{\frac{d}{cnL^2}} \sum_{i=1}^{n} (f(x_i) - \mathbb{E}[f]) z_i \geq t \right) \leq 2 \exp\left(-(t/9)^2 \right) .$$

219 We rewrite the above as:

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^{n} (f(x_i) - \mathbb{E}[f]) z_i \geq \frac{\epsilon}{6} \right) \leq 2 \exp\left(-\frac{\epsilon^2 n d}{8^4 c L^2} \right) . \tag{2.3}$$

220 Since we assumed that the range of the functions is in $[-1,1]$ we have $\mathbb{E}[f] \in [-1,1]$ and hence:

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} \mathbb{E}[f] z_i \geq \frac{\epsilon}{6} \right) \leq \mathbb{P}\left(\left| \frac{1}{n} \sum_{i=1}^{n} z_i \right| \geq \frac{\epsilon}{6} \right) . \tag{2.4}$$

221 (This step is the analog of requiring the labels y_i to be well-balanced in the example of perfect
222 interpolation.) By Hoeffding's inequality, the above quantity is smaller than $2 \exp(-n\epsilon^2/8^3)$ (recall
223 that $|z_i| \leq 2$). Thus we obtain with an union bound:

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} f(x_i) z_i \geq \frac{\epsilon}{3} \right) \quad \leq \quad |\mathcal{F}| \cdot \mathbb{P}\left(\frac{1}{n} \sum_{i=1}^{n} (f(x_i) - \mathbb{E}[f]) z_i \geq \frac{\epsilon}{6} \right) + \mathbb{P}\left(\left| \frac{1}{n} \sum_{i=1}^{n} z_i \right| \geq \frac{\epsilon}{6} \right)$$

$$\leq \quad 2|\mathcal{F}| \cdot \exp\left(-\frac{\epsilon^2 n d}{8^4 c L^2} \right) + 2 \exp\left(-\frac{n \epsilon^2}{8^3} \right) .$$

224 Together with Lemma 2.1 this concludes the proof for $k = 1$.

225

226 We now turn to the case $k > 1$. We first sample the mixture component $\ell_i \in [k]$ for each data point
227 $i \in [n]$, and we now reason conditioned on these mixture components. Let $S_\ell \subset [n]$ be the set of
228 data points sampled from mixture component $\ell \in [k]$, that is $x_i, i \in S_\ell$, is i.i.d. from μ_ℓ. We now
229 have that $\sqrt{\frac{d}{c}} \frac{f(x_i) - \mathbb{E}^{\mu_{\ell_i}}[f]}{L}$ is 1-subgaussian (notice that the only difference is that now we need to
230 center by $\mathbb{E}^{\mu_{\ell_i}}[f]$, which depends on the mixture component). In particular using the same reasoning
231 as for (2.4) we obtain (crucially note that Proposition 1.2 does not require the random variables to
232 be identically distributed):

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^{n} (f(x_i) - \mathbb{E}^{\mu_{\ell_i}}[f]) z_i \geq \frac{\epsilon}{6} \right) \leq 2 \exp\left(-\frac{\epsilon^2 n d}{8^4 c L^2} \right) . \tag{2.5}$$

233 Next we want to appropriately modify (2.4). To do so note that:

$$\max_{m_1, \ldots, m_k \in [-1,1]} \sum_{i=1}^{n} m_{\ell_i} z_i = \sum_{\ell=1}^{k} \left| \sum_{i \in S_\ell} z_i \right| ,$$

234 so that the we can rewrite (2.4) as:

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^{n} \mathbb{E}^{\mu_{\ell_i}}[f] z_i \geq \frac{\epsilon}{6} \right) \leq \mathbb{P}\left(\frac{1}{n} \sum_{\ell=1}^{k} \left| \sum_{i \in S_\ell} z_i \right| \geq \frac{\epsilon}{6} \right) .$$

235 Now note that $\sum_{\ell=1}^{k} \sqrt{|S_\ell|} \leq \sqrt{nk}$ and thus we have:

$$\mathbb{P}\left(\frac{1}{n} \sum_{\ell=1}^{k} \left| \sum_{i \in S_\ell} z_i \right| \geq \frac{\epsilon}{6} \right) \leq \mathbb{P}\left(\sum_{\ell=1}^{k} \left| \sum_{i \in S_\ell} z_i \right| \geq \frac{\epsilon}{6} \sqrt{\frac{n}{k}} \sum_{\ell=1}^{k} \sqrt{|S_\ell|} \right) \leq \sum_{\ell=1}^{k} \mathbb{P}\left(\left| \sum_{i \in S_\ell} z_i \right| \geq \frac{\epsilon}{6} \sqrt{\frac{n}{k}} \sqrt{|S_\ell|} \right) .$$

236 Finally by Hoeffding's inequality, we have for any $\ell \in [k]$, $\mathbb{P}\left(\left|\sum_{i \in S_\ell} z_i\right| \geq t\sqrt{|S_\ell|}\right) \leq$

237 $2\exp\left(-\frac{t^2}{8}\right)$, and thus the last display is bounded from above by $2k\exp\left(-\frac{n\epsilon^2}{8^3 k}\right)$. The proof can

238 now be concluded as we did for $k = 1$. $\qquad\square$

239 Finally we can now state and prove the formal version of the informal theorem stated in the intro-
240 duction.

241 **Theorem 3.** *Let $\mathcal{F}$ be a class of functions from $\mathbb{R}^d \to \mathbb{R}$ and let $(x_i, y_i)_{i=1}^n$ be i.i.d. input-output*
242 *pairs in $\mathbb{R}^d \times [-1, 1]$. Fix $\epsilon, \delta \in (0, 1)$. Assume that:*

    243 *1. The function class can be written as $\mathcal{F} = \{f_{\boldsymbol{w}}, \boldsymbol{w} \in \mathcal{W}\}$ with $\mathcal{W} \subset \mathbb{R}^p$, $\mathrm{diam}(\mathcal{W}) \leq W$*
    244 *and for any $\boldsymbol{w}_1, \boldsymbol{w}_2 \in \mathcal{W}$,*

$$||f_{\boldsymbol{w}_1} - f_{\boldsymbol{w}_2}||_\infty \leq J||\boldsymbol{w}_1 - \boldsymbol{w}_2||.$$

    245 *2. The distribution μ of the covariates x_i can be written as $\mu = \sum_{\ell=1}^k \alpha_\ell \mu_\ell$, where each μ_ℓ*
    246 *satisfies c-isoperimetry, $\alpha_\ell \geq 0$, $\sum_{\ell=1}^k \alpha_\ell = 1$, and k is such that $8^4 k \log(8k/\delta) \leq n\epsilon^2$.*

    247 *3. The expected conditional variance of the output is strictly positive, denoted $\sigma^2 :=$*
    248 $\mathbb{E}^\mu[Var[y|x]] > 0$.

249 *Then, with probability at least $1 - \delta$ with respect to the sampling of the data, one has simultaneously*
250 *for all $f \in \mathcal{F}$:*

$$\frac{1}{n}\sum_{i=1}^n (f(x_i) - y_i)^2 \leq \sigma^2 - \epsilon \implies \mathrm{Lip}(f) \geq \frac{\epsilon}{2^9\sqrt{c}}\sqrt{\frac{nd}{p\log(60WJ\epsilon^{-1}) + \log(4/\delta)}}.$$

251 *Proof.* Define $\mathcal{W}_L \subseteq \mathcal{W}$ by $\mathcal{W}_L = \{\boldsymbol{w} \in \mathcal{W} : \mathrm{Lip}(f_{\boldsymbol{w}}) \leq L\}$. Denote $\mathcal{W}_{L,\epsilon}$ for an $\frac{\epsilon}{6J}$-net of $\mathcal{W}_L$.
252 We have in particular $|\mathcal{W}_\epsilon| \leq (60WJ\epsilon^{-1})^p$. We apply Theorem 2 to $\mathcal{F}_{L,\epsilon} = \{f_{\boldsymbol{w}}, \boldsymbol{w} \in \mathcal{W}_{L,\epsilon}\}$:

$$\mathbb{P}\left(\exists f \in \mathcal{F}_{L,\epsilon} : \frac{1}{n}\sum_{i=1}^n (y_i - f(x_i))^2 \leq \sigma^2 - \frac{\epsilon}{2} \text{ and } \mathrm{Lip}(f) \leq 2L\right)$$

$$\leq 4k\exp\left(-\frac{n\epsilon^2}{8^4 k}\right) + 2\exp\left(p\log(60WJ\epsilon^{-1}) - \frac{\epsilon^2 nd}{8^6 cL^2}\right).$$

253 Observe that if $||f - g||_\infty \leq \frac{\epsilon}{6}$ and $||y||_\infty, ||f||_\infty, ||g||_\infty \leq 1$, then $\frac{1}{n}\sum_{i=1}^n (y_i - f(x_i))^2 \leq \frac{\epsilon}{2} +$
254 $\frac{1}{n}\sum_{i=1}^n (y_i - g(x_i))^2$. (We may again assume without loss of generality that all functions in $\mathcal{F}$ map
255 to $[-1, 1]$.) Thus we obtain for any $L > 0$:

$$\mathbb{P}\left(\exists f \in \mathcal{F} : \frac{1}{n}\sum_{i=1}^n (y_i - f(x_i))^2 \leq \sigma^2 - \epsilon \text{ and } \mathrm{Lip}(f) \leq L\right)$$

$$\leq 4k\exp\left(-\frac{n\epsilon^2}{8^4 k}\right) + 2\exp\left(p\log(60WJ\epsilon^{-1}) - \frac{\epsilon^2 nd}{8^6 cL^2}\right).$$

256 The first assumption ensures that for any $\boldsymbol{w} \in \mathcal{W}_L$, there is $\boldsymbol{w}' \in \mathcal{W}_{L,\epsilon}$ with $||f_{\boldsymbol{w}} -$
257 $f_{\boldsymbol{w}'}||_\infty \leq \frac{\epsilon}{6}$. The second assumption shows the probability just above is at most δ when
258 $L = \frac{\epsilon}{2^9\sqrt{c}}\sqrt{\frac{nd}{p\log(60WJ\epsilon^{-1}) + \log(4/\delta)}}$. This concludes the proof. $\qquad\square$

3 Deep neural networks

260 We now specialize the law of robustness (Theorem 3) to multi-layer neural networks. We consider
261 a rather general class of depth D neural networks described as follows. First, we require that the
262 neurons are partitioned into layers $\mathcal{L}_1, \ldots, \mathcal{L}_D$, and that all connections are from $\mathcal{L}_i \to \mathcal{L}_j$ for
263 some $i < j$. This includes the basic feed-forward case in which only connections $\mathcal{L}_i \to \mathcal{L}_{i+1}$ are
264 used as well as more general skip connections. We specify (in the natural way) a neural network by
265 matrices W_j of shape $|\mathcal{L}_j| \times \sum_{i<j} |\mathcal{L}_i|$ for each $1 \leq j \leq D$, as well as 1-Lipschitz non-linearities

$\sigma_{j,\ell}$ and scalar biases $b_{j,\ell}$ for each (j,ℓ) satisfying $\ell \in |\mathcal{L}_j|$. We use fixed non-linearities $\sigma_{j,\ell}$ as well as a fixed architecture, in the sense that each matrix entry $W_j[k,\ell]$ is either always 0 or else it is variable (and similarly for the bias terms).

To match the notation of Theorem 3, we identify the parametrization in terms of the matrices (W_j) and bias terms $(b_{j,\ell})$ to a single p-dimensional vector $\boldsymbol{w}$ as follows. A variable matrix entry $W_j[k,\ell]$ is set to $w_{a(j,k,\ell)}$ for some fixed index $a(j,k,\ell) \in [p]$, and a variable bias term $b_{j,\ell}$ is set to $w_{a(j,\ell)}$ for some $a(j,\ell) \in [p]$. Thus we now have a parametrization $\boldsymbol{w} \in \mathbb{R}^p \mapsto f_{\boldsymbol{w}}$ where $f_{\boldsymbol{w}}$ is the neural network represented by the parameter vector $\boldsymbol{w}$. Importantly, note that our formulation allows for weight sharing (in the sense that a shared weight is counted only as a single parameter). For example, this is important to obtain an accurate count of the number of parameters in convolutional architectures.

In order to apply Theorem 3 to this class of functions we need to estimate the Lipschitz constant of the parametrization $\boldsymbol{w} \mapsto f_{\boldsymbol{w}}$. To do this we introduce three more quantities. First, we shall assume that all the parameters are bounded in magnitude by W, that is we consider the set of neural networks parametrized by $\boldsymbol{w} \in [-W,W]^p$. Next, for the architecture under consideration, denote Q for the maximum number of matrix entries/bias terms that are tied to a single parameter w_a for some $a \in [p]$. Finally we define

$$B(\boldsymbol{w}) = \max_{i \in \{0,1,\dots,D\}} \prod_{j \in [D], j \neq i} \|W_j\|_{op}, .$$

(The choice $i = 0$ gives the full product while all other choices give the product of $D - 1$ of the D operator norms.) Observe that $B(\boldsymbol{w})$ is an upper bound on the Lipschitz constant of the network itself, i.e., the map $x \mapsto f_{\boldsymbol{w}}(x)$. It turns out that a uniform control on it also controls the Lipschitz constant of the *parametrization* $\boldsymbol{w} \mapsto f_{\boldsymbol{w}}$. Namely we have the following lemma:

Lemma 3.1. *Let $x \in \mathbb{R}^d$ such that $\|x\| \le R$, and $\boldsymbol{w}_1, \boldsymbol{w}_2 \in \mathbb{R}^p$ such that $B(\boldsymbol{w}_1), B(\boldsymbol{w}_2) \le \overline{B}$. Then one has*

$$|f_{\boldsymbol{w}_1}(x) - f_{\boldsymbol{w}_2}(x)| \le \overline{B}QR\sqrt{p}\|\boldsymbol{w}_1 - \boldsymbol{w}_2\|\,.$$

Moreover for any $\boldsymbol{w} \in [-W,W]^p$ with $W \ge 1$, one has

$$B(\boldsymbol{w}) \le (W\sqrt{pQ})^D.$$

Proof. Fix an input x and define g_x by $g_x(\boldsymbol{w}) = f_{\boldsymbol{w}}(x)$. A standard gradient calculation for multi-layer neural networks directly shows that $\|\nabla g_x(\boldsymbol{w})\|_\infty \le B(\boldsymbol{w})QR$ so that $\|\nabla g_x(\boldsymbol{w})\| \le B(\boldsymbol{w})QR\sqrt{p}$. Moreover, the map $\boldsymbol{w} \mapsto B(\boldsymbol{w})$ is convex since the matrix operator norm is convex (and nonnegative), which implies that $B(\boldsymbol{w}) \le \overline{B}$ on the entire segment $[\boldsymbol{w}_1, \boldsymbol{w}_2]$, and thus $\|\nabla g_x(\boldsymbol{w})\| \le \overline{B}QR\sqrt{p}$ on that segment, which concludes the proof of the first claimed inequality. The second claimed inequality follows directly from $\|W_j\|_{op} \le \|W_j\|_2 \le W\sqrt{pQ}$. $\qquad\square$

Lemma 3.1 shows that when applying Theorem 3 to our class of neural networks one can always take $J = R(WQp)^D$ (assuming that the covariate measure μ is supported on the ball of radius R). Thus in this case the law of robustness (under the assumptions of Theorem 3) directly states that with high probability, any neural network in our class that fits the training data well below the noise level must also have:

$$\mathrm{Lip}(f) \ge \tilde{\Omega}\left(\sqrt{\frac{nd}{Dp}}\right), \tag{3.1}$$

where $\tilde{\Omega}$ hides logarithmic factors in W, p, R, Q, and the probability of error δ. Thus we see that the law of robustness, namely that the number of parameters should be at least nd for a smooth model with low training error, remains intact for constant depth neural networks. If taken at face value, the lower bound (3.1) suggests that it is better in practice to distribute the parameters towards *depth* rather than *width*, since the lower bound is decreasing with D. On the other hand, we note that (3.1) can be strengthened to:

$$\mathrm{Lip}(f) \ge \tilde{\Omega}\left(\sqrt{\frac{nd}{p\log(\overline{B})}}\right), \tag{3.2}$$

for the class of neural networks such that $B(\boldsymbol{w}) \leq \overline{B}$. In other words the dependency on the depth all but disappears by simply assuming that the quantity $B(\boldsymbol{w})$ (a natural upper bound on the Lipschitz constant of the network) is polynomially controlled. Interestingly many works have suggested to keep $B(\boldsymbol{w})$ under control, either for regularization purpose (for example [BFT17] relates $B(\boldsymbol{w})$ to the Rademacher complexity of multi-layer neural networks) or to simply control gradient explosion during training, see e.g., [ASB16, CBG$^+$17, MHRB17, MKKY18, JCC$^+$19, YM17]. Moreover, in addition to being well-motivated in practice, the assumption that $\overline{B}$ is polynomially controlled seems also somewhat unavoidable in theory, since $B(\boldsymbol{w})$ is an *upper bound* on the Lipschitz constant $\mathrm{Lip}(f_{\boldsymbol{w}})$. Thus a theoretical construction showing that the lower bound in (3.1) is tight (at some large depth D) would necessarily need to have an exponential gap between $\mathrm{Lip}(f_{\boldsymbol{w}})$ and $B(\boldsymbol{w})$. We are not aware of any such example, and it would be interesting to fully elucidate the role of depth in the law of robustness (particularly if it could give recommendation on how to best distribute parameters in a neural network).

4 Necessity of Polynomially Bounded Weights

In [BLN21] it was conjectured that the law of robustness should hold for the class of *all* two-layer neural networks. In this paper we prove that in fact it holds for arbitrary smoothly parametrized function classes, as long as the parameters are of size at most polynomial. In this section we demonstrate that this polynomial size restriction is in fact necessary for bounded depth neural networks.

First we note that *some* restriction on the size of the parameters is certainly necessary in the most general case. Indeed one can build a single-parameter family, where the single real parameter is used to approximately encode all Lipschitz functions from a compact set in $\mathbb{R}^d$ to $[-1, 1]$, simply by brute-force enumeration. In particular no tradeoff between number of parameters and attainable Lipschitz constant would exist for this function class.

Showing a counter-example to the law of robustness with unbounded parameters and "reasonable" function classes is slightly harder. Here we build a three-layer neural network, with a single fixed nonlinearity $\sigma : \mathbb{R} \to \mathbb{R}$, but the latter is rather complicated and we do not know how to describe it explicitly (it is based on the Kolmogorov-Arnold theorem). It would be interesting to derive similar construction with say ReLU networks.

Theorem 4. *For each $d \in \mathbb{Z}^+$ there is a continuous function $\sigma : \mathbb{R} \to \mathbb{R}$ and a sequence $(b_\ell)_{\ell \leq 2^{2^d}}$ such that the following holds. The function Φ_a defined by*

$$\Phi_a(x) = \sum_{\ell=1}^{2^{2^d}} \sigma(a - \ell) \sum_{i=1}^{2d} \sigma\left(b_\ell + \sum_{j=1}^{d} \sigma(x_j + b_\ell)\right), \qquad |a| \leq 2^{2^d} \tag{4.1}$$

is always $O(d^{3/2})$-Lipschitz, and the parametrization $a \to \Phi_a$ is 1-Lipschitz. Moreover for $n \leq \frac{2^d}{100}$, given i.i.d. uniform points $x_1, \ldots, x_n \in \mathbb{S}^{d-1}$ and random labels $y_1, \ldots, y_n \in \{-1, 1\}$, with probability $1 - e^{-\Omega(d)}$ there exists $\ell \in [2^{2^d}]$ such that $\Phi_\ell(x_i) = y_i$ for at least $\frac{3n}{4}$ of the values $i \in [n]$.

Theorem 4 is proved in the appendix.

Remark 4.1. The representation (4.1) is a three-layer neural network because the $\sigma(a - \ell)$ terms are just matrix entries for the final layer.

References

[ASB16] Martin Arjovsky, Amar Shah, and Yoshua Bengio. Unitary evolution recurrent neural networks. In *International Conference on Machine Learning*, pages 1120–1128. PMLR, 2016.

[BB19] Roland Bauerschmidt and Thierry Bodineau. A very simple proof of the lsi for high temperature spin systems. *Journal of Functional Analysis*, 276(8):2582–2588, 2019.

[BFT17] Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach,

R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017.

[BHMM19] Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019.

[BL97] Sergey Bobkov and Michel Ledoux. Poincaré's inequalities and talagrand's concentration phenomenon for the exponential distribution. *Probability Theory and Related Fields*, 107(3):383–400, 1997.

[BL00] Sergey G Bobkov and Michel Ledoux. From brunn-minkowski to brascamp-lieb and to logarithmic sobolev inequalities. *Geometric & Functional Analysis GAFA*, 10(5):1028–1052, 2000.

[BLLT20] Peter L. Bartlett, Philip M. Long, Gábor Lugosi, and Alexander Tsigler. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020.

[BLN21] Sébastien Bubeck, Yuanzhi Li, and Dheeraj Nagaraj. A law of robustness for two-layers neural networks. *Conference on Learning Theory (COLT)*, 2021.

[CBG⁺17] Moustapha Cisse, Piotr Bojanowski, Edouard Grave, Yann Dauphin, and Nicolas Usunier. Parseval networks: Improving robustness to adversarial examples. In *International Conference on Machine Learning*, pages 854–863. PMLR, 2017.

[CCNW21] Hong-Bin Chen, Sinho Chewi, and Jonathan Niles-Weed. Dimension-free log-sobolev inequalities for mixture distributions. *arXiv preprint arXiv:2102.11476*, 2021.

[FMN16] Charles Fefferman, Sanjoy Mitter, and Hariharan Narayanan. Testing the manifold hypothesis. *Journal of the American Mathematical Society*, 29(4):983–1049, 2016.

[GQU⁺21] Sven Gowal, Chongli Qin, Jonathan Uesato, Timothy Mann, and Pushmeet Kohli. Uncovering the limits of adversarial training against norm-bounded adversarial examples, 2021.

[Gro86] Mikhael Gromov. Isoperimetric inequalities in riemannian manifolds. In *Asymptotic Theory of Finite Dimensional Spaces*, volume 1200, pages 114–129. Springer Berlin, 1986.

[HS89] Trevor Hastie and Werner Stuetzle. Principal curves. *Journal of the American Statistical Association*, 84(406):502–516, 1989.

[JCC⁺19] Haoming Jiang, Zhehui Chen, Minshuo Chen, Feng Liu, Dingding Wang, and Tuo Zhao. On computation and generalization of gans with spectrum control. *Proc. of International Conference on Learning Representation (ICLR)*, 2019.

[KL93] Nanda Kambhatla and Todd K Leen. Fast nonlinear dimension reduction. In *IEEE International Conference on Neural Networks*, pages 1213–1218. IEEE, 1993.

[Led99] Michel Ledoux. Concentration of measure and logarithmic sobolev inequalities. In *Seminaire de probabilites XXXIII*, pages 120–216. Springer, 1999.

[Led01] M. Ledoux. The concentration of measure phenomenon. In *Mathematical Surveys and Monographs*, volume 89. American Mathematical Society, Providence, RI, 2001.

[MHRB17] Zakaria Mhammedi, Andrew Hellicar, Ashfaqur Rahman, and James Bailey. Efficient orthogonal parametrisation of recurrent neural networks using householder reflections. In *International Conference on Machine Learning*, pages 2401–2409. PMLR, 2017.

[MKKY18] Takeru Miyato, Toshiki Kataoka, Masanori Koyama, and Yuichi Yoshida. Spectral normalization for generative adversarial networks. *Proc. of International Conference on Learning Representation (ICLR)*, 2018.

[MM19] Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and double descent curve. *arXiv preprint arXiv:1908.05355*, 2019.

[MMS⁺18] Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. *Proc. of International Conference on Learning Representation (ICLR)*, 2018.

[NKB⁺20] Preetum Nakkiran, Gal Kaplun, Yamini Bansal, Tristan Yang, Boaz Barak, and Ilya Sutskever. Deep double descent: Where bigger models and more data hurt. In *International Conference on Learning Representations*, 2020.

[NM10] Hariharan Narayanan and Sanjoy Mitter. Sample complexity of testing the manifold hypothesis. In *Proceedings of the 23rd International Conference on Neural Information Processing Systems-Volume 2*, pages 1786–1794, 2010.

[RS00] Sam T Roweis and Lawrence K Saul. Nonlinear dimensionality reduction by locally linear embedding. *Science*, 290(5500):2323–2326, 2000.

[RWK20] Leslie Rice, Eric Wong, and Zico Kolter. Overfitting in adversarially robust deep learning. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 8093–8104. PMLR, 2020.

[TDSL00] Joshua B Tenenbaum, Vin De Silva, and John C Langford. A global geometric framework for nonlinear dimensionality reduction. *science*, 290(5500):2319–2323, 2000.

[Ver18] Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.

[vH14] Ramon van Handel. Probability in high dimension. Technical report, Princeton University, 2014.

[XY20] Cihang Xie and Alan Yuille. Intriguing properties of adversarial training at scale. In *International Conference on Learning Representations*, 2020.

[YM17] Yuichi Yoshida and Takeru Miyato. Spectral norm regularization for improving the generalizability of deep learning. *arXiv preprint arXiv:1705.10941*, 2017.