EQUIVARIANT TRANSFORMERS FOR NEURAL NETWORK BASED MOLECULAR POTENTIALS

ABSTRACT

The prediction of quantum mechanical properties is historically plagued by a trade-off between accuracy and speed. Machine learning potentials have previously shown great success in this domain, reaching increasingly better accuracy while maintaining computational efficiency comparable with classical force fields. In this work we propose a novel equivariant Transformer architecture, outperforming state-of-the-art on MD17, ANI-1, and many QM9 targets in both accuracy and computational efficiency. Through an extensive attention weight analysis, we gain valuable insights into the black box predictor and show differences in the learned representation of conformers versus conformations sampled from molecular dynamics or normal modes. Furthermore, we highlight the importance of datasets including off-equilibrium conformations for the evaluation of molecular potentials.

1   INTRODUCTION

Quantum mechanics are essential for the computational analysis and design of molecules and materials. However, the complete solution of the Schrödinger equation is analytically and computationally not practical, which initiated the study of approximations in the past decades (Szabo & Ostlund, 1996). A common quantum mechanics approximation method is to model atomic systems according to density functional theory (DFT), which can provide energy estimates with sufficiently high accuracy for different application cases in biology, physics, chemistry, and materials science. Even more accurate techniques like coupled-cluster exist but both still lack the computational efficiency to be applied on a larger scale, although recent advances are promising in the case of coupled-cluster (Pfau et al., 2020; Hermann et al., 2020). Other methods include force-field and semi-empirical quantum mechanical theories, which provide very efficient estimates but lack accuracy.

The field of machine learning molecular potentials is relatively novel. The first important contributions are rooted in the Behler-Parrinello (BP) representation (Behler, 2011) and the seminal work from Rupp et al. (2012). One of the best transferable machine learning potentials for biomolecules, called ANI (Smith et al., 2017a), is based on BP. A second class of methods, mainly developed in the field of materials science and quantum chemistry, uses more modern graph convolutions (Schütt et al., 2018; Unke & Meuwly, 2019; Qiao et al., 2020; Schütt et al., 2021). SchNet (Schütt et al., 2017b; 2018), for example, uses continuous filter convolutions in a graph network architecture to predict the energy of a system and computes forces by direct differentiation of the neural network against atomic coordinates. Outside of its original use case, this approach has been extended to coupled-cluster solvers (Hermann et al., 2020) and protein folding using coarse-grained systems (Wang et al., 2019; Husic et al., 2020). Recently, other work has shown that a shift towards rotationally equivariant networks (Anderson et al., 2019; Fuchs et al., 2020; Schütt et al., 2021), particularly useful when the predicted quantities are vectors and tensors, can also improve the accuracy on scalars (e.g. energy).

Next to the parametric group of neural network based methods, a nonparametric class of approaches exists. These are usually based on kernel methods, particularly used in materials science. In this work, we will focus on parametric neural network potentials (NNPs) because they have a scaling advantage to large amounts of data, while kernel methods usually work best in a scarce data regime.

Previous deep learning based work in the domain of quantum chemistry focused largely on graph neural network architectures (GNNs) with different levels of handcrafted and learned features

(Schütt et al., 2017b; Qiao et al., 2020; Klicpera et al., 2020b; Unke & Meuwly, 2019; Liu et al., 2020; Schütt et al., 2021). For example, Qiao et al. (2020) first perform a low-cost mean-field electronic structure calculation, from which different quantities are used as input to their neural network. Recently proposed neural network architectures in this context usually include some form of attention (Luong et al., 2015) inside the GNN's message passing step (Qiao et al., 2020; Unke & Meuwly, 2019; Liu et al., 2020).

In this work, we introduce an equivariant Transformer (ET) architecture for the prediction of quantum mechanical properties. By building on top of the Transformer (Vaswani et al., 2017) architecture, we are centering the design around the attention mechanism, achieving state-of-the-art accuracy on multiple benchmarks while relying solely on a learned featurization of atomic types and coordinates. Furthermore, we gain insights into the black box prediction of neural networks by analyzing the Transformer's attention weights and comparing latent representations between different types of data such as energy-minimized (QM9 (Ramakrishnan et al., 2014)), molecular dynamics (MD17 (Chmiela et al., 2017) and normal mode sampled data (ANI-1 (Smith et al., 2017b)).

2 METHODS

The traditional Transformer architecture as proposed by Vaswani et al. (2017) operates on a sequence of tokens. In the context of chemistry, however, the natural data structure for the representation of molecules is a graph. To work on graphs, one can interpret self-attention as constructing a fully connected graph over input tokens and computing interactions between nodes. We leverage this concept and extend it to include information stored in the graph's edges, corresponding to interatomic distances in the context of molecular data. This requires the use of a modified attention mechanism, which we introduce in the following sections, along with the overall architecture of our equivariant Transformer (ET).

The equivariant Transformer is made up of three main blocks. An embedding layer encodes atom types Z and the atomic neighborhood of each atom into a dense feature vector x_i. Then, a series of update layers compute interactions between pairs of atoms through a modified multi-head attention mechanism, with which the latent atomic representations are updated. Finally, an output network computes scalar atomwise predictions using gated equivariant blocks (Schütt et al., 2021), which get aggregated into a single molecular prediction. This can be matched with a scalar target variable or differentiated against atomic coordinates, providing force predictions. An illustration of the architecture is given in Figure 1.

2.1 NOTATION

To differentiate between the concepts of scalar and vector features, this work follows a certain notation. Scalar features are written as $x \in \mathbb{R}^F$, while we refer to vector features as $\vec{v} \in \mathbb{R}^{3 \times F}$. The vector norm $\|\cdot\|$ and scalar product $\langle \cdot, \cdot \rangle$ of vector features are applied to the spatial dimension, while all other operations act on the feature dimension. Upper case letters denote matrices $A \in \mathbb{R}^{N \times M}$.

2.2 EMBEDDING LAYER

The embedding layer assigns two learned vectors to each atom type z_i. One is used to encode information specific to an atom, the other takes the role of a neighborhood embedding. The neighborhood embedding, which is an embedding of the types of neighboring atoms, is multiplied by a distance filter. This operation resembles a continuous-filter convolution (Schütt et al., 2017b) but, as it is used in the first layer, allows the model to store atomic information in two separate weight matrices. These can be thought of as containing information that is intrinsic to an atom versus information about the interaction of two atoms.

The distance filter is generated from expanded interatomic distances using a linear transformation W^F. First, the distance d_{ij} between two atoms i and j is expanded via a set of exponential normal radial basis functions e^{RBF}, defined as

$$e_k^{\mathrm{RBF}}(d_{ij}) = \phi(d_{ij}) \exp(-\beta_k (\exp(-d_{ij}) - \mu_k)^2) \tag{1}$$

where β_k and μ_k are fixed parameters specifying the center and width of radial basis function k. The μ vector is initialized with values equally spaced between $\exp(-d_{\mathrm{cut}})$ and 1, β is initialized

Figure 1: Overview of the equivariant Transformer architecture. Thin lines: scalar features in $\mathbb{R}^F$, thick lines: vector features in $\mathbb{R}^{3 \times F}$, dashed lines: multiple feature vectors. **(a)** Transformer consisting of an embedding layer, update layers and an output network. **(b)** Residual update layer including attention based interatomic interactions and information exchange between scalar and vector features. **(c)** Modified dot-product attention mechanism, scaling values (blue) by the attention weights (red).

as $(2K^{-1}(1 - \exp(-d_{\text{cut}})))^{-2}$ for all k as proposed by Unke & Meuwly (2019). The cutoff distance d_{cut} was set to 5Å. The cosine cutoff $\phi(d_{ij})$ is used to ensure a smooth transition to 0 as d_{ij} approaches d_{cut} in order to avoid jumps in the regression landscape. It is given by

$$\phi(d_{ij}) = \begin{cases} \frac{1}{2}\left(\cos\left(\frac{\pi d_{ij}}{d_{\text{cut}}}\right) + 1\right), & \text{if } d_{ij} \leq d_{\text{cut}} \\ 0, & \text{if } d_{ij} > d_{\text{cut}}. \end{cases} \tag{2}$$

The neighborhood embedding n_i for atom i is then defined as

$$n_i = \sum_{j=1}^{N} a_n(z_j) \odot W^F e^{\text{RBF}}(d_{ij}) \tag{3}$$

with a_n being the neighborhood embedding function and N the number of atoms in the graph. The final atomic embedding x_i is calculated as a linear projection of the concatenated intrinsic embedding and neighborhood embedding $[a_i(z_i), n_i]$, resulting in

$$x_i = W^C[a_i(z_i), n_i] + b^C \tag{4}$$

with a_i being the intrinsic embedding function. The vector features $\vec{v}_i$ are initially set to 0.

2.3 MODIFIED ATTENTION MECHANISM

We use a modified multi-head attention mechanism (Figure 1c), extending dot-product attention, in order to include edge data into the calculation of attention weights. The edge data, i.e. interatomic distances r_{ij}, are projected into two multidimensional filters D^K and D^V, according to

$$\begin{aligned} D^K &= \sigma(W^{D^K} e^{\text{RBF}}(r_{ij}) + b^{D^K}) \\ D^V &= \sigma(W^{D^V} e^{\text{RBF}}(r_{ij}) + b^{D^V}) \end{aligned} \tag{5}$$

The attention weights are computed via an extended dot product, i.e. an elementwise multiplication and subsequent sum over the feature dimension, of the three input vectors: query Q, key K and distance projection D^K:

$$\text{dot}(Q, K, D^K) = \sum_k^F Q_k \odot K_k \odot D_k^K \tag{6}$$

The resulting matrix is passed through a nonlinear activation function and is weighted by a cosine cutoff (see equation 2), ensuring that atoms with a distance larger than d_{cut} do not interact. Traditionally, the resulting attention matrix A is passed through a softmax activation, however, we replace this step with a SiLU function to preserve the distance cutoff. The softmax scaling factor of $\sqrt{d_k}^{-1}$, which normally rescales small gradients from the softmax function, is left out. Work by Choromanski et al. (2021) suggests that replacing the softmax activation function in Transformers with ReLU-like functions might even improve accuracy, supporting the idea of switching to SiLU in this case.

We place a continuous filter graph convolution (Schütt et al., 2017b) in the attention mechanism's value pathway. This enables the model to not only consider interatomic distances in the attention weights but also incorporate this information into the feature vectors directly. The resulting representation is split into three equally sized vectors $s_{ij}^1, s_{ij}^2, s_{ij}^3 \in \mathbb{R}^F$. The vector s_{ij}^3 is scaled by the attention matrix A and aggregated over the value-dimension, leading to an updated list of feature vectors. The linear transformation O is used to combine the attention heads' outputs into a single feature vector $y_i \in \mathbb{R}^{384}$.

$$s_{ij}^1, s_{ij}^2, s_{ij}^3 = \text{split}(V_j \odot D^V{}_{ij})$$

$$y_i = O\left(\sum_j^N A_{ij} \cdot s_{ij}^3\right) \tag{7}$$

The attention mechanism's output, therefore, corresponds to the updated scalar feature vectors y_i and scalar filters s_{ij}^1 and s_{ij}^2, which are used to weight the directional information inside the update layer.

2.4 UPDATE LAYER

The update layer (Figure 1b) is used to compute interactions between atoms (attention block) and exchange information between scalar and vector features. The updated scalar features y_i from the attention block are split up into three feature vectors $q_i^1, q_i^2, q_i^3 \in \mathbb{R}^F$. The first feature vector, q_i^1, takes the role of a residual around the scaled vector features. The resulting scalar feature update Δx_i of this update layer is then defined as

$$\Delta x_i = q_i^1 + q_i^2 \odot \langle U_1 \vec{v}_i, U_2 \vec{v}_i \rangle \tag{8}$$

where $\langle U_1 \vec{v}_i, U_2 \vec{v}_i \rangle$ denotes the scalar product of vector features $\vec{v}_i$, transformed by linear projections U_1 and U_2.

On the side of the vector features, scalar information is introduced through a multiplication between q_i^3 and a linear projection of the vector features $U_3 \vec{v}_i$. The representation is updated with equivariant features using the directional vector between two atoms. The edge-wise directional information is multiplied with scalar filter s_{ij}^2 and added to the rescaled vector features $s_{ij}^1 \cdot \vec{v}_j$. The result is aggregated inside each atom, forming $\vec{w}_i$. The final vector feature update $\Delta \vec{v}_i$ for the current update layer is then produced by adding the weighted scalar features to the equivariant features $\vec{w}_i$.

$$\vec{w}_i = \sum_j^N s_{ij}^1 \odot \vec{v}_j + s_{ij}^2 \odot \frac{\vec{r}_i - \vec{r}_j}{\|\vec{r}_i - \vec{r}_j\|} \tag{9}$$

$$\Delta \vec{v}_i = \vec{w}_i + q_i^3 \odot U_3 \vec{v}_i$$

2.5 TRAINING

Models are trained using mean squared error loss and the Adam optimizer (Kingma & Ba, 2017) with parameters $\beta_1 = 0.9$, $\beta_2 = 0.999$ and $\epsilon = 10^{-8}$. Linear learning rate warm-up is applied as suggested by Vaswani et al. (2017) by scaling the learning rate with $\xi = \frac{\text{step}}{n_{\text{steps}}}$. After the warm-up period, we systematically decrease the learning rate by scaling with a decay factor upon reaching a plateau in validation loss. The learning rate is decreased down to a minimum of 10^{-7}. We found that weight decay and dropout do not improve generalization in this context. When training on energies and forces, we apply exponential smoothing to the energy's train and validation loss. New losses are discounted with a factor of $\alpha = 0.05$. See Table 1 for a summary of hyperparameters.

Table 1: Comparison of various hyperparameters used for QM9, MD17 and ANI-1.

Parameter	QM9	MD17	ANI-1
initial learning rate	$4 \cdot 10^{-4}$	$1 \cdot 10^{-3}$	$7 \cdot 10^{-4}$
lr patience (epochs)	15	30	5
lr decay factor	0.8	0.8	0.5
lr warmup steps	10,000	1,000	10,000
batch size	128	8	2048
no. layers	8	6	6
no. RBFs	64	32	32
feature dimension	256	128	128
no. parameters	6.87M	1.34M	1.34M

3 EXPERIMENTS AND RESULTS

We evaluate the equivariant Transformer on the QM9 (Ramakrishnan et al., 2014), MD17 (Chmiela et al., 2017) and ANI-1 (Smith et al., 2017b) benchmark datasets. QM9 comprises 133,885 small organic molecules with up to nine heavy atoms of type C, O, N, and F. It reports computed geometric, thermodynamic, energetic, and electronic properties for locally optimized geometries. As suggested by the authors, we used a revised version of the dataset, which excludes 3,054 molecules due to failed geometric consistency checks. The remaining molecules were split into a training set with 110,000 and a validation set with 10,000 samples, leaving 10,831 samples for testing.

Table 2 compares the equivariant Transformer's results on QM9 with the invariant architectures SchNet (Schütt et al., 2018), PhysNet (Unke & Meuwly, 2019) and DimeNet++ (Klicpera et al., 2020a), the covariant Cormorant (Anderson et al., 2019) architecture and equivariant PaiNN (Schütt et al., 2021). We use specialized output models for two of the QM9 targets, which add certain features directly to the prediction. For the molecular dipole moment μ, both scalar and vector features are used in the final calculation. The output MLP consists of two gated equivariant blocks (Schütt et al., 2021) with the same layer sizes as in the otherwise used output network. The updated scalar features x_i and vector features $\vec{v}_i$ are then used to compute μ as

$$\mu = \left\| \sum_i^N \vec{v}_i + x_i(\vec{r}_i - \vec{r}) \right\| \tag{10}$$

where $\vec{r}$ is the center of mass of the molecule. For the prediction of the electronic spatial extent $\langle R^2 \rangle$, scalar features are transformed using gated equivariant blocks as described above, yielding scalar atomic predictions x_i, and multiplied by the squared norm of atomic positions

$$\langle R^2 \rangle = \sum_i^N x_i \|\vec{r}_i\|^2 \tag{11}$$

The MD17 dataset consists of molecular dynamics (MD) trajectories of small organic molecules, including both energies and forces. In order to guarantee conservation of energy, forces are predicted using the negative gradient of the energy with respect to atomic coordinates $\vec{F}_i = -\partial \hat{E}/\partial \vec{r}_i$. To evaluate the architecture's performance in a limited data setting, the model is trained on only 1000 samples from which 50 are used for validation. The remaining data is used for evaluation and is the basis for comparison with other work. Separate models are trained for each molecule using a combined loss function for energies and forces where the energy loss is multiplied with a factor of 0.2 and the force loss with 0.8. An overview of the results and comparison to the invariant models SchNet (Schütt et al., 2017b), PhysNet (Unke & Meuwly, 2019) and DimeNet (Klicpera et al., 2020b), as well as the equivariant architectures PaiNN (Schütt et al., 2021) and NequIP (Batzner et al., 2021) can be found in Table 3.

To evaluate the architecture's capabilities on a large collection of off-equilibrium conformations, we train and evaluate the equivariant Transformer on the ANI-1 dataset. It contains 22,057,374 configurations of 57,462 small organic molecules with up to 8 heavy atoms and atomic species H,

Table 2: Results on all QM9 targets and comparison to previous work. Scores are reported as mean absolute errors (MAE).

Target	Unit	SchNet	PhysNet	DimeNet++	Cormorant	PaiNN	ET
μ	D	0.033	0.0529	0.0297	0.038	0.012	**0.002**
α	a_0^3	0.235	0.0615	0.0435	0.085	0.045	**0.010**
ϵ_{HOMO}	meV	41	32.9	24.6	34	27.6	**21.2**
ϵ_{LUMO}	meV	34	24.7	19.5	38	20.4	**17.8**
$\Delta\epsilon$	meV	63	42.5	**32.6**	61	45.7	38.0
$\langle R^2 \rangle$	a_0^2	0.073	0.765	0.331	0.961	0.066	**0.015**
$ZPVE$	meV	1.7	1.39	**1.21**	2.027	1.28	2.12
U_0	meV	14	8.15	6.32	22	**5.85**	6.24
U	meV	19	8.34	6.28	21	**5.83**	6.30
H	meV	14	8.42	6.53	21	**5.98**	6.48
G	meV	14	9.4	7.56	20	**7.35**	7.64
C_v	$\frac{\text{cal}}{\text{mol K}}$	0.033	0.028	**0.023**	0.026	0.024	0.026

Table 3: Results on MD trajectories from the MD17 dataset. Scores are given by the MAE of energy predictions (kcal/mol) and forces (kcal/mol/Å). NequIP does not provide errors on energy, for PaiNN we include the results with lower force error out of training only on forces versus on forces and energy. Benzene corresponds to the dataset originally released in Chmiela et al. (2017), which is sometimes left out from the literature. ET results are averaged over three random splits.

Molecule		SchNet	PhysNet	DimeNet	PaiNN	NequIP	ET
Aspirin	*energy*	0.37	0.230	0.204	0.167	-	**0.124**
	forces	1.35	0.605	0.499	0.338	0.348	**0.255**
Benzene	*energy*	0.08	-	0.078	-	-	**0.056**
	forces	0.31	-	**0.187**	-	**0.187**	0.201
Ethanol	*energy*	0.08	0.059	0.064	0.064	-	**0.054**
	forces	0.39	0.160	0.230	0.224	0.208	**0.116**
Malondialdehyde	*energy*	0.13	0.094	0.104	0.091	-	**0.079**
	forces	0.66	0.319	0.383	0.319	0.337	**0.176**
Naphthalene	*energy*	0.16	0.142	0.122	0.116	-	**0.085**
	forces	0.58	0.310	0.215	0.077	0.097	**0.060**
Salicylic Acid	*energy*	0.20	0.126	0.134	0.116	-	**0.094**
	forces	0.85	0.337	0.374	0.195	0.238	**0.135**
Toluene	*energy*	0.12	0.100	0.102	0.095	-	**0.074**
	forces	0.57	0.191	0.216	0.094	0.101	**0.066**
Uracil	*energy*	0.14	0.108	0.115	0.106	-	**0.096**
	forces	0.56	0.218	0.301	0.139	0.173	**0.094**

C, N, and O. The off-equilibrium data points are generated via exhaustive normal mode sampling of the energy minimized molecules. The model is fitted on DFT energies from 80% of the dataset, while 5% are used as validation and the remaining 15% of the data make up the test set. Figure 2 compares the equivariant Transformer's performance to previous methods DTNN (Schütt et al., 2017a), SchNet (Schütt et al., 2017b), MGCN (Lu et al., 2019) and ANI (Smith et al., 2017a).

Figure 2: Comparison of testing MAE on the ANI-1 dataset in eV. Results for DTNN, SchNet and MGCN are provided by Lu et al. (2019). The ANI method refers to the ANAKIN-ME (Smith et al., 2017a) model used for constructing the ANI-1 dataset.

Figure 3: Depiction of bond probabilities and attention scores extracted from the ET using QM9 (total energy U_0), MD17 (average over 8 discussed molecules) and ANI-1 testing data. Attention scores are given as z_i attending z_j, bond probabilities follow the same idea, showing the conditional probability of a bond between z_i and z_j, given z_i. Darker colors correspond to larger values, element pairs without data are grayed out. See Appendix D for an overview of elemental composition in the respective datasets.

3.1 ATTENTION WEIGHT ANALYSIS

Neural network predictions are notoriously difficult to interpret due to the complex nature of the learned transformations. To shed light into the black box predictor, we extract and analyze the equivariant Transformer's attention weights. We run inference on the ANI-1, QM9, and MD17 test sets for all molecules and extract each sample's attention matrix from all attention heads in all layers. Attention rollout (Abnar & Zuidema, 2020) under the single head assumption is applied during the extraction, resulting in a single attention matrix per sample. Next to the direct visualization of these weights in the form of rendered molecules (Figure 4 and Appendix C), we average attention weights over each unique combination of interacting atom types. This leaves two attention scores for each pair of atom types, one from the perspective of z_1 attending z_2 and vice versa.

The attention scores are compared to bond probabilities extracted from the same molecules to make sure the network does not simply attend interacting atoms proportional to the relative frequency in the dataset. Figure 3 presents a summary of the distilled probabilities and attention scores for QM9, ANI-1, and the average attention scores for all MD17 models. The sum over each row equals one, meaning that the probabilities represent the conditional probability

$$P_{z_i}(\text{bond}_{z_m,z_n}|z_m = z_i) = \frac{N_{\text{bonded}}(z_m, z_n)}{\sum_{z_k \in Z} N_{\text{bonded}}(z_m, z_k)} \tag{12}$$

where $N_{\text{bonded}}(z_i, z_j)$ is the total number of bonds between atom types z_i and z_j found in the collection of molecules. For a separate summary of attention scores for individual molecules in the MD17 dataset, see Appendix E.

Since the equivariant Transformer is trained to predict the energy of a certain molecular conformation, we expect it to pay attention to atoms that are displaced from the equilibrium conformation of the molecule. We test this hypothesis by comparing the attention weights of displaced atoms to those

Figure 4: Visualization of two molecules from the QM9 dataset with attention scores corresponding to models trained on ANI-1, MD17 (uracil) and QM9. Blue and red lines represent negative and positive attention scores respectively.

of the equilibrium conformation. Using the QM9 test set as the source of equilibrium conformations, we displace single atoms by 0.4Å in a random direction and compare the absolute magnitude of attention weights involving the displaced atom to the remaining attention weights. We restrict the molecules to only contain hydrogen, carbon, and oxygen in order to compare results between models trained on QM9, MD17 (aspirin), and ANI-1. We find increased attention for displaced carbon and oxygen atoms in all models, however, only the model trained on ANI-1 attends more to displaced hydrogen atoms than to hydrogen in its equilibrium position. Attention for displaced hydrogen atoms even decreases in the model trained on QM9, which suggests that the energy labels in QM9 do not depend strongly on the exact location of hydrogen atoms. For a detailed overview of the results, see Appendix B.

3.2 ABLATION STUDIES

We quantify the effectiveness of the neighbor embedding layer by the change in accuracy when ablating this architectural component. The neighbor embedding layer is replaced by a regular atom type embedding, which we evaluate by comparing the testing MAE on U_0 from QM9. The ablation causes a drop in accuracy of roughly 6% (from 6.24 to 6.60 meV). We also try replacing the neighbor embedding by an additional update layer as the neighbor embedding resembles a graph convolution operation, which is the equivalent operation to an update layer in graph convolutional networks such as SchNet. Here, the drop in accuracy is even more pronounced with a decrease of around 10% (from 6.24 to 6.85 meV). However, this may also be the result of overfitting as each update layer has about 4.6 times more parameters than a neighbor embedding layer.

3.3 COMPUTATIONAL EFFICIENCY

To assess the computational efficiency of the equivariant Transformer, we measure the inference time of random QM9 batches comprising 50 molecules on an NVIDIA V100 GPU (see Table 4). We report times for different sizes of the ET, differing in the number of update layers, the feature dimension, and the size of the RBF distance expansion. ET-large uses the QM9 hyperparameter set, while ET-small is constructed using MD17/ANI-1 hyperparameters (see Table 1). The measurements were conducted using just-in-time (JIT) compiled models. The JIT-compiled versions of

ET-small and ET-large are 1.5 and 1.8 times respectively more efficient than the raw implementation. We only measure the duration of the forward pass, excluding the backward pass required to predict forces. The force prediction decreases inference speed by approximately 75.9%, resulting in 39.0ms per batch using ET-small and 48.5ms using ET-large.

Table 4: Comparison of computational efficiency between PaiNN, DimeNet++ and different sizes of the ET. The time is measured at inference using random batches of 50 molecules from QM9. Speed of ET models is reported as mean $\pm$ standard deviation over 1000 calls. Values for PaiNN and DimeNet++ are taken from Schütt et al. (2021) so differences in efficiency may to some degree originate from different implementations.

	PaiNN	DimeNet++	ET-small	ET-large
time per batch	13 ms	45 ms	**9.4ms** $\pm$ 3.4ms	11.7ms $\pm$ 4.0ms
no. parameters	600k	1.8M	1.29M	6.70M

4 DISCUSSION

In this work, we introduce a novel attention-based architecture for the prediction of quantum mechanical properties, leveraging the use of rotationally equivariant features. We show a high degree of accuracy on the QM9 benchmark dataset, however, the architecture's effectiveness is particularly clear when looking at the prediction of energies and atomic forces in the context of molecular dynamics. We set a new state-of-the-art on all MD17 targets (except force prediction of the molecule Benzene) and demonstrate the architecture's ability to work in a low data regime. As described in previous work (Schütt et al., 2021; Fuchs et al., 2020), the model's vector features and equivariance can be utilized in the prediction of variables beyond scalars. Here, only the dipole moment is predicted in this fashion, however, the architecture is capable of predicting tensorial properties. By extracting and analyzing the model's attention weights, we gain insights into the molecular representation, which is characterized by the nature of the corresponding training data. We show that the model does not pay much attention to the location of hydrogen when trained only on energy-minimized molecules, while a model trained on data including off-equilibrium conformations focuses to a large degree on hydrogen. Furthermore, we validate the learned representation by analyzing attention weights involving atoms displaced from their equilibrium location. We demonstrate that off-equilibrium conformations in the training data play an important role in the accurate prediction of a molecule's energy. This highlights the importance of configurational diversity in the evaluation of neural network potentials.

SOFTWARE AND DATA

The equivariant Transformer is implemented in PyTorch (Paszke et al., 2019), using PyTorch Geometric (Fey & Lenssen, 2019) as the underlying framework for geometric deep learning. Training is done using pytorch-lightning (Falcon & The PyTorch Lightning team, 2019), a high-level interface for training PyTorch models. The datasets QM9[1], MD17[2] and ANI-1[3] are publicly available and all source code for training, running and analyzing the models presented in this work is available at github.com/xxx.

REFERENCES

Samira Abnar and Willem Zuidema. Quantifying Attention Flow in Transformers. *arXiv:2005.00928 [cs]*, May 2020. URL http://arxiv.org/abs/2005.00928. arXiv: 2005.00928.

---

[1]https://doi.org/10.6084/m9.figshare.c.978904.v5

[2]http://www.quantum-machine.org/gdml/#datasets

[3]https://figshare.com/articles/dataset/ANI-1x_Dataset_Release/10047041/1

Brandon Anderson, Truong-Son Hy, and Risi Kondor. Cormorant: Covariant Molecular Neural Networks. *arXiv:1906.04015 [physics, stat]*, November 2019. URL http://arxiv.org/abs/1906.04015. arXiv: 1906.04015.

Simon Batzner, Albert Musaelian, Lixin Sun, Mario Geiger, Jonathan P. Mailoa, Mordechai Kornbluth, Nicola Molinari, Tess E. Smidt, and Boris Kozinsky. SE(3)-Equivariant Graph Neural Networks for Data-Efficient and Accurate Interatomic Potentials. *arXiv:2101.03164 [cond-mat, physics:physics]*, July 2021. URL http://arxiv.org/abs/2101.03164. arXiv: 2101.03164.

Jörg Behler. Atom-centered symmetry functions for constructing high-dimensional neural network potentials. *The Journal of Chemical Physics*, 134(7):074106, February 2011. ISSN 0021-9606. doi: 10.1063/1.3553717. URL https://aip.scitation.org/doi/10.1063/1.3553717. Publisher: American Institute of Physics.

Stefan Chmiela, Alexandre Tkatchenko, Huziel E. Sauceda, Igor Poltavsky, Kristof T. Schütt, and Klaus-Robert Müller. Machine learning of accurate energy-conserving molecular force fields. *Science Advances*, 3(5):e1603015, May 2017. ISSN 2375-2548. doi: 10.1126/sciadv.1603015. URL https://advances.sciencemag.org/content/3/5/e1603015. Publisher: American Association for the Advancement of Science Section: Research Article.

Krzysztof Choromanski, Valerii Likhosherstov, David Dohan, Xingyou Song, Andreea Gane, Tamas Sarlos, Peter Hawkins, Jared Davis, Afroz Mohiuddin, Lukasz Kaiser, David Belanger, Lucy Colwell, and Adrian Weller. Rethinking Attention with Performers. *arXiv:2009.14794 [cs, stat]*, March 2021. URL http://arxiv.org/abs/2009.14794. arXiv: 2009.14794.

William Falcon and The PyTorch Lightning team. PyTorch Lightning, March 2019. URL https://github.com/PyTorchLightning/pytorch-lightning.

Matthias Fey and Jan Eric Lenssen. Fast Graph Representation Learning with PyTorch Geometric, May 2019. URL https://github.com/rusty1s/pytorch_geometric.

Fabian B. Fuchs, Daniel E. Worrall, Volker Fischer, and Max Welling. SE(3)-Transformers: 3D Roto-Translation Equivariant Attention Networks. *arXiv:2006.10503 [cs, stat]*, November 2020. URL http://arxiv.org/abs/2006.10503. arXiv: 2006.10503.

Jan Hermann, Zeno Schätzle, and Frank Noé. Deep neural network solution of the electronic Schrödinger equation. *Nature Chemistry*, 12(10):891–897, October 2020. ISSN 1755-4330, 1755-4349. doi: 10.1038/s41557-020-0544-y. URL http://arxiv.org/abs/1909.08423. arXiv: 1909.08423.

Brooke E. Husic, Nicholas E. Charron, Dominik Lemm, Jiang Wang, Adrià Pérez, Maciej Majewski, Andreas Krämer, Yaoyi Chen, Simon Olsson, Gianni de Fabritiis, Frank Noé, and Cecilia Clementi. Coarse Graining Molecular Dynamics with Graph Neural Networks. *The Journal of Chemical Physics*, 153(19):194101, November 2020. ISSN 0021-9606, 1089-7690. doi: 10.1063/5.0026133. URL http://arxiv.org/abs/2007.11412. arXiv: 2007.11412.

Diederik P. Kingma and Jimmy Ba. Adam: A Method for Stochastic Optimization. *arXiv:1412.6980 [cs]*, January 2017. URL http://arxiv.org/abs/1412.6980. arXiv: 1412.6980.

Johannes Klicpera, Shankari Giri, Johannes T. Margraf, and Stephan Günnemann. Fast and Uncertainty-Aware Directional Message Passing for Non-Equilibrium Molecules. *arXiv:2011.14115 [physics]*, December 2020a. URL http://arxiv.org/abs/2011.14115. arXiv: 2011.14115.

Johannes Klicpera, Janek Groß, and Stephan Günnemann. Directional Message Passing for Molecular Graphs. *arXiv:2003.03123 [physics, stat]*, March 2020b. URL http://arxiv.org/abs/2003.03123. arXiv: 2003.03123.

Ziteng Liu, Liqiang Lin, Qingqing Jia, Zheng Cheng, Yanyan Jiang, Yanwen Guo, and Jing Ma. Transferable Multi-level Attention Neural Network for Accurate Prediction of Quantum Chemistry Properties via Multi-task Learning. *ChemRxiv*, July 2020. doi: 10.26434/chemrxiv.12588170.v1. URL https://chemrxiv.org/articles/

preprint/Transferable_Multi-level_Attention_Neural_Network_
for_Accurate_Prediction_of_Quantum_Chemistry_Properties_via_
Multi-task_Learning/12588170. Publisher: ChemRxiv.

Chengqiang Lu, Qi Liu, Chao Wang, Zhenya Huang, Peize Lin, and Lixin He. Molecular Property Prediction: A Multilevel Quantum Interactions Modeling Perspective. *Proceedings of the AAAI Conference on Artificial Intelligence*, 33(01):1052–1060, July 2019. ISSN 2374-3468. doi: 10.1609/aaai.v33i01.33011052. URL https://ojs.aaai.org/index.php/AAAI/article/view/3896. Number: 01.

Minh-Thang Luong, Hieu Pham, and Christopher D. Manning. Effective Approaches to Attention-based Neural Machine Translation. *arXiv:1508.04025 [cs]*, September 2015. URL http://arxiv.org/abs/1508.04025. arXiv: 1508.04025.

Adam Paszke, Sam Gross, Francisco Massa, Adam Lerer, James Bradbury, Gregory Chanan, Trevor Killeen, Zeming Lin, Natalia Gimelshein, Luca Antiga, Alban Desmaison, Andreas Köpf, Edward Yang, Zach DeVito, Martin Raison, Alykhan Tejani, Sasank Chilamkurthy, Benoit Steiner, Lu Fang, Junjie Bai, and Soumith Chintala. PyTorch: An Imperative Style, High-Performance Deep Learning Library. *arXiv:1912.01703 [cs, stat]*, December 2019. URL http://arxiv.org/abs/1912.01703. arXiv: 1912.01703.

David Pfau, James S. Spencer, Alexander G. D. G. Matthews, and W. M. C. Foulkes. Ab initio solution of the many-electron Schrödinger equation with deep neural networks. *Physical Review Research*, 2(3):033429, September 2020. doi: 10.1103/PhysRevResearch.2.033429. URL https://link.aps.org/doi/10.1103/PhysRevResearch.2.033429. Publisher: American Physical Society.

Zhuoran Qiao, Matthew Welborn, Animashree Anandkumar, Frederick R. Manby, and Thomas F. Miller III. OrbNet: Deep Learning for Quantum Chemistry Using Symmetry-Adapted Atomic-Orbital Features. *arXiv:2007.08026 [physics]*, September 2020. doi: 10.1063/5.0021955. URL http://arxiv.org/abs/2007.08026. arXiv: 2007.08026.

Raghunathan Ramakrishnan, Pavlo O. Dral, Matthias Rupp, and O. Anatole von Lilienfeld. Quantum chemistry structures and properties of 134 kilo molecules. *Scientific Data*, 1(1):140022, August 2014. ISSN 2052-4463. doi: 10.1038/sdata.2014.22. URL https://www.nature.com/articles/sdata201422. Number: 1 Publisher: Nature Publishing Group.

Matthias Rupp, Alexandre Tkatchenko, Klaus-Robert Müller, and O. Anatole von Lilienfeld. Fast and Accurate Modeling of Molecular Atomization Energies with Machine Learning. *Physical Review Letters*, 108(5):058301, January 2012. doi: 10.1103/PhysRevLett.108.058301. URL https://link.aps.org/doi/10.1103/PhysRevLett.108.058301. Publisher: American Physical Society.

K. T. Schütt, H. E. Sauceda, P.-J. Kindermans, A. Tkatchenko, and K.-R. Müller. SchNet – A deep learning architecture for molecules and materials. *The Journal of Chemical Physics*, 148(24):241722, March 2018. ISSN 0021-9606. doi: 10.1063/1.5019779. URL https://aip.scitation.org/doi/abs/10.1063/1.5019779. Publisher: American Institute of Physics.

Kristof T. Schütt, Farhad Arbabzadah, Stefan Chmiela, Klaus R. Müller, and Alexandre Tkatchenko. Quantum-chemical insights from deep tensor neural networks. *Nature Communications*, 8(1):13890, January 2017a. ISSN 2041-1723. doi: 10.1038/ncomms13890. URL https://www.nature.com/articles/ncomms13890. Publisher: Nature Publishing Group.

Kristof T. Schütt, Pieter-Jan Kindermans, Huziel E. Sauceda, Stefan Chmiela, Alexandre Tkatchenko, and Klaus-Robert Müller. SchNet: A continuous-filter convolutional neural network for modeling quantum interactions. *arXiv:1706.08566 [physics, stat]*, December 2017b. URL http://arxiv.org/abs/1706.08566. arXiv: 1706.08566.

Kristof T. Schütt, Oliver T. Unke, and Michael Gastegger. Equivariant message passing for the prediction of tensorial properties and molecular spectra. *arXiv:2102.03150 [physics]*, June 2021. URL http://arxiv.org/abs/2102.03150. arXiv: 2102.03150.

J. S. Smith, O. Isayev, and A. E. Roitberg. ANI-1: an extensible neural network potential with DFT accuracy at force field computational cost. *Chemical Science*, 8(4):3192–3203, March 2017a. ISSN 2041-6539. doi: 10.1039/C6SC05720A. URL `https://pubs.rsc.org/en/content/articlelanding/2017/sc/c6sc05720a`. Publisher: The Royal Society of Chemistry.

Justin S. Smith, Olexandr Isayev, and Adrian E. Roitberg. ANI-1, A data set of 20 million calculated off-equilibrium conformations for organic molecules. *Scientific Data*, 4(1):170193, December 2017b. ISSN 2052-4463. doi: 10.1038/sdata.2017.193. URL `https://www.nature.com/articles/sdata2017193`. Publisher: Nature Publishing Group.

Attila Szabo and Neil S. Ostlund. *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory*. Dover Books on Chemistry. Dover Publications, July 1996. ISBN 978-0-486-69186-2. Google-Books-ID: 6mV9gYzEkgIC.

Oliver T. Unke and Markus Meuwly. PhysNet: A Neural Network for Predicting Energies, Forces, Dipole Moments and Partial Charges. *arXiv:1902.08408 [physics]*, March 2019. doi: 10.1021/acs.jctc.9b00181. URL `http://arxiv.org/abs/1902.08408`. arXiv: 1902.08408.

Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N. Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention Is All You Need. *arXiv:1706.03762 [cs]*, December 2017. URL `http://arxiv.org/abs/1706.03762`. arXiv: 1706.03762.

Jiang Wang, Simon Olsson, Christoph Wehmeyer, Adrià Pérez, Nicholas E. Charron, Gianni de Fabritiis, Frank Noé, and Cecilia Clementi. Machine Learning of Coarse-Grained Molecular Dynamics Force Fields. *ACS Central Science*, 5(5):755–767, May 2019. ISSN 2374-7943. doi: 10.1021/acscentsci.8b00913. URL `https://doi.org/10.1021/acscentsci.8b00913`. Publisher: American Chemical Society.

A   IMPORTANCE OF HYDROGEN

While the ET trained on QM9 mostly attends to carbon atoms, models trained on molecular dynamics trajectories show a strong focus on carbon-hydrogen interactions. This highlights the different nature of datasets containing only energy minimized conformations in contrast to datasets containing MD data. We further support this hypothesis by comparing the reduction in accuracy for models, which are trained without hydrogen, on the datasets QM9 (total energy U_0) and MD17 (aspirin). We show that the loss in accuracy when predicting the energy using the model trained on MD17 is one order of magnitude larger than when training only on molecules in the ground state. Furthermore, the accuracy of force predictions drops by another 1.5x compared to MD17 energy predictions when excluding hydrogen. A summary of the results is given in Table 5.

Table 5: Test MAE of the ET on QM9 and MD17 (aspirin), trained with and without hydrogen.

Dataset		**with** hydrogen	**without** hydrogen	relative change
QM9		6.37 meV	20.83 meV	227.0%
MD17	*energy*	5.33 meV	137.59 meV	2481.4%
	forces	10.67 meV/Å	433.99 meV/Å	3966.5%

B   ATOM DISPLACEMENT

Figure 5 shows averaged absolute attention scores for molecules where a single atom has been displaced from the equilibrium structure. This reiterates the idea that the model trained only on equilibrium structures (QM9) focuses on carbon and neglects hydrogen. The models trained on off-equilibrium conformations show a higher degree of attention for displaced hydrogen atoms.

Figure 5: Averaged attention weights extracted from the ET on the QM9 test set (molecules consisting of H, C, and O only) with a displacement of 0.4Å in single atoms. Blue bars show attention towards atoms in equilibrium locations, orange bars correspond to attention weights involving the displaced atom. Attention scores are normalized inside each molecule. The black bars show the attention weights' standard deviation.

C   MOLECULAR REPRESENTATION BY DATASET

Figures 6a, 6b and 6c show a three dimensional visualization of three random molecules from the datasets QM9, MD17 (aspirin) and ANI-1 respectively. We extract the attention weights from the best performing equivariant Transformer on the test sets of the three datasets respectively to make sure that no model has seen a visualized conformation during training. The red and blue lines between atoms depict the 10 largest absolute attention weights where the line width and alpha value represent the absolute attention weight. Red lines show positive attention weights, blue lines correspond to negative attention weights, which occur due to using SiLU activations inside the attention mechanism. While the model trained on QM9 focuses largely on carbon-carbon interactions, it is clear that models trained on MD17 and ANI-1 have a strong focus on hydrogen-carbon and hydrogen-oxygen interactions. This corresponds to the results found in the attention weights averaged over pairwise interactions between elements. The attention weights are normalized inside each molecule.

(a) Random molecules from the QM9 test set.

Molecules from MD17 (aspirin)

(b) Random conformations from the MD17 test set of aspirin trajectories.

Molecules from ANI-1

(c) Random molecules from the ANI-1 test set.

Figure 6: Visualization of 10 largest attention weights by absolute value on random molecules from QM9 (a), MD17-aspirin (b) and ANI-1 (c). Each column shows the same molecule, rows correspond to the same ET model trained on QM9, MD17-uracil and ANI-1 respectively.

D FREQUENCY OF ELEMENTS

Figure 7 shows the distribution of elements in the datasets QM9, MD17 and ANI-1 where MD17 corresponds to the combination of all target molecules. It aims to assist with the interpretation of attention scores where certain pairs of elements are largely underrepresented in the model's attention. This corresponds predominantly to nitrogen and fluorine as MD17 contains only a single molecule with nitrogen (uracil) and fluorine is only found in QM9.

Figure 7: Distribution of elements in the datasets QM9, MD17 (combination of all target molecules) and ANI-1.

E MD17 ATTENTION WEIGHTS

Figures 8a and 8b contain the extracted rolled out attention weights for the MD17 target molecules aspirin, benzene, ethanol, malondialdehyde, naphthalene, salicylic acid, toluene and uracil. While the molecules show different attention patterns, a high degree of attention for hydrogen atoms is common for most of the molecules. However, models trained on aspirin, malondialdehyde and uracil additionally exhibit a strong focus on oxygen.

(a) Attention scores and bond probabilities from aspirin, benzene, ethanol and malondialdehyde.

(b) Attention scores and bond probabilities from naphthalene, salicylic acid, toluene and uracil.

Figure 8: Bond probabilities and attention scores extracted from the ET using testing data from all individual molecules in the MD17 dataset. Attention scores are given as z_i attending to z_j, bond probabilities follow the same principle, showing the conditional probability of a bond between z_i and z_j, given z_i. The rightmost subfigure displays the total number of atoms for each type in the data.