SAMPLING IS AS EASY AS LEARNING THE SCORE: THEORY FOR DIFFUSION MODELS WITH MINIMAL DATA ASSUMPTIONS

ABSTRACT

We provide theoretical convergence guarantees for score-based generative models (SGMs) such as denoising diffusion probabilistic models (DDPMs), which constitute the backbone of large-scale real-world generative models such as DALL·E 2. Our main result is that, assuming accurate score estimates, such SGMs can efficiently sample from essentially any realistic data distribution. In contrast to prior works, our results (1) hold for an L^2-accurate score estimate (rather than L^∞-accurate); (2) do not require restrictive functional inequality conditions that preclude substantial non-log-concavity; (3) scale polynomially in all relevant problem parameters; and (4) match state-of-the-art complexity guarantees for discretization of the Langevin diffusion, provided that the score error is sufficiently small. We view this as strong theoretical justification for the empirical success of SGMs. We also examine SGMs based on the critically damped Langevin diffusion (CLD). Contrary to conventional wisdom, we provide evidence that the use of the CLD does *not* reduce the complexity of SGMs.

1 INTRODUCTION

Score-based generative models (SGMs) are a family of generative models which achieve state-of-the-art performance for generating audio and image data (Sohl-Dickstein et al., 2015; Ho et al., 2020; Dhariwal & Nichol, 2021; Kingma et al., 2021; Song et al., 2021a;b; Vahdat et al., 2021); see, e.g., the recent surveys (Cao et al., 2022; Croitoru et al., 2022; Yang et al., 2022). One notable example of an SGM are denoising diffusion probabilistic models (DDPMs) (Sohl-Dickstein et al., 2015; Ho et al., 2020), which are a key component in large-scale generative models such as DALL·E 2 (Ramesh et al., 2022). As the importance of SGMs continues to grow due to newfound applications in commercial domains, it is a pressing question of both practical and theoretical concern to understand the mathematical underpinnings which explain their startling empirical successes.

As we explain in Section 2, at their mathematical core, SGMs consist of two stochastic processes, the forward process and the reverse process. The forward process transforms samples from a data distribution q (e.g., images) into noise, whereas the reverse process transforms noise into samples from q, hence performing generative modeling. Running the reverse process requires estimating the *score function* of the law of the forward process; this is typically done by training neural networks on a score matching objective (Hyvärinen, 2005; Vincent, 2011; Song & Ermon, 2019).

Providing precise guarantees for estimation of the score function is difficult, as it requires an understanding of the non-convex training dynamics of neural network optimization that is currently out of reach. However, given the empirical success of neural networks on the score estimation task, a natural and important question is whether accurate score estimation implies that SGMs provably converge to the true data distribution in realistic settings. This is a surprisingly delicate question, as even with accurate score estimates, as we explain in Section 2.1, there are several other sources of error which could cause the SGM to fail to converge. Indeed, despite a flurry of recent work (Block et al., 2020; De Bortoli et al., 2021; De Bortoli, 2022; Lee et al., 2022a; Pidstrigach, 2022; Liu et al., 2022), prior analyses fall short of answering this question, for (at least) one of three main reasons:

1. **Super-polynomial convergence.** The bounds obtained are not quantitative (e.g., (De Bortoli et al., 2021; Pidstrigach, 2022)), or scale exponentially in the dimension and other problem parameters

like time and smoothness (Block et al., 2020; De Bortoli, 2022; Liu et al., 2022), and hence are typically vacuous for the high-dimensional settings of interest in practice.

2. **Strong assumptions on the data distribution.** The bounds require strong assumptions on the true data distribution, such as a log-Sobolev inequality (LSI) (see, e.g., (Lee et al., 2022a)). While the LSI is slightly weaker than log-concavity, it ultimately precludes the presence of substantial non-convexity, which impedes the application of these results to complex and highly multi-modal real-world data distributions. Indeed, obtaining a polynomial-time convergence analysis for SGMs that holds for multi-modal distributions was posed as an open question in (Lee et al., 2022a).

3. **Strong assumptions on the score estimation error.** The bounds require that the score estimate is L^∞-accurate (i.e., *uniformly* accurate), as opposed to L^2-accurate (see, e.g., (De Bortoli et al., 2021)). This is problematic because the score matching objective is an L^2 loss (see Section A.1 in the supplement), and there are empirical studies suggesting that in practice, the score estimate is not in fact L^∞-accurate (e.g., (Zhang & Chen, 2022)). Intuitively, this is because we cannot expect that the score estimate we obtain will be accurate in regions of space where the true density is very low, simply because we do not expect to see many (or indeed, any) samples from there.

Providing an analysis which goes beyond these limitations is a pressing first step towards theoretically understanding why SGMs actually work in practice.

1.1 OUR CONTRIBUTIONS

In this work, we take a step towards bridging theory and practice by providing a convergence guarantee for SGMs, under realistic (in fact, quite minimal) assumptions, which scales polynomially in all relevant problem parameters. Namely, our main result (Theorem 2) only requires the following assumptions on the data distribution q, which we make more quantitative in Section 3:

A1 The score function of the forward process is L-Lipschitz.

A2 The $(2 + \eta)$-th moment of q is finite, where $\eta > 0$ is an arbitrarily small constant.

A3 The data distribution q has finite KL divergence w.r.t. the standard Gaussian.

We note that all of these assumptions are either standard or, in the case of **A2**, far weaker than what is needed in prior work. Crucially, unlike prior works, we do *not* assume log-concavity, an LSI, or dissipativity; hence, our assumptions cover *arbitrarily non-log-concave* data distributions. Our main result is summarized informally as follows.

Theorem 1 (informal, see Theorem 2). *Under assumptions* **A1-A3***, and if the score estimation error in L^2 is sufficiently polynomially small in $1/d$, $1/L$, and ε, then with an appropriate choice of step size, the SGM outputs a measure which is ε-close in total variation (TV) to q in $\widetilde{O}(L^2 d/\varepsilon^2)$ iterations.*

Our iteration complexity is quite tight: it matches state-of-the-art discretization guarantees for the Langevin diffusion (Vempala & Wibisono, 2019; Chewi et al., 2021a).

We find Theorem 1 surprising, because it shows that SGMs can sample from the data distribution q with polynomial complexity, even when q is highly non-log-concave (a task that is usually intractable), *provided that one has access to an accurate score estimator*. This answers the open question of (Lee et al., 2022a) regarding whether or not SGMs can sample from multimodal distributions, e.g., mixtures of distributions with bounded log-Sobolev constant. In the context of neural networks, our result implies that so long as the neural network succeeds at the score estimation task, the remaining part of the SGM algorithm based on the diffusion model is completely principled, in that it admits a strong theoretical justification.

In general, learning the score function is also a difficult task. Nevertheless, our result opens the door to further investigations, such as: do score functions for real-life data have intrinsic (e.g., low-dimensional) structure which can be exploited by neural networks? A positive answer to this question, combined with our sampling result, would then provide an end-to-end guarantee for SGMs.

More generally, our result can be viewed as a black-box reduction of the task of sampling to the task of learning the score function of the forward process, at least for distributions satisfying our mild assumptions. Existing computational hardness results for learning natural high-dimensional distributions like mixtures of Gaussians (Diakonikolas et al., 2017; Bruna et al., 2021; Gupte et al.,

2022) and pushforwards of Gaussians by shallow ReLU networks (Daniely & Vardi, 2021; Chen et al., 2022a;b) thus immediately imply hardness of score estimation for these distributions. To our knowledge this yields the first known information-computation gaps for this task.

Arbitrary distributions with bounded support. The assumption that the score function is Lipschitz entails in particular that the data distribution has a density w.r.t. Lebesgue measure; in particular, our theorem fails when q satisfies the manifold hypothesis, i.e., is supported on a lower-dimensional submanifold of $\mathbb{R}^d$. But this is for good reason: it is not possible to obtain non-trivial TV guarantees, because the output distribution of the SGM has full support. Instead, we show in Section 3.2 that we can obtain polynomial convergence guarantees in the bounded Lipschitz metric by stopping the SGM algorithm early, under the *sole* assumption that the data distribution q has bounded support, without assuming that q has a density. Since data distributions encountered in real life satisfy this assumption, our results yield the following compelling takeaway:

Given an L^2-accurate score estimate, SGMs can sample from (essentially) any data distribution.

Critically damped Langevin diffusion (CLD). Using our techniques, we also investigate the use of the critically damped Langevin diffusion (CLD) for SGMs, which was proposed in (Dockhorn et al., 2022). Although numerical experiments and intuition from the log-concave sampling literature suggest that the CLD could potentially speed up sampling via SGMs, we provide theoretical evidence to the contrary. Based on this, in Section 3.3, we conjecture that SGMs based on the CLD do not exhibit improved dimension dependence compared to the original DDPM algorithm.

1.2 PRIOR WORK

We now provide a detailed comparison to prior work. By now, there is a vast literature on providing precise complexity estimates for log-concave sampling; see, e.g., (Chewi, 2022) for an exposition on recent developments. The proofs in this work build upon the techniques developed in this literature. However, our work addresses the significantly more challenging setting of *non-log-concave* sampling.

The work of De Bortoli et al. (2021) provides guarantees for the diffusion Schrödinger bridge (Song et al., 2021b). However, as previously mentioned their result is not quantitative, and they require an L^∞-accurate score estimate. The works (Block et al., 2020; Lee et al., 2022a; Liu et al., 2022)[1] instead analyze SGMs under the more realistic assumption of an L^2-accurate score estimate. However, the bounds of (Block et al., 2020; Liu et al., 2022) suffer from exponential dependencies on parameters like dimension and smoothness, whereas the bounds of (Lee et al., 2022a) require q to satisfy an LSI.

The recent work of De Bortoli (2022), motivated by the *manifold hypothesis*, considers a different pointwise assumption on the score estimation error which allows the error to blow up at time 0 and at spatial ∞. We discuss the manifold setting in more detail in Section 3.2. Unfortunately, the bounds of De Bortoli (2022) also scale exponentially in problem parameters such as the manifold diameter.

We also mention that the use of reversed SDEs for sampling is implicit in the interpretation of the proximal sampler (Lee et al., 2021) given by Chen et al. (2022c). Our work can be viewed as expanding upon the theory of Chen et al. (2022c) using a different forward channel (the OU process).

Concurrent work. Very recently, Lee et al. (2022b) independently obtained results similar to our results for DDPM. While our assumptions are technically somewhat incomparable (they assume the score error can vary with time but assume the data is compactly supported), our quantitative bounds are stronger. Additionally, the upper and lower bounds for CLD are unique to our work.

2 BACKGROUND ON SGMs

Throughout this paper, given a probability measure p which admits a density w.r.t. Lebesgue measure, we abuse notation and identify it with its density function. Additionally, we will let q denote the data distribution from which we want to generate new samples. We assume that q is a probability measure

---

[1] Unfortunately, the current analysis of (Block et al., 2020) contains a gap, as Theorem 27 therein is false: an LSI does not imply contraction in the Wasserstein metric for the Langevin diffusion.

on $\mathbb{R}^d$ with full support, and that it admits a smooth density. Except in Section 3.2, we assume that q admits a smooth density written $q = \exp(-U)$, where $U : \mathbb{R}^d \to \mathbb{R}$ is the *potential*.

In this section, we provide a brief exposition to SGMs, following Song et al. (2021b).

2.1 Background on denoising diffusion probabilistic modeling (DDPM)

Forward process. In denoising diffusion probabilistic modeling (DDPM), we start with a forward process, which is a stochastic differential equation (SDE). For clarity, we consider the simplest possible choice, which is the Ornstein–Uhlenbeck (OU) process

$$\mathrm{d}\bar{X}_t = -\bar{X}_t \, \mathrm{d}t + \sqrt{2} \, \mathrm{d}B_t, \qquad \bar{X}_0 \sim q, \tag{1}$$

where $(B_t)_{t \geq 0}$ is a standard Brownian motion in $\mathbb{R}^d$. The OU process is the unique time-homogeneous Markov process which is also a Gaussian process, with stationary distribution equal to the standard Gaussian distribution γ^d on $\mathbb{R}^d$. In practice, it is also common to introduce a positive smooth function $g : \mathbb{R}_+ \to \mathbb{R}$ and consider the time-rescaled OU process

$$\mathrm{d}\bar{X}_t = -g(t)^2 \, \bar{X}_t \, \mathrm{d}t + \sqrt{2} \, g(t) \, \mathrm{d}B_t, \qquad X_0 \sim q, \tag{2}$$

but in this work we stick with the choice $g \equiv 1$.

The forward process has the interpretation of transforming samples from the data distribution q into pure noise. From the well-developed theory of Markov diffusions, it is known that if $q_t := \mathrm{law}(X_t)$ denotes the law of the OU process at time t, then $q_t \to \gamma^d$ exponentially fast in various divergences and metrics such as the 2-Wasserstein metric W_2; see Bakry et al. (2014).

Reverse process. If we reverse the forward process (1) in time, then we obtain a process that transforms noise into samples from q, which is the aim of generative modeling. In general, suppose that we have an SDE of the form

$$\mathrm{d}\bar{X}_t = b_t(\bar{X}_t) \, \mathrm{d}t + \sigma_t \, \mathrm{d}B_t,$$

where $(\sigma_t)_{t \geq 0}$ is a deterministic matrix-valued process. Then, under mild conditions on the process (e.g., (Föllmer, 1985; Cattiaux et al., 2022)), which are satisfied for all processes under consideration in this work, the reverse process also admits an SDE description. Namely, if we fix the terminal time $T > 0$ and set

$$\bar{X}_t^{\leftarrow} := \bar{X}_{T-t}, \qquad \text{for } t \in [0, T],$$

then the process $(\bar{X}_t^{\leftarrow})_{t \in [0,T]}$ satisfies the SDE

$$\mathrm{d}\bar{X}_t^{\leftarrow} = b_t^{\leftarrow}(\bar{X}_t^{\leftarrow}) \, \mathrm{d}t + \sigma_{T-t} \, \mathrm{d}B_t,$$

where the backwards drift satisfies the relation

$$b_t + b_{T-t}^{\leftarrow} = \sigma_t \sigma_t^{\mathsf{T}} \nabla \ln q_t, \qquad q_t := \mathrm{law}(\bar{X}_t). \tag{3}$$

Applying this to the forward process (1), we obtain the reverse process

$$\mathrm{d}\bar{X}_t^{\leftarrow} = \{\bar{X}_t^{\leftarrow} + 2 \nabla \ln q_{T-t}(\bar{X}_t^{\leftarrow})\} \, \mathrm{d}t + \sqrt{2} \, \mathrm{d}B_t, \qquad \bar{X}_0^{\leftarrow} \sim q_T, \tag{4}$$

where now $(B_t)_{t \in [0,T]}$ is the reversed Brownian motion.[2] Here, $\nabla \ln q_t$ is called the *score function* for q_t. Since q (and hence q_t for $t \geq 0$) is not explicitly known, in order to implement the reverse process the score function must be estimated on the basis of samples. The mechanism behind this is the idea of *score matching* which goes back to Hyvärinen (2005); Vincent (2011): roughly speaking, Gaussian integration by parts implies that minimizing the $L^2(q_t)$ loss achieved by an estimate s_t for the score $\nabla \ln q_t$ is *equivalent* to minimizing the $L^2(q_t)$ loss in predicting, given a sample from the forward process at time t, what noise was applied to the corresponding sample at time 0 to obtain it. We defer an exposition of the details of score matching to Sections A.1 and D of the supplement.

In light of this, it is thus most natural to assume an $L^2(q_t)$ error bound $\mathbb{E}_{q_t}[\|s_t - \nabla \ln q_t\|^2] \leq \varepsilon_{\mathrm{score}}^2$ for the score estimator s_t. If s_t is taken to be the empirical risk minimizer for a suitable function class, then guarantees for the $L^2(q_t)$ error can be obtained via standard statistical analysis, see e.g. Block et al. (2020).

---

[2]For ease of notation, we do not distinguish between the forward and the reverse Brownian motions.

Discretization and implementation. We now discuss the final steps required to obtain an implementable algorithm. First, in the learning phase, given samples $\bar{X}_0^{(1)}, \ldots, \bar{X}_0^{(n)}$ from q (e.g., a database of natural images), we train a neural network via score matching, see Song & Ermon (2019). Let $h > 0$ be the step size of the discretization; we assume that we have obtained a score estimate s_{kh} of $\nabla \ln q_{kh}$ for each time $k = 0, 1, \ldots, N$, where $T = Nh$.

In order to approximately implement the reverse SDE (4), we first replace the score function $\nabla \ln q_{T-t}$ with the estimate s_{T-t}. Then, for $t \in [kh, (k+1)h]$ we freeze the value of this coefficient in the SDE at time kh. It yields the new SDE

$$\mathrm{d}X_t^{\leftarrow} = \{X_t^{\leftarrow} + 2\, s_{T-kh}(X_{kh}^{\leftarrow})\}\, \mathrm{d}t + \sqrt{2}\, \mathrm{d}B_t\,, \qquad t \in [kh, (k+1)h]\,. \tag{5}$$

Since this is a linear SDE, it can be integrated in closed form; in particular, conditionally on $X_{kh}^{\leftarrow}$, the next iterate $X_{(k+1)h}^{\leftarrow}$ has an explicit Gaussian distribution.

There is one final detail: although the reverse SDE (4) should be started at q_T, we do not have access to q_T directly. Instead, taking advantage of the fact that $q_T \approx \gamma^d$, we instead initialize the algorithm at $X_0^{\leftarrow} \sim \gamma^d$, i.e., from pure noise.

Let $p_t := \mathrm{law}(X_t^{\leftarrow})$ denote the law of the algorithm at time t. The goal of this work is to bound $\mathsf{TV}(p_T, q)$, taking into account three sources of error: (1) estimation of the score; (2) discretization of the SDE with step size $h > 0$; and (3) initialization of the algorithm at γ^d rather than at q_T.

2.2 BACKGROUND ON THE CRITICALLY DAMPED LANGEVIN DIFFUSION (CLD)

The critically damped Langevin diffusion (CLD) is based on the forward process

$$\begin{aligned}
\mathrm{d}\bar{X}_t &= -\bar{V}_t\, \mathrm{d}t\,, \\
\mathrm{d}\bar{V}_t &= -(\bar{X}_t + 2\,\bar{V}_t)\, \mathrm{d}t + 2\, \mathrm{d}B_t\,.
\end{aligned} \tag{6}$$

Compared to the OU process (1), this is now a coupled system of SDEs, where we have introduced a new variable $\bar{V}$ representing the velocity process. The stationary distribution of the process is γ^{2d}, the standard Gaussian measure on phase space $\mathbb{R}^d \times \mathbb{R}^d$, and we initialize at $\bar{X}_0 \sim q$ and $\bar{V}_0 \sim \gamma^d$.

More generally, the CLD (6) is an instance of what is referred to as the *kinetic Langevin* or the *underdamped Langevin* process in the sampling literature. In the context of log-concave sampling, the smoother paths of $\tilde{X}$ lead to smaller discretization error, thereby furnishing an algorithm with $\tilde{O}(\sqrt{d}/\varepsilon)$ gradient complexity (as opposed to sampling based on the overdamped Langevin process, which has complexity $\tilde{O}(d/\varepsilon^2)$), see Cheng et al. (2018); Shen & Lee (2019); Dalalyan & Riou-Durand (2020); Ma et al. (2021). The recent paper (Dockhorn et al., 2022) proposed to use the CLD as the basis for an SGM and they empirically observed improvements over DDPM.

Applying (3), the corresponding reverse process is

$$\begin{aligned}
\mathrm{d}\bar{X}_t^{\leftarrow} &= -\bar{V}_t^{\leftarrow}\, \mathrm{d}t\,, \\
\mathrm{d}\bar{V}_t^{\leftarrow} &= \big(\bar{X}_t^{\leftarrow} + 2\,\bar{V}_t^{\leftarrow} + 4\,\nabla_v \ln \boldsymbol{q}_{T-t}(\bar{X}_t^{\leftarrow}, \bar{V}_t^{\leftarrow})\big)\, \mathrm{d}t + 2\, \mathrm{d}B_t\,,
\end{aligned} \tag{7}$$

where $\boldsymbol{q}_t := \mathrm{law}(\bar{X}_t, \bar{V}_t)$ is the law of the forward process at time t. Note that the gradient in the score function is only taken w.r.t. the velocity coordinate. Upon replacing the score function with an estimate $\boldsymbol{s}$, we arrive at the algorithm

$$\begin{aligned}
\mathrm{d}X_t^{\leftarrow} &= -V_t^{\leftarrow}\, \mathrm{d}t\,, \\
\mathrm{d}V_t^{\leftarrow} &= \big(X_t^{\leftarrow} + 2\,V_t^{\leftarrow} + 4\,\boldsymbol{s}_{T-kh}(X_{kh}^{\leftarrow}, V_{kh}^{\leftarrow})\big)\, \mathrm{d}t + 2\, \mathrm{d}B_t\,,
\end{aligned}$$

for $t \in [kh, (k+1)h]$. We provide further background on the CLD in Section C.1.

3 RESULTS

We now formally state our assumptions and our main results.

3.1 RESULTS FOR DDPM

For DDPM, we make the following mild assumptions on the data distribution q.

Assumption 1 (Lipschitz score). *For all $t \geq 0$, the score $\nabla \ln q_t$ is L-Lipschitz.*

Assumption 2 (second moment bound). *For some $\eta > 0$, $\mathbb{E}_q[\|\cdot\|^{2+\eta}]$ is finite. We also write $\mathfrak{m}_2^2 \coloneqq \mathbb{E}_q[\|\cdot\|^2]$ for the second moment of q.*

For technical reasons, we need to assume that q has a finite moment of order slightly but strictly bigger than 2, but our quantitative bounds will only depend on the second moment $\mathfrak{m}_2^2$.

Assumption 1 is standard and has been used in the prior works (Block et al., 2020; Lee et al., 2022a). However, unlike Lee et al. (2022a); Liu et al. (2022), we do not assume Lipschitzness of the score estimate. Moreover, unlike Block et al. (2020); De Bortoli et al. (2021); Liu et al. (2022), we do not assume any convexity or dissipativity assumptions on the potential U, and unlike Lee et al. (2022a) we do not assume that q satisfies a log-Sobolev inequality. Hence, our assumptions cover a wide range of highly non-log-concave data distributions. Our proof technique is fairly robust and even Assumption 1 could be relaxed (as well as other extensions, such as considering the time-changed forward process (2)), although we focus on the simplest setting in order to better illustrate the conceptual significance of our results.

We also assume a bound on the score estimation error.

Assumption 3 (score estimation error). *For all $k = 1, \dots, N$, $\mathbb{E}_{q_{kh}}[\|s_{kh} - \nabla \ln q_{kh}\|^2] \leq \varepsilon_{\text{score}}^2$.*

This is the same assumption as in Lee et al. (2022a), and as discussed in Section 2.1, it is a natural and realistic assumption in light of the derivation of the score matching objective.

Our main result for DDPM is the following theorem.

Theorem 2 (DDPM, see Section B of supplement). *Suppose that Assumptions 1, 2, and 3 hold. Let p_T be the output of the DDPM algorithm (Section 2.1) at time T, and suppose that the step size $h \coloneqq T/N$ satisfies $h \lesssim 1/L$, where $L \geq 1$. Then, it holds that*

$$
\mathsf{TV}(p_T, q) \lesssim \underbrace{\sqrt{D_{\text{KL}}(q \parallel \gamma^d) \exp(-T)}}_{\text{convergence of forward process}} + \underbrace{(L\sqrt{dh} + L\mathfrak{m}_2 h)\sqrt{T}}_{\text{discretization error}} + \underbrace{\frac{N^{1/3}\varepsilon_{\text{score}}^{2/3}}{T^{1/6}}}_{\text{score estimation error}} .
$$

To interpret this result, suppose that, e.g., $D_{\text{KL}}(q \parallel \gamma^d) \leq \text{poly}(d)$ and $\mathfrak{m}_2 \leq d$. Choosing $T \asymp \log(D_{\text{KL}}(q \parallel \gamma)/\varepsilon)$ and $h \asymp \frac{\varepsilon^2}{L^2 d}$, and hiding logarithmic factors,

$$
\mathsf{TV}(p_T, q) \leq \widetilde{O}\Big(\varepsilon + \frac{L^{2/3}d^{1/3}\varepsilon_{\text{score}}^{2/3}}{\varepsilon^{2/3}}\Big), \qquad \text{for } N = \widetilde{\Theta}\Big(\frac{L^2 d}{\varepsilon^2}\Big).
$$

In particular, in order to have $\mathsf{TV}(p_T, q) \leq \varepsilon$, it suffices to have score error $\varepsilon_{\text{score}} \leq \widetilde{O}\Big(\frac{\varepsilon^{5/2}}{L d^{1/2}}\Big)$.

We remark that the iteration complexity of $N = \widetilde{\Theta}(\frac{L^2 d}{\varepsilon^2})$ matches state-of-the-art complexity bounds for the Langevin Monte Carlo (LMC) algorithm for sampling under a log-Sobolev inequality (LSI), see Vempala & Wibisono (2019); Chewi et al. (2021a). This provides some evidence that our discretization bounds are of the correct order, at least with respect to the dimension and accuracy parameters, and without higher-order smoothness assumptions.

3.2 CONSEQUENCES FOR ARBITRARY DATA DISTRIBUTIONS WITH BOUNDED SUPPORT

We now elaborate upon the implications of our results under the *sole* assumption that the data distribution q is compactly supported, $\text{supp}\, q \subseteq \mathsf{B}(0, R)$. In particular, we do not assume that q has a smooth density w.r.t. Lebesgue measure, which allows for studying the case when q is supported on a lower-dimensional submanifold of $\mathbb{R}^d$ as in the *manifold hypothesis*. This setting was investigated recently in De Bortoli (2022).

For this setting, our results do not apply directly because the score function of q is not well-defined and hence Assumption 1 fails to hold. Also, the bound in Theorem 2 has a term involving $D_{\text{KL}}(q \parallel \gamma^d)$

which is infinite if q is not absolutely continuous w.r.t. γ^d. As pointed out by De Bortoli (2022), in general we cannot obtain non-trivial guarantees for $\mathsf{TV}(p_T, q)$, because p_T has full support and therefore $\mathsf{TV}(p_T, q) = 1$ under the manifold hypothesis. Nevertheless, we show that we can apply our results using an early stopping technique.

Namely, consider q_t the law of the OU process at a time $t > 0$, initialized at q. We show:

Lemma 3 (Lemma 18 in supplement). *Suppose that* $\operatorname{supp} q \subseteq \mathsf{B}(0, R)$ *where* $R \geq 1$, *and let* q_t *denote the law of the OU process at time* t, *started at* q. *Let* $\varepsilon_{W_2} > 0$ *be such that* $\varepsilon_{W_2} \ll \sqrt{d}$ *and set* $t \asymp \varepsilon_{W_2}^2 / (\sqrt{d} \, (R \vee \sqrt{d}))$. *Then, (1)* $W_2(q_t, q) \leq \varepsilon_{W_2}$, *(2)* q_t *satisfies* $D_{\mathrm{KL}}(q_t \,\|\, \gamma^d) \lesssim \frac{\sqrt{d} \, (R \vee \sqrt{d})^3}{\varepsilon_{W_2}^2}$, *and (3) for every* $t' \geq t$, $q_{t'}$ *satisfies Assumption 1 with* $L \lesssim \frac{dR^2 \, (R \vee \sqrt{d})^2}{\varepsilon_{W_2}^4}$.

By substituting q_t for this choice of t in place of q in Theorem 2, we obtain Corollary 4 below. We remark that taking q_t as the new target corresponds to stopping the algorithm early: instead of running the algorithm backward for a time T, we run the algorithm backward for a time $T - t$ (note that $T - t$ should be a multiple of the step size h).

Corollary 4 (compactly supported data). *Suppose that* q *is supported on the ball of radius* $R \geq 1$. *Let* $t \asymp \varepsilon_{W_2}^2 / (\sqrt{d} \, (R \vee \sqrt{d}))$. *Then, the output* p_{T-t} *of DDPM is* $\varepsilon_{\mathrm{TV}}$-*close in TV to the distribution* q_t, *which is* ε_{W_2}-*close in* W_2 *to* q, *provided that the step size* h *is chosen appropriately according to Theorem 2 and* $N = \widetilde{\Theta}\Big(\frac{d^3 R^4 \, (R \vee \sqrt{d})^4}{\varepsilon_{\mathrm{TV}}^2 \, \varepsilon_{W_2}^8} \Big)$ *and* $\varepsilon_{\mathrm{score}} \leq \widetilde{O}\Big(\frac{\varepsilon_{\mathrm{TV}}^{5/2} \, \varepsilon_{W_2}^4}{d^{3/2} R^2 \, (R \vee \sqrt{d})^2} \Big)$.

Note that the dependencies in this corollary are polynomial in all of the relevant problem parameters, which vastly improves upon the exponential dependencies of De Bortoli (2022).

Remark. The statement of Corollary 4 can be simplified by observing that both TV and W_1 distances are upper bounds for the bounded Lipschitz metric $\mathsf{d}_{\mathrm{BL}}(\mu, \nu) := \sup\{\int f \, \mathrm{d}\mu - \int f \, \mathrm{d}\nu \mid f : \mathbb{R}^d \to [-1, 1] \text{ is } 1\text{-Lipschitz}\}$. Hence, Corollary 4 can be phrased as a convergence guarantee in the bounded Lipschitz metric, which is known to metrize weak convergence (see Dudley (2002, Theorem 11.3.3)).

3.3 Results for CLD

In order to state our results for score-based generative modeling based on the CLD, we must first modify Assumptions 1 and 3 accordingly.

Assumption 4. *For all* $t \geq 0$, *the score* $\nabla_v \ln \boldsymbol{q}_t$ *is* L-*Lipschitz.*

Assumption 5. *For all* $k = 1, \dots, N$, $\mathbb{E}_{\boldsymbol{q}_{kh}}[\|\boldsymbol{s}_{kh} - \nabla_v \ln \boldsymbol{q}_{kh}\|^2] \leq \varepsilon_{\mathrm{score}}^2$.

If we ignore the dependence on L and assume that the score estimate is sufficiently accurate, then the iteration complexity guarantee of Theorem 2 is $N = \widetilde{\Theta}(d/\varepsilon^2)$. On the other hand, recall from Section 2.2 that based on intuition from the literature on log-concave sampling and from empirical findings in Dockhorn et al. (2022), we might expect that SGMs based on the CLD have a smaller iteration complexity than DDPM. We prove the following theorem.

Theorem 5 (CLD, see Section C of supplement). *Suppose that Assumptions 2, 4, and 5 hold. Let* $\boldsymbol{p}_T$ *be the output of the SGM algorithm based on the CLD (Section 2.2) at time* T, *and suppose the step size* $h := T/N$ *satisfies* $h \lesssim 1/L$, *where* $L \geq 1$. *Then, there is a universal constant* $c > 0$ *such that* $\mathsf{TV}(\boldsymbol{p}_T, q \otimes \gamma^d)$ *is bounded, up to a constant factor, by*

$$\underbrace{\sqrt{D_{\mathrm{KL}}(q \,\|\, \gamma^d) + \mathsf{FI}(q \,\|\, \gamma^d)} \exp(-cT)}_{\textit{convergence of forward process}} + \underbrace{(L\sqrt{dh} + L\mathfrak{m}_2 h)\sqrt{T}}_{\textit{discretization error}} + \underbrace{\frac{N^{1/3}\varepsilon_{\mathrm{score}}^{2/3}}{T^{1/6}}}_{\textit{score estimation error}} \,,$$

where $\mathsf{FI}(q \,\|\, \gamma^d)$ *is the relative Fisher information* $\mathsf{FI}(q \,\|\, \gamma^d) := \mathbb{E}_q[\|\nabla \ln(q/\gamma^d)\|^2]$.

Note that the result of Theorem 5 is in fact no better than our guarantee for DDPM in Theorem 2. Although it is possible that this is an artefact of our analysis, we believe that it is in fact fundamental. As we discuss in Remark C.2, from the form of the reverse process (7), the SGM based on CLD lacks

a certain property (that the discretization error should only depend on the size of the increment of the X process, not the increments of both the X and V processes) which is crucial for the improved dimension dependence of the CLD over the Langevin diffusion in log-concave sampling. Hence, in general, we conjecture that under our assumptions, SGMs based on the CLD do not achieve a better dimension dependence than DDPM.

We provide evidence for our conjecture via a lower bound. In our proofs of Theorems 2 and 5, we rely on bounding the KL divergence between certain measures on the path space $\mathcal{C}([0, T]; \mathbb{R}^d)$ via Girsanov's theorem. The following result lower bounds this KL divergence, even for the setting in which the score estimate is perfect ($\varepsilon_{\text{score}} = 0$) and the data distribution q is the standard Gaussian.

Theorem 6 (See Section C.5 of supplement). *Let $\boldsymbol{p}_T$ be the output of the SGM algorithm based on the CLD (Section 2.2) at time T, where the data distribution q is the standard Gaussian γ^d, and the score estimate is exact ($\varepsilon_{\text{score}} = 0$). Suppose that the step size h satisfies $h \leq \frac{1}{10}$. Then, for the path measures $\boldsymbol{P}_T$ and $\boldsymbol{Q}_T^{\leftarrow}$ of the algorithm and the continuous-time process (7) respectively (see Section C for details), it holds that $D_{\text{KL}}(\boldsymbol{Q}_T^{\leftarrow} \parallel \boldsymbol{P}_T) \geq dhT$.*

Theorem 6 shows that in order to make the KL divergence between the path measures small, we must take $h \lesssim 1/d$, which leads to an iteration complexity that scales linearly in the dimension d. Theorem 6 is not a proof that SGMs based on the CLD cannot achieve better than linear dimension dependence, as it is possible that the output $\boldsymbol{p}_T$ of the SGM is close to $q \otimes \gamma^d$ even if the path measures are not close, but it rules out the possibility of obtaining a better dimension dependence via our Girsanov proof technique. We believe that it provides compelling evidence for our conjecture, i.e., that under our assumptions, the CLD does not improve the complexity of SGMs over DDPM.

We remark that in this section, we have only considered the error arising from discretization of the SDE. It is possible that the score function $\nabla_v \ln \boldsymbol{q}_t$ for the SGM with the CLD is easier to estimate than the score function for DDPM, providing a *statistical* benefit of using the CLD. We do not investigate this question here and leave it as future work.

4 TECHNICAL OVERVIEW

We now give a detailed technical overview for the proof for DDPM (Theorem 2). The proof for CLD (Theorem 5) follows along similar lines.

Recall that we must deal with three sources of error: (1) the estimation of the score function; (2) the discretization of the SDE; and (3) the initialization of the reverse process at γ^d rather than at q_T.

First, we ignore the errors (1) and (2), and focus on the error (3). Hence, we consider the continuous-time reverse SDE (4), initialized from γ^d (resp. q_T) and denote by $\tilde{p}_t$ (resp. q_{T-t}) its marginal distributions. Note that $\tilde{p}_0 = \gamma^d$ and that $q_0 = q$, the data distribution. First, using the exponential contraction of the KL divergence along the (forward) OU process, we have $D_{\text{KL}}(q_T \parallel \gamma^d) \leq \exp(-2T) D_{\text{KL}}(q \parallel \gamma^d)$. Then, using the data processing inequality along the backward process, we have $\text{TV}(\tilde{p}_T, q) \leq \text{TV}(\gamma^d, q_T)$. Therefore, using Pinsker inequality, we get

$$\text{TV}(\tilde{p}_T, q) \leq \text{TV}(\gamma^d, q_T) \leq \sqrt{D_{\text{KL}}(q_T \parallel \gamma^d)} \leq \exp(-T)\sqrt{D_{\text{KL}}(q \parallel \gamma^d)}, \tag{8}$$

i.e., the output $\tilde{p}_T$ converges to the data distribution q exponentially fast as $T \to \infty$.

Next, we consider the score estimation error (1) and the discretization error (2). Similarly to (Lee et al., 2022a) we first assume that the score estimate is L^∞-accurate:

$$\max_{k=0,1,\ldots,N-1} \|s_{kh} - \nabla \ln q_{kh}\|_{\text{sup}} \leq \varepsilon_{\text{score}}. \tag{9}$$

Let p_t^∞ denote the law of the algorithm at time t with L^∞-accurate score estimate. Girsanov's theorem lets us bound the discretization error incurred by p_t^∞ by

$$\sum_{k=0}^{N-1} \mathbb{E} \int_{kh}^{(k+1)h} \|s_{T-kh}^\infty(\bar{X}_{kh}^{\leftarrow}) - \nabla \ln q_{T-t}(\bar{X}_t^{\leftarrow})\|^2 \, \mathrm{d}t$$

(see Corollary 9 in the supplement), where the expectation is with respect to the law of the reverse process. To control the integrand in this expression we can decompose

$$\|s^{\infty}_{T-kh}(\bar{X}^{\leftarrow}_{kh}) - \nabla \ln q_{T-t}(\bar{X}^{\leftarrow}_t)\|^2$$
$$\lesssim \|s^{\infty}_{T-kh}(\bar{X}^{\leftarrow}_{kh}) - \nabla \ln q_{T-kh}(\bar{X}^{\leftarrow}_{kh})\|^2 + \|\nabla \ln q_{T-kh}(\bar{X}^{\leftarrow}_{kh}) - \nabla \ln q_{T-t}(\bar{X}^{\leftarrow}_{kh})\|^2$$
$$+ \|\nabla \ln q_{T-t}(\bar{X}^{\leftarrow}_{kh}) - \nabla \ln q_{T-t}(\bar{X}^{\leftarrow}_t)\|^2 .$$

The first term on the right-hand side is bounded by the assumption that p^{∞} is given by an L^{∞}-accurate score estimate. The second term corresponds to the change in the score function along the forward process, which we can bound using tools from Lee et al. (2022a) (see also Lemma 15 for a generalization thereof that we prove for the purposes of analyzing CLD). The final term can be bounded using the fact that the score function is Lipschitz by assumption, together with a bound on how much the forward process moves over a small time period. Putting everything together, we can obtain the bound on the discretization error given by Theorem 10 in the supplement.

Finally, we must replace the assumption (9) of an L^{∞}-accurate score estimate with Assumption 3, for which we take inspiration from (Lee et al., 2022a). For $k = 0, 1, \dots, (N-1)h$, define the bad set

$$B_{kh} := \{\|s_{kh} - \nabla \ln q_{kh}\| \geq \varepsilon_{\text{score},\infty}\} .$$

If the algorithm never enters the bad sets, i.e., $X^{\leftarrow}_{kh} \notin B_{T-kh}$ for $k = 0, 1, \dots, N-1$, then it is as if the algorithm is using an L^{∞}-accurate score estimate, and we can apply the Girsanov discretization analysis. Therefore, if we can control the error probability

$$\mathbb{P}(X^{\leftarrow}_{kh} \in B_{T-kh} \text{ for some } k = 0, 1, \dots, N-1) \leq \dots ,$$

then by a coupling argument, we can bound the overall total variation error $\mathsf{TV}(p_T, q)$.

However, by Chebyshev's, we have $q_{kh}(B_{kh}) \leq \varepsilon^2_{\text{score}}/\varepsilon^2_{\text{score},\infty} =: \delta$, and by a union bound,

$$\mathbb{P}(\bar{X}^{\leftarrow}_{kh} \in B_{T-kh} \text{ for some } k = 0, 1, \dots, N-1) \leq N\delta . \tag{10}$$

In brief, the difficulty now is that our assumption allows us to control the probability of the bad event for the reverse process $\bar{X}^{\leftarrow}$, but we need to control the probability under the algorithm $X^{\leftarrow}$. To control the difference of these probabilities, we notice that Girsanov allows to control the divergence between the *joint distributions* of the two processes. So we first control the difference between the two probabilities by the TV distance between the joint distributions of the process $\bar{X}^{\leftarrow}$ and the process $X^{\leftarrow}$ initialized from q_T. We then pass to KL divergence using Pinsker's inequality and finally we use the bound from our discretization analysis with L^{∞}-accurate score estimate, which reads

$$D_{\text{KL}}\big(\text{law}(\bar{X}^{\leftarrow}_{kh} : k = 0, 1, \dots, N) \,\|\, \text{law}(X^{\leftarrow}_{kh} : k = 0, 1, \dots, N)\big) \leq \varepsilon^2_{\text{Girsanov}} . \tag{11}$$

Putting Eqs. (8), (10), and (11) together, we have

$$\mathbb{P}(X^{\leftarrow}_{kh} \in B_{T-kh} \text{ for some } 0 \leq k < N) \leq \mathsf{TV}(q_T, \gamma^d) + \varepsilon_{\text{Girsanov}} + N\delta , \tag{12}$$

where the three terms correspond to the components of the bound in Theorem 2: the convergence of the forward process, the discretization error, and the probability of failure. For the third term, the parameter δ is balanced in the final bound to optimize a trade-off (a smaller choice of δ leads to a larger value for $\varepsilon_{\text{score},\infty}$, which in turn increases the error $\varepsilon_{\text{Girsanov}}$), ultimately leading to a bound in terms of the score estimation error $\varepsilon_{\text{score}}$.

5 CONCLUSION

In this work, we provided the first convergence guarantees for SGMs which hold under realistic assumptions (namely, L^2-accurate score estimation and arbitrarily non-log-concave data distributions) and which scale polynomially in the problem parameters. Our results take a step towards explaining the remarkable empirical success of SGMs, at least assuming the score is learned with small L^2 error.

The main limitation of this work is that we did not address the question of when the score function can be learned well. In general, studying the non-convex training dynamics of learning the score function via neural networks is challenging, but we believe that the resolution of this problem, even for simple learning tasks, would shed considerable light on SGMs. Together with the results in this paper, it would yield the first end-to-end guarantees for SGMs.

In light of the interpretation of our result as a reduction of the task of sampling to the task of score function estimation, we also ask whether there are interesting situations where it is easier to learn the score function (not necessarily via a neural network) than to (directly) sample.

REFERENCES

Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2014.

Adam Block, Youssef Mroueh, and Alexander Rakhlin. Generative modeling with denoising auto-encoders and Langevin sampling. *arXiv e-prints*, art. arXiv:2002.00107, 2020.

Joan Bruna, Oded Regev, Min Jae Song, and Yi Tang. Continuous LWE. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pp. 694–707, 2021.

Hanqun Cao, Cheng Tan, Zhangyang Gao, Guangyong Chen, Pheng-Ann Heng, and Stan Z. Li. A survey on generative diffusion model. *arXiv e-prints*, art. arXiv:2209.02646, 2022.

Patrick Cattiaux, Giovanni Conforti, Ivan Gentil, and Christian Léonard. Time reversal of diffusion processes under a finite entropy condition. September 2022.

Sitan Chen, Aravind Gollakota, Adam R. Klivans, and Raghu Meka. Hardness of noise-free learning for two-hidden-layer neural networks. *arXiv preprint arXiv:2202.05258*, 2022a.

Sitan Chen, Jerry Li, and Yuanzhi Li. Learning (very) simple generative models is hard. *arXiv preprint arXiv:2205.16003*, 2022b.

Yongxin Chen, Sinho Chewi, Adil Salim, and Andre Wibisono. Improved analysis for a proximal algorithm for sampling. In Po-Ling Loh and Maxim Raginsky (eds.), *Proceedings of Thirty Fifth Conference on Learning Theory*, volume 178 of *Proceedings of Machine Learning Research*, pp. 2984–3014. PMLR, 7 2022c.

Xiang Cheng, Niladri S. Chatterji, Peter L. Bartlett, and Michael I. Jordan. Underdamped Langevin MCMC: a non-asymptotic analysis. In Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet (eds.), *Proceedings of the 31st Conference on Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pp. 300–323. PMLR, 7 2018.

Sinho Chewi. *Log-concave sampling*. 2022. Book draft available at `https://chewisinho.github.io/`.

Sinho Chewi, Murat A. Erdogdu, Mufan B. Li, Ruoqi Shen, and Matthew Zhang. Analysis of Langevin Monte Carlo from Poincaré to log-Sobolev. *arXiv e-prints*, art. arXiv:2112.12662, 2021a.

Sinho Chewi, Chen Lu, Kwangjun Ahn, Xiang Cheng, Thibaut Le Gouic, and Philippe Rigollet. Optimal dimension dependence of the Metropolis-adjusted Langevin algorithm. In Mikhail Belkin and Samory Kpotufe (eds.), *Proceedings of Thirty Fourth Conference on Learning Theory*, volume 134 of *Proceedings of Machine Learning Research*, pp. 1260–1300. PMLR, 8 2021b.

Florinel-Alin Croitoru, Vlad Hondru, Radu Tudor Ionescu, and Mubarak Shah. Diffusion models in vision: a survey. *arXiv e-prints*, art. arXiv:2209.04747, 2022.

Arnak S. Dalalyan and Lionel Riou-Durand. On sampling from a log-concave density using kinetic Langevin diffusions. *Bernoulli*, 26(3):1956–1988, 2020.

Arnak S. Dalalyan, Avetik Karagulyan, and Lionel Riou-Durand. Bounding the error of discretized Langevin algorithms for non-strongly log-concave targets. *arXiv e-prints*, art. arXiv:1906.08530, June 2019.

Amit Daniely and Gal Vardi. From local pseudorandom generators to hardness of learning. In *Conference on Learning Theory*, pp. 1358–1394. PMLR, 2021.

Valentin De Bortoli. Convergence of denoising diffusion models under the manifold hypothesis. *arXiv e-prints*, art. arXiv:2208.05314, 2022.

Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. Diffusion Schrödinger bridge with applications to score-based generative modeling. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, volume 34, pp. 17695–17709. Curran Associates, Inc., 2021.

Prafulla Dhariwal and Alexander Nichol. Diffusion models beat GANs on image synthesis. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, volume 34, pp. 8780–8794. Curran Associates, Inc., 2021.

Ilias Diakonikolas, Daniel M Kane, and Alistair Stewart. Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pp. 73–84. IEEE, 2017.

Tim Dockhorn, Arash Vahdat, and Karsten Kreis. Score-based generative modeling with critically-damped Langevin diffusion. In *International Conference on Learning Representations*, 2022.

Richard M. Dudley. *Real analysis and probability*, volume 74 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Revised reprint of the 1989 original.

Hans Föllmer. An entropy approach to the time reversal of diffusion processes. In *Stochastic differential systems (Marseille-Luminy, 1984)*, volume 69 of *Lect. Notes Control Inf. Sci.*, pp. 156–163. Springer, Berlin, 1985.

Aparna Gupte, Neekon Vafa, and Vinod Vaikuntanathan. Continuous LWE is as hard as LWE & applications to learning Gaussian mixtures. *arXiv preprint arXiv:2204.02550*, 2022.

Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 33:6840–6851, 2020.

Aapo Hyvärinen. Estimation of non-normalized statistical models by score matching. *J. Mach. Learn. Res.*, 6:695–709, 2005.

Diederik Kingma, Tim Salimans, Ben Poole, and Jonathan Ho. Variational diffusion models. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, volume 34, pp. 21696–21707. Curran Associates, Inc., 2021.

Jean-François Le Gall. *Brownian motion, martingales, and stochastic calculus*, volume 274 of *Graduate Texts in Mathematics*. Springer, [Cham], French edition, 2016.

Holden Lee, Jianfeng Lu, and Yixin Tan. Convergence for score-based generative modeling with polynomial complexity. *arXiv e-prints*, art. arXiv:2206.06227, 2022a.

Holden Lee, Jianfeng Lu, and Yixin Tan. Convergence of score-based generative modeling for general data distributions. 2022b.

Yin Tat Lee, Ruoqi Shen, and Kevin Tian. Structured logconcave sampling with a restricted Gaussian oracle. In Mikhail Belkin and Samory Kpotufe (eds.), *Proceedings of Thirty Fourth Conference on Learning Theory*, volume 134 of *Proceedings of Machine Learning Research*, pp. 2993–3050. PMLR, 8 2021.

Xingchao Liu, Lemeng Wu, Mao Ye, and Qiang Liu. Let us build bridges: Understanding and extending diffusion generative models. *arXiv preprint arXiv:2208.14699*, 2022.

Yi-An Ma, Niladri S. Chatterji, Xiang Cheng, Nicolas Flammarion, Peter L. Bartlett, and Michael I. Jordan. Is there an analog of Nesterov acceleration for gradient-based MCMC? *Bernoulli*, 27(3): 1942 – 1992, 2021.

Dan Mikulincer and Yair Shenfeld. On the lipschitz properties of transportation along heat flows. *arXiv preprint arXiv:2201.01382*, 2022.

Felix Otto and Cédric Villani. Comment on: "Hypercontractivity of Hamilton–Jacobi equations", by S. G. Bobkov, I. Gentil and M. Ledoux. *J. Math. Pures Appl. (9)*, 80(7):697–700, 2001.

Jakiw Pidstrigach. Score-based generative models detect manifolds. *arXiv e-prints*, art. arXiv:2206.01018, 2022.

Aditya Ramesh, Prafulla Dhariwal, Alex Nichol, Casey Chu, and Mark Chen. Hierarchical text-conditional image generation with CLIP latents. *arXiv preprint arXiv:2204.06125*, 2022.

Ruoqi Shen and Yin Tat Lee. The randomized midpoint method for log-concave sampling. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (eds.), *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.

Jascha Sohl-Dickstein, Eric Weiss, Niru Maheswaranathan, and Surya Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In Francis Bach and David Blei (eds.), *Proceedings of the 32nd International Conference on Machine Learning*, volume 37 of *Proceedings of Machine Learning Research*, pp. 2256–2265, Lille, France, 7 2015. PMLR.

Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (eds.), *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.

Yang Song, Conor Durkan, Iain Murray, and Stefano Ermon. Maximum likelihood training of score-based diffusion models. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, volume 34, pp. 1415–1428. Curran Associates, Inc., 2021a.

Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2021b.

J. Michael Steele. *Stochastic calculus and financial applications*, volume 45 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 2001.

Arash Vahdat, Karsten Kreis, and Jan Kautz. Score-based generative modeling in latent space. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems*, volume 34, pp. 11287–11302. Curran Associates, Inc., 2021.

Santosh Vempala and Andre Wibisono. Rapid convergence of the unadjusted Langevin algorithm: isoperimetry suffices. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (eds.), *Advances in Neural Information Processing Systems 32*, pp. 8094–8106. Curran Associates, Inc., 2019.

Cédric Villani. Hypocoercivity. *Mem. Amer. Math. Soc.*, 202(950):iv+141, 2009.

Pascal Vincent. A connection between score matching and denoising autoencoders. *Neural Comput.*, 23(7):1661–1674, 2011.

Ling Yang, Zhilong Zhang, Shenda Hong, Wentao Zhang, and Bin Cui. Diffusion models: a comprehensive survey of methods and applications. *arXiv e-prints*, art. arXiv:2209.00796, 2022.

Qinsheng Zhang and Yongxin Chen. Fast sampling of diffusion models with exponential integrator. *arXiv e-prints*, art. arXiv:2204.13902, 2022.

A PRELIMINARIES

In this section, we review the notion of score matching, state and prove the elementary lemma that leads to (12) above, and provide a list of notation for the proofs.

A.1 PRIMER ON SCORE MATCHING

In order to estimate the score function $\nabla \ln q_t$, consider minimizing the $L^2(q_t)$ loss over a function class $\mathscr{F}$,

$$\underset{s_t \in \mathscr{F}}{\text{minimize}} \quad \mathbb{E}_{q_t}[\|s_t - \nabla \ln q_t\|^2], \tag{13}$$

where $\mathscr{F}$ could be, e.g., a class of neural networks. The idea of score matching, which goes back to Hyvärinen (2005); Vincent (2011), is that after applying integration by parts for the Gaussian measure, the problem (13) is *equivalent* to the following problem:

$$\underset{s_t \in \mathscr{F}}{\text{minimize}} \quad \mathbb{E}\left[\left\|s_t(\bar{X}_t) + \frac{1}{\sqrt{1 - \exp(-2t)}} Z_t\right\|^2\right], \tag{14}$$

where $Z_t \sim \text{normal}(0, I_d)$ is independent of $\bar{X}_0$ and $\bar{X}_t = \exp(-t)\,\bar{X}_0 + \sqrt{1 - \exp(-2t)}\,Z_t$, in the sense that (13) and (14) share the same minimizers. We give a self-contained derivation in Appendix D for the sake of completeness. Unlike (13), however, the objective in (14) can be replaced with an empirical version and estimated on the basis of samples $\bar{X}_0^{(1)}, \ldots, \bar{X}_0^{(n)}$ from q, leading to the finite-sample problem

$$\underset{s_t \in \mathscr{F}}{\text{minimize}} \quad \frac{1}{n} \sum_{i=1}^{n} \left\|s_t(\bar{X}_t^{(i)}) + \frac{1}{\sqrt{1 - \exp(-2t)}} Z_t^{(i)}\right\|^2,$$

where $(Z_t^{(i)})_{i \in [n]}$ are i.i.d. standard Gaussians independent of the data $(\bar{X}_0^{(i)})_{i \in [n]}$. Moreover, if we parameterize the score function as $s_t = -\frac{1}{\sqrt{1 - \exp(-2t)}} \widehat{z}_t$, then the empirical problem is equivalent to

$$\underset{\widehat{z}_t \in -\sqrt{1 - \exp(-2t)}\,\mathscr{F}}{\text{minimize}} \quad \frac{1}{n} \sum_{i=1}^{n} \left\|\widehat{z}_t(\bar{X}_t^{(i)}) - Z_t^{(i)}\right\|^2,$$

which has the illuminating interpretation of predicting the added noise $Z_t^{(i)}$ from the noised data $\bar{X}_t^{(i)}$.

A.2 TRUNCATION LEMMA

Lemma 7. *Suppose that $(\bar{X}_{kh}^{\leftarrow})_{k \in \mathbb{N}}$, $(X_{kh}^{\leftarrow})_{k \in \mathbb{N}}$, and $(X_{kh}^{\leftarrow,\infty})_{k \in \mathbb{N}}$ are random processes defined on the same probability space. For each $k \in \mathbb{N}$, let $q_{kh}^{\leftarrow}$, p_{kh}, and p_{kh}^{∞} denote the laws of $\bar{X}_{kh}^{\leftarrow}$, $X_{kh}^{\leftarrow}$, and $X_{kh}^{\leftarrow,\infty}$ respectively.*

Suppose that $(B_{kh})_{k \in \mathbb{N}}$ are measurable subsets of $\mathbb{R}^d$ and that the following hold.

1. *If $X_{kh}^{\leftarrow,\infty} \notin B_{T-kh}$ for $k = 0, 1, \ldots, N-1$, then $X_{kh}^{\leftarrow} = X_{kh}^{\leftarrow,\infty}$ for all $k = 0, 1, \ldots, N$.*

2. *It holds that*

$$\mathsf{TV}\big(\text{law}(X_{kh}^{\leftarrow,\infty} : k = 0, 1, \ldots, N-1),\ \text{law}(\bar{X}_{kh}^{\leftarrow} : k = 0, 1, \ldots, N-1)\big) \le \varepsilon\,.$$

3. *$\mathbb{P}[\bar{X}_{kh}^{\leftarrow} \in B_{T-kh}] \le \delta$ for all $k = 0, 1, \ldots, N-1$.*

Then, the following holds:

$$\mathsf{TV}(p_{Nh}, q_{Nh}^{\leftarrow}) \le 2\varepsilon + N\delta\,.$$

Proof. By triangle inequality,

$$\mathsf{TV}(p_{Nh}, q_{Nh}^\leftarrow) \le \mathsf{TV}(p_{Nh}, p_{Nh}^\infty) + \mathsf{TV}(p_{Nh}^\infty, q_{Nh}^\leftarrow) \le \mathsf{TV}(p_{Nh}, p_{Nh}^\infty) + \varepsilon \,,$$

where we used the data processing inequality. Then, by the union bound

$$\begin{aligned}
\mathsf{TV}(p_{Nh}, p_{Nh}^\infty) &\le \mathbb{P}(X_{Nh}^\leftarrow \ne X_{Nh}^{\leftarrow,\infty}) \le \mathbb{P}(X_{kh}^{\leftarrow,\infty} \in B_{T-kh} \text{ for some } k = 0, 1, \dots, N-1) \\
&\le \varepsilon + \mathbb{P}(\bar{X}_{kh}^\leftarrow \in B_{T-kh} \text{ for some } k = 0, 1, \dots, N-1) \\
&\le \varepsilon + \sum_{k=0}^{N-1} \mathbb{P}(\bar{X}_{kh}^\leftarrow \in B_{kh}) \le \varepsilon + N\delta \,.
\end{aligned}$$

This proves the lemma. □

A.3 NOTATION

L^∞**-accurate score estimate.** The bad set is defined to be

$$B_t := \{\|s_t - \nabla \ln q_t\| \ge \varepsilon_{\mathrm{score},\infty}\} \,,$$

where $\varepsilon_{\mathrm{score},\infty} > 0$ is a parameter to be chosen later. We define the L^∞-accurate score estimate to be

$$s_t^\infty := s_t \mathbb{1}_{B_t^c} + \nabla \ln q_t \mathbb{1}_{B_t} \,. \tag{15}$$

We note that $\|s_t^\infty - \nabla \ln q_t\| \le \varepsilon_{\mathrm{score},\infty}$.

Stochastic processes and their laws.

- The data distribution is $q = q_0$.
- The forward process (1) is denoted $(\bar{X}_t)_{t \in [0,T]}$, and $\bar{X}_t \sim q_t$.
- The reverse process (4) is denoted $(\bar{X}_t^\leftarrow)_{t \in [0,T]}$, where $\bar{X}_t^\leftarrow := \bar{X}_{T-t} \sim q_{T-t}$.
- The SGM algorithm (5) is denoted $(X_t^\leftarrow)_{t \in [0,T]}$, and $X_t^\leftarrow \sim p_t$. Recall that we initialize at $p_0 = \gamma^d$, the standard Gaussian measure.
- The SGM algorithm with L^∞-accurate score, denoted $(X_t^{\leftarrow,\infty})_{t \in [0,T]}$, is the same as (5) except that we use s_{T-kh}^∞ defined in (15) instead of s_{T-kh} for $k = 0, 1, \dots, N-1$. We write $X_t^{\leftarrow,\infty} \sim p_t^\infty$, and we initialize the process at $p_0^\infty = \gamma^d$.
- The process $(X_t^{\leftarrow,\infty,q_T})_{t \in [0,T]}$ is the same as $(X_t^{\leftarrow,\infty})_{t \in [0,T]}$, except that we initialize this process at q_T rather than at γ^d. We write $X_t^{\leftarrow,\infty,q_T} \sim p_t^{\infty,q_T}$.

Conventions for Girsanov's theorem. When we apply Girsanov's theorem, it is convenient to instead think about a single stochastic process, which for ease of notation we denote simply via $(X_t)_{t \in [0,T]}$, and we consider different measures over the path space $\mathcal{C}([0,T]; \mathbb{R}^d)$.

The two measures we consider over path space are:

- $Q_T^\leftarrow$, under which $(X_t)_{t \in [0,T]}$ has the law of the reverse process (4); and
- P_T^{∞,q_T}, under which $(X_t)_{t \in [0,T]}$ has the law of the SGM algorithm with L^∞-accurate score initialized at q_T (corresponding to the process $(X_t^{\leftarrow,\infty,q_T})_{t \in [0,T]}$ defined above).

Other parameters. We recall that $T > 0$ denotes the total time for which we run the forward process; $h > 0$ is the step size of the discretization; $L \ge 1$ is the Lipschitz constant of the score function; $\mathfrak{m}_2^2 := \mathbb{E}_q[\|\cdot\|^2]$ is the second moment under the data distribution; and $\varepsilon_{\mathrm{score}}$ is the L^2 score estimation error.

Notation for CLD. The notational conventions for the CLD are similar; however, we must also consider a velocity variable V. When discussing quantities which involve both position and velocity (e.g., the joint distribution $\boldsymbol{q}_t$ of $(\bar{X}_t, \bar{V}_t)$), we typically use boldface fonts.

B PROOFS FOR DDPM

B.1 PRELIMINARIES ON GIRSANOV'S THEOREM

First, we recall the statement of Girsanov's theorem.

Theorem 8 (Girsanov's theorem, (Le Gall, 2016, Theorem 5.22)). *Let P_T and Q_T be two probability measures on path space $\mathcal{C}([0, T]; \mathbb{R}^d)$. Suppose that under P_T, the process $(X_t)_{t \in [0,T]}$ follows*

$$\mathrm{d}X_t = \tilde{b}_t \, \mathrm{d}t + \sigma_t \, \mathrm{d}\tilde{B}_t \,,$$

where $\tilde{B}$ is a P_T-Brownian motion, and under Q_T, the process $(X_t)_{t \in [0,T]}$ follows

$$\mathrm{d}X_t = b_t \, \mathrm{d}t + \sigma_t \, \mathrm{d}B_t \,,$$

where B is a Q_T-Brownian motion. We assume that for each $t > 0$, σ_t is a $d \times d$ symmetric positive definite matrix. Then, provided that Novikov's condition holds,

$$\mathbb{E}_{Q_T} \exp\left(\frac{1}{2} \int_0^T \|\sigma_t^{-1} (\tilde{b}_t - b_t)\|^2 \, \mathrm{d}t\right) < \infty \,, \tag{16}$$

we have that

$$\frac{\mathrm{d}P_T}{\mathrm{d}Q_T} = \exp\left(\int_0^T \sigma_t^{-1} (\tilde{b}_t - b_t) \, \mathrm{d}B_t - \frac{1}{2} \int_0^T \|\sigma_t^{-1} (\tilde{b}_t - b_t)\|^2 \, \mathrm{d}t\right) \,.$$

We would like to apply Girsanov's theorem with $P_T = P_T^{\infty, q_T}$, $Q_T = Q_T^{\leftarrow}$, $\tilde{b}_t = X_t + 2 \, s_{T-kh}^\infty(X_{kh})$ (for $t \in [kh, (k+1)h]$), $b_t = X_t + 2 \nabla \ln q_{T-t}(X_t)$, and $\sigma_t = \sqrt{2} \, I_d$. Unfortunately, under Assumptions 1, 2, and 3 alone, Novikov's condition need not hold.[3] In Appendix F we give a workaround for this issue by defining a sequence of suitable truncations of the reverse and SGM processes for which Novikov's condition does hold and showing that these truncations converge in an appropriate sense to the processes we care about. In this section however, we will work under the assumption that P_T^{∞, q_T} and $Q_T^{\leftarrow}$ do satisfy Novikov's condition as it leads to a considerably simpler proof.

Corollary 9. *Assuming that P_T^{∞, q_T} and Q_T satisfy Novikov's condition (16), it holds that*

$$D_{\mathrm{KL}}(Q_T^{\leftarrow} \| P_T^{\infty, q_T}) = \mathbb{E}_{Q_T^{\leftarrow}} \ln \frac{\mathrm{d}Q_T^{\leftarrow}}{\mathrm{d}P_T^{\infty, q_T}} = \sum_{k=0}^{N-1} \mathbb{E}_{Q_T^{\leftarrow}} \int_{kh}^{(k+1)h} \|s_{T-kh}^\infty(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2 \, \mathrm{d}t \,.$$

B.2 GIRSANOV DISCRETIZATION ARGUMENT

We now apply Corollary 9 and bound the discretization error.

Theorem 10 (discretization error for DDPM). *Suppose that Assumptions 1, 2, and 3 hold. Let $Q_T^{\leftarrow}$ and P_T^{∞, q_T} denote the measures on path space corresponding to the reverse process (4) and the SGM algorithm with L^∞-accurate score estimate initialized at q_T. Assume that $L \geq 1$ and $h \lesssim 1/L$. Then,*

$$\mathsf{TV}(P_T^{\infty, q_T}, Q_T^{\leftarrow})^2 \lesssim (\varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 \mathfrak{m}_2^2 h^2) \, T \,.$$

We first give a proof under the assumption that P_T^{∞, q_T} and $Q_T^{\leftarrow}$ satisfy Novikov's condition (16). In Appendix F, we show how to lift this assumption and give a fully rigorous proof of this theorem.

Proof assuming that Novikov's condition holds. For $t \in [kh, (k+1)h]$, we can decompose

$$\|s_{T-kh}^\infty(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2$$
$$\lesssim \|s_{T-kh}^\infty(X_{kh}) - \nabla \ln q_{T-kh}(X_{kh})\|^2 + \|\nabla \ln q_{T-kh}(X_{kh}) - \nabla \ln q_{T-t}(X_{kh})\|^2$$
$$\quad + \|\nabla \ln q_{T-t}(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2$$
$$\lesssim \varepsilon_{\mathrm{score},\infty}^2 + \left\|\nabla \ln \frac{q_{T-kh}}{q_{T-t}}(X_{kh})\right\|^2 + L^2 \|X_{kh} - X_t\|^2 \,. \tag{17}$$

---

[3] In order to check Novikov's condition, we would want X_0 to have sub-Gaussian tails for instance.

We must bound the change in the score function along the forward process. If $S : \mathbb{R}^d \to \mathbb{R}^d$ is the mapping $S(x) := \exp(-(t-kh))\, x$, then $q_{T-kh} = S_\# q_{T-t} * \mathsf{normal}(0, 1 - \exp(-2\,(t-kh)))$. We can then use (Lee et al., 2022a, Lemma C.12) (or the more general Lemma 15 that we prove in Section C.4) with $\alpha = \exp(t - kh) = 1 + O(h)$ and $\sigma^2 = 1 - \exp(-2\,(t-kh)) = O(h)$ to obtain

$$
\begin{aligned}
\left\| \nabla \ln \frac{q_{T-kh}}{q_{T-t}}(X_{kh}) \right\|^2 &\lesssim L^2 dh + L^2 h^2 \|X_{kh}\|^2 + (1+L^2)\, h^2 \|\nabla \ln q_{T-t}(X_{kh})\|^2 \\
&\lesssim L^2 dh + L^2 h^2 \|X_{kh}\|^2 + L^2 h^2 \|\nabla \ln q_{T-t}(X_{kh})\|^2
\end{aligned}
\tag{18}
$$

where the last line uses $L \geq 1$.

For the last term,

$$
\begin{aligned}
\|\nabla \ln q_{T-t}(X_{kh})\|^2 &\lesssim \|\nabla \ln q_{T-t}(X_t)\|^2 + \|\nabla \ln q_{T-t}(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2 \\
&\lesssim \|\nabla \ln q_{T-t}(X_t)\|^2 + L^2 \|X_{kh} - X_t\|^2,
\end{aligned}
\tag{19}
$$

where the second term above is absorbed into the third term of the decomposition (17). Hence,

$$
\begin{aligned}
\|s^\infty_{T-kh}(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2 &\lesssim \varepsilon^2_{\text{score},\infty} + L^2 dh + L^2 h^2 \|X_{kh}\|^2 \\
&\quad + L^2 h^2 \|\nabla \ln q_{T-t}(X_t)\|^2 + L^2 \|X_{kh} - X_t\|^2.
\end{aligned}
$$

Using the fact that under $Q^\leftarrow_T$, the process $(X_t)_{t \in [0,T]}$ is the time reversal of the forward process $(\bar{X}_t)_{t \in [0,T]}$, we can apply the moment bounds in Lemma 11 and the movement bound in Lemma 12 to obtain

$$
\begin{aligned}
\mathbb{E}_{Q^\leftarrow_T}&[\|s^\infty_{T-kh}(X_{kh}) - \nabla \ln q_{T-t}(X_t)\|^2] \\
&\lesssim \varepsilon^2_{\text{score},\infty} + L^2 dh + L^2 h^2\,(d + \mathfrak{m}_2^2) + L^3 dh^2 + L^2\,(\mathfrak{m}_2^2 h^2 + dh) \\
&\lesssim \varepsilon^2_{\text{score},\infty} + L^2 dh + L^2 \mathfrak{m}_2^2 h^2.
\end{aligned}
$$

The result now follows from Corollary 9 and Pinsker's inequality. $\qquad\square$

B.3 Proof of Theorem 2

We can now conclude our main result, namely a convergence guarantee when we only have an L^2-accurate score estimate. To prove this, we use the following truncation lemma.

Proof of Theorem 2. We recall the notation from Section 4. By the data processing inequality, Theorem 10 implies

$$
\begin{aligned}
\mathsf{TV}\big(\mathsf{law}(X^{\leftarrow,\infty,q_T}_{kh} : k = 0, 1, \dots, N-1),\ \mathsf{law}(\bar{X}^\leftarrow_{kh} : k = 0, 1, \dots, N-1)\big) \\
\lesssim (\varepsilon_{\text{score},\infty} + L\sqrt{dh} + L\mathfrak{m}_2 h)\,\sqrt{T}.
\end{aligned}
$$

By the data processing inequality and Pinsker's inequality,

$$
\begin{aligned}
\mathsf{TV}\big(\mathsf{law}(X^{\leftarrow,\infty,q_T}_{kh} : k = 0, 1, \dots, N-1),\ \mathsf{law}(X^{\leftarrow,\infty}_{kh} : k = 0, 1, \dots, N-1)\big) \\
\leq \mathsf{TV}(q_T, \gamma^d) \leq \sqrt{D_{\mathrm{KL}}(q_T \,\|\, \gamma^d)} \leq \sqrt{D_{\mathrm{KL}}(q \,\|\, \gamma^d)}\, \exp(-T),
\end{aligned}
$$

where we used the convergence of the OU process in KL divergence (see, e.g., (Bakry et al., 2014, Theorem 5.2.1)). Hence, we can apply Lemma 7. Here, we take the bad set $B_t := \{\|s_t - \nabla \ln q_t\| \geq \varepsilon_{\text{score},\infty}\}$, so by Chebyshev's inequality and Assumption 3, we can take $\delta = (\varepsilon_{\text{score}}/\varepsilon_{\text{score},\infty})^2$. It yields

$$
\mathsf{TV}(p_T, q) \lesssim \sqrt{D_{\mathrm{KL}}(q \,\|\, \gamma^d)}\, \exp(-T) + (\varepsilon_{\text{score},\infty} + L\sqrt{dh} + L\mathfrak{m}_2 h)\,\sqrt{T} + \frac{N\varepsilon^2_{\text{score}}}{\varepsilon^2_{\text{score},\infty}}.
$$

Theorem 2 follows after optimizing over the choice of $\varepsilon_{\text{score},\infty}$. $\qquad\square$

B.4 Auxiliary lemmas

In this section, we prove some auxiliary lemmas which are used in the proof of Theorem 2.

Lemma 11 (moment bounds for DDPM). *Suppose that Assumptions 1 and 2 hold. Let $(\bar{X}_t)_{t\in[0,T]}$ denote the forward process* (1).

1. *(moment bound) For all $t \geq 0$,*

$$\mathbb{E}[\|\bar{X}_t\|^2] \leq d \vee \mathrm{m}_2^2 \,.$$

2. *(score function bound) For all $t \geq 0$,*

$$\mathbb{E}[\|\nabla \ln q_t(\bar{X}_t)\|^2] \leq Ld \,.$$

Proof. 1. Along the OU process, we have $\bar{X}_t \stackrel{\mathrm{d}}{=} \exp(-t)\,\bar{X}_0 + \sqrt{1 - \exp(-2t)}\,\xi$, where $\xi \sim \mathsf{normal}(0, I_d)$ is independent of $\bar{X}_0$. Hence,

$$\mathbb{E}[\|\bar{X}_t\|^2] = \exp(-2t)\mathbb{E}[\|X\|^2] + \{1 - \exp(-2t)\}\,d \leq d \vee \mathrm{m}_2^2 \,. \qquad \square$$

2. This follows from the L-smoothness of $\ln q_t$ (see, e.g., (Vempala & Wibisono, 2019, Lemma 9)). We give a short proof for the sake of completeness.

If $\mathscr{L}_t f := \Delta f - \langle \nabla U_t, \nabla f \rangle$ is the generator associated with $q_t \propto \exp(-U_t)$, then

$$0 = \mathbb{E}_{q_t}\mathscr{L}U_t = \mathbb{E}_{q_t}\Delta U_t - \mathbb{E}_{q_t}[\|\nabla U_t\|^2] \leq Ld - \mathbb{E}_{q_t}[\|\nabla U_t\|^2] \,.$$

Lemma 12 (movement bound for DDPM). *Suppose that Assumption 2 holds. Let $(\bar{X}_t)_{t\in[0,T]}$ denote the forward process* (1). *For $0 \leq s < t$ with $\delta := t - s$, if $\delta \leq 1$, then*

$$\mathbb{E}[\|\bar{X}_t - \bar{X}_s\|^2] \lesssim \delta^2 \mathrm{m}_2^2 + \delta d \,.$$

Proof. We can write

$$\begin{aligned}
\mathbb{E}[\|\bar{X}_t - \bar{X}_s\|^2] &= \mathbb{E}\Big[\Big\|-\int_s^t \bar{X}_r \,\mathrm{d}r + \sqrt{2}\,(B_t - B_s)\Big\|^2\Big] \\
&\lesssim \delta \int_s^t \mathbb{E}[\|\bar{X}_r\|^2] \,\mathrm{d}r + \delta d \lesssim \delta^2\,(d + \mathrm{m}_2^2) + \delta d \\
&\lesssim \delta^2 \mathrm{m}_2^2 + \delta d \,,
\end{aligned}$$

where we used Lemma 11. $\qquad \square$

C Proofs for CLD

C.1 Background on the CLD process

More generally, for the forward process we can introduce a *friction parameter* $\gamma > 0$ and consider

$$\begin{aligned}
\mathrm{d}\bar{X}_t &= \bar{V}_t \,\mathrm{d}t \,, \\
\mathrm{d}\bar{V}_t &= -\bar{X}_t \,\mathrm{d}t - \gamma\,\bar{V}_t \,\mathrm{d}t + \sqrt{2\gamma}\,\mathrm{d}B_t \,.
\end{aligned}$$

If we write $\bar{\boldsymbol{\theta}}_t := (\bar{X}_t, \bar{V}_t)$, then the forward process satisfies the linear SDE

$$\mathrm{d}\bar{\boldsymbol{\theta}}_t = \boldsymbol{A}_\gamma \bar{\boldsymbol{\theta}}_t \,\mathrm{d}t + \Sigma_\gamma \,\mathrm{d}B_t \,, \qquad \text{where } \boldsymbol{A}_\gamma := \begin{bmatrix} 0 & 1 \\ -1 & -\gamma \end{bmatrix} \text{ and } \Sigma_\gamma := \begin{bmatrix} 0 \\ \sqrt{2\gamma} \end{bmatrix} \,.$$

The solution to the SDE is given by

$$\bar{\boldsymbol{\theta}}_t = \exp(t\boldsymbol{A}_\gamma)\,\bar{\boldsymbol{\theta}}_0 + \int_0^t \exp\{(t - s)\,\boldsymbol{A}_\gamma\}\,\Sigma_\gamma \,\mathrm{d}B_s \,, \tag{20}$$

which means that by the Itô isometry,

$$\mathrm{law}(\bar{\boldsymbol{\theta}}_t) = \exp(t\boldsymbol{A}_\gamma)_\# \mathrm{law}(\bar{\boldsymbol{\theta}}_0) * \mathrm{normal}\Big(0, \int_0^t \exp\{(t-s)\,\boldsymbol{A}_\gamma\}\,\Sigma_\gamma\Sigma_\gamma^\mathsf{T}\exp\{(t-s)\,\boldsymbol{A}_\gamma^\mathsf{T}\}\,\mathrm{d}s\Big).$$

Since $\det \boldsymbol{A}_\gamma = 1$, $\boldsymbol{A}_\gamma$ is always invertible. Moreover, from $\mathrm{tr}\,\boldsymbol{A}_\gamma = -\gamma$, one can work out that the spectrum of $\boldsymbol{A}_\gamma$ is

$$\mathrm{spec}(\boldsymbol{A}_\gamma) = \Big\{-\frac{\gamma}{2} \pm \sqrt{\frac{\gamma^2}{4}-1}\Big\}.$$

However, $\boldsymbol{A}_\gamma$ is not diagonalizable. The case of $\gamma = 2$ is special, as it corresponds to the case when the spectrum is $\{-1\}$, and it corresponds to the *critically damped case*. Following (Dockhorn et al., 2022), which advocated for setting $\gamma = 2$, we will also only consider the critically damped case. This also has the advantage of substantially simplifying the calculations.

C.2 GIRSANOV DISCRETIZATION ARGUMENT

In order to apply Girsanov's theorem, we introduce the path measures $\boldsymbol{P}_T^{\infty,\boldsymbol{q}_T}$ and $\boldsymbol{Q}_T^{\leftarrow}$, under which

$$\begin{aligned}
\mathrm{d}X_t &= -V_t\,\mathrm{d}t\,, \\
\mathrm{d}V_t &= \{X_t + 2\,V_t + 4\,\boldsymbol{s}_{T-kh}(X_{kh},V_{kh})\}\,\mathrm{d}t + 2\,\mathrm{d}B_t\,,
\end{aligned}$$

for $t \in [kh, (k+1)h]$, and

$$\begin{aligned}
\mathrm{d}X_t &= -V_t\,\mathrm{d}t\,, \\
\mathrm{d}V_t &= \{X_t + 2\,V_t + 4\,\nabla_v \ln \boldsymbol{q}_{T-t}(X_t,V_t)\}\,\mathrm{d}t + 2\,\mathrm{d}B_t\,,
\end{aligned}$$

respectively. We can apply Girsanov's theorem with

$$\tilde{b}_t = \begin{bmatrix} -V_t \\ X_t + 2\,V_t + 4\,\boldsymbol{s}_{T-kh}(X_{kh},V_{kh}) \end{bmatrix},\quad b_t = \begin{bmatrix} -V_t \\ X_t + 2\,V_t + 4\,\nabla_v \ln \boldsymbol{q}_{T-t}(X_t,V_t) \end{bmatrix},\quad \sigma_t = \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix}.$$

Here, σ_t is not invertible, so strictly speaking Theorem 8 does not apply; however, since $\tilde{b}_t - b_t \in \mathrm{range}\,\sigma_t$, the expression $\sigma_t^{-1}\,(\tilde{b}_t - b_t)$ is still well-defined, and the following corollary can be justified, provided that Novikov's condition holds.

Corollary 13. *Suppose that Novikov's condition* (16) *holds. Then,*

$$\begin{aligned}
D_{\mathrm{KL}}(\boldsymbol{Q}_T^{\leftarrow} \,\|\, \boldsymbol{P}_T^{\infty,\boldsymbol{q}_T}) &= \mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}} \ln \frac{\mathrm{d}\boldsymbol{Q}_T^{\leftarrow}}{\mathrm{d}\boldsymbol{P}_T^{\infty,\boldsymbol{q}_T}} \\
&= 2\sum_{k=0}^{N-1} \mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}} \int_{kh}^{(k+1)h} \|\boldsymbol{s}_{T-kh}(X_{kh},V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t,V_t)\|^2\,\mathrm{d}t\,.
\end{aligned}$$

Using this, we now aim to prove the following theorem.

Theorem 14 (discretization error for CLD). *Suppose that Assumptions 2, 4, and 5 hold. Let $\boldsymbol{Q}_T^{\leftarrow}$ and $\boldsymbol{P}_T^{\infty,\boldsymbol{q}_T}$ denote the measures on path space corresponding to the reverse process* (7) *and the SGM algorithm with L^∞-accurate score estimate initialized at $\boldsymbol{q}_T$. Assume that $L \geq 1$ and $h \lesssim 1/L$. Then,*

$$\mathsf{TV}(\boldsymbol{P}_T^{\infty,\boldsymbol{q}_T}, \boldsymbol{Q}_T^{\leftarrow})^2 \lesssim (\varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 \mathfrak{m}_2^2 h^2)\,T\,.$$

Similarly to Section B.2, we prove the theorem assuming that Novikov's condition (16) holds. This assumption can be lifted using similar arguments to the ones in Appendix F, and we omit the details.

Proof assuming that Novikov's condition holds. For $t \in [kh, (k+1)h]$, we can decompose

$$\begin{aligned}
&\|\boldsymbol{s}_{T-kh}^\infty(X_{kh},V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t,V_t)\|^2 \\
&\qquad\lesssim \|\boldsymbol{s}_{T-kh}^\infty(X_{kh},V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-kh}(X_{kh},V_{kh})\|^2 \\
&\qquad\quad + \|\nabla_v \ln \boldsymbol{q}_{T-kh}(X_{kh},V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_{kh},V_{kh})\|^2 \\
&\qquad\quad + \|\nabla_v \ln \boldsymbol{q}_{T-t}(X_{kh},V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t,V_t)\|^2 \\
&\qquad\lesssim \varepsilon_{\mathrm{score},\infty}^2 + \Big\|\nabla_v \ln \frac{\boldsymbol{q}_{T-kh}}{\boldsymbol{q}_{T-t}}(X_{kh},V_{kh})\Big\|^2 + L^2\,\|(X_{kh},V_{kh}) - (X_t,V_t)\|^2\,. \qquad (21)
\end{aligned}$$

The change in the score function is bounded by Lemma 15, which generalizes (Lee et al., 2022a, Lemma C.12). From the representation (20) of the solution to the CLD, we note that

$$\boldsymbol{q}_{T-kh} = (\boldsymbol{M}_0)_\# \boldsymbol{q}_{T-t} * \mathsf{normal}(0, \boldsymbol{M}_1)$$

with

$$\boldsymbol{M}_0 = \exp\bigl((t - kh)\,\boldsymbol{A}_2\bigr),$$
$$\boldsymbol{M}_1 = \int_0^{t-kh} \exp\{(t - kh - s)\,\boldsymbol{A}_2\}\,\Sigma_2 \Sigma_2^\mathsf{T} \exp\{(t - kh - s)\,\boldsymbol{A}_2^\mathsf{T}\}\,\mathrm{d}s\,.$$

In particular, since $\|\boldsymbol{A}_2\|_{\mathrm{op}} \lesssim 1$, $\|\boldsymbol{A}_2^{-1}\|_{\mathrm{op}} \lesssim 1$, and $\|\Sigma_2\|_{\mathrm{op}} \lesssim 1$ it follows that $\|\boldsymbol{M}_0\|_{\mathrm{op}} = 1 + O(h)$ and $\|\boldsymbol{M}_1\|_{\mathrm{op}} = O(h)$. Substituting this into Lemma 15, we deduce that if $h \lesssim 1/L$, then

$$\left\| \nabla_v \ln \frac{\boldsymbol{q}_{T-kh}}{\boldsymbol{q}_{T-t}}(X_{kh}, V_{kh}) \right\|^2$$
$$\leq \left\| \nabla \ln \frac{\boldsymbol{q}_{T-kh}}{\boldsymbol{q}_{T-t}}(X_{kh}, V_{kh}) \right\|^2$$
$$\lesssim L^2 dh + L^2 h^2\,(\|X_{kh}\|^2 + \|V_{kh}\|^2) + (1 + L^2)\,h^2\,\|\nabla \ln \boldsymbol{q}_{T-t}(X_{kh}, V_{kh})\|^2$$
$$\lesssim L^2 dh + L^2 h^2\,(\|X_{kh}\|^2 + \|V_{kh}\|^2) + L^2 h^2\,\|\nabla \ln \boldsymbol{q}_{T-t}(X_{kh}, V_{kh})\|^2\,,$$

where in the last step we used $L \geq 1$.

For the last term,

$$\|\nabla \ln \boldsymbol{q}_{T-t}(X_{kh}, V_{kh})\|^2 \lesssim \|\nabla \ln \boldsymbol{q}_{T-t}(X_t, V_t)\|^2 + L^2\,\|(X_{kh}, V_{kh}) - (X_t, V_t)\|^2\,,$$

where the second term above is absorbed into the third term of the decomposition (21). Hence,

$$\|\boldsymbol{s}_{T-kh}^\infty(X_{kh}, V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t, V_t)\|^2$$
$$\lesssim \varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 h^2\,(\|X_{kh}\|^2 + \|V_{kh}\|^2)$$
$$+ L^2 h^2\,\|\nabla \ln \boldsymbol{q}_{T-t}(X_t, V_t)\|^2 + L^2\,\|(X_{kh}, V_{kh}) - (X_t, V_t)\|^2\,.$$

By applying the moment bounds in Lemma 16 together with Lemma 17 on the movement of the CLD,

$$\|\boldsymbol{s}_{T-kh}^\infty(X_{kh}, V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t, V_t)\|^2$$
$$\lesssim \varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 h^2\,(d + \mathfrak{m}_2^2) + L^3 dh^2 + L^2\,(dh + \mathfrak{m}_2^2 h^2)$$
$$\lesssim \varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 \mathfrak{m}_2^2 h^2\,.$$

Together with Corollary 13 and Pinsker's inequality, it completes the proof. $\square$

Remark. We now pause to discuss why the discretization bound above does not improve upon the result for DDPM (Theorem 10). In the context of log-concave sampling, one instead considers the underdamped Langevin process

$$\mathrm{d}X_t = V_t\,,$$
$$\mathrm{d}V_t = -\nabla U(X_t)\,\mathrm{d}t - \gamma\,V_t\,\mathrm{d}t + \sqrt{2\gamma}\,\mathrm{d}B_t\,,$$

which is discretized to yield the algorithm

$$\mathrm{d}X_t = V_t\,,$$
$$\mathrm{d}V_t = -\nabla U(X_{kh})\,\mathrm{d}t - \gamma\,V_t\,\mathrm{d}t + \sqrt{2\gamma}\,\mathrm{d}B_t\,,$$

for $t \in [kh, (k+1)h]$. Let $\boldsymbol{P}_T$ denote the path measure for the algorithm, and let $\boldsymbol{Q}_T$ denote the path measure for the continuous-time process. After applying Girsanov's theorem, we obtain

$$D_{\mathrm{KL}}(\boldsymbol{Q}_T \parallel \boldsymbol{P}_T) \asymp \frac{1}{\gamma} \sum_{k=0}^{N-1} \mathbb{E}_{\boldsymbol{Q}_T} \int_{kh}^{(k+1)h} \|\nabla U(X_t) - \nabla U(X_{kh})\|^2\,\mathrm{d}t\,.$$

In this expression, note that ∇U depends only on the position coordinate. Since the X process is smoother (as we do not add Brownian motion directly to X), the error $\|\nabla U(X_t) - \nabla U(X_{kh})\|^2$ is of size $O(dh^2)$, which allows us to take step size $h \lesssim 1/\sqrt{d}$. This explains why the use of the underdamped Langevin diffusion leads to improved dimension dependence for log-concave sampling.

In contrast, consider the reverse process, in which

$$D_{\mathrm{KL}}(\boldsymbol{Q}_T^{\leftarrow} \| \boldsymbol{P}_T^{\infty, \boldsymbol{q}_T}) = 2 \sum_{k=0}^{N-1} \mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}} \int_{kh}^{(k+1)h} \|\boldsymbol{s}_{T-kh}(X_{kh}, V_{kh}) - \nabla_v \ln \boldsymbol{q}_{T-t}(X_t, V_t)\|^2 \,\mathrm{d}t\,.$$

Since discretization of the reverse process involves the score function, which depends on both X and V, the error now involves controlling $\|V_t - V_{kh}\|^2$, which is of size $O(dh)$ (the process V is not very smooth because it includes a Brownian motion component). Therefore, from the form of the reverse process, we may expect that SGMs based on the CLD do not improve upon the dimension dependence of DDPM.

In Section C.5, we use this observation in order to prove a rigorous lower bound against discretization of SGMs based on the CLD.

C.3 PROOF OF THEOREM 5

Proof of Theorem 5. Similarly to the proof of Theorem 2 in Section B.3, Theorem 14 together with the data processing inequality yield

$$\mathsf{TV}\big(\mathrm{law}(X_{kh}^{\leftarrow,\infty}, V_{kh}^{\leftarrow,\infty} : k = 0, 1, \ldots, N-1), \,\mathrm{law}(\bar{X}_{kh}^{\leftarrow}, \bar{V}_{kh}^{\leftarrow} : k = 0, 1, \ldots, N-1)\big)$$
$$\lesssim \sqrt{D_{\mathrm{KL}}(\boldsymbol{q}_T \| \boldsymbol{\gamma}^{2d})} + (\varepsilon_{\mathrm{score},\infty} + L\sqrt{dh} + L\mathfrak{m}_2 h)\sqrt{T}\,.$$

In (Ma et al., 2021), following the entropic hypocoercivity approach of (Villani, 2009), Ma et al. consider a Lyapunov functional $\mathcal{L}$ which is equivalent to the sum of the KL divergence and the Fisher information,

$$\mathcal{L}(\boldsymbol{\mu} \| \boldsymbol{\gamma}^{2d}) \asymp D_{\mathrm{KL}}(\boldsymbol{\mu} \| \boldsymbol{\gamma}^{2d}) + \mathsf{FI}(\boldsymbol{\mu} \| \boldsymbol{\gamma}^{2d})\,,$$

which decays exponentially fast in time: there exists a universal constant $c > 0$ such that for all $t \geq 0$,

$$\mathcal{L}(\boldsymbol{q}_t \| \boldsymbol{\gamma}^{2d}) \leq \exp(-ct)\,\mathcal{L}(\boldsymbol{q}_0 \| \boldsymbol{\gamma}^{2d})\,.$$

Since $\boldsymbol{q}_0 = q \otimes \gamma^d$ and $\boldsymbol{\gamma}^{2d} = \gamma^d \otimes \gamma^d$, then $\mathcal{L}(\boldsymbol{q}_0 \| \boldsymbol{\gamma}^{2d}) \lesssim D_{\mathrm{KL}}(q \| \gamma^d) + \mathsf{FI}(q \| \gamma^d)$. Applying Lemma 7 and Pinsker's inequality, we deduce that

$$\mathsf{TV}(\boldsymbol{p}_T, \boldsymbol{q}_0) \lesssim \sqrt{D_{\mathrm{KL}}(q \| \gamma^d) + \mathsf{FI}(q \| \gamma^d)} \exp(-cT) + (\varepsilon_{\mathrm{score},\infty} + L\sqrt{dh} + L\mathfrak{m}_2 h)\sqrt{T} + \frac{N\varepsilon_{\mathrm{score}}^2}{\varepsilon_{\mathrm{score},\infty}^2}\,.$$

Theorem 5 follows from optimizing over the choice of $\varepsilon_{\mathrm{score},\infty}$. $\qquad\square$

C.4 AUXILIARY LEMMAS

We begin with the perturbation lemma for the score function.

Lemma 15 (score perturbation lemma)**.** *Let $0 < \zeta < 1$. Suppose that $\boldsymbol{M}_0, \boldsymbol{M}_1 \in \mathbb{R}^{2d \times 2d}$ are two matrices, where $\boldsymbol{M}_1$ is symmetric. Also, assume that $\|\boldsymbol{M}_0 - \boldsymbol{I}_{2d}\|_{\mathrm{op}} \leq \zeta$, so that $\boldsymbol{M}_0$ is invertible. Let $\boldsymbol{q} = \exp(-\boldsymbol{H})$ be a probability density on $\mathbb{R}^{2d}$ such that $\nabla \boldsymbol{H}$ is L-Lipschitz with $L \leq \frac{1}{4\|\boldsymbol{M}_1\|_{\mathrm{op}}}$. Then, it holds that*

$$\left\|\nabla \ln \frac{(\boldsymbol{M}_0)_\# \boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{\boldsymbol{q}}(\boldsymbol{\theta})\right\| \lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}} d} + L\zeta \|\boldsymbol{\theta}\| + (\zeta + L\|\boldsymbol{M}_1\|_{\mathrm{op}}) \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|\,.$$

Proof. The proof follows along the lines of (Lee et al., 2022a, Lemma C.12). First, we show that when $\boldsymbol{M}_0 = \boldsymbol{I}_{2d}$, if $L \leq \frac{1}{2\|\boldsymbol{M}_1\|_{\mathrm{op}}}$ then

$$\left\|\nabla \ln \frac{\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{\boldsymbol{q}}(\boldsymbol{\theta})\right\| \lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}} d} + L\|\boldsymbol{M}_1\|_{\mathrm{op}} \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|\,. \qquad (22)$$

Let $\mathcal{S}$ denote the subspace $\mathcal{S} := \operatorname{range} \boldsymbol{M}_1$. Then, since

$$\big(\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)\big)(\boldsymbol{\theta}) = \int_{\boldsymbol{\theta}+\mathcal{S}} \exp\big(-\frac{1}{2}\,\langle \boldsymbol{\theta} - \boldsymbol{\theta}', \boldsymbol{M}_1^{-1}\,(\boldsymbol{\theta} - \boldsymbol{\theta}')\rangle\big)\,\boldsymbol{q}(\mathrm{d}\boldsymbol{\theta}')\,,$$

where $\boldsymbol{M}_1^{-1}$ is well-defined on $\mathcal{S}$, we have

$$\Big\|\nabla \ln \frac{\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{\boldsymbol{q}}(\boldsymbol{\theta})\Big\| = \Big\|\frac{\int_{\boldsymbol{\theta}+\mathcal{S}} \nabla \boldsymbol{H}(\boldsymbol{\theta}') \exp(-\frac{1}{2}\,\langle \boldsymbol{\theta} - \boldsymbol{\theta}', \boldsymbol{M}_1^{-1}\,(\boldsymbol{\theta} - \boldsymbol{\theta}')\rangle)\,\boldsymbol{q}(\mathrm{d}\boldsymbol{\theta}')}{\int_{\boldsymbol{\theta}+\mathcal{S}} \exp(-\frac{1}{2}\,\langle \boldsymbol{\theta} - \boldsymbol{\theta}', \boldsymbol{M}_1^{-1}\,(\boldsymbol{\theta} - \boldsymbol{\theta}')\rangle)\,\boldsymbol{q}(\mathrm{d}\boldsymbol{\theta}')} - \nabla \boldsymbol{H}(\boldsymbol{\theta})\Big\|$$

$$= \|\mathbb{E}_{\boldsymbol{q_\theta}} \nabla \boldsymbol{H} - \nabla \boldsymbol{H}(\boldsymbol{\theta})\|\,.$$

Here, $\boldsymbol{q_\theta}$ is the measure on $\boldsymbol{\theta} + \mathcal{S}$ such that

$$\boldsymbol{q_\theta}(\mathrm{d}\boldsymbol{\theta}') \propto \exp\big(-\frac{1}{2}\,\langle \boldsymbol{\theta} - \boldsymbol{\theta}', \boldsymbol{M}_1^{-1}\,(\boldsymbol{\theta} - \boldsymbol{\theta}')\rangle\big)\,\boldsymbol{q}(\mathrm{d}\boldsymbol{\theta}')\,.$$

Note that since $L \le \frac{1}{2\,\|\boldsymbol{M}_1\|_{\mathrm{op}}}$, then if we write $\boldsymbol{q_\theta}(\boldsymbol{\theta}') \propto \exp(-\boldsymbol{H_\theta}(\boldsymbol{\theta}'))$, we have

$$\nabla^2 \boldsymbol{H_\theta} \succeq \big(\frac{1}{\|\boldsymbol{M}_1\|_{\mathrm{op}}} - L\big)\,I_d \succeq \frac{1}{2\,\|\boldsymbol{M}_1\|_{\mathrm{op}}}\,I_d \qquad \text{on } \boldsymbol{\theta} + \mathcal{S}\,.$$

Let $\boldsymbol{\theta}_\star \in \arg\min \boldsymbol{H_\theta}$ denote a mode. We bound

$$\|\mathbb{E}_{\boldsymbol{q_\theta}} \nabla \boldsymbol{H} - \nabla \boldsymbol{H}(\boldsymbol{\theta})\| \le L\,\mathbb{E}_{\boldsymbol{\theta}' \sim \boldsymbol{q_\theta}} \|\boldsymbol{\theta}' - \boldsymbol{\theta}\| \le L\,\mathbb{E}_{\boldsymbol{\theta}' \sim \boldsymbol{q_\theta}} \|\boldsymbol{\theta}' - \boldsymbol{\theta}_\star\| + L\,\|\boldsymbol{\theta}_\star - \boldsymbol{\theta}\|\,.$$

For the first term, (Dalalyan et al., 2019, Proposition 2) yields

$$\mathbb{E}_{\boldsymbol{\theta}' \sim \boldsymbol{q_\theta}} \|\boldsymbol{\theta}' - \boldsymbol{\theta}_\star\| \le \sqrt{2\,\|\boldsymbol{M}_1\|_{\mathrm{op}}\,d}\,.$$

For the second term, since the mode satisfies $\nabla \boldsymbol{H}(\boldsymbol{\theta}_\star) + \boldsymbol{M}_1^{-1}\,(\boldsymbol{\theta}_\star - \boldsymbol{\theta}) = 0$, we have

$$\|\boldsymbol{\theta}_\star - \boldsymbol{\theta}\| \le \|\boldsymbol{M}_1\|_{\mathrm{op}} \|\nabla \boldsymbol{H}(\boldsymbol{\theta}_\star)\| \le L\,\|\boldsymbol{M}_1\|_{\mathrm{op}} \|\boldsymbol{\theta}_\star - \boldsymbol{\theta}\| + \|\boldsymbol{M}_1\|_{\mathrm{op}} \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|$$

which is rearranged to yield

$$\|\boldsymbol{\theta}_\star - \boldsymbol{\theta}\| \le 2\,\|\boldsymbol{M}_1\|_{\mathrm{op}} \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|\,.$$

After combining the bounds, we obtain the claimed estimate (22).

Next, we consider the case of general $\boldsymbol{M}_0$. We have

$$\Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{\boldsymbol{q}}(\boldsymbol{\theta})\Big\| \le \Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{(\boldsymbol{M}_0)_{\#}\boldsymbol{q}}(\boldsymbol{\theta})\Big\| + \Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q}}{\boldsymbol{q}}(\boldsymbol{\theta})\Big\|\,.$$

We can apply (22) with $(\boldsymbol{M}_0)_{\#}\boldsymbol{q}$ in place of $\boldsymbol{q}$, noting that $(\boldsymbol{M}_0)_{\#}\boldsymbol{q} \propto \exp(-\boldsymbol{H}')$ for $\boldsymbol{H}' := \boldsymbol{H} \circ \boldsymbol{M}_0$ which is L'-smooth for $L' := L\,\|\boldsymbol{M}_0\|_{\mathrm{op}}^2 \lesssim L$, to get

$$\Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{(\boldsymbol{M}_0)_{\#}\boldsymbol{q}}(\boldsymbol{\theta})\Big\| \lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}}\,d} + L\,\|\boldsymbol{M}_1\|_{\mathrm{op}} \|\boldsymbol{M}_0 \nabla \boldsymbol{H}(\boldsymbol{M}_0\boldsymbol{\theta})\|$$

$$\lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}}\,d} + L\,\|\boldsymbol{M}_1\|_{\mathrm{op}} \|\nabla \boldsymbol{H}(\boldsymbol{M}_0\boldsymbol{\theta})\|\,.$$

Note that

$$\|\nabla \boldsymbol{H}(\boldsymbol{M}_0\boldsymbol{\theta})\| \le \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\| + L\,\|(\boldsymbol{M}_0 - \boldsymbol{I}_{2d})\,\boldsymbol{\theta}\| \lesssim \|\nabla \boldsymbol{H}(\boldsymbol{\theta})\| + L\zeta\,\|\boldsymbol{\theta}\|\,.$$

We also have

$$\Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q}}{\boldsymbol{q}}(\boldsymbol{\theta})\Big\| = \|\boldsymbol{M}_0 \nabla \boldsymbol{H}(\boldsymbol{M}_0\boldsymbol{\theta}) - \nabla \boldsymbol{H}(\boldsymbol{\theta})\|$$

$$\le \|\boldsymbol{M}_0 \nabla \boldsymbol{H}(\boldsymbol{M}_0\boldsymbol{\theta}) - \boldsymbol{M}_0 \nabla \boldsymbol{H}(\boldsymbol{\theta})\| + \|\boldsymbol{M}_0 \nabla \boldsymbol{H}(\boldsymbol{\theta}) - \nabla \boldsymbol{H}(\boldsymbol{\theta})\|$$

$$\lesssim L\,\|(\boldsymbol{M}_0 - \boldsymbol{I}_{2d})\,\boldsymbol{\theta}\| + \zeta\,\|\nabla \boldsymbol{H}(\boldsymbol{\theta})\| \lesssim L\zeta\,\|\boldsymbol{\theta}\| + \zeta\,\|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|\,.$$

Combining the bounds,

$$\Big\|\nabla \ln \frac{(\boldsymbol{M}_0)_{\#}\boldsymbol{q} * \mathsf{normal}(0, \boldsymbol{M}_1)}{\boldsymbol{q}}(\boldsymbol{\theta})\Big\|$$

$$\lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}}\,d} + L\zeta\,(1 + L\,\|\boldsymbol{M}_1\|_{\mathrm{op}})\,\|\boldsymbol{\theta}\| + (\zeta + L\,\|\boldsymbol{M}_1\|_{\mathrm{op}})\,\|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|$$

$$\lesssim L\sqrt{\|\boldsymbol{M}_1\|_{\mathrm{op}}\,d} + L\zeta\,\|\boldsymbol{\theta}\| + (\zeta + L\,\|\boldsymbol{M}_1\|_{\mathrm{op}})\,\|\nabla \boldsymbol{H}(\boldsymbol{\theta})\|$$

so the lemma follows. $\qquad\square$

Next, we prove the moment and movement bounds for the CLD.

Lemma 16 (moment bounds for CLD). *Suppose that Assumptions 2 and 4 hold. Let $(\bar{X}_t, \bar{V}_t)_{t\in[0,T]}$ denote the forward process* (6).

1. *(moment bound) For all $t \geq 0$,*
$$\mathbb{E}[\|(\bar{X}_t, \bar{V}_t)\|^2] \lesssim d + \mathfrak{m}_2^2.$$

2. *(score function bound) For all $t \geq 0$,*
$$\mathbb{E}[\|\nabla \ln \boldsymbol{q}_t(\bar{X}_t, \bar{V}_t)\|^2] \leq Ld.$$

Proof. 1. We can write
$$\mathbb{E}[\|(\bar{X}_t, \bar{V}_t)\|^2] = W_2^2(\boldsymbol{q}_t, \delta_{\mathbf{0}}) \lesssim W_2^2(\boldsymbol{q}_t, \boldsymbol{\gamma}^{2d}) + W_2^2(\boldsymbol{\gamma}^{2d}, \delta_{\mathbf{0}}) \lesssim d + W_2^2(\boldsymbol{q}_t, \boldsymbol{\gamma}^{2d}).$$
Next, the coupling argument of (Cheng et al., 2018) shows that the CLD converges exponentially fast in the Wasserstein metric associated to a twisted norm $\|\|\cdot\|\|$ which is equivalent (up to universal constants) to the Euclidean norm $\|\cdot\|$. It implies the following result, see, e.g., (Cheng et al., 2018, Lemma 8):
$$W_2^2(\boldsymbol{q}_t, \boldsymbol{\gamma}^{2d}) \lesssim W_2^2(\boldsymbol{q}, \boldsymbol{\gamma}^{2d}) \lesssim W_2^2(\boldsymbol{q}, \delta_{\mathbf{0}}) + W_2^2(\delta_{\mathbf{0}}, \boldsymbol{\gamma}^{2d}) \lesssim d + \mathfrak{m}_2^2.$$

2. The proof is the same as in Lemma 11.

$\square$

Lemma 17 (movement bound for CLD). *Suppose that Assumptions 2 holds. Let $(\bar{X}_t, \bar{V}_t)_{t\in[0,T]}$ denote the forward process* (6). *For $0 < s < t$ with $\delta := t - s$, if $\delta \leq 1$,*
$$\mathbb{E}[\|(\bar{X}_t, \bar{V}_t) - (\bar{X}_s, \bar{V}_s)\|^2] \lesssim \delta^2 \mathfrak{m}_2^2 + \delta d.$$

Proof. First,
$$\mathbb{E}[\|\bar{X}_t - \bar{X}_s\|^2] = \mathbb{E}\Big[\Big\|\int_s^t \bar{V}_r \, \mathrm{d}r\Big\|^2\Big] \leq \delta \int_s^t \mathbb{E}[\|\bar{V}_r\|^2] \, \mathrm{d}r \lesssim \delta^2 \, (d + \mathfrak{m}_2^2),$$
where we used the moment bound in Lemma 16. Next,
$$\mathbb{E}[\|\bar{V}_t - \bar{V}_s\|^2] = \mathbb{E}\Big[\Big\|\int_s^t (-\bar{X}_r - 2\bar{V}_r) \, \mathrm{d}r + 2 \, (B_t - B_s)\Big\|^2\Big] \lesssim \delta \int_s^t \mathbb{E}[\|\bar{X}_r\|^2 + \|\bar{V}_r\|^2] \, \mathrm{d}r + \delta d$$
$$\lesssim \delta^2 \, (d + \mathfrak{m}_2^2) + \delta d,$$
where we used Lemma 16 again. $\square$

C.5 LOWER BOUND AGAINST CLD

Proof of Theorem 6. Since $\boldsymbol{q}_0 = \gamma^d \otimes \gamma^d = \boldsymbol{\gamma}^{2d}$ is stationary for the forward process (6), we have $\boldsymbol{q}_t = \boldsymbol{\gamma}^{2d}$ for all $t \geq 0$. In this proof, since the score estimate is perfect and $\boldsymbol{q}_T = \boldsymbol{\gamma}^{2d}$, we simply denote the path measure for the algorithm as $\boldsymbol{P}_T = \boldsymbol{P}_T^{\infty, \boldsymbol{q}_T}$. From Girsanov's theorem in the form of Corollary 13 and from $\boldsymbol{s}_{T-kh}(x, v) = \nabla_v \ln \boldsymbol{q}_{T-kh}(x, v) = -v$, we have
$$D_{\mathrm{KL}}(\boldsymbol{Q}_T^{\leftarrow} \| \boldsymbol{P}_T) = 2 \sum_{k=0}^{N-1} \mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}} \int_{kh}^{(k+1)h} \|V_{kh} - V_t\|^2 \, \mathrm{d}t. \tag{23}$$
To lower bound this quantity, we use the inequality $\|x + y\|^2 \geq \frac{1}{2} \|x\|^2 - \|y\|^2$ to write, for $t \in [kh, (k+1)h]$
$$\mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}}[\|V_{kh} - V_t\|^2] = \mathbb{E}[\|\bar{V}_{T-kh} - \bar{V}_{T-t}\|^2] = \mathbb{E}\Big[\Big\|\int_{T-t}^{T-kh} \{-\bar{X}_s - 2\bar{V}_s\} \, \mathrm{d}s + 2 \, (B_{T-kh} - B_{T-t})\Big\|^2\Big]$$
$$\geq 2\mathbb{E}[\|B_{T-kh} - B_{T-t}\|^2] - \mathbb{E}\Big[\Big\|\int_{T-t}^{T-kh} \{-\bar{X}_s - 2\bar{V}_s\} \, \mathrm{d}s\Big\|^2\Big]$$
$$\geq 2d \, (t - kh) - (t - kh) \int_{T-t}^{T-kh} \mathbb{E}[\|\bar{X}_s + 2\bar{V}_s\|^2] \, \mathrm{d}s$$
$$\geq 2d \, (t - kh) - (t - kh) \int_{T-t}^{T-kh} \mathbb{E}[2 \|\bar{X}_s\|^2 + 8 \|\bar{V}_s\|^2] \, \mathrm{d}s.$$

Using the fact that $\bar{X}_s \sim \gamma^d$ and $\bar{V}_s \sim \gamma^d$ for all $s \in [0, T]$, we can then bound

$$\mathbb{E}_{\boldsymbol{Q}_T^{\leftarrow}}[\|V_{kh} - V_t\|^2] \geq 2d\,(t - kh) - 10d\,(t - kh)^2 \geq d\,(t - kh)\,,$$

provided that $h \leq \frac{1}{10}$. Substituting this into (23),

$$D_{\mathrm{KL}}(\boldsymbol{Q}_T^{\leftarrow} \,\|\, \boldsymbol{P}_T) \geq 2d \sum_{k=0}^{N-1} \int_{kh}^{(k+1)h} (t - kh)^2 \,\mathrm{d}t = dh^2 N = dhT\,.$$

This proves the result. $\qquad\square$

This lower bound shows that the Girsanov discretization argument of Theorem 14 is essentially tight (except possibly the dependence on L).

D   DERIVATION OF THE SCORE MATCHING OBJECTIVE

In this section, we present a self-contained derivation of the score matching objective (14) for the reader's convenience. See also (Hyvärinen, 2005; Vincent, 2011; Song & Ermon, 2019).

Recall that the problem is to solve

$$\underset{s_t \in \mathscr{F}}{\text{minimize}} \quad \mathbb{E}_{q_t}[\|s_t - \nabla \ln q_t\|^2]\,.$$

This objective cannot be evaluated, even if we replace the expectation over q_t with an empirical average over samples from q_t. The trick is to use an integration by parts identity to reformulate the objective. Here, C will denote any constant that does not depend on the optimization variable s_t. Expanding the square,

$$\mathbb{E}_{q_t}[\|s_t - \nabla \ln q_t\|^2] = \mathbb{E}_{q_t}[\|s_t\|^2 - 2\,\langle s_t, \nabla \ln q_t\rangle] + C\,.$$

We can rewrite the second term using integration by parts:

$$\int \langle s_t, \nabla \ln q_t\rangle \,\mathrm{d}q_t = \int \langle s_t, \nabla q_t\rangle = -\int (\mathrm{div}\, s_t) \,\mathrm{d}q_t$$
$$= -\iint (\mathrm{div}\, s_t)\big(\exp(-t)\,x_0 + \sqrt{1 - \exp(-2t)}\,z_t\big) \,\mathrm{d}q(x_0)\,\mathrm{d}\gamma^d(z_t)\,,$$

where $\gamma^d = \mathrm{normal}(0, I_d)$ and we used the explicit form of the law of the OU process at time t. Recall the Gaussian integration by parts identity: for any vector field $v : \mathbb{R}^d \to \mathbb{R}^d$,

$$\int (\mathrm{div}\, v) \,\mathrm{d}\gamma^d = \int \langle x, v(x)\rangle \,\mathrm{d}\gamma^d(x)\,.$$

Applying this identity,

$$\int \langle s_t, \nabla \ln q_t\rangle \,\mathrm{d}q_t = -\frac{1}{\sqrt{1 - \exp(-2t)}} \int \langle z_t, s_t(x_t)\rangle \,\mathrm{d}q(x_0)\,\mathrm{d}\gamma^d(z_t)$$

where $x_t = \exp(-t)\,x_0 + \sqrt{1 - \exp(-2t)}\,z_t$. Substituting this in,

$$\mathbb{E}_{q_t}[\|s_t - \nabla \ln q_t\|^2] = \mathbb{E}\Big[\|s_t(X_t)\|^2 + \frac{2}{\sqrt{1 - \exp(-2t)}}\,\langle Z_t, s_t(X_t)\rangle\Big] + C$$
$$= \mathbb{E}\Big[\Big\|s(X_t) + \frac{1}{\sqrt{1 - \exp(-2t)}}\,Z_t\Big\|^2\Big] + C\,,$$

where $X_0 \sim q$ and $Z_t \sim \gamma^d$ are independent, and $X_t := \exp(-t)\,X_0 + \sqrt{1 - \exp(-2t)}\,Z_t$.

E   DEFERRED PROOFS

Lemma 18. *Suppose that* supp $q \subseteq \mathsf{B}(0, R)$ *where* $R \geq 1$, *and let* q_t *denote the law of the OU process at time t, started at q. Let* $\varepsilon > 0$ *be such that* $\varepsilon \ll \sqrt{d}$ *and set* $t \asymp \varepsilon^2/(\sqrt{d}\,(R \vee \sqrt{d}))$. *Then,*

1. $W_2(q_t, q) \leq \varepsilon$.

2. q_t *satisfies*

$$D_{\mathrm{KL}}(q_t \parallel \gamma^d) \lesssim \frac{\sqrt{d}\,(R \vee \sqrt{d})^3}{\varepsilon^2}\,.$$

3. *For every* $t' \geq t$, $q_{t'}$ *satisfies Assumption 1 with*

$$L \lesssim \frac{dR^2\,(R \vee \sqrt{d})^2}{\varepsilon^4}\,.$$

Proof.   1. For the OU process (1), we have $\bar{X}_t := \exp(-t)\,\bar{X}_0 + \sqrt{1 - \exp(-2t)}\,Z$, where $Z \sim \mathsf{normal}(0, I_d)$ is independent of $\bar{X}_0$. Hence, for $t \lesssim 1$,

$$\begin{aligned} W_2^2(q, q_t) &\leq \mathbb{E}\big[\big\|\big(1 - \exp(-t)\big)\,\bar{X}_0 + \sqrt{1 - \exp(-2t)}\,Z\big\|^2\big] \\ &= \big(1 - \exp(-t)\big)^2 \mathbb{E}[\|\bar{X}_0\|^2] + \big(1 - \exp(-2t)\big)\,d \lesssim R^2 t^2 + dt\,. \end{aligned}$$

We now take $t \lesssim \min\{\varepsilon/R, \varepsilon^2/d\}$ to ensure that $W_2^2(q, q_t) \leq \varepsilon^2$. Since $\varepsilon \ll \sqrt{d}$, it suffices to take $t \asymp \varepsilon^2/(\sqrt{d}\,(R \vee \sqrt{d}))$.

2. For this, we use the short-time regularization result in (Otto & Villani, 2001, Corollary 2), which implies that

$$D_{\mathrm{KL}}(q_t \parallel \gamma^d) \leq \frac{W_2^2(q, \gamma^d)}{4t} \lesssim \frac{W_2^2(q, \delta_0) + W_2^2(\gamma^d, \delta_0)}{t} \lesssim \frac{\sqrt{d}\,(R \vee \sqrt{d})^3}{\varepsilon^2}\,.$$

3. Using (Mikulincer & Shenfeld, 2022, Lemma 4), along the OU process,

$$\frac{1}{1 - \exp(-2t)}\,I_d - \frac{\exp(-2t)\,R^2}{(1 - \exp(-2t))^2}\,I_d \preccurlyeq -\nabla^2 \ln q_t(x) \preccurlyeq \frac{1}{1 - \exp(-2t)}\,I_d\,.$$

With our choice of t, it implies

$$\|\nabla^2 \ln q_{t'}\|_{\mathrm{op}} \lesssim \frac{1}{1 - \exp(-2t')} \vee \frac{\exp(-2t')\,R^2}{(1 - \exp(-2t'))^2} \lesssim \frac{1}{t} \vee \frac{R^2}{t^2} \lesssim \frac{dR^2\,(R \vee \sqrt{d})^2}{\varepsilon^4}\,.$$

$\square$

F   NOVIKOV'S CONDITION FOR DDPM

In Section B.2, we proved the Girsanov discretization bound for DDPM (Theorem 10) under the assumption that the path measures P_T^{∞,q_T} and $Q_T^{\leftarrow}$ satisfy Novikov's condition (16). Unfortunately, Novikov's condition fails under Assumptions 1, 2, and 3 alone. In this section, we remedy this by instead applying the Girsanov discretization argument to a sequence of pairs of auxiliary processes defined in place of P_T^{∞,q_T} and $Q_T^{\leftarrow}$.

F.1   TRUNCATION ARGUMENT

We first construct a smooth cutoff function for truncating the drift terms in the processes.

Lemma 19. *For any* $R > 0$, *there is a smooth function* $\phi_R : \mathbb{R}^d \to [0, 1]$ *satisfying:*

1. $\phi_R(x) = 1$ *for all* $\|x\| \leq R$,

2. $\phi_R(x) = 0$ *for all* $\|x\| \geq 2R$,

3. ϕ_R *is* $O(1/R)$*-Lipschitz.*

Proof. Let $\psi : \mathbb{R} \to [0,1]$ be an $O(1)$-Lipschitz and smooth function for which $\psi(z) = 1$ for all $|z| \leq 1$, $\psi(z) = 0$ for all $|z| \geq 2$. This can be constructed, for instance, by taking the piecewise linear function

$$\widetilde{\psi}(z) \triangleq \begin{cases} 1, & \text{if } |z| \leq 1 + c, \\ 0, & \text{if } |z| \geq 2 - c, \\ 1 - \frac{1}{1-2c} \,\big|\, |z| - (1+c) \,\big|, & \text{otherwise}, \end{cases}$$

for any small constant $c \in (0, \frac{1}{2})$ and convolving $\widetilde{\psi}$ with $z \mapsto c^{-1}\phi(z/c)$ for any mollifier ϕ supported on $(-1, 1)$. Now define $\phi_R(x) \triangleq \psi(\|x\|/R)$. Parts 1 and 2 of the lemma are immediate. For Lipschitzness, we can calculate for $x \neq 0$

$$\nabla \phi_R(x) = \frac{1}{R} \, \psi'\big(\frac{\|x\|}{R}\big) \, \frac{x}{\|x\|} \,,$$

which has norm at most $|\psi'(\|x\|/R)|/R = O(1/R)$ as desired, whereas $\nabla \phi_R(0) = 0$. $\qquad \square$

Lemma 20. *Suppose that* $v : \mathbb{R}^d \to \mathbb{R}^d$ *is an* L*-Lipschitz vector field. Then,* $x \mapsto \phi_R(x) \, v(x)$ *is* L'*-Lipschitz for* $L' \lesssim L + \frac{1}{R} \|v(0)\|$.

Proof. Let $x, y \in \mathbb{R}^d$. If $\|y\| \leq 2R$, then

$$\|\phi_R(x) \, v(x) - \phi_R(y) \, v(y)\| \leq \phi_R(x) \, \|v(x) - v(y)\| + \|v(y)\| \, |\phi_R(x) - \phi_R(y)|$$

$$\leq L \, \|x - y\| + O\big(\frac{1}{R}\big) \, (\|v(0)\| + 2LR) \, \|x - y\|$$

$$\leq O\big(L + \frac{1}{R} \|v(0)\|\big) \, \|x - y\| \,.$$

If $\|y\| > 2R$, then because $\phi_R(y) = 0$,

$$\|\phi_R(x) \, v(x) - \phi_R(y) \, v(y)\| = \phi_R(x) \, \|v(x)\| \leq (\|v(0)\| + 2LR) \, |\phi_R(x) - \phi_R(y)|$$

$$\leq O\big(L + \frac{1}{R} \|v(0)\|\big) \, \|x - y\| \,,$$

as claimed. $\qquad \square$

Now define the following modified processes: for $t \in [kh, (k+1)h]$,

$$\mathrm{d}\bar{X}_t^{\mathsf{tr},\leftarrow} = \phi_R(\bar{X}_t^{\mathsf{tr},\leftarrow}) \, \{\bar{X}_t^{\mathsf{tr},\leftarrow} + 2 \, \nabla \ln q_{T-t}(\bar{X}_t^{\mathsf{tr},\leftarrow})\} \, \mathrm{d}t + \sqrt{2} \, \mathrm{d}B_t \,, \tag{24}$$

$$\mathrm{d}X_t^{\mathsf{tr},\leftarrow,\infty,q_T} = \phi_R(X_t^{\mathsf{tr},\leftarrow,\infty,q_T}) \, \{X_t^{\mathsf{tr},\leftarrow,\infty,q_T} + 2 \, s_{T-kh}^\infty(X_{kh}^{\mathsf{tr},\leftarrow,\infty,q_T})\} \, \mathrm{d}t + \sqrt{2} \, \mathrm{d}B_t \,, \tag{25}$$

where $\bar{X}_0^{\mathsf{tr},\leftarrow} = X_0^{\mathsf{tr},\leftarrow,\infty,q_T}$ is obtained by sampling $\bar{X}_T \sim q_T$ and setting $\bar{X}_0^{\mathsf{tr},\leftarrow} = \bar{X}_T$ if $\|\bar{X}_T\| \leq R$ and setting $\bar{X}_0^{\mathsf{tr},\leftarrow} = 0$ otherwise. For clarity of presentation, we suppress the dependence on R in the notation.

Let $q_t^{\mathsf{tr},\leftarrow} := \mathrm{law}(\bar{X}_t^{\mathsf{tr},\leftarrow})$ and $p_t^{\mathsf{tr},\infty,q_T} := \mathrm{law}(X_t^{\mathsf{tr},\leftarrow,\infty,q_T})$. Likewise, denote the corresponding measures over path space by $Q_T^{\mathsf{tr},\leftarrow}$ and $P_T^{\mathsf{tr},\infty,q_T}$. As it is convenient to think about a single sochastic process, for ease of notation we denote this by $(X_t^{\mathsf{tr}})_{t \in [0,T]}$ so that under $Q_T^{\mathsf{tr},\leftarrow}$ (resp. $P_T^{\mathsf{tr},\infty,q_T}$), $(X_t^{\mathsf{tr}})_{t \in [0,T]}$ has the law of the modified reverse process (24) (resp. the modified SGM process (25)).

We first verify that these modified processes satisfy Novikov's condition.

Lemma 21. *It holds that*

$$\mathbb{E}_{Q_T^{\mathsf{tr},\leftarrow}} \exp\bigg(\sum_{k=0}^{N-1} \int_{kh}^{(k+1)h} \|\phi_R(X_{kh}^{\mathsf{tr}}) \, s_{T-kh}^\infty(X_{kh}^{\mathsf{tr}}) - \phi_R(X_t^{\mathsf{tr}}) \, \nabla \ln q_{T-t}(X_t^{\mathsf{tr}})\|^2 \, \mathrm{d}t\bigg) < \infty \,.$$

Proof. For any x,

$$\|\phi_R(x)\, s^\infty_{T-kh}(x)\| \le \sup_{x^*\in\mathsf{B}(0,R),\ k^*\in\{0,\dots,N-1\}} \|s^\infty_{T-k^*h}(x^*)\| =: A < \infty\,,$$

$$\|\phi_R(x)\,\nabla\ln q_{T-t}(x)\| \le \sup_{x^*\in\mathsf{B}(0,R),\ t^*\in[0,T]} \|\nabla\ln q_{T-t^*}(x^*)\| =: B < \infty\,.$$

So the quantity in the lemma is at most $\exp(2T\,(A^2+B^2)) < \infty$ as claimed. $\qquad\square$

We can thus apply Girsanov's theorem (Theorem 8), with the choices $P_T = P^{\mathsf{tr},\infty,q_T}_T$, $Q_T = Q^{\mathsf{tr},\leftarrow}_T$, $\tilde b_t = \phi_R(X^{\mathsf{tr}}_t)\,\{X^{\mathsf{tr}}_t + 2\,s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh})\}$, $b_t = \phi_R(X^{\mathsf{tr}}_t)\,\{X^{\mathsf{tr}}_t + 2\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\}$, and $\sigma_t = \sqrt{2}\,I_d$. It yields the following corollary.

Corollary 22. *It holds that*

$$D_{\mathrm{KL}}(Q^{\mathsf{tr},\leftarrow}_T \parallel P^{\mathsf{tr},\infty,q_T}_T) = \mathbb{E}_{Q^{\mathsf{tr},\leftarrow}_T}\ln\frac{\mathrm{d}Q^{\mathsf{tr},\leftarrow}_T}{\mathrm{d}P^{\mathsf{tr},\infty,q_T}_T}$$

$$= \sum_{k=0}^{N-1}\mathbb{E}_{Q^{\mathsf{tr},\leftarrow}_T}\int_{kh}^{(k+1)h}\|\phi_R(X^{\mathsf{tr}}_{kh})\,s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh}) - \phi_R(X^{\mathsf{tr}}_t)\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2\,\mathrm{d}t\,.$$

Our analysis for the quantity on the right-hand side of Corollary 22 proceeds along similar lines to the argument in Section B.2, though the moment and movement bounds require some additional work as we must work with $\bar X^{\mathsf{tr},\leftarrow}_t$ instead of $\bar X^{\leftarrow}_t$, the former of which is not the reverse of the OU process.

Theorem 23 (discretization error for modified process). *Suppose that Assumptions 1, 2, and 3 hold. For $R>0$, let $Q^{\mathsf{tr},\leftarrow}_T$ and $P^{\mathsf{tr},\infty,q_T}_T$ denote the measures on path space corresponding to the modified processes* (24) *and* (25) *respectively. Assume that $L\ge 1$ and $h\lesssim 1/L$. Then,*

$$\limsup_{R\to\infty} D_{\mathrm{KL}}(Q^{\mathsf{tr},\leftarrow}_T \parallel P^{\mathsf{tr},\infty,q_T}_T) \lesssim (\varepsilon^2_{\mathrm{score},\infty} + L^2 dh + L^2\mathfrak{m}^2_2 h^2)\,T\,.$$

Proof. First, as before, we can decompose

$$\|\phi_R(X^{\mathsf{tr}}_{kh})\,s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh}) - \phi_R(X^{\mathsf{tr}}_t)\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2$$

$$\lesssim \|\phi_R(X^{\mathsf{tr}}_{kh})\,\{s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh}) - \nabla\ln q_{T-kh}(X^{\mathsf{tr}}_{kh})\}\|^2$$

$$+ \|\phi_R(X^{\mathsf{tr}}_{kh})\,\{\nabla\ln q_{T-kh}(X^{\mathsf{tr}}_{kh}) - \nabla\ln q_{T-t}(X^{\mathsf{tr}}_{kh})\}\|^2$$

$$+ \|\phi_R(X^{\mathsf{tr}}_{kh})\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_{kh}) - \phi_R(X^{\mathsf{tr}}_t)\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2$$

$$\lesssim \varepsilon^2_{\mathrm{score},\infty} + \left\|\phi_R(X^{\mathsf{tr}}_{kh})\,\nabla\ln\frac{q_{T-kh}}{q_{T-t}}(X^{\mathsf{tr}}_{kh})\right\|^2 + L'^2\,\|X^{\mathsf{tr}}_{kh} - X^{\mathsf{tr}}_t\|^2\,,$$

where in the last step we used the fact that $\phi_R(X^{\mathsf{tr}}_{kh})\in[0,1]$ together with Lemma 20 applied to the vector field $v = \nabla\ln q_{T-t}$ (L' is the quantity defined therein).

Because (18) and (19) hold pointwise, we can bound the second term above by

$$\left\|\phi_R(X^{\mathsf{tr}}_{kh})\,\nabla\ln\frac{q_{T-kh}}{q_{T-t}}(X^{\mathsf{tr}}_{kh})\right\|^2 \lesssim \phi_R(X^{\mathsf{tr}}_{kh})^2\,(L^2 dh + L^2 h^2\,\|X^{\mathsf{tr}}_{kh}\|^2 + L^2 h^2\,\|\nabla\ln q_{T-t}(X^{\mathsf{tr}}_{kh})\|^2)$$

$$\lesssim L^2 dh + \phi_R(X^{\mathsf{tr}}_{kh})^2\,L^2 h^2\,\|X^{\mathsf{tr}}_{kh}\|^2 + \phi_R(X^{\mathsf{tr}}_t)^2\,L^2 h^2\,\|\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2 + L'^2\,\|X^{\mathsf{tr}}_{kh} - X^{\mathsf{tr}}_t\|^2\,,$$

where in the last line we applied Lemma 20 again. Hence,

$$\|\phi_R(X^{\mathsf{tr}}_{kh})\,s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh}) - \phi_R(X^{\mathsf{tr}}_t)\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2$$

$$\lesssim \varepsilon^2_{\mathrm{score},\infty} + L^2 dh + \phi_R(X^{\mathsf{tr}}_{kh})^2\,L^2 h^2\,\|X^{\mathsf{tr}}_{kh}\|^2 + \phi_R(X^{\mathsf{tr}}_t)^2\,L^2 h^2\,\|\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2 + L'^2\,\|X^{\mathsf{tr}}_{kh} - X^{\mathsf{tr}}_t\|^2\,.$$

By taking $R\to\infty$ above, noting that $\limsup_{R\to\infty} L' \lesssim L$, and applying Lemmas 25, 26, and 27, we get

$$\limsup_{R\to\infty}\mathbb{E}_{Q^{\mathsf{tr},\leftarrow}_T}[\|\phi_R(X^{\mathsf{tr}}_{kh})\,s^\infty_{T-kh}(X^{\mathsf{tr}}_{kh}) - \phi_R(X^{\mathsf{tr}}_t)\,\nabla\ln q_{T-t}(X^{\mathsf{tr}}_t)\|^2]$$

$$\lesssim \varepsilon^2_{\mathrm{score},\infty} + L^2 dh + L^2 h^2\mathbb{E}_{Q^{\leftarrow}_T}[\|X_t\|^2] + L^2 h^2\mathbb{E}_{Q^{\leftarrow}_T}[\|\nabla\ln q_{T-t}(X_t)\|^2] + L^2\,(\mathfrak{m}^2_2 h^2 + dh)\,.$$

We can bound $\mathbb{E}_{Q_T^\leftarrow}[\|X_t\|^2] \leq d \vee \mathfrak{m}_2^2$ and $\mathbb{E}_{Q_T^\leftarrow}[\|\nabla \ln q_{T-t}(X_t)\|^2] \leq Ld$ via the moment bounds of Lemma 11.

Putting everything together, we get

$$\limsup_{R\to\infty} \mathbb{E}[\|\phi_R(X_{kh}^{\mathsf{tr}})\, s_{T-kh}^\infty(X_{kh}^{\mathsf{tr}}) - \phi_R(X_t^{\mathsf{tr}})\, \nabla \ln q_{T-t}(X_t^{\mathsf{tr}})\|^2] \lesssim \varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 \mathfrak{m}_2^2 h^2\,.$$

The result now follows from Corollary 22. $\qquad\square$

We are now ready to bound the discretization error for DDPM in total variation.

Full proof of Theorem 10. Consider a coupling of the laws of $(X_t)_{t\in[0,T]}$ and $(X_t^{\mathsf{tr}})_{t\in[0,T]}$ under $Q_T^\leftarrow$ (resp. P_T^{∞,q_T}) and $Q_T^{\mathsf{tr},\leftarrow}$ (resp. $P_T^{\mathsf{tr},\infty,q_T}$) such that if $X_t \in \mathsf{B}(0,R)$ for all $t \in [0,T]$, then $X_t^{\mathsf{tr}} = X_t$ for $t \in [0,T]$. Denote this event by $\mathcal{E}^R$ (resp. $\mathcal{E}'^R$). Then

$$\mathsf{TV}(Q_T^{\mathsf{tr},\leftarrow}, Q_T^\leftarrow) \leq \mathbb{P}[\mathcal{E}^R] \qquad \text{and} \qquad \mathsf{TV}(P_T^{\mathsf{tr},\infty,q_T}, P_T^{\infty,q_T}) \leq \mathbb{P}[\mathcal{E}'^R]\,.$$

Clearly $\lim_{R\to\infty} \mathbb{P}[\mathcal{E}^R] = 0$ and $\lim_{R\to\infty} \mathbb{P}[\mathcal{E}'^R] = 0$. By Theorem 23 and Pinsker's inequality,

$$\limsup_{R\to\infty} \mathsf{TV}(Q_T^{\mathsf{tr},\leftarrow}, P_T^{\mathsf{tr},\infty,q_T})^2 \lesssim (\varepsilon_{\mathrm{score},\infty}^2 + L^2 dh + L^2 \mathfrak{m}_2^2 h^2)\,T\,.$$

Hence, by the triangle inequality,

$$\mathsf{TV}(Q_T^\leftarrow, P_T^{\infty,q_T}) \leq \limsup_{R\to\infty} \{\mathsf{TV}(Q_T^{\mathsf{tr},\leftarrow}, Q_T^\leftarrow) + \mathsf{TV}(P_T^{\mathsf{tr},\infty,q_T}, P_T^{\infty,q_T}) + \mathsf{TV}(Q_T^{\mathsf{tr},\leftarrow}, P_T^{\mathsf{tr},\infty,q_T})\}$$
$$\lesssim (\varepsilon_{\mathrm{score},\infty} + L\sqrt{dh} + L\mathfrak{m}_2 h)\,\sqrt{T}\,.$$

This is the claimed result. $\qquad\square$

F.2 Auxiliary lemmas

In this section, we prove some auxiliary lemmas which were used in the proof of Theorem 23 above.

Lemma 24 (escape probability for reverse process). *Under Assumption 2,*

$$\mathbb{P}\{\bar{X}_t^\leftarrow \in \mathsf{B}(0,R)\ \text{for all } t \in [0,T]\} \geq 1 - o\big(\frac{1}{R^2}\big) \qquad \text{as } R \to \infty\,.$$

Proof. As $\bar{X}^\leftarrow$ is the reverse of the OU process, we can equivalently bound $\mathbb{P}\{\bar{X}_t \in \mathsf{B}(0,R)$ for all $t \in [0,T]\}$. From the explicit solution of the OU process, $\bar{X}_t = \exp(-t)\,\bar{X}_0 + \sqrt{2} \int_0^t \exp(-(t-s))\,\mathrm{d}B_s$. Let $\tilde{T} := \int_0^t \exp(-2(t-s))\,\mathrm{d}s$. By the representation of stochastic integrals as time changes of Brownian motion (see, e.g., (Steele, 2001, Theorem 12.4)), it follows that $\sup_{t\in[0,T]} \|\int_0^t \exp(-(t-s))\,\mathrm{d}B_s\| = \sup_{t\in[0,\tilde{T}]} \|\tilde{B}_t\|$, where $\tilde{B}$ is another standard Brownian motion. Then, we can bound

$$\sup_{t\in[0,T]} \|\bar{X}_t\| \leq \|\bar{X}_0\| + \sqrt{2} \sup_{t\in[0,\tilde{T}]} \|\tilde{B}_t\|\,.$$

By Assumption 2, we have $\mathbb{P}\{\|\bar{X}_0\| > R/2\} \leq \mathbb{E}[\|\bar{X}_0\|^{2+\eta}]/(R/2)^{2+\eta} = o(1/R^2)$. On the other hand, by (Chewi et al., 2021b, Lemma 23) applied to $\lambda = 1/(4\tilde{T})$,

$$\mathbb{P}\{\sup_{t\in[0,\tilde{T}]} \|\tilde{B}_t\| > \frac{R}{2\sqrt{2}}\} \leq 3^d \exp\big(-\frac{R^2}{32\tilde{T}}\big) = o\big(\frac{1}{R^2}\big)\,.$$

The lemma follows from a union bound over the two events. $\qquad\square$

Lemma 25 (effect of truncation on the moment bound). *Under Assumption 2, we have*

$$\limsup_{R\to\infty} \mathbb{E}_{Q_T^{\mathsf{tr},\leftarrow}}[\phi_R(X_t^{\mathsf{tr}})^2 \|X_t^{\mathsf{tr}}\|^2] \leq \mathbb{E}_{Q_T^\leftarrow}[\|X_t\|^2]\,.$$

Proof. Consider a coupling of the laws of $(X_t)_{t\in[0,T]}$ and $(X_t^{\text{tr}})_{t\in[0,T]}$ such that if $X_{t'} \in \mathsf{B}(0,R)$ for all $0 \le t' < t$, then $X_t^{\text{tr}} = X_t$. Denote the complement of this event by $\mathcal{E}_t$. Then we have

$$\mathbb{E}[\phi_R(X_t^{\text{tr}})^2 \|X_t^{\text{tr}}\|^2] \le \mathbb{E}[\|X_t\|^2] + \mathbb{E}[\phi_R(X_t^{\text{tr}})^2 \|X_t^{\text{tr}}\|^2 \mathbb{1}_{\mathcal{E}_t}].$$

Because $\phi_R(X_t^{\text{tr}}) = 0$ if $\|X_t^{\text{tr}}\| \ge 2R$, we can bound the latter term via Hölder's inequality by

$$(2R)^2 \cdot \mathbb{P}[\mathcal{E}_t] = o(1),$$

from Lemma 24. As this tends to 0 as $R \to \infty$, the lemma follows. $\qquad\square$

Lemma 26 (effect of truncation on the score function bound). *Under Assumptions 1 and 2,*

$$\limsup_{R\to\infty} \mathbb{E}_{Q_T^{\text{tr},\leftarrow}}[\phi_R(X_t^{\text{tr}})^2 \|\nabla \ln q_{T-t}(X_t^{\text{tr}})\|^2] \le \mathbb{E}_{Q_T^{\leftarrow}}[\|\nabla \ln q_{T-t}(X_t)\|^2]$$

for any $t \in [0,T]$.

Proof. Consider the same coupling of the laws of $(X_t)_{t\in[0,T]}$ and $(X_t^{\text{tr}})_{t\in[0,T]}$ from the proof of Lemma 25, and let $\mathcal{E}_t$ again denote the event that $X_{t'} \in \mathsf{B}(0,R)$ for all $0 \le t' < t$. Then we have

$$\mathbb{E}[\phi_R(X_t^{\text{tr}})^2 \|\nabla \ln q_{T-t}(X_t^{\text{tr}})\|^2] \le \mathbb{E}[\|\nabla \ln q_{T-t}(X_t)\|^2] + \mathbb{E}[\phi_R(X_t^{\text{tr}})^2 \|\nabla \ln q_{T-t}(X_t^{\text{tr}})\|^2 \mathbb{1}_{\mathcal{E}_t}].$$

Because $\phi_R(X_t^{\text{tr}}) = 0$ if $\|X_t^{\text{tr}}\| \ge 2R$, we can bound the latter term via Hölder's inequality by

$$\sup_{x\in\mathsf{B}(0,2R)} \|\nabla \ln q_{T-t}(x)\|^2 \cdot \mathbb{P}[\mathcal{E}_t].$$

By Assumption 1, if $\|x\| \le 2R$, then $\|\nabla \ln q_{T-t}(x)\|^2 \lesssim \|\nabla \ln q_{T-t}(0)\|^2 + L^2R^2$, so the above quantity is bounded via Lemma 24 by $o(1)$ as $R \to \infty$. $\qquad\square$

Lemma 27 (movement bound for modified process). *Suppose that Assumptions 1 and 2 hold. Then for $0 \le s < t$ satisfying $t - s \le h$, if $h \lesssim 1/L$, then*

$$\limsup_{R\to\infty} \mathbb{E}_{Q_T^{\text{tr},\leftarrow}}[\|\bar{X}_t^{\text{tr},\leftarrow} - \bar{X}_s^{\text{tr},\leftarrow}\|^2] \lesssim \mathfrak{m}_2^2 h^2 + dh.$$

Proof. In the proof, we drop the subscript $Q_T^{\text{tr},\leftarrow}$. We can write

$$\mathbb{E}[\|\bar{X}_t^{\text{tr},\leftarrow} - \bar{X}_s^{\text{tr},\leftarrow}\|^2] = \mathbb{E}\left[\left\|\int_s^t \phi_R(\bar{X}_r^{\text{tr},\leftarrow}) \{\bar{X}_r^{\text{tr},\leftarrow} + 2\nabla \ln q_{T-r}(\bar{X}_r^{\text{tr},\leftarrow})\} \, dr + \sqrt{2}\,(B_t - B_s)\right\|^2\right]$$

$$\lesssim h \int_s^t \mathbb{E}[\phi_R(\bar{X}_r^{\text{tr},\leftarrow})^2 \|\bar{X}_r^{\text{tr},\leftarrow}\|^2] \, dr$$

$$+ h \int_s^t \mathbb{E}[\phi_R(\bar{X}_r^{\text{tr},\leftarrow})^2 \|\nabla \ln q_{T-r}(\bar{X}_r^{\text{tr},\leftarrow})\|^2] \, dr + dh.$$

Taking the $\limsup$ on both sides as $R \to \infty$, we can apply Lemmas 25 and 26 to get

$$\limsup_{R\to\infty} \mathbb{E}[\|\bar{X}_t^{\text{tr},\leftarrow} - \bar{X}_s^{\text{tr},\leftarrow}\|^2] \lesssim h \int_s^t \mathbb{E}[\|\bar{X}_r^{\leftarrow}\|^2] \, dr + h \int_s^t \mathbb{E}[\|\nabla \ln q_{T-r}(\bar{X}_r^{\leftarrow})\|^2] \, dr + dh.$$

The claimed bound then follows from the moment and score function bounds of Lemma 11. $\qquad\square$