SUPPLEMENTARY MATERIAL FOR "IMPROVED ALGO-RITHM AND BOUNDS FOR SUCCESSIVE PROJECTION"

Jiashun Jin & Gabriel Moryoussef

Department of Statistics Carnegie Mellon University Pittsburgh, PA 15213, USA {jiashun, gmoryous}@andrew.cmu.edu

Zheng Tracy Ke & Jiajun Tang & Jingming Wang Department of Statistics Harvard University Cambridge, MA 02138, USA {zke, jiajuntang, jingmingwang}@fas.harvard.edu

CONTENTS

A	Proof of preliminary lemmas		
	A.1	Proof of Lemma 1	2
	A.2	Proof of Lemma 3	2
	A.3	Proof of Lemma 4	3
	A.4	Proof of Lemma 5	4
B	Ana	lysis of the SPA algorithm	5
	B.1	Some preliminary lemmas in linear algebra	6
	B.2	The simplicial neighborhoods and a key lemma	6
	B.3	Proof of Theorem A (Theorem 1 in the main paper)	6
	B.4	Proof of the supplementary lemmas	9
		B.4.1 Proof of Lemma A	9
		B.4.2 Proof of Lemma B	10
		B.4.3 Proof of Lemma C	10
		B.4.4 Proof of Lemma D	11
		B.4.5 Proof of Lemma E	11
С	Proof of the main theorems		
	C.1	Proof of Theorems 2 and 3	13
	C.2	Proof of Theorems B and C.	14
	C.3	Proof of Lemma F	17
D	Nun	nerical simulation for Theorem A	18

A **PROOF OF PRELIMINARY LEMMAS**

A.1 PROOF OF LEMMA 1

This is a quite standard result, which can be found at tutorial materials (e.g., https://people.math.wisc.edu/~roch/mmids/roch-mmids-llssvd-6svd.pdf). We include a proof here only for convenience of readers.

We start by introducing some notation. Let $Z_i = X_i - \overline{X}$ and let $Z = [Z_1, \ldots, Z_n] \in \mathbb{R}^{d,n}$. Suppose the singular value decomposition of Z is given by $Z = U_Z D_Z V'_Z$. Since H is a rank-(K-1) projection matrix, we have H = QQ', where $Q \in \mathbb{R}^{d,K-1}$ is such that $Q'Q = I_{K-1}$. Hence, we rewrite the optimization in (3) as follows:

minimize
$$\sum_{i=1}^{n} (X_i - x_0)' (I_d - QQ') (X_i - x_0)$$
, subject to $Q'Q = I_{K-1}$.

For $\lambda \in \mathbb{R}$, consider the Lagrangian objective function

$$\widetilde{S}(x_0, Q, \lambda) = \sum_{i=1}^n (X_i - x_0)' (I_d - QQ') (X_i - x_0) + \lambda (Q'Q - I_{K-1}).$$
(A.1)

Setting its gradients w.r.t. x_0 and Q to be 0 yields

$$\nabla_{x_0} \widetilde{S}(x_0, Q, \lambda) = -2(I_d - QQ') \sum_{i=1}^n (X_i - x_0) = 0,$$
(A.2)

$$\nabla_Q \widetilde{S}(x_0, Q, \lambda) = -2Q' \sum_{i=1}^n (X_i - x_0)(X_i - x_0)' + 2\lambda Q' = 0.$$
(A.3)

Firstly, we deduce from (A.2) that $\hat{x}_0 = \bar{X}$, which in view of (A.3) implies that $Q'(ZZ' - \lambda I_d) = 0$. The above equations also implies that the (K - 1) columns of \hat{Q} should be the distinct columns of U_Z . Now, the objective function in (A.1) is given by

$$\widetilde{S}(x_0, Q, \lambda) = \sum_{i=1}^{n} Z'_i (I_d - QQ') Z_i = \operatorname{tr}[(I_d - QQ') ZZ'] = \operatorname{tr}[(I_d - QQ') U_Z D_Z^2 U'_Z]$$

= $\operatorname{tr}(D_Z)^2 - \operatorname{tr}[Q' U_Z D_Z^2 U'_Z Q] = \operatorname{tr}(D_Z^2) - \|D_Z U'_Z Q\|_{\mathrm{F}}^2.$ (A.4)

Note that for each column of $U'_Z Q \in \mathbb{R}^{d,K-1}$, it has exactly one entry being 1 and its other entries are all 0. Therefore, taking $\hat{Q} = U$ maximizes $\|D_Z U'_Z Q\|_{\mathrm{F}}^2$ and hence minimizes the objective function \tilde{S} in (A.1), that is, $\hat{H} = UU'$. The proof is complete.

A.2 PROOF OF LEMMA 3

For the simplex formed by $V \in \mathbb{R}^{d \times K}$, we can always find an orthogonal matrix $O \in \mathbb{R}^{d \times d}$ and a scalar a such that

$$OV = \begin{pmatrix} x_1 & x_2 & \dots & x_K \\ a & a & \dots & a \\ 0 & 0 & \dots & 0 \end{pmatrix}, \quad \text{where} \quad x_k \in \mathbb{R}^{K-1} \text{ for } k = 1, \dots, K.$$

Denote $\bar{x} = K^{-1} \sum_{k=1}^{K} x_k$. Further we can represent

$$O\tilde{V} = \begin{pmatrix} x_1 - \bar{x} & x_2 - \bar{x} & \dots & x_K - \bar{x} \\ 0 & 0 & \dots & 0 \end{pmatrix}$$

We write $\tilde{X} := (x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_K - \bar{x})$. Since rotation and location do not change the volume,

$$\operatorname{Volume}(\mathcal{S}_0) = \operatorname{Volume}(\mathcal{S}(X)).$$

where $\mathcal{S}(\tilde{X})$ represents the simplex formed by \tilde{X} . By Stein (1966), we have

$$\operatorname{Volume}(\mathcal{S}_0) = \frac{\det(\tilde{A})}{(K-1)!}, \quad \text{with} \quad \tilde{A} = \begin{bmatrix} 1 & (x_1 - \bar{x})' \\ 1 & (x_2 - \bar{x})' \\ \vdots & \vdots \\ 1 & (x_K - \bar{x})' \end{bmatrix}$$

We also define

$$A = \begin{bmatrix} 1 & (v_1 - \bar{v})' \\ 1 & (v_2 - \bar{v})' \\ \vdots & \vdots \\ 1 & (v_K - \bar{v})' \end{bmatrix} = [\mathbf{1}_K, \tilde{V}'],$$

Since $(\tilde{A}, 0) = A \begin{pmatrix} 1 & 0 \\ 0 & O \end{pmatrix}$, it follows that $\tilde{A}\tilde{A}' = AA'$ and $\operatorname{Volume}(S_0) = \frac{\sqrt{\det(AA')}}{(K-1)!} = \frac{\sqrt{\det(A'A)}}{(K-1)!}$. Note that $A'A = \begin{pmatrix} K & 0 \\ 0 & \tilde{V}\tilde{V}' \end{pmatrix}$ by the fact that $\tilde{V}\mathbf{1}_K = 0$. Then $\det(A'A) = K\det(\tilde{V}\tilde{V}')$. Further notice that $\operatorname{rank}(\tilde{V}\tilde{V}') = K - 1$. We thus conclude that

Volume
$$(\mathcal{S}_0) = \frac{\sqrt{K}}{(K-1)!} \prod_{k=1}^{K-1} s_k(\tilde{V}).$$

This proves the first claim.

For the second and last claims, we first notice that $V = \tilde{V} - \bar{v} \mathbf{1}'_K$. Then $VV' = \tilde{V}\tilde{V}' + K\bar{v}\bar{v}'$ again by $\tilde{V}\mathbf{1}_K = 0$. Because both $\tilde{V}\tilde{V}'$ and $K\bar{v}\bar{v}'$ are positive semi-definite, by Weyl's inequality (see, for example Horn & Johnson (1985)), it follows that $s_{K-1}(V) \ge s_{K-1}(\tilde{V})$ and $s_K(V) = \sqrt{\lambda_{\min}(VV')} \le \sqrt{K\|\bar{v}\|^2} = \sqrt{K}\|\bar{v}\|$.

A.3 PROOF OF LEMMA 4

We first prove claim (a). Let $\Pi = [\pi_1 - \bar{\pi}, \dots, \pi_n - \bar{\pi}] \in \mathbb{R}^{K,n}$. Recalling the definitions of G and V, we have $G = n^{-1}\Pi\Pi'$ and $R = n^{-1/2}V\Pi$, so that $RR' = n^{-1}V\Pi\Pi'V' = VGV'$.

Next, we prove claim (b). Recall that $\tilde{V} = V - \bar{v}\mathbf{1}'_K$, so that $\tilde{V}\tilde{V}' = (V - \bar{v}\mathbf{1}'_K)(V - \bar{v}\mathbf{1}'_K)' = VV' - K\bar{v}\bar{v}'$. Note that Since $\pi'_i\mathbf{1}_K = \bar{\pi}'\mathbf{1}_K = 1$, we have $\Pi'\mathbf{1}_K = 0$, which implies that $G\mathbf{1}_K = n^{-1}\Pi(\Pi'\mathbf{1}_K) = 0$. We deduce from this observation that $\lambda_K(G) = 0$ and its associated eigenvector is $K^{-1/2}\mathbf{1}_K$. Therefore, $G - \lambda_{K-1}(G)\mathbf{1}_K + K^{-1}\lambda_{K-1}(G)\mathbf{1}_K\mathbf{1}'_K$ is a positive semi-definite matrix, so that

$$VGV' - \lambda_{K-1}(G)\tilde{V}\tilde{V}' = VGV' - \lambda_{K-1}(G)VV' + \lambda_{K-1}(G)K\bar{v}\bar{v}'$$
$$= V[G - \lambda_{K-1}(G)I_K + K^{-1}\lambda_{K-1}(G)\mathbf{1}_K\mathbf{1}'_K]V' \ge 0.$$

In addition, observing that $\Pi' 1_K = 0$ due to the fact that $\|\pi_i\|_1 = \|\bar{\pi}\|_1 = 1$, we obtain that

$$\tilde{V}G\tilde{V}' = (V - \bar{v}1'_K)G(V - \bar{v}1'_K)' = n^{-1}(V - \bar{v}1'_K)\Pi\Pi'(V - \bar{v}1'_K)' = VGV'.$$

Therefore,

$$\lambda_1(G)\tilde{V}\tilde{V}' - VGV' = \lambda_1(G)\tilde{V}\tilde{V}' - \tilde{V}G\tilde{V}' = \tilde{V}[\lambda_1(G)I_K - G]\tilde{V}' \ge 0,$$

which completes the proof of claim (b).

Finally, for claim (c), we obtain from (a) that $\sigma_{K-1}^2(R) = \lambda_{K-1}(RR') = \lambda_{K-1}(VGV')$, which by Weyl's inequality (see, for example, Horn & Johnson (1985)) and in view of claim (b) implies that $\lambda_{K-1}(G)\lambda_{K-1}(\tilde{V}\tilde{V}') \leq \sigma_{K-1}^2(R) \leq \lambda_1(G)\lambda_{K-1}(\tilde{V}\tilde{V}')$. The proof is therefore complete.

A.4 PROOF OF LEMMA 5

Recall that $z_1 \sim \chi^2_d(0)$. Let b_n be the value such that

$$\mathbb{P}(z_1 \ge b_n) = 1/n.$$

By basic extreme value theory, it is known that

$$\frac{\max_{1 \le i \le n} \{z_i\}}{b_n} \to 1, \qquad \text{in probability.}$$

We now solve for b_n . It is seen that $b_n \ge d$. Recall that the density of $\chi^2_d(0)$ is

$$\frac{1}{2^{d/2}\Gamma(d/2)}x^{d/2-1}e^{-x/2}, \qquad x > 0.$$

Note that for any $x_0 \ge d$,

$$\int_{x_0}^{\infty} x^{d/2-1} e^{-x/2} dx = 2x_0^{d/2-1} e^{-x_0/2} + \int_{x_0}^{\infty} (d-2) x^{d/2-2} e^{-x/2} dx$$
(A.5)

where the RHS is no greater than

$$\leq 2x_0^{d/2-1}e^{-x_0/2} + \frac{(d-2)}{x_0} \int_{x_0}^\infty x^{d/2-1}e^{-x/2}dx.$$

It follows that for all $x_0 \ge d$,

$$2x_0^{d/2-1}e^{-x_0/2} \le \int_{x_0}^{\infty} x^{d/2-1}e^{-x/2}dx \le x_0 \cdot x_0^{d/2-1}e^{-x_0/2},\tag{A.6}$$

where we have used

$$\frac{x_0}{x_0 - d + 2} \le x_0/2.$$

It now follows that there is a term a(x) such that when $x \ge d$,

$$1 \le a(x) \le x/2$$

and

$$\mathbb{P}(z_1 \ge x) = a(x) \frac{1}{2^{d/2} \gamma(d/2)} 2x^{d/2 - 1} e^{-x/2}.$$

Combining these, b_n is the solution of

$$a(x)\frac{1}{2^{d/2}\gamma(d/2)}2x^{d/2-1}e^{-x/2} = \frac{1}{n}.$$
(A.7)

We now solve the equation in (A.7). Consider the case d is even. The case where d is odd is similar, so we omit it. When d is even, using

$$\Gamma(d/2) = (d/2 - 1)! = (2/d)(d/2)! = (2/d)\theta(\frac{d}{2e})^{d/2},$$

where θ is the factor in the Stirling's formula which is $\leq C\sqrt{\log(d)}$. Plugging this into the left hand side of (A.7) and re-arrange, we have

$$\log(d/x) + (d/2)\log(\frac{ex}{d}) - x/2 = -\log(n) + o(\log(n)).$$
(A.8)

We now consider three cases below separately.

- Case 1. $d \ll \log(n)$.
- Case 2. $d = a_0 \log(n)$ for a constant $a_0 > 0$.
- Case 3. $d \gg \log(n)$.

Consider Case 1. In this case, it is seen that when

$$x = O(\log(n)),$$

the LHS of (A.8) is

$$-x/2 + o(\log(n)).$$

Therefore, the solution of (A.8) is seen to be

$$b_n = (1 + o(1)) \cdot 2\log(n).$$

Consider Case 2. In this case, $d = a_0 \log(n)$. Let $x = b_1 \log(n)$. Plugging these into (A.8) and rearranging,

$$a_1 - a_0 \log(a_1) = 2 + a_0 - a_0 \log(a_0) + o(1).$$
 (A.9)

Now, consider the equation

$$a_1 - a_0 \log(a_1) = 2 + a_0 - a_0 \log(a_0).$$

It is seen that the equation has a unique solution (denoted by b_0) that is bigger than 2. Therefore, in this case,

$$b_n = (1 + o(1))b_0,$$

Consider Case 3. In this case, $d \gg \log(n)$. Consider again the equation

$$\log(d/x) + (d/2)\log(\frac{ex}{d}) - x/2 = -\log(n) + o(\log(n)).$$

Letting y = x/d and rearranging, it follows that

$$y - \log(y) - 1 = o(1),$$
 (A.10)

where for sufficiently large n, o(1) > 0 and $o(1) \to 0$. Note that the function $g(y) = y - \log(y) - 1$ is a convex function with a minimum of 0 reached at y = 1, it follows

$$y = 1 + o(1)$$

Recalling y = x/d, this shows

 $b_n = (1 + o(1))d.$

This completes the proof of Lemma 5.

B ANALYSIS OF THE SPA ALGORITHM

Fix $d \ge K - 1$. For any $V = [v_1, v_2, \dots, v_K] \in \mathbb{R}^{d \times K}$, let $\sigma_k(V)$ denote the kth singular value of V, and define

$$\gamma(V) = \min_{v_0 \in \mathbb{R}^d} \max_{1 \le k \le K} ||v_k - v_0||, \qquad d_{\max}(V) = \max_{x \in S} ||x||.$$

To capture the error bound for SPA, we introduce a useful quantity

$$\beta(X,V) := \max\left\{\max_{1 \le i \le n} \operatorname{Dist}(X_i, \mathcal{S}), \quad \max_{1 \le k \le K} \min_{i:r_i = v_k} \|X_i - v_k\|\right\}.$$
 (B.11)

In this quantity, the first term is the maximum distance from any outside X_i to the simplex, and the second term is maximum radius of the neighborhood of a vertex containing no X_i . When $\beta(V, X)$ is small, it implies: (a) there is at least one data point near each true vertex, and (b) no point is too far away from the true simplex.

Theorem A. Suppose for each $1 \le k \le K$, there exists $1 \le i \le n$ such that $\pi_i = e_k$. Suppose $\beta(X, V)$ satisfies that $450d_{\max} \max\{1, \frac{d_{\max}}{\sigma_*}\}\beta \le \sigma_*^2$. Let $\hat{v}_1, \hat{v}_2, \ldots, \hat{v}_r$ be the output of SPA. Up to a permutation of these r vectors,

$$\max_{1 \le k \le r} \|\hat{v}_k - v_k\| \le \left(1 + \frac{30\gamma}{\sigma_*} \max\left\{1, \frac{d_{\max}}{\sigma_*}\right\}\right) \beta(X, V).$$

B.1 SOME PRELIMINARY LEMMAS IN LINEAR ALGEBRA

The following lemmas are proved in later subsections.

Lemma A. Let $S^* \subset \mathbb{R}^K$ be the standard probability simplex consisting of all weight vectors. Let $F: S^* \to S$ be the mapping with $F(\pi) = V\pi$. For any π and $\tilde{\pi}$ in S^* ,

$$\sigma_{K-1}(V) \cdot \|\pi - \tilde{\pi}\| \le \|F(\pi) - F(\tilde{\pi})\| \le \gamma(V) \cdot \|\pi - \tilde{\pi}\|_1.$$
(B.12)

Fix $1 \le s \le K - 2$. If π and $\tilde{\pi}$ share at least s common entries, then

$$||F(\pi) - F(\tilde{\pi})|| \ge \sigma_{K-1-s}(V)||\pi - \tilde{\pi}||.$$
 (B.13)

Lemma B. Fix $m \ge 2$ and $x_1, ..., x_m \in \mathbb{R}^d$. Let $a = \min_{i \ne j} ||x_i - x_j||$ and $b = \max_{i \ne j} ||x_i|| - ||x_j|||$. For any $w_1, ..., w_m \ge 0$ such that $\sum_{i=1}^m w_i = 1$,

$$\left\|\sum_{i=1}^{m} w_i x_i\right\| \le L - \frac{a^2 - b^2}{4L} \sum_{i=1}^{m} w_i (1 - w_i), \quad \text{with } L := \sum_{i=1}^{m} w_i \|x_i\|.$$
(B.14)

Lemma C. Fix $1 \le s \le K - 2$. For any projection matrix $H \in \mathbb{R}^d$ with rank s,

$$\sigma_{K-1-s}((I_d - H)V) \ge \sigma_{K-1}(V).$$
 (B.15)

Lemma D. Fix $0 \le s \le K-2$. Suppose there are at least s indices, $\{k_1, \ldots, k_s\} \subset \{1, 2, \ldots, K\}$, such that $||v_k|| \leq \delta$. If $\sigma_{K-1-s}^2(V) \geq 2(K-2)\delta^2$, then

$$\max_{1 \le k \le K} \|v_k\| \ge \frac{\sqrt{K-s-1}}{\sqrt{2(K-s)}} \,\sigma_{K-1-s}(V) \ge \frac{1}{2} \sigma_{K-1-s}(V). \tag{B.16}$$

B.2 THE SIMPLICIAL NEIGHBORHOODS AND A KEY LEMMA

We need some preparations. Write $\beta = \beta(X, V), \sigma_* = \sigma_{K-1}(V), \gamma = \gamma(V)$, and $d_{\max} = d_{\max}(V)$ for short. Let S^* denote the standard probability simplex, and let F be as in Lemma A. Note that $v_k = F(e_k)$, where $e_k \in \mathbb{R}^K$ is a standard basis vector. By Lemma A,

$$\|v_k - v_\ell\| \ge \sigma_* \|e_k - e_\ell\| \ge \sqrt{2\sigma_*}, \quad \text{for any } 1 \le k \ne \ell \le K.$$
(B.17)

Let $J_k = \{1 \le i \le n : \pi_i(k) = 1\}$, for $1 \le k \le K$. By definition,

$$\max_{1 \le i \le n} \operatorname{Dist}(X_i, \mathcal{S}) \le \beta, \qquad \max_{1 \le k \le K} \min_{i \in J_k} \|X_i - v_k\| \le \beta.$$
(B.18)

Given $\epsilon \in (0, 1/2)$, we introduce K local neighborhoods, one for each v_k :

$$\mathcal{V}_k(\epsilon) = \{F(\pi) : \pi \in \mathcal{S}^*, \pi_i(k) \ge 1 - \epsilon\} \subset \mathbb{R}^d, \qquad 1 \le k \le K.$$
(B.19)

When $x \in \mathcal{V}_k$, the kth entry of $\pi := F^{-1}(x)$ is at least $1 - \epsilon \leq 1/2$. Since each $\pi \in \mathcal{S}^*$ cannot have two entries larger than 1/2, these neighborhoods are disjoint:

$$\mathcal{V}_k(\epsilon) \cap \mathcal{V}_\ell(\epsilon) = \emptyset, \quad \text{for any } 1 \le k \ne \ell \le K.$$
 (B.20)

We verify that each $\mathcal{V}_k(\epsilon)$ is indeed a "local" neighborhood. Let $e_k \in \mathbb{R}^K$ be the kth standard basis vector. For any $\pi \in S^*$, $\|\pi - e_k\|_1 = 2[1 - \pi(k)]$. It follows from Lemma A that

$$\max_{x \in \mathcal{V}_k(\epsilon)} \|x - v_k\| = \max_{\pi \in \mathcal{S}^* : \pi(k) \le 1 - \epsilon} \|F(\pi) - F(e_k)\| \le 2\gamma\epsilon.$$
(B.21)

In the first iteration, i_1 is selected to maximize $||X_i||$. To study X_{i_1} , the key is to study the maximum Euclidean norm for different regions of the true simplex S. Recall that $d_{\max} = \max_{x \in S} ||x||$. The maximum is attained at one or multiple vertices. Let

$$\mathcal{K}^* = \{k : \|v_k\| = d_{\max}\}, \quad \text{where} \quad d_{\max} := \max_{x \in S} \|x\| = \max_k \|v_k\|$$

Given any $h_0 > 0$ and $\epsilon_0 > 0$, define an index set $\mathcal{K}(h_0)$ and a region $\mathcal{V}(\epsilon_0, h_0) \subset \mathcal{S}$ as follows:

$$\mathcal{K}(h_0) = \{k : \|v_k\| \ge d_{\max} - h_0\}, \qquad \mathcal{V}(\epsilon_0, h_0) = \bigcup_{k \in \mathcal{K}(h_0)} \mathcal{V}_k(\epsilon_0), \tag{B.22}$$

where $\mathcal{V}_k(\epsilon_0)$ is the same as defined in (B.19). The following lemma is a key technical lemma: **Lemma E.** Let $\mathcal{K}(h_0)$ and $\mathcal{V}(\epsilon_0, h_0)$ be as defined in (B.22). Suppose $d_{\max} \geq \sigma_*/2$. For any $C_0 > 0$, if we set $h_0 = \sigma_*/3$ and $1/2 > \epsilon_0 \ge 6C_0\sigma_*^{-1} \max\{1, d_{\max}/\sigma_*\}\beta$, then ||x||

$$\leq d_{\max} - C_0\beta$$
, for any $x \in S \setminus \mathcal{V}(\epsilon_0, h_0)$.

B.3 PROOF OF THEOREM A (THEOREM 1 IN THE MAIN PAPER)

We now show the claim. The analysis consists of three steps. In Step 1, we study the first iteration of SPA and show that \hat{v}_1 falls in the neighborhood of a true vertex. In Steps 2-3, we recursively study the remaining iterations and show that, if $\hat{v}_1, \ldots, \hat{v}_{k-1}$ fall into the neighborhoods of (k-1) true vertices, one per each, then \hat{v}_k will also fall into the neighborhood of another true vertex. For clarity, we first study the second iteration in Step 2 (for which the notations are simpler), and then study the *s*th iteration for a general *s* in Step 3.

Step 1: Analysis of the first iteration of SPA.

Applying Lemma D with s = 0, we have $d_{\max} \ge \sigma_*/2$. We then apply Lemma E. Let $C_0 = 7/3$, $h_0 = \sigma_*/3$, and $\epsilon_0 = 15 \max\{\sigma_*, \sigma_*^{-2}d_{\max}\}$ (our assumption guarantees $\epsilon_0 < 1/2$). It follows by Lemma E that

$$\max_{x \in \mathcal{S} \setminus \mathcal{V}(\epsilon_0, h_0)} \|x\| \le d_{\max} - 7\beta/3.$$
(B.23)

At the same time, for any $k \in \mathcal{K}^*$, it follows by (B.18) that there exists at least one $i^* \in J_k$ such that $||X_i - v_k|| \le \beta$. It follows by triangle inequality that $||X_{i^*}|| \ge d_{\max} - \beta$. Hence,

$$||X_{i_1}|| \ge ||X_{i^*}|| \ge d_{\max} - \beta.$$
(B.24)

Combining (B.23) and (B.24), we conclude that X_{i_1} can only be inside $\mathcal{V}(\epsilon_0, h_0)$ or outside S. Suppose X_{i_1} is outside S. It follows by (B.18) that

$$||r_{i_1}|| \ge ||X_{i_1}|| - \beta \ge d_{\max} - 2\beta.$$

Combining it with (B.23), we conclude that r_{i_1} must fall into $\mathcal{V}(\epsilon_0, h_0)$. So far, we have shown that one of the following cases must happen:

Case 1:
$$X_{i_1} \in \mathcal{V}(\epsilon_0, h_0)$$
. Case 2: $X_{i_1} \notin \mathcal{S}$ but $r_{i_1} \in \mathcal{V}(\epsilon_0, h_0)$. (B.25)

Recall that $\mathcal{V}(\epsilon_0, h_0)$ is as defined in (B.22). In Case 1, since $\mathcal{V}_1(\epsilon_0), \ldots, \mathcal{V}_K(\epsilon_0)$ are disjoint, there is only one $k_1 \in \mathcal{K}(h_0)$ such that $X_{i_1} \in \mathcal{V}_{k_1}(\epsilon_0)$. It follows by (B.21) that $||X_{i_1} - v_{k_1}|| \leq 2\gamma\epsilon_0$. In Case 2, we similarly have $||r_{i_1} - v_{k_1}|| \leq 2\gamma\epsilon_0$. In addition, since $X_{i_1} \notin S$ in this case, we have $||X_{i_1} - r_{i_1}|| \leq \beta$. Combining these arguments gives

$$\|X_{i_1} - v_{k_1}\| \le (1 + 2\gamma\epsilon_0)\beta \le \left(1 + \frac{30\gamma}{\sigma_*}\max\{1, \frac{d_{\max}}{\sigma_*}\}\right)\beta, \quad \text{for some } k_1. \quad (B.26)$$

Step 2: Analysis of the second iteration of SPA.

Let $H_1 = I_d - \frac{1}{\|X_{i_1}\|^2} X_{i_1} X_{i_1}$ and $\widetilde{X}_i = H_1 X_i$, for $1 \le i \le n$. The second iteration operates on the data points $\widetilde{X}_1, \ldots, \widetilde{X}_n \in \mathbb{R}^d$. Write $\widetilde{r}_i = H_1 r_i$, $\widetilde{\epsilon}_1 = H_1 \epsilon_i$, $\widetilde{v}_k = H_1 v_k$, and $\widetilde{V} = [\widetilde{v}_1, \widetilde{v}_2, \ldots, \widetilde{v}_K]$. It follows that

$$\widetilde{X}_i = \widetilde{V}\pi_i + \widetilde{\epsilon}_i, \qquad 1 \le i \le n.$$
(B.27)

Let $\widetilde{S} \subset \mathbb{R}^d$ denote the projected simplex, whose vertices are $\tilde{v}_1, \ldots, \tilde{v}_K$. Let \widetilde{F} denote the mapping from the standard probability simplex \mathcal{S}^* to the projected simplex \widetilde{S} (note that F_1 is not necessarily a one-to-one mapping). We introduce the neighborhoods similar to those in (B.19)

$$\widetilde{\mathcal{V}}_k(\epsilon) = \left\{ \widetilde{F}(\pi) : \pi \in \mathcal{S}^*, \pi_i(k) \ge 1 - \epsilon \right\} \subset \mathbb{R}^d, \qquad 1 \le k \le K.$$
(B.28)

Let k_1 be as in (B.26). Let $\tilde{d}_{\max} := \max_{x \in \tilde{S}} ||x||$. The maximum distance \tilde{d}_{\max} is attained at one or multiple vertices. Same as before, let $\tilde{\mathcal{K}}^*$ be the index set of k at which $||\tilde{v}_k|| = \tilde{d}_{\max}$. We similarly define

$$\widetilde{\mathcal{K}}(h_0) = \{k : \|\widetilde{v}_k\| \ge \widetilde{d}_{\max} - h_0\}, \qquad \widetilde{\mathcal{V}}(\epsilon_0, h_0) = \bigcup_{k \in \widetilde{\mathcal{K}}(h_0)} \widetilde{\mathcal{V}}_k(\epsilon_0).$$
(B.29)

At the same time, let $\tilde{\beta} = \beta(\tilde{X}, \tilde{V})$. It is easy to see that for any points x and y, $||H_1x - H_1y|| \le ||x - y||$. Hence, $\tilde{\beta} \le \beta$. It follows that

$$\max_{1 \le i \le n} \operatorname{Dist}(\widetilde{X}_i, \widetilde{S}) \le \beta, \qquad \max_{1 \le k \le K} \min_{i \in J_k} \|\widetilde{X}_i - \widetilde{v}_k\| \le \beta.$$
(B.30)

Given (B.27)-(B.30), we can extend Lemma E. A quick look of the proof of this lemma suggests that the same conclusion holds as long as we can show the following:

$$\tilde{d}_{\max} \ge \sigma_*/2, \quad \min_{\substack{(k,\ell): k \neq k_1, \\ \ell \neq k_1, k \neq \ell}} \|\tilde{v}_k - \tilde{v}_\ell\| \ge \sqrt{2}\sigma^*, \quad \text{and} \quad k_1 \notin \widetilde{\mathcal{K}}(h_0). \tag{B.31}$$

When (B.31) holds, we can similarly prove Lemma E. It is worth mentioning that d_{\max} is changed to a smaller number \tilde{d}_{\max} now, so the choice of (h_0, ϵ_0) will change accordingly. Similar to (B.23), by choosing $h_0 = \sigma_*$ and $\epsilon_1 = 15 \max\{\sigma_*, \sigma_*^{-2}\tilde{d}_{\max}\}$, we get $\max_{x \in \tilde{S} \setminus \tilde{V}(\epsilon_1, h_0)} ||x|| \le \tilde{d}_{\max} - 7\beta/3$. Note that $\epsilon_1 \le \epsilon_0$, and the set $\tilde{S} \setminus \tilde{V}(\epsilon, h_0)$ gets smaller as ϵ increases. We immediately have

$$\max_{x \in \tilde{\mathcal{S}} \setminus \tilde{\mathcal{V}}(\epsilon_0, h_0)} \|x\| \le \tilde{d}_{\max} - 7\beta/3.$$
(B.32)

At the same time, by (B.31), it is easy to get $\|\tilde{X}_{i_2}\| \ge \tilde{d}_{\max} - \beta$. We can mimic the analysis between (B.24) and (B.26) to get

$$\|\widetilde{X}_{i_2} - \widetilde{v}_{k_2}\| \le \left(1 + \frac{30\gamma}{\sigma_*} \max\left\{1, \frac{\widetilde{d}_{\max}}{\sigma_*}\right\}\right)\beta, \quad \text{for some } k_2 \ne k_1.$$
(B.33)

Note that $d_{\max} \leq d_{\max}$. Hence, the above bound for is no larger than the one in (B.33).

It remains to show (B.31). Without loss of generality, we assume $k_1 = 1$. By definition, $\tilde{V} = H_1 V$, where H_1 is a rank-1 projection matrix. It follows by Lemma C that

$$\sigma_{K-2}(V) \ge \sigma_{K-1}(V) = \sigma_*$$

Note that $\tilde{d}_{\max} \ge \max_{k \ne 1} \|\tilde{v}_k\|$. We apply Lemma D with s = 1 to get $\tilde{d}_{\max} \ge \frac{1}{2}\sigma_{K-2}(\tilde{V}) \ge \frac{1}{2}\sigma_*$. This proves the first claim in (B.31). For any k, $\tilde{v}_k = \tilde{V}e_k$, where $e_k \in \mathbb{R}^K$ is a standard basis vector. For any $2 \le k \ne \ell \le K$, e_k and e_ℓ both have a zero at the first coordinate; and we apply Lemma A with s = 1 to get $\|v_k - v_\ell\| \ge \sigma_{K-2}(\tilde{V})\|e_k - e_\ell\| \ge \sqrt{2}\sigma_*$. This proves the second claim in (B.31). Last, we show the third claim. Note that

$$\tilde{v}_1 = H_1 v_1 = v_1 - \frac{v_1' X_{i_1}}{\|X_{i_1}\|^2} X_{i_1} = \frac{X_{i_1}' (X_{i_1} - v_1)}{\|X_{i_1}\|^2} v_1 - \frac{v_1' X_{i_1}}{\|X_{i_1}\|^2} (X_{i_1} - v_1).$$

Here, $||v_1|| \le d_{\max}$, and by (B.24), $||X_{i_1}|| \ge d_{\max} - \beta$. Since $|X'_{i_1}(X_{i_1} - v_1)| \le ||X_{i_1}|| \cdot ||X_{i_1} - v_1||$, we have $|\frac{X'_{i_1}(X_{i_1} - v_1)}{||X_{i_1}||^2}|||v_1|| \le \frac{||v_1||}{||X_{i_1}||} ||X_{i_1} - v_1|| \le \frac{d_{\max}}{d_{\max} - \beta} ||X_{i_1} - v_1||$, and $\frac{v'_1 X_{i_1}}{||X_{i_1}||^2} \le \frac{||v_1||}{||X_{i_1}||} \le \frac{d_{\max}}{d_{\max} - \beta}$. Plugging these inequalities into the above equation and applying (B.26), we obtain:

$$\|\tilde{v}_{1}\| \leq \frac{2d_{\max}}{d_{\max} - \beta} \|X_{i_{1}} - r_{i_{1}}\| \leq \frac{2d_{\max}}{d_{\max} - \beta} \Big(\beta + \frac{30\gamma}{\sigma_{*}} \max\{1, \frac{d_{\max}}{\sigma_{*}}\}\beta\Big).$$
(B.34)

By our assumption, $\frac{30d_{\max}}{\sigma_*} \max\{1, \frac{d_{\max}}{\sigma_*}\}\beta \leq \sigma_*/15$. Moreover, we have shown $d_{\max} \geq \tilde{d}_{\max} \geq \sigma_*/2$. It further implies $\beta \leq \frac{\sigma_*^2}{450d_{\max}} \leq \frac{1}{225}\sigma_* \leq \frac{1}{100}\tilde{d}_{\max}$. As a result,

$$\|\tilde{v}_1\| \le \frac{200}{99} (\beta + \frac{\sigma_*}{15}) \le \frac{3}{10} \tilde{d}_{\max} \le \tilde{d}_{\max} - \frac{7}{20} \sigma_*.$$
(B.35)

At the same time, $h_0 = \sigma_*/3$. Hence, $\|\tilde{v}_1\|$ is strictly smaller than $\tilde{d}_{\max} - h_0$. This shows that 1 cannot belong to the set $\tilde{\mathcal{K}}(h_0)$ as defined in (B.29).

Step 3: Analysis of the remaining iterations of SPA.

Fix $3 \le s \le K - 1$. We now study the *s*th iteration, for $3 \le s \le K - 1$. Let i_1, \ldots, i_K denote the sequentially selected indices in SPA. Let $X_i^{(1)} = X_i$ and H_1 be the same as in Step 1 of this proof. We define $X_i^{(s)}$ and H_s recursively to describe the iterations in SPA:

$$\hat{y}_{s-1} = \frac{X_{i_{s-1}}^{(s-1)}}{\|X_{i_{s-1}}^{(s-1)}\|}, \qquad H_s = (I_d - \hat{y}_{s-1}\hat{y}_{s-1})H_{s-1}, \qquad X_i^{(s)} = H_s X_i^{(s-1)}.$$

It is seen that $H_{s-1} = \prod_{m=1}^{s-1} (I_d - \hat{y}_s \hat{y}_s)$. Note that each \hat{y}_s is orthogonal to $\hat{y}_1, \ldots, \hat{y}_{s-1}$. As a result, H_{s-1} is a projection matrix with rank (s-1). We apply Lemma C to obtain that

$$\sigma_{K-s}(H_{s-1}V) \ge \sigma_{K-1}(V) \ge \sigma_*, \quad \text{for } 3 \le s \le K-1.$$
 (B.36)

Let $\widetilde{X}_i = H_{k-1}X_i$ and $\widetilde{V} = H_{k-1}V$. We will follow similarsteps as in (B.27)-(B.33). Suppose there exist distinct $k_1, k_2, \ldots, k_{s-1}$ such that (B.33) holds for each of them. Let $\mathcal{M}_{s-1} = \{k_1, k_2, \ldots, k_{s-1}\}$. The analysis is very similar to (B.27)-(B.33), except that (B.31) is replaced by

$$\tilde{d}_{\max} \ge \sigma_*/2, \quad \min_{\substack{\{k,\ell\} \cap \mathcal{M}_{s-1} = \emptyset, \\ k \neq \ell}} \|\tilde{v}_k - \tilde{v}_\ell\| \ge \sqrt{2}\sigma^*, \quad \text{and} \quad \mathcal{M}_{s-1} \cap \widetilde{\mathcal{K}}(h_0) = \emptyset.$$
(B.37)

Once this is proved, we will follow similar arguments as in the text around (B.32) to get (B.33).

It remains to show (B.37). Suppose we have already obtained (B.37) and (B.33) for each $1 \le j \le s - 1$, and we would like to show (B.37) for s. First, consider the second claim in (B.37). For each $k \notin \mathcal{M}_{s-1}$, it has (s-1) zeros in the coordinates in \mathcal{M}_{s-1} . We apply Lemma A and (B.37) to obtain that

$$\|\tilde{v}_k - \tilde{v}_\ell\| \ge \sqrt{2}\sigma_{K-s}(\tilde{V}) \ge \sqrt{2}\sigma_*, \quad \text{for all } k \ne \ell \text{ in } \{1, \dots, K\} \setminus \mathcal{M}_{s-1}$$

Next, consider the third claim in (B.37). Note that $\mathcal{M}_{s-1} = \{k_1, k_2, \dots, k_{s-1}\}$. For each $1 \leq j \leq s-1$, by definition, $\tilde{v}_{k_j} = \left[\prod_{m>j} (I_d - \hat{y}_m \hat{y}_m')\right] \cdot (I_d - \hat{y}_j \hat{y}_j) H_{j-1} v_{k_j}$. It follows that

$$\|\tilde{v}_{k_j}\| \le \|(I_d - \hat{y}_j \hat{y}_j) H_{j-1} v_{k_j}\|, \quad \text{where} \quad \hat{y}_j = \frac{H_{j-1} X_{i_j}}{\|H_{j-1} X_{i_j}\|}.$$
(B.38)

Here, $||H_{j-1}X_{i_j}||$ is the maximum Euclidean distance attained in the (j-1)th iteration. Since we have already established (B.37) for j, we immediately have $||H_{j-1}X_{i_j}|| \ge \sigma_*/2$. In addition, we have shown (B.33) for j, which implies that

$$\|H_{j-1}v_{k_j} - H_{j-1}X_{i_j}\| \le \left(1 + \frac{30\gamma}{\sigma_*} \max\{1, \frac{d_{\max}}{\sigma_*}\}\right)\beta.$$

Mimicking the proof of (B.34), we can get a similar bound as

$$\|\tilde{v}_{j}\| \leq \|(I_{d} - \hat{y}_{j}\hat{y}_{j})H_{j-1}v_{k_{j}}\| \leq \left(1 + \frac{30\gamma}{\sigma_{*}}\max\{1, \frac{\hat{d}_{\max}}{\sigma_{*}}\}\right)\beta.$$
 (B.39)

We then mimic the argument in (B.34) to show that $||v_{k_j}|| \leq \tilde{\beta}_{\max} - 7\sigma_*/20 < \tilde{\beta} - h_0$. This implies that $j \notin \tilde{\mathcal{K}}(h_0)$. Last, consider the first argument in (B.37). Let Δ denote the right hand side of (B.39). We have shown that $||\tilde{v}_k|| \leq \Delta$, for all $k \in \mathcal{M}_{s-1}$. By our assumption, we can easily conclude that $\sigma_*^2 \geq 2(K-2)\Delta$. We then apply Lemma D to get $\tilde{d}_{\max} \geq \sigma_*/2$. So far, we have proved all three arguments in (B.37). This completes the proof of this theorem.

B.4 PROOF OF THE SUPPLEMENTARY LEMMAS

B.4.1 PROOF OF LEMMA A

By definition, $F(\pi) = \sum_{k=1}^{K} \pi(k)v_k$. Since $\sum_{k=1}^{K} \pi(k) = 1$, for any $v_0 \in \mathbb{R}^d$, we can re-express $F(\pi)$ as $F(\pi) = v_0 + \sum_{k=1}^{K} \pi(k)(v_k - v_0)$. It follows immediately that

$$\|F(\pi) - F(\tilde{\pi})\| = \left\|\sum_{k=1}^{K} [\pi(k) - \tilde{\pi}(k)](v_k - v_0)\right\| \le \|\pi - \tilde{\pi}\|_1 \cdot \max_k \|v_k - v_0\|.$$

At the same time, since $\mathbf{1}'_K(\pi - \tilde{\pi}) = 0$, the vector $\pi - \tilde{\pi}$ is an (K-1)-dimensional linear subspace. It follows by basic properties of singular values that

$$||F(\pi) - F(\tilde{\pi})|| \ge \sigma_{K-1}(V) \cdot ||\pi - \tilde{\pi}||.$$

Combining the above gives (B.12).

Suppose there are $1 \le k_1 < k_2 < \ldots < k_s \le K$ such that $\pi(k_j) = \tilde{\pi}(k_j)$, for $1 \le j \le s$. Then, the vector $\delta = \pi - \tilde{\pi}$ satisfies (s+1) constraints: $\mathbf{1}'_K \delta = 0$, $\delta(k_j) = 0$, for $1 \le j \le s$. In other words, δ lives in a (K-1-s)-dimensional linear space. It follows by properties of singular values that

$$\|F(\pi) - F(\tilde{\pi})\| \ge \sigma_{K-1-s}(V) \cdot \|\pi - \tilde{\pi}\|.$$

This proves (B.13)

B.4.2 PROOF OF LEMMA B

Write for short $x = \sum_{i=1}^{m} \pi_i x_i \in \mathbb{R}^d$ and $L = \sum_{i=1}^{m} w_i ||x_i||$. By concavity of the norm function, $||x|| \leq L$.

In this lemma, we would like to get a lower bound for L - ||x||. By definition,

$$\|x\|^{2} = \sum_{i} w_{i}^{2} \|x_{i}\|^{2} + \sum_{i \neq j} w_{i} w_{j} x_{i}' x_{j}$$
(B.40)

For any vectors $u, v \in \mathbb{R}^d$, we have a universal equality: $2u'v = 2||u|| ||v|| + (||u|| - ||v||)^2 - ||u-v||^2$. By our assumption, $||x_i - x_j|| \ge a$ and $(||x_i|| - ||x_j||)^2 \le b^2$, for all $i \ne j$. It follows that

$$x'_{i}x_{j} \le ||x_{i}|| ||x_{j}|| - (a^{2} - b^{2})/2, \qquad 1 \le i \ne j \le m.$$
 (B.41)

We plug (B.41) into (B.42) to get

$$|x||^{2} \leq \sum_{i} w_{i}^{2} ||x_{i}||^{2} + \sum_{i \neq j} w_{i} w_{j} ||x_{i}|| ||x_{j}|| - \frac{1}{2} (a^{2} - b^{2}) \sum_{i \neq j} w_{i} w_{j}$$

= $L^{2} - \frac{1}{2} (a^{2} - b^{2}) \sum_{i \neq j} w_{i} w_{j}.$ (B.42)

Note that $\sum_{i \neq j} w_i w_j = \sum_i \sum_{j:i \neq j} w_j = \sum_i w_i (1 - w_i)$. Combining it with (B.42) gives

$$||x||^{2} \le L^{2} - \frac{1}{2}(a^{2} - b^{2})\sum_{i} w_{i}(1 - w_{i}).$$
(B.43)

At the same time, $L + ||x|| \le 2L$. It follows that

$$L - \|x\| = \frac{L^2 - \|x\|^2}{L + \|x\|} \frac{L^2 - \|x\|^2}{2L} \ge \frac{a^2 - b^2}{4L} \sum_i w_i (1 - w_i).$$
(B.44)

This proves the claim.

B.4.3 PROOF OF LEMMA C

Since H is a projection matrix, there exists $Q_1 \in \mathbb{R}^s$ and $Q_2 \in \mathbb{R}^{d-s}$ such that $Q = [Q_1, Q_2]$ is an orthogonal matrix, $H = Q_1Q'_1$, and $I_d - H = Q_2Q'_2$. It follows that

$$(I_d - H)VV'(I_d - H) = Q_2(Q'_2VV'Q_2)Q'_2$$

Since Q_2 has orthonormal columns, for any symmetric matrix $M \in \mathbb{R}^{(d-s) \times (d-s)}$, M and $Q_2 M Q'_2$ have the same set of nonzero eigenvalues. Hence,

$$\sigma_{K-1-s}^2((I_d - H)V) = \lambda_{K-1-s}(Q'_2 V V' Q_2).$$

We note that $Q'_2 V V' Q_2 \in \mathbb{R}^{(d-s) \times (d-s)}$ is a principal submatrix of $Q' V V' Q \in \mathbb{R}^{d \times d}$. Using the eigenvalue interlacing theorem (Horn & Johnson, 1985, Theorem 4.3.28),

$$\lambda_{K-1-s}(Q_2'VV'Q_2) \ge \lambda_{K-1}(Q'VV'Q).$$

The claim follows immediately by noting that $\lambda_{K-1}(Q'VV'Q) = \lambda_{K-1}(VV') = \sigma_{K-1}^2(V)$.

B.4.4 PROOF OF LEMMA D

Write $\ell_{\max} = \max_{1 \le k \le K} ||v_k||$. We target to show

$$\ell_{\max}^2 \ge \frac{K - s - 1}{2(K - s)} \sigma_*^2, \quad \text{with } \sigma_* := \sigma_{K - 1 - s}(V).$$
 (B.45)

The right hand side of (B.45) is minimized at s = K - 2, at which $\ell_{\max}^2 \ge \sigma_*^2/4$. We now show (B.45). When s = 0, it is seen that

$$K\ell_{\max}^2 \ge \operatorname{trace}(V'V) = \sum_k ||v_k||^2 \ge (K-1)\sigma_*^2(V).$$

Therefore, $\ell_{\max}^2 \ge \frac{K-1}{K}\sigma_*^2$, which implies (B.16) for s = 0. When $1 \le s \le K - 2$. it is seen that $s\delta^2 + (K-s)\ell_{\max}^2 \ge \operatorname{trace}(V'V) \ge (K-1-s)\sigma_*^2$.

As a result,

$$\ell_{\max}^2 \ge \frac{(K-s-1)\sigma_*^2 - s\delta^2}{K-s}.$$
(B.46)

Note that $\frac{s}{K-s-1}$ is a monotone increasing function of s. Hence, $\frac{s}{K-s-1} \leq K-2$. The assumption of $2(K-2)\delta^2 \leq \sigma_*^2$ implies that $\frac{2s}{K-s-1}\delta^2 \leq \sigma_*^2$, or equivalently, $s\delta^2 \leq \frac{K-s-1}{2}\sigma_*^2$. We plug it into (B.46) to get $\ell_{\max}^2 \geq \frac{K-s-1}{2}\sigma_*^2$. This proves (B.16) for $1 \leq s \leq K-2$.

B.4.5 PROOF OF LEMMA E

Write $\mathcal{K} = \mathcal{K}(h_0)$, $\mathcal{V}_k = \mathcal{V}_k(\epsilon_0)$, and $\mathcal{V} = \mathcal{V}(\epsilon_0, h_0)$ for short. By definition of \overline{K} ,

 $d_{\max} - h_0 \leq ||v_k|| \leq d_{\max}$, for $k \in \mathcal{K}$, $||v_k|| \leq d_{\max} - h_0$, for $k \notin \mathcal{K}$. (B.47) Same as before, let \mathcal{S} denote the true simplex. Fix a point $x \in \mathcal{S} \setminus \mathcal{V}$, and let $\pi = F^{-1}(x)$ be its barycentric coordinate in the simplex, where F is the mapping in Lemma A. By definition of \mathcal{V} ,

$$\max_{k \in \mathcal{K}} \pi(k) \le 1 - \epsilon_0, \qquad \text{whenever } x := F(\pi) \text{ is in } \mathcal{S} \setminus \mathcal{V}. \tag{B.48}$$

We now show the claim. Define

$$\rho := \sum_{k \in \mathcal{K}} \pi_i(k), \qquad \eta := \begin{cases} \rho^{-1} \sum_{k \in \mathcal{K}} \pi(k) v_k, & \text{if } \rho \neq 0, \\ \mathbf{0}_d, & \text{otherwise.} \end{cases}$$
(B.49)

By the triangle inequality,

$$\|x\| = \left\|\rho\eta + \sum_{k \notin \mathcal{K}} \pi(k)v_k\right\| \le \rho\|\eta\| + \sum_{k \notin \mathcal{K}} \pi(k)\|v_k\| \le \rho\|\eta\| + (1-\rho)(d_{\max} - h_0).$$
(B.50)

We consider two different cases. In Case 1, $1 - \rho \ge \epsilon_0/2$. Since $\|\eta\| \le d_{\max}$, it follows from (B.50) that

$$\|x\| \le d_{\max} - (1-\rho)h_0 \le d_{\max} - \frac{h_0\epsilon_0}{2} \quad \text{if } 1-\rho \ge \frac{\epsilon_0}{2} \quad (\text{Case 1}). \tag{B.51}$$

In Case 2, $1 - \rho \le \epsilon_0/2$. If $\mathcal{K} = \{k^*\}$ is a singleton, then $\rho = \pi(k^*)$. It follows by (B.48) that $1 - \rho = 1 - \pi(k^*) \le 1 - \epsilon_0$, which yields a contradiction. Hence, it must hold that $|\mathcal{K}| \ge 2$. Then, η is a convex combination of more than one point in $\{v_k : k \in \mathcal{K}\}$. We hope to apply Lemma B. By (B.47), for each $k \in \mathcal{K}$, $||v_k||$ is in the interval $[d_{\max} - h_0, d_{\max}]$. Hence, we can take $b = h_0$ in Lemma B. In addition, by (B.17), $||v_k - v_\ell|| \ge \sqrt{2}\sigma_*$. Hence, we set $a = \sqrt{2}\sigma_*$ in Lemma B. We apply this lemma to the vector η in (B.49). It yields

$$\|\eta\| \le L - \frac{(2\sigma_*^2 - h_0^2)}{4L} \sum_{k \in \mathcal{K}} \frac{\pi(k)[\rho - \pi(k)]}{\rho^2}, \quad \text{with} \quad L := \sum_{k \in \mathcal{K}} \frac{\pi(k)}{\rho} \|v_k\|.$$

Since $L \leq d_{\max}$, the above inequality yields $\|\eta\| \leq d_{\max} - \frac{2\sigma_*^2 - h_0^2}{4\rho d_{\max}} \sum_{k \in \mathcal{K}} \pi(k) [1 - \rho^{-1}\pi(k)]$. In addition, $1 - \rho^{-1}\pi(k) = \rho^{-1}[1 - \pi(k)] - (\rho^{-1} - 1) \geq \rho^{-1}\epsilon_0 - (\rho^{-1} - 1)$, where the last inequality is from (B.48). Combining these arguments and using the fact that $\sum_{k \in \mathcal{K}} \pi(k) = \rho$, we have

$$\|\eta\| \le d_{\max} - \frac{(2\sigma_*^2 - h_0^2)[\rho^{-1}\epsilon_0 - (\rho^{-1} - 1)]}{4\rho d_{\max}} \sum_{k \in \mathcal{K}} \pi(k) \le d_{\max} - \frac{(2\sigma_*^2 - h_0^2)[\epsilon_0 - (1 - \rho)]}{4\rho d_{\max}}$$

Since $1 - \rho \le \epsilon_0/2$, we immediately have $\|\eta\| \le d_{\max} - \frac{2\sigma_*^2 - h_0^2}{4\rho d_{\max}}$. We plug it into (B.50) to get

$$||x|| \le d_{\max} - \frac{(2\sigma_*^2 - h_0^2)\epsilon_0}{4d_{\max}} \quad \text{if } 1 - \rho \le \frac{\epsilon_0}{2} \quad (\text{Case 2}).$$
 (B.52)

We combine the above two cases. By setting $h_0 = \sigma_*/3$, we have a unified expression:

$$||x|| \le d_{\max} - \min\left\{\frac{\sigma_*}{6}, \frac{4\sigma_*^2}{9d_{\max}}\right\}\epsilon_0.$$

Consequently, a sufficient condition for $||x|| \leq d_{\max} - C_0\beta$ to hold is $\epsilon_0 \geq \max\{\frac{6C_0}{\sigma_*}, \frac{9C_0d_{\max}}{4\sigma_*^2}\}\beta$. For notation simplicity, we simply set $\epsilon_0 \geq 6C_0\sigma_*^{-1}\max\{1, \sigma_*^{-1}d_{\max}\}$. This proves the claim.

C PROOF OF THE MAIN THEOREMS

We recall our pp-SPA procedure. On the hyperplane, we obtained the projected points

$$\tilde{X}_i := H(X_i - \bar{X}) + \bar{X} = (I_d - H)\bar{X} + Hr_i + H\epsilon_i$$

after rotation by U, they become $Y_i = U'\tilde{X}_i = U'r_i + U'\epsilon_i = U'X_i \in \mathbb{R}^{K-1}$. Denote $\tilde{Y}_i = U'_0X_i = U'_0r_i + U'_0\epsilon_i \in \mathbb{R}^{K-1}$. In particular, $U'_0\epsilon_i \sim N(0, \sigma^2 I_{K-1})$. Then, without loss of generality, the vertex hunting analysis on \tilde{Y}_i is equivalent to that of $X_i = r_i + \epsilon_i \in \mathbb{R}^p$, where $\epsilon_i \sim N(0, \sigma^2 I_p)$ with p = K - 1. We provide the following theorems for the rate by applying D-SPA on the aforementioned low dimension p = K - 1 space. The proof of these two theorems are postponed to Section C.2.

Theorem B. Consider $X_i = r_i + \epsilon_i \in \mathbb{R}^p$, where $\epsilon_i \sim N(0, \sigma^2 I_p)$ for $1 \leq i \leq n$. Suppose $m \geq c_1 n$ for a constant $c_1 > 0$ and $p \ll \log(n)/\log\log(n)$. Let $p/\log(n) \ll \delta_n \ll 1$. Let $c_2^* = 0.9(2e^2)^{-1/p}\sqrt{(2/p)}(\Gamma(p/2+1))^{1/p}$. Then, $c_2 \to 0.9e^{-1/2}$ as $p \to \infty$. We apply D-SPA to X_1, X_2, \ldots, X_n and output X_1^*, \cdots, X_n^* where some X_i^* may be NA owing to the pruning. If we choose $N = \log(n)$ and

$$\Delta = c_3 \sigma \sqrt{p} \left(\frac{\log(n)}{n^{1-\delta_n}} \right)^{1/p} \text{for a constant } c_3 \le c_2^*,$$

Then,

$$\beta_{new}(X^*) \le \sqrt{\delta_n} \cdot \sigma \cdot \sqrt{2\log(n)}$$

If the last inequality of (8) and (9) hold, then up to a permutation in the columns,

$$\max_{1 \le k \le K} \|\hat{v}_k - v_k\| \le g_{new}(V) \cdot \sqrt{\delta_n} \cdot \sigma \cdot \sqrt{2\log(n)}.$$

The second theorem discuss the case there a fewer pure nodes.

Theorem C. Consider $X_i = r_i + \epsilon_i \in \mathbb{R}^p$, where $\epsilon_i \sim N(0, \sigma^2 I_p)$ for $1 \le i \le n$. Fix $0 < c_0 < 1$ and assume that $m \ge n^{1-c_0+\delta}$ for a sufficiently small constant $0 < \delta < c_0$. Suppose $p \ll \log(n)/\log\log(n)$. Let $c_2^* = 0.9(2e^{2-c_0})^{-1/p}\sqrt{(2/p)}(\Gamma(p/2+1))^{1/p}$. Then $c_2 \to 0.9e^{-1/2}$ as $p \to \infty$. Suppose we apply D-SPA to X_1, X_2, \ldots, X_n and output X_1^*, \cdots, X_n^* where some X_i^* may be NA owing to the pruning. If we choose $N = \log(n)$ and

$$\Delta = c_3 \sigma \sqrt{p} \left(\frac{\log(n)}{n^{1-c_0}} \right)^{1/p} \text{ for a constant } c_3 \le c_2^*.$$

Then,

$$\beta_{new}(X^*) \le \sqrt{c_0} \cdot \sigma \cdot \sqrt{2\log(n)}$$

If the last inequality of (8) and (9) hold, then up to a permutation in the columns,

$$\max_{1 \le k \le K} \|\hat{v}_k - v_k\| \le g_{new}(V) \cdot \sqrt{c_0} \cdot \sigma \sqrt{2\log(n)}.$$

for any arbitrary small constant $\delta < 0$.

Based on the above two theorem, we have the results on $\{\tilde{Y}_i\}'s$. However, what we really care about is on $\{Y_i\}'s$ which differ from $\{\tilde{Y}_i\}'s$ by the rotation matrix. To bridge the gap, we need the following Lemma.

Lemma F. Suppose that $s_{K-1}^2(R) \gg \max\{\sqrt{\sigma^2 d/n}, \sigma^2 d/n\}$ and $\sigma = O(1)$. Then, with probability 1 - o(1),

$$\|U - U_0\| \asymp \|H - H_0\| \le \frac{C}{s_{K-1}^2(R)} \max\{\sqrt{\sigma^2 d/n}, \sigma^2 d/n\}$$
(C.53)

C.1 PROOF OF THEOREMS 2 AND 3

With the help of Theorems B, C and Lemma F, we now prove Theorems 2 and 3. We will present the detailed proof for Theorem 3. The proof of Theorem 2 is nearly identical to that of Theorem 3 with the only difference in employing Theorem C, and we refrain ourselves from repeated details.

Proof of Theorem 3. Recall that $Y_i = U'X_i = U'r_i + U'\epsilon_i$ and $\tilde{Y}_i = U'_0r_i + U'_0\epsilon_i$. Theorem B indicates that applying D-SPA on \bar{Y}_i improves the rate to $\sigma(1+o(1))\sqrt{2c_0\log(n)}$. Note that $||r_i|| \leq 1$. Also, by Lemma 5, $||\epsilon_i|| \leq (1+o(1))\sigma(\sqrt{\max\{d, 2\log(n)\}})$ simultaneously for all *i*, with high probability. Under the assumption $\alpha_n = o(1)$ for both cases and $s_{K-1}^2(R) \approx s_{K-1}^2(\tilde{V})$ by Lemma 4, the first condition in Lemma F is valid. By the last inequality in (8), we have the norm of r_i should be upper bounded for all $1 \leq i \leq n$ and therefore $s_{K-1}(\tilde{V}) \leq C \max_{k \neq l} ||\tilde{v}_k - \tilde{v}_l|| \leq C$. Further with the condition (9), we obtain that $\sigma = O(1)$. Therefore, the conditions in Lemma F are both valid. Then by employing Lemma F, we can derive that

$$\|Y_i - \tilde{Y}_i\| = O_{\mathbb{P}}\left(\frac{\sigma\sqrt{d}}{\sqrt{n}s_{K-1}^2(R)}(1 + \sigma\sqrt{\max\{d, 2\log(n)\}})\right) = O_{\mathbb{P}}(\sigma\alpha_n)$$

where the last step is due to Lemma 4 under the condition (8).

Consider the first case that $\alpha_n \ll t_n^*$. And we choose $\Delta = c_3 t_n^* \sigma$. It is seen that $\sigma \alpha_n \ll \Delta$. We will prove by contradiction that applying pp-SPA with $(\Delta, \log(n))$ on $\{Y_i\}$, the denoise step can remove outlying points whose distance to the underlying simplex larger than $\sigma[\sqrt{2c_0 \log(n)} + C\alpha_n]$ for some C > 0.

First, suppose that with probability c for a small constant c > 0, there is one point Y_{i_0} away from the underlying simplex by a distance larger than $\sigma[\sqrt{2c_0 \log(n)} + C\alpha_n]$ and it is not pruned out. Since $\sigma\alpha_n \ll \Delta$, we see that \tilde{Y}_{i_0} is faraway to the simplex with distance $\sigma\sqrt{2c_0 \log(n)}$ for certain large C and it cannot be pruned out by $(1.5\Delta, \log(n))$. However, by employing Theorem C on $\{\tilde{Y}_i\}$ with p = K - 1 and noticing $c_2^* = 1.8c_2$ with c_2 defined in the manuscript, it can only happen with negligible probability. This leads to a contradiction.

Second, suppose that with probability c for a small constant c > 0, all outliers can be removed but a vertex v_1 is also removed (which means all points near it are removed). Then, $N(\mathcal{B}(v_1, \Delta)) < \log(n)$. For the corresponding vertex for $\{\tilde{Y}_i\}$, denoted by \tilde{v}_1 , it holds that $N(\mathcal{B}(\tilde{v}_i, \Delta/2)) < \log(n)$ which means the vertex \tilde{v}_1 for $\{\tilde{Y}_i\}$ is also pruned. However, again by Theorem C, this can only happen with probability o(1). This leads to another contradiction.

Let us denote by $\beta(Y^*, U'_0 V)$ the maximal distance of points in Y^* to the simplex formed by $U'_0 V$. By the above two contradictions, we conclude that with high probability,

$$\beta(Y^*, U_0'V) \le \sigma[\sqrt{2c_0 \log(n)} + C\alpha_n].$$

where U'_0V is the underlying simplex of $\{Y_i\}$. It is worth noting that $\alpha_n = o(1)$. Then, under the assumptions of the theorem, we can apply Theorem A (Theorem 1 in the manuscript). It gives that

$$\max_{1 \le k \le K} \|\hat{v}_k^* - U_0' v_k\| \le \sigma g_{new}(V) [\sqrt{2c_0 \log(n)} + C\alpha_n]$$

where we use $(\hat{v}_1^*, \cdots, \hat{v}_K^*)$ to denote the output vertices by applying SP on $\{Y_i\}$. Eventually, we output each vertex $\hat{v}_k = (I_K - UU')\bar{X} + U\hat{v}_k^*$. It follows that up to a permutation of the K vectors,

$$\max_{1 \le k \le K} \|\hat{v}_k - v_k\| \le \max_{1 \le k \le K} \|U\hat{v}_k^* - U_0'v_k\| + \|(I_d - UU')\bar{X} - (I_d - U_0U_0')\bar{r}\|$$
$$\le \max_{1 \le k \le K} \|\hat{v}_k^* - v_k\| + \|U - U_0\| + \|(I_d - UU')\bar{X} - (I_d - U_0U_0')\bar{r}\|$$

Further we can derive

$$\begin{aligned} \|(I_d - UU')\bar{X} - (I_d - U_0U'_0)\bar{r}\| &\leq \|H - H_0\| + \|\bar{X} - \bar{r}\| \\ &\leq \sigma\alpha_n + \|\bar{\epsilon}\| \\ &\leq \sigma\alpha_n + \frac{2\sigma\sqrt{\max\{d, 2\log(n)\}}}{\sqrt{n}} \end{aligned}$$

this together with Lemma F, give rise to

$$\max_{1 \le k \le K} \|\hat{v}_k - v_k\| \le \sigma g_{new}(V) [\sqrt{2c_0 \log(n)} + C\alpha_n] + \frac{2\sigma \sqrt{\max\{d, 2\log(n)\}}}{\sqrt{n}}$$

Consider the second case that $\alpha_n \gg t_n^*$ where we choose $\Delta = \sigma \alpha_n$. By Lemma 5, it is observed that with high probability, $\max_{1 \le i \le n} d(\tilde{Y}_i, S) < (1 + o(1))\sigma \sqrt{2\log(n)}$. Notice that $||Y_i - \tilde{Y}_i|| \le C\sigma\alpha_n$ with high probability. For Y_i , if its distance to the underlying simplex is larger than $\sigma[(1 + o(1))\sqrt{2\log(n)} + C_1\alpha_n]$ for a sufficiently large $C_1 > 3C + 1$, then $d(\tilde{Y}_i, S) \ge d(Y_i, S) - C\sigma\alpha_n > \sigma[(1 + o(1))\sqrt{2\log(n)} + (2C + 1)\alpha_n]$. Hence, $\mathbb{B}(\tilde{Y}_i, (2C + 1)\Delta))$ is away from the simplex by a distance larger than $\sigma(1+o(1))\sqrt{2\log(n)}$. It follows that $N(\mathbb{B}(Y_i, \Delta)) \le N(\mathbb{B}(\tilde{Y}_i, (2C+1)\Delta)) < \log(n)$. This is equivalent to say that we do not prune out any points there. Consequently, with high probability,

$$\beta(Y^*, U_0'V) \le \sigma[(1 + o_{\mathbb{P}}(1))\sqrt{2\log(n)} + C_1\alpha_n]$$

and further by Theorem A (Theorem 1 in the manuscript),

$$\max_{1 \le k \le K} \|\hat{v}_k^* - U_0' v_k\| \le \sigma g_{new}(V) [\sqrt{2\log(n)} + C\alpha_n]$$

Next, replicate the proof for $\max_{1 \le k \le K} \|\hat{v}_k - v_k\|$ in the former case, we can conclude that

$$\max_{1 \le k \le K} \|\hat{v}_k - v_k\| \le \sigma g_{new}(V)[(1 + o_{\mathbb{P}}(1))\sqrt{2\log(n)} + C_1\alpha_n] + \frac{2\sigma\sqrt{\max\{d, 2\log(n)\}}}{\sqrt{n}}$$
$$= \sigma g_{new}(V)(1 + o_{\mathbb{P}}(1))\sqrt{2\log(n)}.$$

This concludes our proof.

C.2 PROOF OF THEOREMS B AND C.

In the subsection, we provide the proofs of Theorems B and C. We show the proof of Theorem C in detail and briefly present the proof of Theorems B as it is similar to that of Theorem C.

Proof of Theorem B. We first claim the limit of $c_2^* = 0.9(2e^{2-c_0})^{-1/p}\sqrt{(2/p)}(\Gamma(p/2+1))^{1/p}$. Note that $\Gamma(p/2+1) = (p/2)!$ if p is even and $\Gamma(p/2+1) = \sqrt{\pi}(p+1)!/(2^{p+1}(\frac{p+1}{2})!)$ if p is odd. Using Stirling's approximation, it is elementary to deduce that

$$c_2^* = e^{O(1/p) - (1 - \log(p+1))(p+1)/2p - \log(p)/2} \to e^{-1/2}.$$

Define the radius $\Delta \equiv \Delta_n = c_3 \sigma \sqrt{p} \left(\frac{\log(n)}{n^{1-c_0}}\right)^{1/p}$ for a constant $c_3 \leq c_2$. In the sequel, we will prove that applying D-SPA to X_1, \dots, X_n with (Δ, N) , we can prune out the points whose distance to the underlying true simplex are larger than the rate in the theorem, while the points around vertices are captured.

Denote d(x, S), the distance of x to the simplex S. Let

$$\mathcal{R}_f := \{ x \in \mathbb{R}^p : d(x, \mathcal{S}) \ge 2\sigma \sqrt{\log(n)} \}$$

We first claim that the number of points in \mathcal{R}_f , denoted by $N(\mathcal{R}_f)$, is bounded with probability 1 - o(1). By definition, we deduce

$$N(\mathcal{R}_f) = \sum_{i=1}^n \mathbf{1}(x_i \in \mathcal{R}_f) \le \sum_{i=1}^n \mathbf{1}(\|\varepsilon_i\| \ge 2\sigma\sqrt{\log n})$$

The mean on the RHS is given by $n\mathbb{P}(\|\varepsilon_i\| \ge 2\sigma\sqrt{\log n}) = n\mathbb{P}(\chi_p^2 \ge 4\log n) \le ne^{-1.5\log(n)} = n^{-1/2}$. By similar computations, the order of the variance is again $n^{-1/2}$. By Chebyshev's inequality, we conclude that $N(\mathcal{R}_f) = O_{\mathbb{P}}(1)$.

In the sequel, we use the notation $\mathbb{B}(x, r)$ to represent a ball centered at x with radius r and denote $N(\mathbb{B}(x, r))$ the number of points falling into this ball. And we also denote S the true underlying simplex.

Based on these notation, we introduce

 $P := \mathbb{P}(\exists X_i \text{ satisfying } \sigma \sqrt{2c_0 \log(n)} \le d(X_i, S) \le 2\sigma \sqrt{\log(n)} \text{ cannot be pruned out })$ We aim to show that P = o(1). To see this, we first derive

$$P = \binom{n}{N} N \cdot \mathbb{P}(X_1, \dots X_N \in B(X_1, \Delta) \text{ s.t. } \sigma \sqrt{2c_0 \log(n)} \le d(X_1, \mathcal{S}) \le 2\sigma \sqrt{\log(n)})$$
$$\le \binom{n}{N} N \cdot \int_{a_n \le d(x, \mathcal{S}) \le b_n} f_{X_1}(x) \mathbb{P}(X_2, \dots, X_N \in \mathcal{B}(x, \Delta)) dx$$
$$\le \binom{n}{N} N \cdot \int_{a_n \le d(x, \mathcal{S}) \le b_n} f_{X_1}(x) \prod_{t=2}^N \mathbb{P}(X_t \in \mathcal{B}(x, \Delta)) dx$$

where $a_n := \sigma \sqrt{2c_0 \log(n)}$ and $b_n := 2\sigma \sqrt{\log(n)}$ for simplicity. We can compute that for any $2 \le t \le N$,

$$\mathbb{P}(X_t \in \mathcal{B}(x, \Delta)) = (2\pi\sigma^2)^{-\frac{p}{2}} \int_{\|y-x\| \le \Delta} \exp\{-\|y-r_t\|^2 / 2\sigma^2\} dy$$

$$\leq \frac{(\Delta/\sigma)^p}{2^{p/2}\Gamma(p/2+1)} \exp\{-\frac{(\|x-r_t\| - \Delta)^2}{2\sigma^2}\}$$

$$\leq (\Delta/\sigma)^p C_p \exp\{-\frac{\|x-r_t\|^2}{2\sigma^2}\}$$

where we used $||x - r_t|| \Delta/\sigma \leq 2\sqrt{\log n} \Delta/\sigma = o(1)$ due to our choice of Δ so that Δ/σ is sufficiently small; and we write $C_p := 2^{1-p/2}/\Gamma(p/2+1)$. The above equation, together with $f_{X_1}(x) = (2\pi\sigma^2)^{-\frac{p}{2}} \exp\{-||x - r_1||^2/(2\sigma^2)\}$, leads to

$$P \le \binom{n}{N} N C_p^{N-1} (\Delta/\sigma)^{p(k-1)} \cdot \int_{a_n \le d(x,\mathcal{S}) \le b_n} (2\pi\sigma^2)^{-\frac{p}{2}} \exp\left\{-\frac{\sum_{t=1}^N \|x - r_t\|^2}{2\sigma^2}\right\} \mathrm{d}x$$

Also, notice that $\sum_{t=1}^{N} \|x - r_t\|^2 \ge N \|x - \bar{r}\|^2$ where $\bar{r} = N^{-1} \sum_{t=1}^{N} r_t$. Then,

$$P \leq \binom{n}{N} N C_p^{N-1} (\Delta/\sigma)^{p(N-1)} \cdot \int_{a_n \leq d(x,\mathcal{S}) \leq b_n} (2\pi\sigma^2)^{-\frac{p}{2}} \exp\left\{-\frac{N\|x-\bar{r}\|^2}{2\sigma^2}\right\} \mathrm{d}x$$

$$\leq \binom{n}{N} N C_p^{N-1} (\Delta/\sigma)^{p(N-1)} \int_{\|x-\bar{r}\| \geq a_n} (2\pi\sigma^2)^{-\frac{p}{2}} \exp\left\{-\frac{N\|x-\bar{r}\|^2}{2\sigma^2}\right\} \mathrm{d}x$$

$$\leq \binom{n}{N} N C_p^{N-1} (\Delta/\sigma)^{p(N-1)} N^{-p/2} \cdot \mathbb{P}(\chi_p^2 \geq 2Nc_0 \log n)$$

where we used the fact that $||x - \bar{r}|| \ge d(x, S)$ in the second step and we did change of variables so that the integral reduces to the tail probability of χ_p^2 distribution. By Mills ratio, the tail probability of χ_p^2 is given by

$$\mathbb{P}(\chi_p^2 \ge 2Nc_0 \log n) \le Cn^{-Nc_0} (2Nc_0 \log n)^{p/2-1},$$

we obtain

$$P \le C\binom{n}{N} N C_p^{N-1} (\Delta/\sigma)^{p(N-1)} N^{-p/2} n^{-Nc_0} (2Nc_0 \log n)^{p/2-1}.$$

Using the approximation $\binom{n}{k} \leq C(en/k)^k$, we deduce that

$$P \leq C \left[e(2Nc_0 \log n)^{(p-2)/(2N)} C_p^{1-1/N} N^{(1-p/2)/N} \cdot \frac{n^{1-c_0} (\Delta/\sigma)^{p(1-1/N)}}{N} \right]^N$$
$$=: C \left[A(n,p,N) \cdot \frac{n^{1-c_0} (\Delta/\sigma)^{p(1-1/N)}}{N} \right]^N$$

Now we plug in $N = \log(n)$ and $\Delta = c_3 \sigma \sqrt{p} \left(\frac{\log(n)}{n^{1-c_0}}\right)^{1/p}$ for a constant $c_3 \leq c_2$ where $c_2 = 0.9(2e^{2-c_0})^{-1/p} \sqrt{(2/p)} (\Gamma(p/2+1))^{1/p} = 0.9e^{-(2-c_0)/p} C_p^{-1/p} / \sqrt{p}$ with $C_p = 2^{1-p/2} / \Gamma(p/2+1)$. It is straightforward to compute that

$$\begin{aligned} A(n,p,N) \cdot \frac{n^{1-c_0} (\Delta/\sigma)^{p(1-1/N)}}{N} \\ &\leq e^{1-(2-c_0)(1-1/\log(n))} 2^{\frac{p-2}{2\log(n)}} (c_0 \log(n))^{\frac{p-2}{2\log(n)}} (0.9)^{p(1-1/\log(n))} \Big(\frac{n^{1-c_0}}{\log(n)}\Big)^{1/\log(n)} \\ &\leq e^{o(1)} (0.9)^p < 1.01 \cdot 0.9 < 1 \end{aligned}$$

under the condition that $p \ll \log(n) / \log \log(n)$. This implies $P \le C(0.909)^{\log(n)} = o(1)$. In the mean time, for each vertex v_k ,

$$N(\mathcal{B}(v_k, \Delta/2)) \ge \sum_{i \in \mathcal{M}_k} \mathbf{1}(x_i \in \mathcal{B}(v_k, \Delta/2)) = \sum_{i \in \mathcal{M}_k} \mathbf{1}(\|\varepsilon_i\| \le \Delta/2) \ge mp_\Delta - C\sqrt{mp_\Delta \log \log(n)}.$$

with probability 1 - o(1), and

$$p_{\Delta} := \mathbb{P}(\|\varepsilon_i\| \le \Delta/2) = \mathbb{P}(\chi_p^2 \le 4^{-1}(\Delta/\sigma)^2) \ge \frac{e^{-(\Delta/\sigma)^2/8}2^{-p}}{2^{p/2}\Gamma(p/2+1)}(\Delta/\sigma)^p$$

Recall the condition that $m \ge n^{\delta} n^{1-c_0}$. It follows that

$$mp_{\Delta} \ge n^{\delta} \frac{e^{-(\Delta/\sigma)^{2}/8} 2^{-p}}{2^{p/2} \Gamma(p/2+1)} n^{1-c_{0}} (\Delta/\sigma)^{p} = n^{\delta} \frac{e^{-(\Delta/\sigma)^{2}/8}}{2^{p/2} \Gamma(p/2+1)} \cdot \frac{c \log(n)}{C_{p}} 2^{-p} (c_{3}/c_{2})^{p} \\ \ge cn^{\delta} 2^{-p} (c_{3}/c_{2})^{p} \log(n) \gg \log(n)$$

where c > 0 is some small constant. The last step is due to the fact that $n^{\delta}2^{-p}(c_3/c_2)^p = e^{\delta \log(n) - p \log(2c_2/c_3)} \gg 1$ as $2c_2/c_3 \ge 2$ is a constant and $p \ll \log(n) / \log \log(n)$. Thus, with probability 1 - o(1), $N(\mathcal{B}(v_k, \Delta/2)) \gg \log(n)$. Under this event, for any point $X_{i_0} \in \mathcal{B}(v_k, \Delta/2)$, immediately $\mathcal{B}(v_k, \Delta/2) \subset \mathcal{B}(X_{i_0}, \Delta)$ and further $N(\mathcal{B}(X_{i_0}, \Delta)) \gg \log(n)$. Combining this, with P = o(1) and $N(\mathcal{R}_f) = O_{\mathbb{P}}(1)$, we conclude that we can prune out all points with a distance to the simplex larger than $\sigma \sqrt{2c_0 \log(n)}$ while preserve those points near vertices, with high probability. Thus we finish the claim for $\beta_{new}(X^*)$.

The last claim follows directly from Theorem A (Theorem 1 in the manuscript) under condition (9). We therefore conclude the proof.

We briefly present the proof of Theorem B below.

Proof. The proof strategy is roughly the same as that of Theorem C When $m > c_1 n$, we take $\Delta = c_3 \sigma \sqrt{p} \left(\frac{\log(n)}{n^{1-\delta_n}}\right)^{1/p}$ where $p/\log(n) \ll \delta_n \ll 1$ and $c_3 \leq c_2$, then similarly we can derive that $N(\mathcal{B}(v_k, \Delta/2)) \geq c\log(n)n^{\delta_n}a^p = c\log(n)e^{\delta_n\log(n)-p\log(1/a)} \gg \log(n)$ where c > 0 is a small constant and $0 < a \leq 1$. This gives rise to the conclusion that with high probability, $N(\mathcal{B}(X_{i_0}, \Delta)) \gg \log(n)$ for any $X_{i_0} \in N(\mathcal{B}(v_k, \Delta/2))$. Moreover, in the same manner to the above derivations, replacing c_0 by δ_n , we can claim again that $N(\mathcal{R}_f) = O_{\mathbb{P}}(1)$ and

$$P \le C \left(A(n, p, \log(n)) \cdot \frac{n^{1-\delta_n} (\Delta/\sigma)^{p(1-1/\log(n))}}{\log(n)} \right)^{\log(n)} = o(1).$$

Consequently, all the claims follow from the same reasoning as the proof of Theorem C. We therefore omit the details and conclude the proof . $\hfill \Box$

C.3 PROOF OF LEMMA F

Recall that $R = n^{-1/2}[r_1 - \bar{r}, \dots, r_n - \bar{r}]$. Let $R = U_0 D_0 V_0$ be its singular value decomposition and let $H_0 = U_0 U'_0$. Denote $\epsilon = [\epsilon_1, \dots, \epsilon_n] \in \mathbb{R}^{d,n}$. We start by analyzing the convergence rate of $\|ZZ' - nRR' - n\sigma^2 I_d\|$. Recall that $\bar{X} = \bar{r} + \bar{\epsilon}$, where $\bar{\epsilon} = n^{-1} \sum_{i=1}^n \epsilon_i$. We obtain

$$Z = X_i - \bar{X} = r_i + \epsilon_i - \bar{r} - \bar{\epsilon}, \qquad Z = \sqrt{nR} + \epsilon - \bar{\epsilon}\mathbf{1}'_n.$$
(C.54)

Observing the fact that $R1_n = 0$, we deduce

$$ZZ' - nRR' - n\sigma^{2}I_{d} = (\sqrt{n}R + \epsilon - \bar{\epsilon}1'_{n})(\sqrt{n}R + \epsilon - \bar{\epsilon}1'_{n})' - nRR' - n\sigma^{2}I_{d}$$

$$= \sqrt{n}(\epsilon - \bar{\epsilon}1'_{n})R' + \sqrt{n}R(\epsilon - 1_{n}\bar{\epsilon}')' + (\epsilon - \bar{\epsilon}1'_{n})(\epsilon - \bar{\epsilon}1'_{n})' - n\sigma^{2}I_{d}$$

$$= \sqrt{n}\epsilon R' + \sqrt{n}R\epsilon' + (\epsilon\epsilon' - n\sigma^{2}I_{d}) - n\bar{\epsilon}\bar{\epsilon}'.$$
(C.55)

The above equation implies that

$$||ZZ' - nRR' - n\sigma^2 I_d|| \le 2\sqrt{n} ||\epsilon R'|| + ||\epsilon \epsilon' - n\sigma^2 I_d|| + n||\bar{\epsilon}||^2.$$
(C.56)

We proceed to bound the three terms $\|\epsilon R'\|$, $\|\epsilon\epsilon' - n\sigma^2 I_d\|$ and $n\|\bar{\epsilon}\|^2$ respectively. First, notice that $\epsilon R' \in \mathbb{R}^{d \times d}$ is a Gaussian random matrix with independent rows which follow N(0, RR'). By Theorem 5.39 and Remark 5.40 in Vershynin (2010), we can deduce that with probability 1 - o(1),

$$n \|R\epsilon' \epsilon R'\| \le d\sigma^2 [s_1^2(R) + n/d + c].$$

This, together with the fact that $s_1(R) \leq c$ gives that

$$\sqrt{n} \|\epsilon R' + R\epsilon'\| \le 2\sigma \sqrt{nd}. \tag{C.57}$$

Second, by Bai-Yin law (Bai & Yin (2008)), we can estimate the bound of $\|\mathcal{E}\mathcal{E}' - n\sigma^2 I_d\|$ as follows.

$$\|\epsilon\epsilon' - n\sigma^2 I_d\| \le n\sigma^2 (2\sqrt{d/n} + d/n) \le \sigma^2 (2\sqrt{nd} + d), \tag{C.58}$$

with probability 1 - o(1). Third, observe that $\bar{\epsilon} \sim N(0, \sigma^2/nI_d)$. We therefore obtain that with probability 1 - o(1),

$$n\|\bar{\epsilon}\|^2 \le \sigma^2[d + C\sqrt{d\log(n)}]$$

By applying the condition that $\sigma = O(1)$, combining the above equation with (C.56), (C.57) and (C.58) yields that, with probability at least 1 - o(1),

$$||ZZ' - nRR' - n\sigma^2 I_d|| \le 2\sigma\sqrt{nd} + \sigma^2[d + C\sqrt{d\log(n)}] + \sigma^2(2\sqrt{nd} + d)$$

$$\le C(\sigma\sqrt{nd} + \sigma^2 d).$$
(C.59)

Now, we compute the bound for $\|\hat{H} - H_0\|$. Let $U^{\perp}, U_0^{\perp} \in \mathbb{R}^{d,d-K+1}$ such that their columns are the last (d - K + 1) columns of U and U_0 , respectively. It follows from direct calculations that

$$\begin{aligned} \|\widehat{H} - H_0\| &= \|U_0 U_0' - UU'\| \le \|U_0^{\perp} (U_0^{\perp})' (U_0 U_0' - UU')\| + \|U_0 U_0' (U_0 U_0' - UU')\| \\ &= \|U_0^{\perp} (U_0^{\perp})' UU'\| + \|U_0 U_0' U^{\perp} (U^{\perp})'\| \le \|(U_0^{\perp})' U\| + \|U_0' U^{\perp}\| = 2\|\sin\Theta(U_0, U)\|. \end{aligned}$$

Notably, U_0, U_0^{\perp} is also the eigen-space of $ZZ' - n\sigma^2 I_d$. By Weyl's inequality (see, for example, Horn & Johnson (1985)),

$$\max_{1 \le i \le d} \left| \lambda_i (ZZ' - n\sigma^2 I_d) - \lambda_i (ZZ') \right| \le C \|ZZ' - n\sigma^2 I_d - nRR'\|$$

Under the condition that $s_{K-1}^2(R) \gg \max\{\sqrt{\sigma^2 d/n}, \sigma^2 d/n\}$, by Davis-Kahan Theorem (Davis & Kahan (1970)), we deduce that, with probability at least 1 - o(1),

$$\|\widehat{H} - H_0\| \le 2\|\sin\Theta(U_0, U)\| \le \frac{2\|ZZ' - nRR' - n\sigma^2 I_d\|}{\lambda_{K-1}(RR')}$$
$$\le C \frac{\max\{\sqrt{\sigma^2 d/n}, \sigma^2 d/n\}}{s_{K-1}^2(R)}.$$
(C.60)

The proof is complete.

D NUMERICAL SIMULATION FOR THEOREM A

In this short section, we want to provide a better sense of our bound derived in Theorem A and how it compares with the one from the orthodox SPA. To make it easier for the reader to see the difference between the two bounds, we consider toy example where we fix (K, d) = (3, 3) and

$$V = \{(20, 20, 0), (20, 30, 0), (30, 20, 0)\}$$

while we let

 $V = \widetilde{V} + a \cdot (0, 0, 1).$

We consider 50 different values for a ranging from 10 to 1000. It is not surprising to see that when a is close to 0 the bound of the orthodox SPA goes to infinity whereas as the simplex is bounded far away from the origin, the K^{th} singular value will be bounded away from 0. However, our bound still outperforms the traditional SPA bound even for very large values of a. Looking at two specific values of a we have the following. For a = 10,

$$\beta_{new} = 0.03, \qquad \beta(V) = 0.05$$

Moreover, as a changes, the Figure 1 below illustrate how much the ratio of

our whole bound Gillis bound

changes as the parameter a changes. For example, when a = 10.

$$\frac{g_{new}(V)}{g(V)} = 0.015,$$

and so

$$\frac{\text{our whole bound}}{\text{Gillis bound}} = 0.009$$

so we reduce the bound by 111 . Similarly, when a = 1000,

$$\frac{g_{new}(V)}{g(V)} = 0.19,$$
 our whole bound
Gillis bound = 0.105

so we have reduced the bound by 9.5.



Figure 1: Factor of improvement of our bound over orthodox spa as the true simplex moves away from origin by a distance a.

REFERENCES

- Zhi-Dong Bai and Yong-Qua Yin. Limit of the smallest eigenvalue of a large dimensional sample covariance matrix. In *Advances In Statistics*, pp. 108–127. World Scientific, 2008.
- Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM J. Numer. Anal.*, 7(1):1–46, 1970.
- Roger Horn and Charles Johnson. Matrix Analysis. Cambridge University Press, 1985.
- P Stein. A note on the volume of a simplex. *The American Mathematical Monthly*, 73(3):299–301, 1966.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *ArXiv.1011.3027*, 2010.