


Estimating kinetic constants in the Michaelis–Menten model from one enzymatic assay using Approximate Bayesian Computation

Jakub M. Tomczak¹  and Ewelina Węglarz-Tomczak² 

¹ Institute of Informatics, Faculty of Science, University of Amsterdam, The Netherlands

² Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam, The Netherlands

Correspondence

E. Węglarz-Tomczak, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam, Science Park 904, room: C2.104, 1098 XH Amsterdam, The Netherlands
Tel: +31 (0)20 525 7931
E-mail: ewelina.weglarz.tomczak@gmail.com

(Received 1 January 2019, revised 5 June 2019, accepted 27 June 2019, available online 21 July 2019)

doi:10.1002/1873-3468.13531

Edited by Alfonso Valencia

The Michaelis–Menten equation is one of the most extensively used models in biochemistry for studying enzyme kinetics. However, this model requires at least a couple (e.g., eight or more) of measurements at different substrate concentrations to determine kinetic parameters. Here, we report the discovery of a novel tool for calculating kinetic constants in the Michaelis–Menten equation from only a single enzymatic assay. As a consequence, our method leads to reduced costs and time, primarily by lowering the amount of enzymes, since their isolation, storage and usage can be challenging when conducting research.

Keywords: Approximate Bayesian Computation; Bayesian statistics; enzymology; likelihood-free; Michaelis–Menten kinetics

In biochemistry, the Michaelis–Menten model [1–3] is one of the best-known and useful approaches to enzyme kinetics [4–7]. It takes the form of an equation describing the rate of enzymatic reaction by relating a rate of formation of the product to a substrate concentration. The system involves two reactions where a substrate binds reversibly to the enzyme to form an enzyme–substrate complex, which then reacts irreversibly to generate a product and to regenerate the original enzyme. In the beginning of the early 20th century, Victor Henri discovered that enzyme reactions are initiated by a bond between the enzyme and the substrate [8]. Later, his work was taken up by Leonor Michaelis and Maud Menten [3], who measured the initial velocity as a function of sucrose concentration and derived an equation that approximates the modern version of the Michaelis–Menten equation, which is widely used in enzyme–substrate interaction study [9]. Moreover, this approach is

applied to other biochemical problems, including antigen–antibody binding, protein–protein interaction [2] as well as pharmacokinetics [10], and to areas outside biochemical interaction like clearance of blood alcohol [11], alveolar clearance of dusts [12], and bacteriophage infection [13]. The model provides valuable knowledge for researchers in the form of kinetic parameters that explain the properties of a particular enzyme. The standard method for determining the kinetic parameters involves running a series of enzyme assays at multiple substrate concentrations, and measuring the initial rate of the reaction [2]. By plotting reaction rate against substrate concentration, and applying a nonlinear regression to the Michaelis–Menten equation, the kinetic parameters could be obtained. Until recently, the graphical methods, such as the one described by Lineweaver and Burk, which utilized the double reciprocal plot, were used for this purpose [2]. Nowadays, various software programs

Abbreviations

ABC, Approximate Bayesian Computation; *h*APN, human aminopeptidase; *h*ERAP2, human endoplasmic reticulum aminopeptidase 2; *ss*APN, *Sus scrofa* APN.

provide fast tools to obtain kinetic parameters from nonlinear regression of the Michaelis–Menten equation.

In order to minimize experimental work that involves several enzymatic assays, we propose to use the Approximate Bayesian Computation (ABC) techniques [14–18] to determine kinetic constant in the Michaelis–Menten model. Commonly, biochemical systems could be described using differential equations or a set of nondifferentiable set of rules with parameters having physical interpretation [6,19]. Then, for given initial conditions and specific values of parameters it is possible to generate data, that is, provide a solution of the set of differential equations. However, an analytical form of the distribution of solutions is highly complex and, thus, it is impossible to formulate the likelihood function. ABC techniques offer an almost automated solution in situations where evaluation of the posterior is computationally prohibitive, or whenever suitable likelihoods are not available. The main idea behind ABC is to utilize a mathematical model for a biochemical system and apply an approximate inference (e.g., Monte Carlo methods) to calculate the posterior over parameters. A classical example of applying ABC is a model of spreading tuberculosis [19] that contains a set of probabilistic rules of events, such as, illness transmission, mutation, death, and recovery. The model is relatively simple yet it is analytically intractable. Since the probabilistic inference is infeasible due to lack of an analytical form of the model, a rejection sampling procedure with a prespecified acceptance threshold was used to infer about parameters. Namely, values of parameters are first sampled from a prior distribution and then data are generated by the model. If the generated data are close to the real data in terms of a chosen distance measure (i.e., the distance is smaller than the threshold ϵ), the parameters' values are accepted and further used to approximate the posterior. Later, the ABC techniques were widely used in other applications, such as, biology [20,21], evolution and ecology [15,22], the evolution of genomes [23], the dynamics of gene regulation [18], the demographic spread of species [24–27], or mRNA self-regulation [28]. It is worth mentioning that [28] used Parallel Tempering and Metropolis–Hastings sampling techniques to determine parameters of the Michaelis–Menten model. However, they used a synthetic likelihood function [29,30] assuming 1% Gaussian error while our approach is likelihood-free.

In this paper, we present a novel application of the ABC computational tool for calculating kinetic constants in the Michaelis–Menten equation using one

enzymatic assay at a single substrate concentration. This extremely useful framework gives wider possibilities in determination of kinetic parameters by immense reduction of the cost, primarily by lowering the amount of enzyme, substrate and other reagents as well as time.

Materials and methods

Implementation

The proposed method was implemented in PYTHON using the NumPy package (www.numpy.org). In all experiments, we used our own implementation of the Runge-Kutta (RK4) solver. The code is available at: <https://github.com/e-weglarz-tomczak/mmabc>.

In the implementation of our method, we used two heuristics to obtain samples from the posterior distribution over parameters:

- For computational convenience, the sampling procedure is run 100 times to sample 10 000 parameters.
- If the procedure is stopped before reaching 100 iterations (runs), then we increase the threshold of the rejection sampling ϵ and repeat the procedure.

We noticed that such approach worked very well in practice.

Parameter estimation from synthetic data

In order to verify the usefulness of the proposed approach, we first evaluate our method on synthetic data. We simulate 6000 different synthetic measurements using the following setting:

- We sampled the initial concentration of the enzyme E_0 uniformly from the set of the following values: {0.0005, 0.001, 0.005} [μM].
- The step-size h (a time between two consecutive measurement points) is sampled uniformly from the following set: {8, 10, 12}. The time step is expressed in seconds.
- The measurement duration is set to 15 min.
- The values of k_{cat} (the catalytic rate constant) and K_M (the Michaelis constant) are sampled uniformly from [0.1, 100] and [1, 200] in the linear scale, respectively. Further, the sampled values are used to generate synthetic measurements by solving the Michaelis–Menten equation using the RK4 method.
- During the simulation, we use the following values of the threshold: $\epsilon \in \{0.0001, 0.0002, 0.0003, 0.0005, 0.0007, 0.001, 0.002, 0.003\}$. We start with the smallest value of ϵ and if at the end of the procedure < 5 samples are accepted, we increase ϵ and repeat the procedure.

• In the experiment, we run our procedure for different values of S_0 , namely, $S_0 \in \{10, 50, 100, 200, 300, 400\}$ [μM]. For a single value of S_0 we generate 1000 simulations.

Parameter estimation from real data

The utility of the proposed method is further demonstrated on kinetic parameters measurement data for three real enzymes, namely human aminopeptidase (*hAPN*), *Sus scrofa* APN (*ssAPN*), and human endoplasmic reticulum aminopeptidase 2 (*hERAP2*) [31,32]. For each enzyme eight measurements were collected at different levels of substrate.

Concentration, $S_0 \in \{3.125, 6.25, 12.5, 25, 50, 100, 200, 400\}$ [μM]. In our experiments a single measurement for $S_0 = 200$ is taken, the following settings are used:

- *hAPN*: $E_0 = 0.0005 \mu\text{M}$, $h = 8$ s, one measurement lasted 15 min;
- *ssAPN*: $E_0 = 0.0005 \mu\text{M}$, $h = 12$ s, one measurement lasted 15 min;
- *hERAP2*: $E_0 = 0.001 \mu\text{M}$, $h = 10$ s, one measurement lasted 15 min.

In order to obtain the values of the kinetic constants, the standard method was used [2]. Further details how the experiments were carried out are outlined in Refs [31,32].

Metrics

The following metrics are used to quantify the performance of the presented approach:

- Deviation specifies how the mean estimated value, θ_{est} , deviates from the real one, θ_{real} :

$$\text{Deviation} = \begin{cases} 1 - \frac{\theta_{\text{real}}}{\theta_{\text{est}}}, & \theta_{\text{est}} > \theta_{\text{real}} \\ 1 - \frac{\theta_{\text{est}}}{\theta_{\text{real}}}, & \text{otherwise} \end{cases} \quad (1)$$

- Accuracy indicates how often the real value θ_{real} is within one standard deviation interval from the mean estimated across all simulations.

Results and Discussion

The Michaelis–Menten model

The Michaelis–Menten model takes the form of an equation describing the rate of enzymatic reactions, by relating reaction rate to the concentration of a substrate. The enzymatic reaction in this model involves reversible reaction where an enzyme, E, binds to a

substrate, S, to form a complex, ES, and irreversible releasing a product, P, and the free enzyme. This may be represented schematically as follows:



By applying the law of mass action, which states that the rate of any chemical reaction is directly proportional to the product of the masses/concentrations of the reacting substances [33], system of four ordinary differential equations that define the rate of change of reactants over time are obtained, namely:

$$\frac{d[E]}{dt} = -k_f[E][S] + k_r[ES] + k_{\text{cat}}[ES] \quad (3)$$

$$\frac{d[S]}{dt} = -k_f[E][S] + k_r[ES] \quad (4)$$

$$\frac{d[ES]}{dt} = -k_f[E][S] - k_r[ES] - k_{\text{cat}}[ES] \quad (5)$$

$$\frac{d[P]}{dt} = k_{\text{cat}}[ES] \quad (6)$$

In their original analysis, Michaelis and Menten assumed that the substrate is in instantaneous chemical equilibrium with the complex enzyme–substrate [3]. An alternative analysis was proposed 10 years later by Briggs and Haldane, known as quasi-steady-state of the system, that assumed the concentration of the intermediate complex is approximately steady during progression of the reaction [34]. Both approximations assume irreversible dissociation of ES complex to the product and free enzyme that yields the following equation of the velocity of the reaction:

$$v = \frac{dP}{dt} = \frac{V_{\text{max}}S}{K_M + S} = \frac{E_0k_{\text{cat}}S}{K_M + S}, \quad (7)$$

where P is the concentration of the product, S denotes the concentration of the substrate, V_{max} is the maximal velocity, E is the initial concentration of the enzyme, k_{cat} denotes the constant of the conversion to the product (the catalytic rate constant), and K_M is called the Michaelis constant. Additionally, the initial value of the substrate, S_0 is known.

The model provides crucial information about the nature of the enzyme in the form of the kinetic parameters. The Michaelis constant (K_M) is the concentration of substrate at which the reaction rate is half of the maximum rate. This is an inverse measure of the substrate's affinity for the enzyme. The smaller the value

of K_M , the higher the affinity; hence, the rate will approach maximal velocity with lower S than reactions with a lower K_M . The catalytic rate (k_{cat}) shows how fast reaction is. The k_{cat}/K_M constant provides the knowledge how efficiently an enzyme converts a substrate into a product. The curve describing the relationship between the velocity and the substrate concentration in the Michaelis–Menten kinetics can be formulated as a nonlinear regression model. The commonly used model is expressed using the exponential function:

$$v = V_{\text{max}}(1 - \exp(-bS)), \quad (8)$$

where b is a real-valued parameter, or by using a polynomial of the k -th order:

$$v = a_1S + a_2S^2 + \dots + a_kS^k + a_0, \quad (9)$$

where a_0, a_1, \dots, a_k are constants.

The standard approach to Michaelis–Menten kinetics

The typical method for determining the constants involves performing a series of in vitro experiments at varying substrate concentrations. Next, from enzyme assays, the initial reaction rates are obtained. The initial reaction rate is taken to mean because the equilibrium or quasi-steady-state approximation remains valid. By plotting reaction rate against concentration of the substrate, and using a nonlinear regression of the Michaelis–Menten equation, the kinetic parameters can be calculated. Nowadays, available software programs for enzyme kinetics calculations allow to fit the exponential Eqn (8) and/or the polynomial (9) to measured values. Before computational era, graphical methods involving linearization of the equation were used to perform nonlinear regression, such as, the Eadie–Hofstee diagram, Hanes–Woolf plot, and Lineweaver–Burk plot as most commonly used [2].

The standard approach for finding k_{cat} and K_M assumes the following three steps [2]:

- Calculate the initial rate of the reaction measured at varying substrate concentration from the linear portion of the reaction progress curve (product vs. time) (see step 1 in Figure 1).
- Fit the Eqn (8) or the polynomial (9) to the plotted reaction rate against concentration of the substrate (see step 2 in Figure 1).
- Calculate the kinetic parameters K_M and k_{cat} from the fitted curve (see step 3 in Figure 1).

The procedure is straightforward and rather accurate [2]; however, it is prone to noise in measurements and requires multiple (e.g., eight or more) measurements at different substrate concentration level in order to properly fit the curve and, eventually, determine the kinetic constants. Since carrying out several enzymatic assays is a time-consuming process, the whole procedure becomes rather slow. Additionally, performing all experiments requires substantial amount of enzyme that is typically costly and time-demanding to obtain. As a result, costs and time needed to determine the kinetic constants are usually high and become a considerable bottleneck in the whole research process.

ABC approach to the Michaelis–Menten kinetics

One of the main inference frameworks is the Bayesian approach where the prior knowledge about parameters, θ , is further updated by observations v (the posterior) [35]. According to Bayes' rule the posterior distribution over the model parameters, $p(\theta|v)$ is proportional to the product of the likelihood function of the observed data, $p(v|\theta)$, and the prior distribution over the parameters, $p(\theta)$, namely:

$$p(\theta|v) \propto p(v|\theta)p(\theta). \quad (10)$$

Typically, the likelihood function and the prior are chosen from a known family of distributions. However, in many cases, the form of the distribution $p(v|\theta)$ remains unknown, but we can sample from it (i.e., generate new data). In the ABC literature, such model is called a simulator [15].

In the considered case, we treat the Michaelis–Menten model in (7) as the simulator of enzyme kinetics with parameters $\theta = \{k_{\text{cat}}, K_M\}$. For given values of the kinetic parameters k_{cat} and K_M sampled from some assumed prior (e.g., the uniform prior), the ODE can be solved using a numerical ODE solver (e.g., RK4 solver). As a result, simulated data v_{sim} can be further compared with the real measurements v_{real} . If a distance (e.g., the Euclidean distance, $\|\bullet\|_2^2$) between the simulated data and the real observations is smaller than a prespecified threshold ε (e.g., $\varepsilon = 0.001$), then we can keep the values of parameters. Otherwise, we disregard them. Eventually, we could use all the accepted parameters to estimate the posterior distribution. This procedure is called the ABC rejection algorithm [21].

Since the initial conditions are known (E_0 and S_0), we could easily run the simulator multiple times in order to estimate the posterior distribution over the




Fig. 1. The three steps of the standard approach: (1) Calculate reaction rates at different substrate concentrations. (2) Fit a curve (e.g., a polynomial) to the calculated reaction rates. (3) Calculate k_{cat} as the maximum rate divided by the initial concentration of the enzyme, and K_M as the substrate concentration at which the curve attains half of the maximum rate.

kinetic constants. Moreover, the measurement could be noisy and we can still estimate the parameters by enlarging the threshold ε . Therefore, we propose to utilize the ABC rejection method to determine the kinetic constants of the Michaelis–Menten model. For given ε , E_0 and S_0 , the procedure is the following (Figure 2):

- Sample $\theta = \{k_{\text{cat}}, K_M\}$ from the prior $p(\theta)$ being the uniform distribution over a prespecified closed region (e.g., $k_{\text{cat}} \in [0, 100]$ and $K_M \in [0, 200]$).
- Simulate data v_{sim} using the Michaelis–Menten model and sampled parameters θ in Step 1 by running the RK4 solver.
- For all simulated data in Step 2, if $\|v_{\text{real}} - v_{\text{sim}}\|_2^2 < \varepsilon$, then accept θ that was used to simulate v_{sim} .
- Approximate the posterior distribution over the kinetic constants using all accepted θ s.

The benefit of the proposed approach is apparent for two practical reasons. First, instead of collecting multiple measurements, only one enzymatic assay is sufficient. As a result, the whole process can be sped up a couple of times. Second, there is a significant saving of used enzyme and substrate for determining the kinetic constants. This results in lower costs and

shorter time spent on performing experiments. Further, our approach is based on a probabilistic method; therefore, we are able to provide the mean value with an uncertainty estimate, for example, by using the standard deviation.

Discussion

In our first experiment, we aimed at verifying whether it is possible to correctly determine the kinetic constants using a single observation. For this purpose, we generated synthetic data as described in Materials and methods. In order to quantify the performance of our approach, we used the deviation and accuracy. We expressed both metrics as a percentage. Ideally, the deviation should be equal 0% while the accuracy should be equal 100%.

The results of simulations (1000 simulations per single value of S_0) are presented in Figure 3 and Table A1 (see Appendix 1). First we noticed that the average deviation for k_{cat} is low ($\sim 1\text{--}2\%$). Similarly, for K_M , the deviation is also low ($\sim 5\text{--}6\%$). These results indicate that the proposed probabilistic approach allows to obtain good estimates of the real parameters. Moreover, in more than 90% of cases, the




Fig. 2. The three steps of the proposed approach: (1) Sample parameters (k_{cat} and K_M) from the uniform prior. (2) Simulate data using the Michaelis–Menten model and sampled parameters. If the distance from the simulated data, v_{sim} , to the real data, v_{real} , is smaller than a prespecified threshold, ε , accept the parameters. (3) Calculate the approximate posterior using the accepted samples of the parameters.

real value is within one standard deviation from the estimated mean value. The remaining cases might be concerning; however, the deviation metric indicates that the uncertainty interval was simply very narrow and this could be alleviated by taking slightly larger ε . We also experimented with taking three standard deviations and then the accuracy was 100%. The obtained results confirm that it is indeed possible to find very accurate estimate of the kinetic constants using a single enzymatic assay. Our second observation from this experiment is that there are preferred values of S_0 between and where our method attains the best scores of both metrics for k_{cat} and K_M . We believe that this effect could be explained as follows. If we look at the

Michaelis–Menten plot (Figure 1), taking too large value of S would result in a value of the rate close to the maximal rate V_{max} . In such case we can easily determine k_{cat} , but we have less information about K_M . On the other hand, taking too small value of S_0 would be not sufficiently informative in respect to k_{cat} that is determined by the $V_{\text{max}}/2$. The ideal value would be around the substrate concentration for which the rate is close to V_{max} . This tells us most about K_M and k_{cat} . Obviously, finding such value of S might be challenging; however, the values around 100–200 seem to be sufficient in most cases. We will take advantage of this fact in the experiment on real enzymes.




Fig. 3. The results (as a percentage) of the accuracy and deviation metrics on the synthetic data. The solid and dashed lines denote mean values and the shaded area corresponds to the standard error. (left) The accuracy metric for k_{cat} and varying S_0 . (middle) The accuracy metric for K_M and varying S_0 . (right) The deviation metric for k_{cat} and K_M , and varying S_0 .




Fig. 4. The estimation of the kinetic constants using our approach (the circle represents the mean value and bars corresponds to one standard deviation calculated using the ensemble of parameter samples returned by the ABC procedure) and the standard approach (the red rectangle). The results for k_{cat} (left). The results for K_M (right).

We also inspected whether there is a dependency between a range of values of k_{cat} and K_M and the considered metric. We divided values of both parameters into four bins each and checked whether obtained values of metrics depend on the values of the parameters. Our conclusion is that there is no evident dependency between the values of parameters and the values of the metrics.

In the second experiment, we used data containing the progression of reaction catalyzed by enzyme from our previous study [31,32]. We considered a single substrate concentration assay for the ABC. The data involved experiments on isolated proteolytic enzymes for which the assumption of irreversibility of released product is fulfilled. We evaluated our approach on data from three real enzymes: *hAPN*, *ssAPN*, and *hERAP2* [31,32].

The results for our method and the standard approach are presented in Figure 4 and Table A2 (see Appendix 1). In our method, we utilized a single enzymatic assay for $S_0 = 200$ since for this value we got the best results in the previous experiment. Interestingly, the results obtained using the standard approach and our approach do not differ too much. However, it is a known fact that the standard approach tends to under- or overestimate the true value [2] and, thus, it should be treated as a reference point rather than a real value. Nevertheless, the differences between both methods are rather small that allows us to conclude that our approach provides good estimates. Importantly, our method needs only one enzymatic assay while the standard procedure requires at least couple (e.g., eight) of them.

Conclusions

Application of statistical computational methods to biochemistry, biology, and chemistry allows to understand the underlying phenomena but they could also assist researchers in their daily routine. In this paper, we proposed a probabilistic method for determining the kinetic constants in the Michaelis–Menten model.

The advantage of applying the ABC rejection algorithm to this biochemistry relevant model is the possibility of using only one enzymatic assay. The single enzymatic assay provides information about progression of the reaction catalyzed by enzyme at one initial substrate concentration and we claim that it is sufficient to properly estimate the kinetic parameters. In comparison to the standard procedure that requires at least several measurements, our method notably reduces the whole research process. The obtained results on both synthetic and real data provide empirical evidence in favor of our claims. In our future works, we aim at focusing on more sophisticated ABC procedures [16,17,36] that could improve uncertainty estimation and sampling efficiency. Another interesting topic for future research is to develop new quality measures of the ABC methods. Finally, encouraged by the obtained results, we find investigating other kinetics models as a challenging future research direction.

Acknowledgements

EW-T is financed by a grant within Mobilnosc Plus V from the Polish Ministry of Science and Higher Education (Grant No. 1639/MOB/V/2017/0).

Author contributions

JMT designed and implemented the method, and carried out experiments *in silico*. EW-T formulated the problem, proposed and designed the project, and prepared the data. JMT and EW-T wrote the manuscript.

References

- 1 Ariyawansa R, Basnayake B, Karunarathna A and Mowjood M (2018) Extensions to Michaelis–Menten kinetics for single parameters. *Sci Rep* **8**, 16586.

- 2 Copeland RA (2004) *Enzymes: A Practical Introduction to Structure, Mechanism, and Data Analysis*. John Wiley & Sons, Hoboken, NJ.
- 3 Menten L and Michaelis M (1913) Die Kinetik der Invertinwirkung. *Biochem Z* **49**, 333–369.
- 4 Drag M and Salvesen GS (2010) Emerging principles in protease-based drug discovery. *Nat Rev Drug Discov* **9**, 690.
- 5 Qian H and Elson EL (2002) Single-molecule enzymology: stochastic Michaelis-Menten kinetics. *Biophys Chem* **101**, 565–576.
- 6 Penkler G, Du Toit F, Adams W, Rautenbach M, Palm DC, Van Niekerk DD and Snoep JL (2015) Construction and validation of a detailed kinetic model of glycolysis in *Plasmodium falciparum*. *FEBS J* **282**, 1481–1511.
- 7 Teusink B, Passarge J, Reijenga CA, Esgalhado E, Van derWeijden CC, Schepper M, Walsh MC, Bakker BM, vanDam K, Westerhoff HV et al. (2000) Can yeast glycolysis be understood in terms of in vitro kinetics of the constituent enzymes? Testing biochemistry. *Eur J Biochem* **267**, 5313–5329.
- 8 Henri V (1903) *Lois générales de l'action des diastases*. Librairie Scientifique A. Hermann, Paris.
- 9 Johnson KA and Goody RS (2011) The original Michaelis constant: translation of the 1913 Michaelis–Menten paper. *Biochemistry* **50**, 8264–8269.
- 10 Lietman PS, Chiba K, Ishizaki T, Miura H and Minagawa K (1980) Michaelis–Menten pharmacokinetics of diphenylhydantoin and application in the pediatric age patient. *J Pediatr* **96**, 479–484.
- 11 Jones AW (2010) Evidence-based survey of the elimination rates of ethanol from blood with applications in forensic casework. *Forensic Sci Int* **200**, 1–20.
- 12 Yu RC and Rappaport SM (1997) A lung retention model based on Michaelis-Menten-like kinetics. *Environ Health Perspect* **105**, 496–503.
- 13 Abedon ST (2009) Kinetics of phage-mediated biocontrol of bacteria. *Foodborne Pathog Dis* **6**, 807–815.
- 14 Liepe J, Kirk P, Filippi S, Toni T, Barnes CP and Stumpf MP (2014) A framework for parameter estimation and model selection from experimental data in systems biology using approximate Bayesian computation. *Nat Protoc* **9**, 439–456.
- 15 Lintusaari J, Gutmann MU, Dutta R, Kaski S and Corander J (2017) Fundamentals and recent developments in approximate Bayesian computation. *Syst Biol* **66**, e66–e82.
- 16 Meeds E and Welling M (2014) GPS-ABC: Gaussian process surrogate approximate Bayesian computation. Uncertainty in Artificial Intelligence. AUAI Press. pp. 593–602.
- 17 Park M, Jitkrittum W and Sejdinovic D (2016) K2-ABC: Approximate Bayesian Computation with Kernel embeddings, Proceedings of the 19th International Conference on Artificial Intelligence and Statistics, PMLR 51: 398–407.
- 18 Toni T, Welch D, Strelkowa N, Ipsen A and Stumpf MP (2008) Approximate Bayesian computation scheme for parameter inference and model selection in dynamical systems. *J R Soc Interface* **6**, 187–202.
- 19 Tanaka MM, Francis AR, Luciani F and Sisson S (2006) Using Approximate Bayesian computation to estimate tuberculosis transmission parameters from genotype data. *Genetics* **173**, 1511–1520.
- 20 Liepe J, Barnes C, Cule E, Erguler K, Kirk P, Toni T and Stumpf MP. (2010) ABC-SysBio—approximate Bayesian computation in Python with GPU support. *Bioinformatics* **26**, 1797–1799.
- 21 Tavaré S, Balding DJ, Griffiths RC and Donnelly P (1997) Inferring coalescence times from DNA sequence data. *Genetics* **145**, 505–518.
- 22 Beaumont MA (2010) Approximate Bayesian computation in evolution and ecology. *Annu Rev Ecol Evol Syst* **41**, 379–406.
- 23 Marttinen P, Croucher NJ, Gutmann MU, Corander J and Hanage WP (2015) Recombination produces coherent bacterial species clusters in both core and accessory genomes. *Microbial Genomics* **1**, e000038.
- 24 Currat M and Excoffier L (2004) Modern humans did not admix with Neanderthals during their range expansion into Europe. *PLoS Biol* **2**, e421.
- 25 Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC and Foll M (2013) Robust demographic inference from genomic and SNP data. *PLoS Genet* **9**, e1003905.
- 26 Fagundes NJ, Ray N, Beaumont M, Neuenschwander S, Salzano FM, Bonatto SL and Excoffier L (2007) Statistical evaluation of alternative models of human evolution. *Proc Natl Acad Sci USA* **104**, 17614–17619.
- 27 Itan Y, Powell A, Beaumont MA, Burger J and Thomas MG (2009) The origins of lactase persistence in Europe. *PLoS Comput Biol* **5**, e1000491.
- 28 Gupta S, Hainsworth L, Hogg J, Lee R and Faeder J (2018) Evaluation of parallel tempering to accelerate Bayesian parameter estimation in systems biology. In the 26th Euromicro International Conference on Parallel, Distributed and Network-based Processing (PDP) (Merelli I, Lio P and Kotenko I eds), pp. 690–697. IEEE Computer Society, Cambridge.
- 29 Drovandi CC, Grazian C, Mengersen K and Robert C (2018) Approximating the likelihood in approximate Bayesian computation. *arXiv* 180306645 [PREPRINT].
- 30 Ong VM, Nott DJ, Tran MN, Sisson SA and Drovandi CC (2018) Variational Bayes with synthetic likelihood. *Stat Comput* **28**, 971–988.

- 31 Vassiliou S, Węglarz-Tomczak E, Berlicki L, Pawelczak M, Nocek B, Mulligan R, Joachimiak A and Mucha A (2014) Structure-guided, single-point modifications in the phosphinic dipeptide structure yield highly potent and selective inhibitors of neutral aminopeptidases. *J Med Chem* **57**, 8140–8151.
- 32 Węglarz-Tomczak E, Vassiliou S and Mucha A (2016) Discovery of potent and selective inhibitors of human aminopeptidases ERAP1 and ERAP2 by screening libraries of phosphorus-containing amino acid and dipeptide analogues. *Bioorg Med Chem Lett* **26**, 4122–4126.
- 33 Koudriavtsev AB, Jameson RF and Linert W (2011) *The Law of Mass Action*. Springer Science & Business Media, Berlin.
- 34 Briggs GE and Haldane JBS (1925) A note on the kinetics of enzyme action. *Biochem J* **19**, 338.
- 35 Barnes CP, Silk D, Sheng X and Stumpf MP (2011) Bayesian design of synthetic biological systems. *Proc Natl Acad Sci USA* **108**, 15190–15195.
- 36 Lintusaari J, Vuollekoski H, Kangasrääsio A, Skytén K, Järvenpää M, Marttinen P, Gutmann MU, Vehtari A, Corander J and Kaski S (2018) ELFI: engine for likelihood-free inference. *J Mach Learn Res* **19**, 643–649.

Appendix 1

Table A1. Detailed results of Deviation (\downarrow) and Accuracy (\uparrow) for the synthetic data. An average and a standard error are provided for different values of the initial substrate concentration.

S_0	Deviation		Accuracy	
	k_{cat}	K_M	k_{cat}	K_M
5	5.98 ± 0.30	6.39 ± 0.31	85.20 ± 1.12	85.40 ± 1.12
10	5.20 ± 0.31	5.80 ± 0.35	85.10 ± 1.13	85.70 ± 1.11
50	2.64 ± 0.21	4.61 ± 0.37	90.40 ± 0.93	90.60 ± 0.92
100	1.99 ± 0.16	5.26 ± 0.41	90.10 ± 0.94	90.50 ± 0.93
200	1.42 ± 0.11	5.68 ± 0.43	90.90 ± 0.91	92.10 ± 0.85
300	1.24 ± 0.10	6.44 ± 0.47	89.60 ± 0.97	90.30 ± 0.94
400	1.38 ± 0.09	8.94 ± 0.56	87.20 ± 1.06	88.50 ± 1.01

Table A2. Detailed results of estimated k_{cat} and K_M for the real data. An average and one standard deviation are provided for the standard method and our approach.

	k_{cat}		K_M	
	Standard	Ours	Standard	Ours
hERAP2	0.388	0.426 ± 0.078	91	110.796 ± 56.927
hAPN	5.5	6.353 ± 1.165	88	111.137 ± 56.692
ssAPN	34	41.118 ± 7.863	120	102.252 ± 56.918