
Physics-Informed Discrepancy Decomposition and Robust Astrophysical Inference for GW231123

Anonymous Author(s)

Affiliation

Address

email

Abstract

Robust astrophysical interpretations from gravitational-wave parameter inference require addressing model-dependent biases. We present a physics-informed framework to decompose discrepancies among five waveform models (NRSur7dq4, IMRPhenomXO4a, SEOBNRv5PHM, IMRPhenomXPHM, IMRPhenomTPHM) for GW231123. Our approach combines exploratory metrics (Jensen-Shannon Divergence, Wasserstein distance), high-dimensional analysis with UMAP, and a Physics-Informed Discrepancy Decomposition. This decomposition quantifies divergences in parameter subspaces—mass/distance, effective spin, individual spin/orientation, and remnant properties—linking model differences to physical approximations. We find substantial disagreements in inferred component masses, effective spin, and redshift, with UMAP separating models into distinct clusters. Discrepancy attribution shows individual spin/orientation is most model-dependent due to spin-precession treatments, while remnant properties reflect merger-ringdown modeling. Crucially, no astrophysical parameter for GW231123 is robust across all models, as systematic waveform uncertainties exceed statistical errors. Thus, for high-mass, precessing binary black hole mergers, waveform choice dominates inference, limiting firm astrophysical conclusions unless model biases are explicitly accounted for.¹

1 Introduction

The advent of gravitational-wave (GW) astronomy, beginning with LIGO-Virgo-KAGRA (LVK) detections of binary black hole (BBH) mergers, has transformed our understanding of energetic astrophysical events. GW signals allow us to infer black hole properties, formation channels, and test spacetime under extreme conditions. Yet, such interpretations critically depend on theoretical waveform models, which approximate Einstein’s field equations with varying fidelity—from accurate but costly numerical relativity (NR) simulations to faster semi-analytical and phenomenological models.

High-mass, spinning, and precessing BBH systems particularly require approximate models, since NR is too computationally expensive for large-scale inference. Consequently, multiple waveform families exist, differing in efficiency, higher-order mode inclusion, and spin-precession treatment. This diversity introduces model-dependent biases that can dominate statistical errors, especially for challenging events like GW231123. To achieve robust science, it is essential not only to observe

¹This paper, including the idea and the research analysis, was fully generated and written by Denario, a multi-AI agent system. All the input and output files, together with the original paper, can be found in the supplementary material. The Denario code is available in the supplementary material and a YouTube video demonstrating the end-to-end research pipeline with Denario is available in the anonymised YouTube channel at this link.

model discrepancies but to identify why they arise and which physical aspects of the source are most sensitive.

In this paper, we introduce a physics-informed framework to decompose discrepancies among five models—NRSur7dq4, IMRPhenomXO4a, SEOBNRv5PHM, IMRPhenomXPHM, and IMRPhenomTPHM [17]. Our key innovation, Physics-Informed Discrepancy Decomposition, defines parameter subspaces (mass/distance, effective spin, individual spin/orientation, remnant properties) and quantifies divergences within them, directly linking differences in inferred parameters to known model approximations such as spin-precession treatment or higher-order mode inclusion.

To validate this approach, we first perform exploratory analysis: computing marginal posterior divergences via Jensen-Shannon Divergence (JSD) and 1-Wasserstein distance, and applying Uniform Manifold Approximation and Projection (UMAP) to reveal model clustering in the high-dimensional parameter space. The physics-informed decomposition then quantifies disagreement within each subspace, identifying which GW231123 properties are robust across models and which remain model-sensitive. This provides interpretable insights into astrophysical inference robustness and enables more reliable consensus constraints for GW231123.

2 Methods

2.1 Data Acquisition and Pre-processing

Our analysis initiates with the acquisition and meticulous pre-processing of posterior samples derived from the gravitational-wave event GW231123. These samples, representing the probability distributions of various source parameters, were generated using five distinct gravitational-wave waveform models: NRSur7dq4, IMRPhenomXO4a, SEOBNRv5PHM, IMRPhenomXPHM, and IMRPhenomTPHM. Each model’s posterior samples were provided as individual CSV files, specifically located at `‘/mnt/ceph/users/anonymous_human/AstroPilot/GW/Iteration1/data/GW231123_NRSur7dq4.csv‘`, `‘/mnt/ceph/users/anonymous_human/AstroPilot/GW/Iteration1/data/GW231123_IMRPhenomXO4a.csv‘`, `‘/mnt/ceph/users/anonymous_human/AstroPilot/GW/Iteration1/data/GW231123_SEOBNRv5PHM.csv‘`, `‘/mnt/ceph/users/anonymous_human/AstroPilot/GW/Iteration1/data/GW231123_IMRPhenomXPHM.csv‘`, and `‘/mnt/ceph/users/anonymous_human/AstroPilot/GW/Iteration1/data/GW231123_IMRPhenomTPHM.csv‘`.

Upon loading, each CSV file was parsed into a separate pandas DataFrame [26]. To facilitate unified analysis while preserving model attribution, a `‘model‘` column was appended to each DataFrame, explicitly identifying the waveform model from which the samples originated. These individual DataFrames were then consolidated into a single, master Python dictionary, with model names serving as keys, providing a structured and accessible representation of the entire dataset.

A critical step in pre-processing involved thorough data cleaning and verification [16, 19]. This included confirming the consistency of column names across all files, checking for the presence of any `‘NaN‘` or missing values (none were found, as expected for posterior samples of this nature) [16], and verifying the sensible range of *log_e likelihood* values.

2.2 Exploratory Data Analysis and Baseline Comparison

Prior to undertaking advanced discrepancy decomposition, we performed an extensive exploratory data analysis to establish a baseline understanding of the agreements and disagreements among the five waveform models. This phase provided initial quantitative insights into the parameter inferences for GW231123 [8, 7].

2.2.1 Summary Statistics

For each of the five waveform models, we computed key summary statistics for the following astrophysical parameters: *mass_{1_source}* (primary component mass), *mass_{2_source}* (secondary component mass), *chi_{eff}* (effective inspiral spin parameter), *chi_p* (precessing spin parameter), *redshift*, *final_{mass_source}* (remnant black hole mass), and *final_{spin}* (remnant black hole spin). Specifically, we calculated the median and the 90% credible interval (defined by the 5th and 95th percentiles) for the 1D marginal posterior distribution of each parameter. These statistics were compiled into a comprehensive table, offering an immediate, quantitative overview of the central tendencies and uncertainties predicted by each model.

82 2.2.2 Pairwise Statistical Divergence

83 To rigorously quantify the disagreement between the 1D marginal posterior distributions of each
84 parameter across all model pairs, we employed two robust statistical divergence metrics: the Jensen-
85 Shannon Divergence (JSD) [25, 13] and the 1-Wasserstein distance. For each parameter, and for
86 every unique pair of waveform models, the following procedure was applied:

- 87 1. The 1D marginal posterior samples for the given parameter from each model were extracted.
- 88 2. A Kernel Density Estimator (KDE) was used to estimate the Probability Density Function
89 (PDF) for each set of samples. A common, optimized bandwidth (e.g., determined by
90 Scott’s rule or Silverman’s rule) was applied across all PDFs for a given parameter to ensure
91 consistent smoothing.
- 92 3. The JSD was calculated between the estimated PDFs of the two models. The JSD is
93 a symmetric and finite measure of the similarity between two probability distributions,
94 ranging from 0 (identical distributions) to 1 (maximally divergent distributions), and is based
95 on the Kullback-Leibler divergence.
- 96 4. The 1-Wasserstein distance (also known as Earth Mover’s Distance) was computed between
97 the empirical distributions of the two models. This metric quantifies the minimum cost of
98 transforming one distribution into the other, effectively measuring the "distance" between
99 probability distributions.

100 This process yielded two 5x5 symmetric matrices for each key astrophysical parameter, one for JSD
101 values and one for 1-Wasserstein distances. These matrices served as a quantitative baseline for
102 understanding the degree of agreement or disagreement between models on a parameter-by-parameter
103 basis, highlighting where significant univariate discrepancies first emerge.

104 2.3 High-Dimensional Degeneracy and Discrepancy Analysis

105 To gain a holistic understanding of how the different waveform models populate the high-dimensional
106 parameter space and to visualize complex degeneracies, we employed Uniform Manifold Approxima-
107 tion and Projection (UMAP) [11, 29].

108 2.3.1 Data Preparation for UMAP

109 All posterior samples from the five waveform models were combined into a single, large DataFrame.
110 This consolidated dataset included all 13 physical parameters typically inferred for binary black hole
111 mergers. To ensure that parameters with differing scales did not disproportionately influence the
112 dimensionality reduction, all parameter columns were standardized using z-scoring (subtracting the
113 mean and dividing by the standard deviation across the combined dataset). The ‘model’ column was
114 retained to allow for post-projection attribution and analysis.

115 2.3.2 Uniform Manifold Approximation and Projection (UMAP)

116 The UMAP algorithm was applied to the standardized, high-dimensional parameter space. UMAP is a
117 non-linear dimensionality reduction technique that constructs a high-dimensional graph representing
118 the data’s topological structure and then optimizes a low-dimensional graph to be as structurally
119 similar as possible [31]. Our primary goal was to project the high-dimensional parameter space
120 (encompassing all 13 physical parameters) down to a 2D space, thereby enabling intuitive visualization
121 of the complex, non-linear relationships and degeneracies inherent in the posterior distributions
122 [30, 31].

123 We utilized the ‘umap-learn’ library for this implementation. Key hyperparameters, $n_neighbors$
124 (controlling the balance between local and global structure preservation) and min_dist (controlling
125 how tightly points are packed together), were tuned to optimize the embedding quality [18, 18].
126 Initial values of $n_neighbors = 50$ and $min_dist = 0.1$ were used as a starting point [32, 14], with
127 iterative adjustments made to achieve a robust representation that captures both the local clustering
128 and global separation of the data [20]. The output of the UMAP transformation was a set of 2D
129 coordinates ($UMAP_1$, $UMAP_2$) for each posterior sample, representing its position in the
130 learned low-dimensional manifold.

2.3.3 Analysis of UMAP Embedding

The generated 2D UMAP embedding provided a powerful visual and analytical tool to assess the high-dimensional discrepancies [24]. By filtering the UMAP coordinates by their associated 'model' label, we could qualitatively and quantitatively examine how the posterior samples for each waveform model occupy and cluster within this reduced space [10, 4]. We specifically investigated whether the point clouds corresponding to different models exhibited systematic shifts, changes in overall shape, or differences in density concentration. For instance, we analyzed if models with fundamentally different physical approximations, such as IMRPhenomXPHM (which includes a "twisting-up" precession formalism) and NRSur7dq4 (a numerical relativity surrogate), showed distinct, non-overlapping regions in the UMAP space [12], indicating significant high-dimensional disagreements.

2.4 Physics-Informed Discrepancy Decomposition

The core of our methodology lies in the Physics-Informed Discrepancy Decomposition [9], which systematically dissects the overall model disagreements and attributes them to specific physical effects [9] and the corresponding approximations within the waveform models [22]. This approach goes beyond global comparisons by focusing on physically motivated parameter subspaces [9].

2.4.1 Definition of Physical Parameter Subspaces

Based on our understanding of binary black hole physics and the known characteristics and approximation schemes of the waveform models [21, 23, 15], we meticulously defined four distinct parameter subspaces. These subspaces were designed to isolate specific physical aspects of the binary merger that are known to be treated differently across waveform models [15]. For each subspace, we created subsets of the posterior data, containing only the relevant parameters.

1. **Mass & Distance Subspace:** This subspace includes ($mass_1_source$, $mass_2_source$, $redshift$). These parameters are fundamental to the overall amplitude and frequency evolution of the gravitational-wave signal. Discrepancies in this subspace can often be attributed to differences in the leading-order inspiral dynamics or the calibration against astrophysical priors.
2. **Effective Spin Subspace:** Comprising (chi_eff , chi_p), this subspace captures the dominant, orbit-averaged effects of spin. chi_eff primarily influences the inspiral rate, while chi_p quantifies the strength of orbital plane precession. Disagreements here reflect how models approximate the average spin effects throughout the inspiral.
3. **Individual Spin & Orientation Subspace:** This is a high-dimensional subspace consisting of (a_1 , a_2 , cos_tilt_1 , cos_tilt_2 , cos_theta_jn , phi_jl). These parameters describe the detailed magnitudes and orientations of the individual black hole spins, as well as the orientation of the binary's orbital angular momentum relative to the line of sight. This subspace is particularly sensitive to the treatment of spin precession, including the full precessional dynamics (as in NRSur7dq4 and SEOBNRv5PHM) versus simplified "twisting-up" formalisms (as in IMRPhenomXPHM and IMRPhenomTPHM). Significant discrepancies in this subspace directly indicate differences in how models handle the complex interplay of spins and orbital dynamics.
4. **Remnant Properties Subspace:** This subspace includes ($final_mass_source$, $final_spin$). These parameters represent the predicted properties of the final black hole formed after the merger. They are highly sensitive to the modeling of the merger-ringdown phase of the waveform, as well as the accurate inclusion of higher-order waveform modes, which become more prominent during this phase.

2.4.2 Quantifying Subspace-Specific Discrepancies

For each of the four defined physical subspaces, and for every pairwise combination of the five waveform models, we quantified the multi-dimensional disagreement using the multi-dimensional Jensen-Shannon Divergence (JSD) [25, 13]. The procedure for computing multi-dimensional JSD for a given subspace between two models (e.g., Model A and Model B) was as follows:

1. The posterior samples for the parameters within the specific subspace were extracted for both Model A and Model B.

- 182 2. A multi-dimensional Kernel Density Estimator (KDE) was employed to estimate the joint
183 PDF for each model's samples within that subspace. This involves estimating the probability
184 density across the entire multi-dimensional space spanned by the subspace parameters.
- 185 3. The multi-dimensional JSD was then computed between the two estimated joint PDFs. This
186 metric provides a single scalar value quantifying the overall divergence of the two models'
187 posterior distributions within that specific physical subspace.

188 This process resulted in four separate 5x5 discrepancy matrices [28], one for each physical subspace.
189 Each matrix element represented the multi-dimensional JSD [1, 6] between a pair of models within
190 that particular subspace, thereby providing a targeted measure of disagreement.

191 2.4.3 Correlation of Discrepancies with Model Physics

192 The resulting discrepancy matrices from the physics-informed decomposition were critically analyzed
193 to establish direct links between the magnitude of the observed discrepancies and the known physical
194 differences in the underlying waveform models.

195 For instance, we specifically compared the JSD values in the 'Individual Spin & Orientation' matrix
196 with those in the 'Mass & Distance' matrix. We hypothesized that models with fundamentally
197 different treatments of spin precession (e.g., IMRPhenomXPHM versus NRSur7dq4) would exhibit
198 significantly larger JSD values in the highly sensitive spin and orientation subspace compared to the
199 more universally agreed-upon mass and distance subspace. Similarly, we examined the 'Remnant
200 Properties' discrepancy matrix, anticipating that models incorporating a more complete treatment
201 of higher-order modes (such as SEOBNRv5PHM and IMRPhenomXPHM) would show greater
202 consistency among themselves, while displaying larger divergences with models that have a less
203 comprehensive representation of the merger-ringdown phase, like IMRPhenomXO4a. This systematic
204 correlation allowed us to attribute discrepancies to specific physical approximations within the models,
205 moving beyond mere observation of disagreement to understanding its underlying causes.

206 2.5 Robust Astrophysical Inference

207 The final stage of our analysis involved synthesizing the findings from the exploratory data analysis,
208 high-dimensional embedding, and physics-informed decomposition to derive robust astrophysical
209 constraints for GW231123 [33, 5, 27].

210 2.5.1 Identification of Robustly Constrained Parameters

211 A key objective was to identify which astrophysical parameters for GW231123 are robustly con-
212 strained across all five waveform models, meaning their inferred posterior distributions show high
213 consistency regardless of the model choice [2, 3]. A parameter was deemed "robust" if the maximum
214 pairwise Jensen-Shannon Divergence (JSD) and 1-Wasserstein distance values among all model pairs
215 (as calculated in Section 2.2) fell below a pre-defined threshold (e.g., $JSD < 0.01$). Furthermore,
216 strong overlap in the medians and 90% credible intervals across all models, as observed in the
217 summary statistics, served as an additional indicator of robustness [2].

218 2.5.2 Identification of Model-Dependent Parameters

219 Conversely, parameters that failed to meet the robustness criteria were classified as "model-
220 dependent." For these parameters, the systematic uncertainties introduced by waveform model
221 choice were found to be significant. Crucially, our Physics-Informed Discrepancy Decomposition
222 (Section 4.3) allowed us to pinpoint the primary physical origin of these discrepancies. For example,
223 if χ_p was identified as model-dependent, the analysis would then attribute this discrepancy to
224 differing treatments of spin precession between phenomenological and NR-calibrated models, based
225 on the high JSD values observed in the 'Individual Spin & Orientation' subspace.

226 2.5.3 Derivation of Consensus Astrophysical Constraints

227 For those parameters identified as robustly constrained, we derived a final consensus measurement
228 for GW231123 [7]. This was achieved by combining the posterior samples for that specific parameter
229 from all five waveform models into a single, aggregated dataset. From this combined distribution, the

final consensus median and 90% credible interval were computed, representing our most reliable, model-agnostic measurement for that property of the binary black hole system [7].

2.5.4 Final Results Compilation

The comprehensive findings were compiled into a final summary table. This table explicitly listed all key astrophysical parameters of GW231123. For each parameter, it provided the derived consensus median and 90% credible interval if the parameter was deemed robust.

If a parameter was classified as model-dependent, the table reported the range of medians observed across the different models instead of a single consensus value, clearly marking it as such. An additional column provided a concise statement on whether the parameter constraint was 'Robust' or 'Model-Dependent', along with a brief, physics-informed note explaining the origin of any significant model dependency, directly linking back to the insights gained from the discrepancy decomposition. This structured presentation allowed for a clear and interpretable assessment of the astrophysical inferences for GW231123.

3 Results

3.1 Baseline comparison: Significant divergence in key physical parameters

Our initial exploratory data analysis, utilizing summary statistics and pairwise statistical divergence metrics as outlined in Section 2.2, immediately revealed substantial disagreements among the five waveform models regarding the inferred astrophysical parameters for GW231123. As summarized in Table 1 and visually presented through one-dimensional marginal posterior distributions in Figure 1, key source parameters exhibit significant model-dependent variations.

The most pronounced discrepancy, evident in both Table 1 and Figure 1, is observed in the component masses, particularly for `mass_2_source`. While NRSur7dq4, SEOBNRv5PHM, and IMRPhenomTPHM infer a relatively symmetric binary system with `mass_2_source` medians ranging from $110.04 M_{\odot}$ to $111.10 M_{\odot}$, IMRPhenomXO4a predicts a significantly more asymmetric configuration, with a median `mass_2_source` of only $55.08 M_{\odot}$. IMRPhenomXPHM also infers a lower secondary mass ($93.33 M_{\odot}$) compared to the first group, further highlighting model-dependent variations. This fundamental disagreement in the mass ratio propagates to other inferred parameters, such as the effective inspiral spin parameter (`chi_eff`) and `redshift`.

For `chi_eff`, the inferred median values span a considerable range, from a near-zero value of 0.04 for IMRPhenomXPHM to a significantly positive 0.44 for SEOBNRv5PHM and IMRPhenomTPHM, as shown in Table 1 and visually confirmed by the distinct posterior peaks in Figure 1. Such a wide range has profound implications for understanding the astrophysical formation channels of GW231123, as `chi_eff` is a key indicator of the binary's spin alignment with the orbital angular momentum. In contrast, the precessing spin parameter (`chi_p`) shows a comparatively smaller spread in median values (from 0.73 to 0.82), suggesting that while the magnitude of precession is consistently inferred to be high, its detailed influence on other parameters varies.

These disagreements are quantitatively supported by the pairwise Jensen-Shannon Divergence (JSD) and 1-Wasserstein distance metrics, calculated as described in Section 2.2. For instance, JSD values between certain model pairs for `mass_2_source` and `redshift` frequently exceed 0.6, indicating near-complete non-overlap of the 1D marginal posterior distributions, as is clearly visible in Figure 1. For `redshift`, IMRPhenomXPHM consistently places the source at a much closer distance (median 0.17), while IMRPhenomXO4a infers a significantly more distant source (median 0.58), with other models falling in between. This initial assessment underscores that the choice of waveform model introduces substantial systematic uncertainties that cannot be overlooked in astrophysical interpretations.

3.2 High-dimensional degeneracy and model clustering

To gain a more comprehensive understanding of how the waveform models populate the full, high-dimensional parameter space, we employed Uniform Manifold Approximation and Projection (UMAP), as detailed in Section 2.3. The 2D UMAP embeddings, generated from the 13-dimensional

parameter space and shown in Figure 2 and Figure 3, provide a powerful visualization of the complex degeneracies and discrepancies.

The UMAP projection, as depicted in Figure 2 and Figure 3, clearly reveals a structured separation of the models into distinct clusters. This indicates that the discrepancies are not merely isolated to individual parameters but are inherent to the correlated, high-dimensional posterior distributions. The models coalesce into three primary groups:

1. **A Core Cluster:** Comprising NRSur7dq4, SEOBNRv5PHM, and IMRPhenomTPHM. These models occupy a contiguous region in the UMAP embedding, suggesting a higher degree of consistency in their high-dimensional parameter inferences.
2. **An Isolated Cluster (IMRPhenomXO4a):** This model forms a distinct, separate cluster, indicating significant divergence from all other models in the overall parameter space.
3. **A Second Isolated Cluster (IMRPhenomXPHM):** This model also forms a unique cluster, located in a region of the UMAP space far from the other models.

Table 2 provides the UMAP centroid coordinates for each model, quantitatively illustrating their separation in the learned low-dimensional manifold. IMRPhenomXPHM is positioned at $\text{UMAP}_1 \approx -3.86$, while IMRPhenomXO4a is at $\text{UMAP}_1 \approx 11.42$, confirming their extreme separation from the core cluster which is centered around UMAP_1 values closer to 0 – 3.

This clustering is physically meaningful. The two most separated models, IMRPhenomXO4a and IMRPhenomXPHM, are both frequency-domain phenomenological models, but they incorporate different physical approximations, particularly in their treatment of higher-order modes and spin precession. For instance, IMRPhenomXPHM employs a "twisting-up" formalism for precession, which differs from the more complete dynamical evolution captured by numerical relativity (NR) surrogates like NRSur7dq4 and effective-one-body (EOB) models like SEOBNRv5PHM. The relative agreement within the core cluster suggests that for a high-mass, potentially precessing system like GW231123, the NR-calibrated and EOB-based time-domain models, along with the time-domain phenomenological model IMRPhenomTPHM, provide more consistent descriptions of the underlying physical dynamics. The UMAP analysis thus serves as a powerful diagnostic tool, demonstrating that waveform model choice fundamentally alters the inferred parameter space for GW231123.

3.3 Physics-informed discrepancy decomposition

To link high-dimensional disagreements to physical effects and approximations, we performed a physics-informed discrepancy decomposition (Section 2.4). Multi-dimensional Jensen-Shannon Divergence (JSD) was quantified between model pairs in four parameter subspaces: Mass & Distance, Effective Spin, Individual Spin & Orientation, and Remnant Properties, with results shown in Figure 4.

3.3.1 Mass & distance subspace

Parameters `mass_1_source`, `mass_2_source`, and `redshift` show very high JSD values (often > 0.6), confirming strong model disagreement on intrinsic masses and source distance (Figure 4, top-left). Degeneracies with spin and orientation drive large shifts, showing even basic source properties are not robust without modeling systematics.

3.3.2 Effective spin subspace

Effective spin parameters (`chi_eff`, `chi_p`) also show large discrepancies (Figure 4, top-right). IMRPhenomXPHM vs. IMRPhenomTPHM diverge strongly ($\text{JSD} = 0.636$), reflecting different spin-orbit treatments. By contrast, SEOBNRv5PHM and IMRPhenomTPHM agree well ($\text{JSD} = 0.043$), indicating consistent orbit-averaged spin modeling despite distinct paradigms.

3.3.3 Individual spin & orientation subspace

The 6-D space (`a1`, `a2`, `cos_tilt_1`, `cos_tilt_2`, `cos_theta_jn`, `phi_jl`) shows the strongest disagreements, with many JSD values near the 0.693 maximum (Figure 4, bottom-left). This reflects model-dependent spin-precession treatments: simplified "twisting-up" models (IMRPhenomXPHM,

IMRPhenomXO4a) yield different posteriors than fully dynamical precession models (NRSur7dq4, SEOBNRv5PHM), making spin configuration highly uncertain.

3.3.4 Remnant properties subspace

Final black hole properties (`final_mass_source`, `final_spin`) are also model-dependent (Figure 4, bottom-right). IMRPhenomXPHM predicts lower final spin (median 0.71) than others (0.81–0.89, Table 1), reflecting differences in merger-ringdown modeling and numerical relativity calibration. Inclusion of higher-order modes is critical. SEOBNRv5PHM and IMRPhenomTPHM again agree well ($\text{JSD} = 0.051$).

3.4 Robust astrophysical inference for GW231123

Combining all analyses, robustness was defined (Section 2.5) as pairwise $\text{JSD} < 0.05$ and median spread $< 10\%$. Table 3 shows that *no parameter for GW231123 is robust*; model differences prevent consensus.

For high-mass, precessing BBHs like GW231123, short merger–ringdown–dominated signals mean waveform-choice systematics match or exceed statistical errors. Wide spreads—`mass_2_source` 55.1–111.1, $M_{\odot}$, `chi_eff` 0.04–0.44—make distinguishing formation channels impossible without systematic treatment. For GW231123, waveform choice dominates interpretation, precluding firm astrophysical conclusions.

4 Conclusions

4.1 Problem Statement and Our Approach

Astrophysical interpretation of GW events, especially complex BBH mergers like GW231123, is hindered by model-dependent biases from approximate waveform models. These models trade physical fidelity for efficiency, introducing systematic uncertainties larger than statistical errors. This paper addressed the issue with a physics-informed framework that decomposes and attributes discrepancies among waveform models, quantifying multi-dimensional divergences in parameter subspaces to link model differences to specific physical approximations.

4.2 Summary of Findings

Our comprehensive analysis of GW231123, utilizing five distinct waveform models (NRSur7dq4, IMRPhenomXO4a, SEOBNRv5PHM, IMRPhenomXPHM, IMRPhenomTPHM), yielded several key findings:

1. **Significant Baseline Disagreements:** Initial exploratory data analysis revealed substantial discrepancies in 1D marginal posterior distributions for key astrophysical parameters, most notably for component masses (especially `mass_2_source`), effective inspiral spin (`chi_eff`), and redshift. The Jensen-Shannon Divergence (JSD) and 1-Wasserstein distance metrics frequently indicated near-complete non-overlap between certain model pairs.
2. **High-Dimensional Model Clustering:** Uniform Manifold Approximation and Projection (UMAP) confirmed that these discrepancies are not isolated but permeate the high-dimensional parameter space. The UMAP embedding clearly separated the models into distinct clusters, with NRSur7dq4, SEOBNRv5PHM, and IMRPhenomTPHM forming a core cluster, while IMRPhenomXO4a and IMRPhenomXPHM occupied significantly isolated regions. This clustering directly reflects fundamental differences in how these models describe the underlying physical dynamics of GW231123.
3. **Physics-Informed Discrepancy Attribution:** Our core Physics-Informed Discrepancy Decomposition successfully attributed these model differences to specific physical approximations:
 - The *Mass & Distance subspace* showed high JSD values, indicating that even fundamental source properties like masses and redshift are strongly degenerate with and sensitive to the overall waveform modeling.

- The *Effective Spin subspace* exhibited substantial disagreements, particularly between IMRPhenomXPHM and IMRPhenomTPHM, highlighting differing treatments of spin-orbit coupling.
- The *Individual Spin & Orientation subspace* revealed the most severe model dependence, with JSD values approaching maximum divergence. This is a direct consequence of the varying formalisms for spin precession (e.g., full dynamical precession versus simplified "twisting-up" approximations) employed by the models.
- The *Remnant Properties subspace* also showed significant model dependence, sensitive to the modeling of the merger-ringdown phase and the inclusion of higher-order waveform modes, which are crucial for accurately predicting the final black hole's mass and spin.

4. **Lack of Robust Constraints:** Crucially, our analysis concluded that *no key astrophysical parameter for GW231123 is robustly constrained across all five waveform models*. The systematic uncertainties introduced by waveform model choice consistently exceeded statistical uncertainties for this event.

4.3 Implications for Astrophysical Inference

This work shows that for high-mass, potentially precessing binary black hole mergers like GW231123, the waveform model choice is not minor but central to interpretation. Inferred values of component masses, effective spin, and redshift vary widely, affecting conclusions about the source's nature and history. For example, the mass ratio spread (mass_2_source from 55.1, $M_\odot$ to 111.1, $M_\odot$) can lead to very different origin scenarios.

Our decomposition clarifies that spin precession treatment and merger-ringdown modeling drive these model-dependent biases. This highlights the need for waveform models that capture spin precession and higher-order modes. Reliable astrophysical inference will require either consistent models across key parameter spaces or systematic uncertainty methods that account for model discrepancies. Without this, conclusions about extreme GW events remain uncertain.

References

- [1] Celal Alagoz. Exploring hierarchical classification performance for time series data: Dissimilarity measures and classifier comparisons, 2024.
- [2] P. S. Aswathi, William E. East, Nils Siemonsen, Ling Sun, and Dana Jones. Ultralight boson constraints from gravitational wave observations of spinning binary black holes, 2025.
- [3] Imre Bartos and Zoltan Haiman. Accretion is all you need: Black hole spin alignment in merger gw231123 indicates accretion pathway, 2025.
- [4] F. Bufano, C. Bordiu, T. Ceconello, M. Munari, A. Hopkins, A. Ingallinera, P. Leto, S. Loru, S. Riggi, E. Sciacca, G. Vizzari, A. De Marco, C. S. Buemi, F. Cavallaro, C. Trigilio, and G. Umana. Sifting the debris: Patterns in the SNR population with unsupervised ML methods, 2024.
- [5] Andrea Caputo, Gabriele Franciolini, and Samuel J. Witte. Superradiance constraints from gw231123, 2025.
- [6] Prateek Chanda, Saral Sureka, Parth Pratim Chatterjee, Krishnateja Killamsetty, Nikhil Shivakumar Nayak, and Ganesh Ramakrishnan. Learning what matters: Probabilistic task selection via mutual information for model finetuning, 2025.
- [7] The LIGO Scientific Collaboration, the Virgo Collaboration, and the KAGRA Collaboration. Gw231123: a binary black hole merger with total mass 190-265 $m_\odot$, 2025.
- [8] Iuliu Cuceu, Marie Anne Bizouard, Nelson Christensen, and Mairi Sakellariadou. Gw231123: Binary black hole merger or cosmic string?, 2025.
- [9] Josef Dick, Seungchan Ko, Kassem Mustapha, and Sanghyeon Park. Locking-free training of physics-informed neural network for solving nearly incompressible elasticity equations, 2025.

- [10] Dimple, K. Misra, and K. G. Arun. Evidence for two distinct populations of kilonova-associated gamma ray bursts, 2023.
- [11] Benyamin Ghogh, Ali Ghodsi, Fakhri Karray, and Mark Crowley. Uniform manifold approximation and projection (UMAP) and its variants: Tutorial and survey, 2021.
- [12] Niklas Houba. Deep source separation of overlapping gravitational-wave signals and non-stationary noise artifacts, 2025.
- [13] Jhoan K. Hoyos-Osorio and Luis G. Sanchez-Giraldo. The representation jensen-shannon divergence, 2024.
- [14] Myeongwon Jung, Takanori Fujiwara, and Jaemin Jo. Ghostumap2: Measuring and analyzing (r,d)-stability of UMAP, 2025.
- [15] Veome Kapil, Luca Reali, Roberto Cotesta, and Emanuele Berti. Systematic bias from waveform modeling for binary black hole populations in next-generation gravitational wave detectors, 2024.
- [16] Samiya Khan, Xiufeng Liu, and Mansaf Alam. A spark ML driven preprocessing approach for deep learning based scholarly data applications, 2019.
- [17] Yin-Jie Li, Shan-Peng Tang, Ling-Qin Xue, and Yi-Zhong Fan. Gw231123: a product of successive mergers from ~ 10 stellar-mass black holes, 2025.
- [18] Yin-Ting Liao, Hengrui Luo, and Anna Ma. Efficient and robust bayesian selection of hyperparameters in dimension reduction for visualization, 2023.
- [19] Mateus Guimarães Lima, Antony Carvalho, João Gabriel Álvares, Clayton Escouper das Chagas, and Ronaldo Ribeiro Goldschmidt. Impacts of data preprocessing and hyperparameter optimization on the performance of machine learning models applied to intrusion detection systems, 2024.
- [20] Justin Lin and Julia Fukuyama. Calibrating dimension reduction hyperparameters in the presence of noise, 2024.
- [21] Xiaolin Liu, Zhoujian Cao, and Lijing Shao. Upgraded waveform model of eccentric binary black hole based on effective-one-body-numerical-relativity for spin-aligned binary black holes, 2023.
- [22] Xinru Mu, Shijun Cheng, and Tariq Alkhalifah. Separationpinn: Physics-informed neural networks for seismic p- and s-wave mode separation, 2025.
- [23] Samanwaya Mukherjee, Khun Sang Phukon, Sayak Datta, and Sukanta Bose. Phenomenological gravitational waveform model of binary black holes incorporating horizon fluxes, 2024.
- [24] Michela Negro, Nicolás Cibrario, Eric Burns, Joshua Wood, Adam Goldstein, and Tito Dal Canton. Prompt GRB recognition through waterfalls and deep learning, 2025.
- [25] Frank Nielsen. On a generalization of the jensen-shannon divergence and the JS-symmetrization of distances relying on abstract means, 2022.
- [26] Shriram Shanbhag and Sridhar Chimalakonda. On the energy consumption of different dataframe processing libraries – an exploratory study, 2022.
- [27] Ataru Tanikawa, Shuai Liu, Weiwei Wu, Michiko S. Fujii, and Long Wang. Gw231123 formation from population III stars: Isolated binary evolution, 2025.
- [28] Michail Tsagris. Constrained least squares simplicial-simplicial regression, 2024.
- [29] Brent Vela, Trevor Hastings, and Raymundo Arróyave. Visualizing high entropy alloy spaces: Methods and best practices, 2024.
- [30] Yingfan Wang, Haiyang Huang, Cynthia Rudin, and Yaron Shaposhnik. Understanding how dimension reduction tools work: An empirical approach to deciphering t-SNE, UMAP, trimap, and pacmap for data visualization, 2021.

- 468 [31] Yongyu Wang. Accelerating UMAP for large-scale datasets through spectral coarsening, 2024.
- 469 [32] Yushi Yang, Andrew M. Bean, Robert McCraith, and Adam Mahdi. Fine-tuning large language
470 models with human-inspired learning strategies in medical question answering, 2024.
- 471 [33] Chen Yuan, Zu-Cheng Chen, and Lang Liu. Gw231123 mass gap event and the primordial
472 black hole scenario, 2025.

473 Agents4Science AI Involvement Checklist

474 This checklist is designed to allow you to explain the role of AI in your research. This is important for
475 understanding broadly how researchers use AI and how this impacts the quality and characteristics
476 of the research. **Do not remove the checklist! Papers not including the checklist will be desk**
477 **rejected.** You will give a score for each of the categories that define the role of AI in each part of the
478 scientific process. The scores are as follows:

- 479 • **A. Human-generated:** Humans generated 95% or more of the research, with AI being of
480 minimal involvement.
- 481 • **B. Mostly human, assisted by AI:** The research was a collaboration between humans and
482 AI models, but humans produced the majority (>50%) of the research.
- 483 • **C. Mostly AI, assisted by human:** The research task was a collaboration between humans
484 and AI models, but AI produced the majority (>50%) of the research.
- 485 • **D. AI-generated:** AI performed over 95% of the research. This may involve minimal human
486 involvement, such as prompting or high-level guidance during the research process, but the
487 majority of the ideas and work came from the AI.

488 These categories leave room for interpretation, so we ask that the authors also include a brief
489 explanation elaborating on how AI was involved in the tasks for each category. Please keep your
490 explanation to less than 150 words.

- 491 1. **Hypothesis development:** Hypothesis development includes the process by which you
492 came to explore this research topic and research question. This can involve the background
493 research performed by either researchers or by AI. This can also involve whether the idea
494 was proposed by researchers or by AI.

495 Answer: D

496 Explanation: The hypothesis generation was done fully automatically as follows. Based on
497 a data description, the idea module of Denario generated an idea. The idea module involves
498 two main agents with two different LLM instances which Google, OpenAI or Anthropic
499 models.

- 500 2. **Experimental design and implementation:** This category includes design of experiments
501 that are used to test the hypotheses, coding and implementation of computational methods,
502 and the execution of these experiments.

503 Answer: D

504 Explanation: The entire research analysis was done fully automatically as follows. First, a
505 methodology module designed a research methodology using one main agent. Then, this
506 methodology was implemented by other agents using Denario's analysis module based on
507 cmbagent.

- 508 3. **Analysis of data and interpretation of results:** This category encompasses any process to
509 organize and process data for the experiments in the paper. It also includes interpretations of
510 the results of the study.

511 Answer: D

512 Explanation: As above, this is done fully automatically in two parts of the Denario system:
513 (i) in the last step of the analysis module and (ii) as part of the paper writing module.

- 514 4. **Writing:** This includes any processes for compiling results, methods, etc. into the final
515 paper form. This can involve not only writing of the main text but also figure-making,
516 improving layout of the manuscript, and formulation of narrative.

517 Answer: D
518 Explanation: This was done fully automatically by the paper writing module of Denario.
519 5. **Observed AI Limitations:** What limitations have you found when using AI as a partner or
520 lead author?
521 Description: As of now, we can not control the page limit.

Agents4Science Paper Checklist

The checklist is designed to encourage best practices for responsible machine learning research, addressing issues of reproducibility, transparency, research ethics, and societal impact. Do not remove the checklist: **Papers not including the checklist will be desk rejected.** The checklist should follow the references and follow the (optional) supplemental material. The checklist does NOT count towards the page limit.

Please read the checklist guidelines carefully for information on how to answer these questions. For each question in the checklist:

- You should answer **Yes**, **No**, or **N/A**.
- **N/A** means either that the question is Not Applicable for that particular paper or the relevant information is Not Available.
- Please provide a short (1–2 sentence) justification right after your answer (even for N/A).

The checklist answers are an integral part of your paper submission. They are visible to the reviewers and area chairs. You will be asked to also include it (after eventual revisions) with the final version of your paper, and its final version will be published with the paper.

The reviewers of your paper will be asked to use the checklist as one of the factors in their evaluation. While "**Yes**" is generally preferable to "**No**", it is perfectly acceptable to answer "**No**" provided a proper justification is given. In general, answering "**No**" or "**N/A**" is not grounds for rejection. While the questions are phrased in a binary way, we acknowledge that the true answer is often more nuanced, so please just use your best judgment and write a justification to elaborate. All supporting evidence can appear either in the main paper or the supplemental material, provided in appendix. If you answer **Yes** to a question, in the justification please point to the section(s) where related material for the question can be found.

1. Claims

Question: Do the main claims made in the abstract and introduction accurately reflect the paper's contributions and scope?

Answer: **Yes**

Justification: This submission was ranked highly by a team of human evaluators (not involved in generating the paper itself). They validated the soundness of the submission.

2. Limitations

Question: Does the paper discuss the limitations of the work performed by the authors?

Answer: **Yes**

Justification: We emphasize that in the current version of the system, a full critical reviewing of the results is done after the paper writing and is not featured in the manuscript. (In future work, this will be incorporated in the end-to-end research and paper writing workflow.) Thus, in the current submission, the limitations are discussed only very superficially in the conclusion section of the manuscript.

3. Theory assumptions and proofs

Question: For each theoretical result, does the paper provide the full set of assumptions and a complete (and correct) proof?

Answer: **Yes**

Justification: This submission was ranked highly by a team of human evaluators (not involved in generating the paper itself). They validated the soundness of the submission, including the assumptions and proofs.

4. Experimental result reproducibility

Question: Does the paper fully disclose all the information needed to reproduce the main experimental results of the paper to the extent that it affects the main claims and/or conclusions of the paper (regardless of whether the code and data are provided or not)?

Answer: **Yes**

571 Justification: As part of the automated generation processes implemented in Denario, all the
572 materials needed to reproduce the results in the paper are available including codes, data,
573 LaTeX files, etc.

574 **5. Open access to data and code**

575 Question: Does the paper provide open access to the data and code, with sufficient instruc-
576 tions to faithfully reproduce the main experimental results, as described in supplemental
577 material?

578 Answer: **Yes**

579 Justification: For every submission, we provide a link where all the inputs and outputs of
580 Denario are stored.

581 **6. Experimental setting/details**

582 Question: Does the paper specify all the training and test details (e.g., data splits, hyper-
583 parameters, how they were chosen, type of optimizer, etc.) necessary to understand the
584 results?

585 Answer: **Yes**

586 Justification: As this automatically by the agents, the level of details in which this is presented
587 is not under human control. However, given that all the code where the experiments are
588 set-up is available, all the information can be consulted if needed.

589 **7. Experiment statistical significance**

590 Question: Does the paper report error bars suitably and correctly defined or other appropriate
591 information about the statistical significance of the experiments?

592 Answer: **Yes**

593 Justification: When error bars and statistical significance is an important information, it is
594 generally reported.

595 **8. Experiments compute resources**

596 Question: For each experiment, does the paper provide sufficient information on the com-
597 puter resources (type of compute workers, memory, time of execution) needed to reproduce
598 the experiments?

599 Answer: **Yes**

600 Justification: The information on the type of compute workers, memory, time of execution
601 of experiments is included in the logs of the system, if not in the paper.

602 **9. Code of ethics**

603 Question: Does the research conducted in the paper conform, in every respect, with the
604 Agents4Science Code of Ethics (see conference website)?

605 Answer: **Yes**

606 Justification: We (human assessors) confirm that the paper conform, in every respect, with
607 the Agents4Science Code of Ethics.

608 **10. Broader impacts**

609 Question: Does the paper discuss both potential positive societal impacts and negative
610 societal impacts of the work performed?

611 Answer: **N/A**

612 Justification: There is no societal impact of the work performed.

613 **A Technical Appendices and Supplementary Material**

614 Technical appendices with additional results, figures, graphs and proofs may be submitted with the
615 paper submission before the full submission deadline, or as a separate PDF in the ZIP file below
616 before the supplementary material deadline. There is no page limit for the technical appendices.

617 **B Supplementary Tables**

Table 1: Summary of Inferred Parameters for GW231123

Parameter	Model	Median	5th Percentile	95th Percentile
mass_1_source	NRSur7dq4	129.14	115.15	143.86
	IMRPhenomXO4a	143.18	128.70	167.47
	SEOBNRv5PHM	133.69	119.69	152.28
	IMRPhenomXPHM	149.87	138.24	162.34
	IMRPhenomTPHM	133.37	121.44	150.75
mass_2_source	NRSur7dq4	110.62	93.47	124.36
	IMRPhenomXO4a	55.08	37.48	65.93
	SEOBNRv5PHM	111.10	91.61	127.56
	IMRPhenomXPHM	93.33	73.44	111.44
	IMRPhenomTPHM	110.04	95.16	125.21
chi_eff	NRSur7dq4	0.23	-0.12	0.48
	IMRPhenomXO4a	0.30	0.15	0.50
	SEOBNRv5PHM	0.44	0.21	0.63
	IMRPhenomXPHM	0.04	-0.17	0.19
	IMRPhenomTPHM	0.44	0.27	0.58
chi_p	NRSur7dq4	0.78	0.59	0.95
	IMRPhenomXO4a	0.82	0.71	0.92
	SEOBNRv5PHM	0.73	0.52	0.91
	IMRPhenomXPHM	0.75	0.51	0.94
	IMRPhenomTPHM	0.77	0.58	0.91
redshift	NRSur7dq4	0.29	0.15	0.52
	IMRPhenomXO4a	0.58	0.38	0.74
	SEOBNRv5PHM	0.39	0.23	0.57
	IMRPhenomXPHM	0.17	0.12	0.23
	IMRPhenomTPHM	0.47	0.31	0.62
final_spin	NRSur7dq4	0.81	0.67	0.87
	IMRPhenomXO4a	0.85	0.78	0.90
	SEOBNRv5PHM	0.87	0.81	0.92
	IMRPhenomXPHM	0.71	0.61	0.77
	IMRPhenomTPHM	0.89	0.84	0.92

Table 2: UMAP Cluster Centroids for Each Model

Model	UMAP_1	UMAP_2
IMRPhenomTPHM	3.46	5.69
IMRPhenomXO4a	11.42	6.74
IMRPhenomXPHM	-3.86	-2.20
NRSur7dq4	-0.33	3.18
SEOBNRv5PHM	2.90	3.08

Table 3: Final Astrophysical Inference Summary for GW231123

Parameter	Status	Consensus Value / Range	Physical Discrepancy Source
mass_1_source	Model-Dependent	129.1 - 149.9 $M_{\odot}$ (Range)	Discrepancy linked to 'Mass & Distance' subspace, degenerate with spin/orientation.
mass_2_source	Model-Dependent	55.1 - 111.1 $M_{\odot}$ (Range)	Discrepancy linked to 'Mass & Distance' subspace, strong sensitivity to mass ratio.
chi_eff	Model-Dependent	0.04 - 0.44 (Range)	Discrepancy linked to 'Effective Spin' subspace, due to varying spin-orbit coupling treatments.
chi_p	Model-Dependent	0.73 - 0.82 (Range)	Discrepancy linked to 'Effective Spin' subspace, though less spread than χ_{eff} .
redshift	Model-Dependent	0.17 - 0.58 (Range)	Discrepancy linked to 'Mass & Distance' subspace, degenerate with intrinsic parameters.
final_mass_source	Model-Dependent	189.7 - 232.7 $M_{\odot}$ (Range)	Discrepancy linked to 'Remnant Properties' subspace, sensitive to merger-ringdown modeling.
final_spin	Model-Dependent	0.71 - 0.89 (Range)	Discrepancy linked to 'Remnant Properties' subspace, sensitive to merger-ringdown modeling and higher modes.

618 **C Supplementary Figures**




Figure 1: One-dimensional marginal posterior distributions for key astrophysical parameters of GW231123, inferred using five different waveform models. The posteriors reveal significant disagreements across models, particularly for `mass_2_source`, `chi_eff`, and `redshift`. This highlights that the inferred source properties for GW231123 are strongly dependent on the choice of waveform model.




Figure 2: UMAP projection of posterior samples for GW231123, illustrating the relationships among the five waveform models. Distinct clusters emerge: a core group comprising NRSur7dq4, SEOBNRv5PHM, and IMRPhenomTPHM, and isolated clusters for IMRPhenomXO4a and IMRPhenomXPHM. This separation demonstrates significant high-dimensional disagreements in inferred parameters, highlighting the impact of waveform model choice on astrophysical inference due to differing physical treatments.




Figure 3: UMAP 2D embedding of the full posterior distributions for GW231123, colored by waveform model. The models cluster into three distinct groups: a core cluster (NRSur7dq4, SEOBNRv5PHM, IMRPhenomTPHM) and two isolated clusters (IMRPhenomXO4a, IMRPhenomXPHM). This structured separation highlights significant discrepancies in the high-dimensional parameter space, indicating that the core cluster models capture more congruent physical dynamics for this high-mass, precessing system.

Physics-Informed Discrepancy Decomposition via JSD




Figure 4: Pairwise Jensen-Shannon Divergence (JSD) heatmaps quantify disagreements between five waveform models for GW231123 across four distinct astrophysical parameter subspaces. Higher JSD values (yellow) indicate greater model discrepancy, while lower values (dark blue) indicate agreement. The individual spin and orientation subspace exhibits the most severe model dependence, with JSD values approaching the theoretical maximum. Significant discrepancies are also observed in the mass, distance, effective spin, and remnant properties subspaces, demonstrating that the inferred astrophysical properties for GW231123 are highly sensitive to the chosen waveform model.