

NETS: A NON-EQUILIBRIUM TRANSPORT SAMPLER

Anonymous authors

Paper under double-blind review

ABSTRACT

We propose an algorithm, termed the Non-Equilibrium Transport Sampler (NETS), to sample from unnormalized probability distributions. NETS can be viewed as a variant of annealed importance sampling (AIS) based on Jarzynski’s equality, in which the stochastic differential equation used to perform the non-equilibrium sampling is augmented with an additional learned drift term that lowers the impact of the unbiasing weights used in AIS. We show that this drift is the minimizer of a variety of objective functions, which can all be estimated in an unbiased fashion without backpropagating through solutions of the stochastic differential equations governing the sampling. We also prove that some these objectives control the Kullback-Leibler divergence of the estimated distribution from its target. NETS is shown to be unbiased and, in addition, has a tunable diffusion coefficient which can be adjusted post-training to maximize the effective sample size. We demonstrate the efficacy of the method on standard benchmarks, high-dimensional Gaussian mixture distributions, and a model from statistical lattice field theory, for which it surpasses the performances of related work and existing baselines.

1 INTRODUCTION

The aim of this paper is to sample probability distributions supported on \mathbb{R}^d and known only up to a normalization constant. This problem arises in a wide variety in applications, ranging from statistical physics (Faulkner & Livingstone, 2023; Wilson, 1974; Hénin et al., 2022) to Bayesian inference (Neal, 1993), and it is known to be challenging when the target distribution is not log-concave. In this situation, vanilla methods based on ergodic sampling using Markov Chain Monte Carlo (MCMC) or stochastic differential equations (SDE) such as the Langevin dynamics typically have very slow convergence rates, making them inefficient in practice.

To overcome these difficulties, a variety of more sophisticated methods have been introduced, based e.g. on importance sampling. Here we will be interested in a class of methods of this type which involve non-equilibrium sampling, by which we mean algorithms that attempt to sample from a non-stationary probability distribution. The idea is to first generate samples from a simple base distribution (e.g. a normal distribution) then push them in finite time unto samples from the target. Traditionally, this aim has been achieved by combining some transport using e.g. the Langevin dynamics with reweighting, so as to remove the bias introduced by the non-equilibrium quench. Neal’s Annealed Importance Sampling (AIS) (Neal, 2001), Sequential Monte Carlo (SMC) methods (Del Moral, 1997; Doucet et al., 2001), or continuous-time variants thereof based on Jarzynski’s equality (Hartmann et al., 2018) are ways to implement this idea in practice. While these methods work better than ergodic sampling in many instances, they too can fail when the variance of the un-biasing weights become too large compared to their mean: this arises when the samples end up being too far from the target distribution at the end of the non-equilibrium quench.

This problem suggests to modify the dynamics of the samples to help them evolves towards the target during the quench. Here we propose a way to achieve this by learning an *additional drift* to include in the Langevin SDE. As we will show below, there exists a drift that removes the need for the un-biasing weights altogether, and it is the minimizer of a variety of objective functions that are amenable to empirical estimation. In practice, this offers the possibility to estimate this drift using deep learning methods. We exploit this idea here, showing that it results in an unbiased sampling strategy in which importance weights can still be used to correct the samples exactly, but the variance of these weights can be made much smaller due to the additional transport.

To this end, our work makes the following **main contributions**:

- We present a sampling algorithm which combines annealed Langevin dynamics with learnable additional transport. The algorithm, which we call the Non-Equilibrium Transport Sampler (NETS), is shown to be unbiased through a generalization of the Jarzynski equality.
- We show that the drift coefficient contributing this additional transport is the minimizer of two separate objective functions, which can be learned without backpropagating through solutions of the SDE used for sampling.
- We show that one of these objectives, a physics informed neural network (PINN) loss, is also an *off-policy* objective, meaning that it does not need samples from the target density. In addition, this objective controls the KL-divergence between the model and its target.
- The resultant samplers can be adapted *after training* by tuning the integration time-step as well as the diffusivity to improve the sample quality, which we demonstrate on high-dimensional numerical experiments below.

1.1 RELATED WORK

Dynamical Measure Transport. Most contemporary generative models for continuous data are built upon dynamical measure transport, in which samples from a base density are mapped to samples from a target density by means of solving ordinary or stochastic differential equations (ODE/SDE) whose drift coefficients are estimable. Initial attempts to do this started with (Chen et al., 2018; Grathwohl et al., 2019). Recent work built upon these ideas by recasting the challenge of estimating the drift coefficients in these dynamical equations as a problem of quadratic regression, most notably with score-based diffusion models (Ho et al., 2020; Song et al., 2020), and since then with more general frameworks (Albergo & Vanden-Eijnden, 2022; Lipman et al., 2022; Albergo et al., 2023; Liu et al., 2022; De Bortoli et al., 2021; Neklyudov et al., 2023). Importantly, these methods work because samples from the base and the target are readily available. Here, we show that analogous equations governing those systems can lead to objective functions that can be minimized with no initial data but still allow us to exploit the expressivity of models built on dynamical transport.

Augmenting sampling with learning. Augmenting MCMC and importance sampling procedures with transport has been an active area of research for the past decade. Early work makes use of the independence Metropolis algorithm (Hastings, 1970; Liu, 1996), in which proposals come from a transport map (Parno & Marzouk, 2018; Noé et al., 2019; Albergo et al., 2019; Gabrié et al., 2022) that are accepted or rejected based off their likelihood ratio with the target. These methods were further improved by combining them with AIS and SMC perspectives, learning incremental maps that connect a sequence of interpolating densities between the base and target (Arbel et al., 2021; Matthews et al., 2022; Midgley et al., 2023). Similar works in the high-energy physics community posit that interleaving stochastic updates within a sequence of maps can be seen as a form of non-equilibrium sampling (Caselle et al., 2022; Bonanno et al., 2024). Following the success of generative models built out of dynamical transport, there has been a surge of interest in applying these perspectives to sampling: Vargas et al. (2023); Berner et al. (2024) translate ideas from diffusion models to minimize the KL divergence between the model and the target, and Zhang & Chen (2021) as well as Behjoo & Chertkov (2024) reformulate sampling as a stochastic optimal control (SOC) problem. These approaches require backpropagating through the solution of an SDE, which is too costly in high dimensions. In addition, the methods based on SOC must start with samples from a point mass, which may be far from the target. Akhound-Sadegh et al. (2024) avoid the need to backpropagate through an SDE, but in the process introduce a bias into their objective function. An alternative perspective was also recently given in (Bruna & Han, 2024) by using denoising oracles to turn the original sampling problem into an easier one. [A final line of work \(Malkin et al., 2023\) shows how ideas used for modeling distributions on graphs can be repurposed as tools for sampling, including with off-policy training \(Sendera et al., 2024\).](#)

Vargas et al. (2024) establish an unbiased sampler with added transport called Controlled Monte Carlo Diffusions (CMCD) that is similar to ours. The main difference is how we learn the drift. In Vargas et al. (2024) an objective for this drift in gradient form is derived through the use of path integrals and Girsanov’s theorem. This objective either needs backpropagating through the SDE or has to be computed with a reference measure, and is done on a fixed grid. In practice the latter can be

numerically unstable. Here, through simple manipulations of Fokker-Planck equations, we propose a variety of new objective functions for the additional drift, none of which require backpropagating through the simulation. In addition, our learning can be done in an optimize-then-discretize fashion so that the sampling can be done with arbitrary step size and time-dependent diffusion after learning the model. This gives us an adaptive knob to increase performance, which we demonstrate below. **One of the objectives we propose is a Physics-Informed Neural Network (PINN) loss that has recently appeared elsewhere in the literature for sampling (Máté & Fleuret, 2023; Tian et al., 2024; Fan et al., 2024). Importantly, here we establish that this objective is valid in the context of annealed Langevin dynamics, and, moreover, that this PINN objective *directly controls* the KL divergence as well as the importance weights that emerge through the Jarzynski equality.**

2 METHODS

2.1 SETUP AND NOTATIONS

We assume that the target distribution is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d , with probability density function (PDF) $\rho_1(x) = Z_1^{-1}e^{-U_1(x)}$: here $x \in \mathbb{R}^d$, $U : \mathbb{R}^d \rightarrow \mathbb{R}$ is a known energy potential, assumed twice differentiable and bounded below, and $Z_1 = \int_{\mathbb{R}^d} e^{-U_1(x)} dx < \infty$ is an unknown normalization constant, referred to as the partition function in physics and the evidence in statistics. Our aim is to generate samples from $\rho_1(x)$ so as to be able to estimate expectations with respect to this density. Additionally we wish to estimate Z_1 .

To this end we will use a series of time-dependent potentials $U_t(x)$ which connects some simple $U_0(x)$ at $t = 0$ (e.g. $U_0(x) = \frac{1}{2}|x|^2$) to $U_1(x)$ at $t = 1$. For example we could use linear interpolation:

$$U_t(x) = (1 - t)U_0(x) + tU_1(x), \quad (1)$$

but other choices are possible as long as $U_{t=0} = U_0$, $U_{t=1} = U_1$, and $U_t(x)$ is twice differentiable in $(t, x) \in [0, 1] \times \mathbb{R}^d$, which we explore below. We assume that the time-dependent PDF associated with this potential $U_t(x)$ is normalizable for all $t \in [0, 1]$ and denote it as

$$\rho_t(x) = Z_t^{-1}e^{-U_t(x)}, \quad Z_t = \int_{\mathbb{R}^d} e^{-U_t(x)} dx < \infty, \quad (2)$$

so that $\rho_{t=0}(x) = \rho_0(x)$ and $\rho_{t=1}(x) = \rho_1(x)$; we also assume that $\rho_0(x)$ is simple to sample (either directly or via MCMC or Langevin dynamics) and that its partition function Z_0 is known. To simplify the notations we also introduce the free energy

$$F_t = -\log Z_t. \quad (3)$$

Since $\partial_t F_t = -\partial_t \log \int_{\mathbb{R}^d} e^{-U_t(x)} dx = \int_{\mathbb{R}^d} \partial_t U_t(x) e^{-U_t(x)} dx / \int_{\mathbb{R}^d} e^{-U_t(x)} dx$ we have the useful identity

$$\partial_t F_t = \int_{\mathbb{R}^d} \partial_t U_t(x) \rho_t(x) dx. \quad (4)$$

2.2 NONEQUILIBRIUM SAMPLING WITH IMPORTANCE WEIGHTS

Annealed importance sampling uses a finite sequence of MCMC moves that satisfy detailed-balance locally in time but not globally, thereby introducing a bias that can be corrected with weights. Here we present a time-continuous variant of AIS based on Jarzynski equality that will be more useful for our purpose.

By definition of the PDF in (2), $\nabla \rho_t(x) = -\nabla U_t(x) \rho_t(x)$ and hence, for any $\varepsilon_t \geq 0$, we have

$$0 = \varepsilon_t \nabla \cdot (\nabla U_t \rho_t + \nabla \rho_t). \quad (5)$$

Since we also have

$$\partial_t \rho_t = -(\partial_t U_t - \partial_t F_t) \rho_t, \quad (6)$$

we can combine these last two equations to deduce that

$$\partial_t \rho_t = \varepsilon_t \nabla \cdot (\nabla U_t \rho_t + \nabla \rho_t) - (\partial_t U_t - \partial_t F_t) \rho_t. \quad (7)$$

The effect of the last term at the right hand-side of this equation can be accounted for by using weights. To see how, notice that if we extend the phase space to $(x, a) \in \mathbb{R}^{d+1}$ and introduce the PDF $f_t(x, a)$ solution to the Fokker-Planck equation (FPE)

$$\partial_t f_t = \varepsilon_t \nabla \cdot (\nabla U_t f_t + \nabla f_t) + \partial_t U_t \partial_a f_t, \quad f_{t=0}(x, a) = \delta(a) \rho_0(x), \quad (8)$$

then a direct calculation using (4) (for details see Appendix 4.1) shows that

$$\rho_t(x) = \frac{\int_{\mathbb{R}} e^a f_t(x, a) da}{\int_{\mathbb{R}^{d+1}} e^a f_t(y, a) dady}. \quad (9)$$

Therefore we can use the solution to SDE associated with the FPE (8) in the extended space to estimate expectations with respect to $\rho_t(x)$:

Proposition 1 (Jarzynski equality). *Let (X_t, A_t) solve the coupled system of SDE/ODE*

$$dX_t = -\varepsilon_t \nabla U_t(X_t) dt + \sqrt{2\varepsilon_t} dW_t, \quad X_0 \sim \rho_0, \quad (10)$$

$$dA_t = -\partial_t U_t(X_t) dt, \quad A_0 = 0, \quad (11)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \frac{\mathbb{E}[e^{A_t} h(X_t)]}{\mathbb{E}[e^{A_t}]}, \quad Z_t/Z_0 = e^{-F_t+F_0} = \mathbb{E}[e^{A_t}], \quad (12)$$

where the expectations are taken over the law of (X_t, A_t) .

The proof of this proposition is given in Appendix 4.1 and it relies on the identity $\int_{\mathbb{R}^d} h(x) \rho_t(x) = \int_{\mathbb{R}^{d+1}} e^a h(x) f_t(x, a) dadx / \int_{\mathbb{R}^{d+1}} f_t(x, a) dadx$ which follows from (9). The second equation in (12) for the free energy F_t is what is referred to as Jarzynski's equality, and was originally surmised in the context of non-equilibrium thermodynamics (Jarzynski, 1997).

Remark 1. *We stress that it is key to use the weights e^{A_t} in (12) because $\rho_t(x)$ is not the PDF of X_t in general. Indeed, if we denote by $\tilde{\rho}_t(x)$ the PDF of X_t , it satisfies the FPE*

$$\partial_t \tilde{\rho}_t = \varepsilon_t \nabla \cdot (\nabla U_t \tilde{\rho}_t + \nabla \tilde{\rho}_t), \quad \tilde{\rho}_{t=0} = \rho. \quad (13)$$

This FPE misses the term $-(\partial_t U_t - \partial_t F_t) \rho_t$ at the right hand-side of (7), and as a result $\tilde{\rho}_t(x) \neq \rho_t(x)$ in general – intuitively, $\tilde{\rho}_t(x)$ lags behind $\rho_t(x)$ when the potential $U_t(x)$ evolves and this lag is what the weights in (12) correct for.

It is important to realize that, while the relation (12) can be used to compute unbiased estimators of expectations, this estimator on its own can be high variance if the lag between the PDF $\tilde{\rho}_t(x)$ of X_t and $\rho_t(x)$ is too pronounced. This issue can be alleviated by using resampling methods as is done in sequential Monte Carlo (Doucet et al., 2001). Here we will solve it by adding some additional drift in (10) that will compensate for this lag and reduce the effect of the weights.

2.3 NONEQUILIBRIUM SAMPLING WITH PERFECT ADDITIONAL TRANSPORT

To see how we can add a transport term to eliminate the need of the weights, let us introduce a velocity field $b_t(x) \in \mathbb{R}^d$ which at all times $t \in [0, 1]$ satisfies

$$\nabla \cdot (b_t \rho_t) = -\partial_t \rho_t. \quad (14)$$

We stress that this is an equation for $b_t(x)$ in which $\rho_t(x)$ is fixed and given by (2): In Appendix 4.3 we show how to express the solution to (14) via Feynman-Kac formula. If (14) is satisfied, then we can combine this equation with (5) and (6) to arrive at

$$\partial_t \rho_t = \varepsilon_t \nabla \cdot (\nabla U_t \rho_t + \nabla \rho_t) - \nabla \cdot (b_t \rho_t), \quad (15)$$

which is a standard FPE. Therefore the solution to the SDE associated with (15) allows us to sample $\rho_t(x)$ directly (without weights). We phrase this result as:

Proposition 2 (Sampling with perfect additional transport.). *Let $b_t(x)$ be a solution to (14) and let X_t^b satisfy the SDE*

$$dX_t^b = -\varepsilon_t \nabla U_t(X_t^b) dt + b_t(X_t^b) dt + \sqrt{2\varepsilon_t} dW_t, \quad X_0^b \sim \rho_0, \quad (16)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then $\rho_t(x)$ is the PDF of X_T^b , i.e. for all $t \in [0, 1]$ and, given any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \mathbb{E}[h(X_t^b)], \quad (17)$$

where the expectation at the right-hand side is taken over the law of (X_t^b) .

This proposition is proven in Appendix 4.1 and it shows that we can in principle get rid of the weights altogether by adding the drift $b_t(x)$ in the Langevin SDE. This possibility was first noted in Vaikuntanathan & Jarzynski (2008) and is also exploited in Tian et al. (2024) for deterministic dynamics (i.e. setting $\varepsilon_t = 0$ in (16)) and in Vargas et al. (2024) using the SDE (16). Of course, in practice we need to estimate this drift, and also correct for sampling errors if this drift is imperfectly learned. Let us discuss this second question first, and defer the derivation of objectives to learn $b_t(x)$ to Secs. 2.5 and 2.6. In Appendix 4.3 we show how to express the solution to (14) via Feynman-Kac formula.

2.4 NON-EQUILIBRIUM TRANSPORT SAMPLER

Let us now show that we can combine the approaches discussed in Secs. 2.2 and 2.3 to design samplers in which we use an added transport, possibly imperfect, and importance weights.

To this end, suppose that we wish to add an additional transport term $-\nabla \cdot (\hat{b}_t \rho_t)$ in (7), where $\hat{b}_t(x) \in \mathbb{R}^d$ is some given velocity that does not necessarily solve (14). Using the expression in (2) for $\rho_t(x)$, we have the identity

$$-\nabla \cdot (\hat{b}_t \rho_t) = -\nabla \cdot \hat{b}_t \rho_t + \nabla U_t \cdot \hat{b}_t \rho_t \quad (18)$$

Therefore we can rewrite (5) equivalently as

$$\partial_t \rho_t = \varepsilon_t \nabla \cdot (\nabla U_t \rho_t + \nabla \rho_t) - \nabla \cdot (\hat{b}_t \rho_t) + (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t + \partial_t F_t) \rho_t. \quad (19)$$

We can now proceed as we did with (5) and extend state space to account for the effect of the terms $(\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t + \partial_t F_t) \rho_t$ in this equation via weights, while having the term $-\nabla \cdot (\hat{b}_t(x) \rho_t(x))$ contribute to some additional transport. This leads us to a result originally obtained in Vaikuntanathan & Jarzynski (2008) and recently re-derived in Vargas et al. (2024):

Proposition 3 (Nonequilibrium Transport Sampler (NETS)). *Let $(X_t^{\hat{b}}, A_t^{\hat{b}})$ solve the coupled system of SDE/ODE*

$$dX_t^{\hat{b}} = -\varepsilon_t \nabla U_t(X_t^{\hat{b}}) dt + \hat{b}_t(X_t^{\hat{b}}) dt + \sqrt{2\varepsilon_t} dW_t, \quad X_0^{\hat{b}} \sim \rho_0, \quad (20)$$

$$dA_t^{\hat{b}} = \nabla \cdot \hat{b}_t(X_t^{\hat{b}}) dt - \nabla U_t(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}}) dt - \partial_t U_t(X_t^{\hat{b}}) dt, \quad A_0^{\hat{b}} = 0, \quad (21)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \frac{\mathbb{E}[e^{A_t^{\hat{b}}} h(X_t^{\hat{b}})]}{\mathbb{E}[e^{A_t^{\hat{b}}}]}, \quad Z_t/Z_0 = e^{-F_t+F_0} = \mathbb{E}[e^{A_t^{\hat{b}}}], \quad (22)$$

where the expectations are taken over the law of $(X_t^{\hat{b}}, A_t^{\hat{b}})$.

A simple proof of this proposition using simple manipulations of the FPE is given in Appendix 4.1, which will allow us to write down a variety of new loss functions for learning \hat{b}_t ; for an alternative proof using Girsanov theorem, see Vargas et al. (2024). For completeness, in Appendix 4.2 we also give a time-discretized version of Proposition 3, and in Appendix 4.4 we generalize it in two ways: to include inertia and to turn t into a vector coordinate for multimarginal sampling. We also discuss the connection between NETS and the method of Vargas et al. (2024) in Appendix 4.8.

Notice that, if $\hat{b}_t(x) = 0$, the equations in Proposition 3 simply reduce to those in Proposition 1, whereas if $\hat{b}_t(x) = b_t(x)$ solves (14) we can show that

$$A_t^b = -F_t + F_0, \quad (23)$$

i.e. the weights have zero variance and give the free energy difference. Indeed, by expanding both sides of (14) and dividing them by $\rho_t(x) > 0$, this equation can equivalently be written as

$$\nabla \cdot b_t - \nabla U_t \cdot b_t = \partial_t U_t - \partial_t F_t. \quad (24)$$

As a result, when $\hat{b}_t(x) = b_t(x)$, (21) reduces to

$$dA_t^b = -\partial_t F_t dt, \quad A_0^b = 0, \quad (25)$$

and the solution to this equation is (23). In practice, achieving zero variance of the weights by estimating $b_t(x)$ exactly is not generally possible, but having a good approximation of $b_t(x)$ can help reducing this variance dramatically, as we will illustrate below via experiments.

2.5 ESTIMATING THE DRIFT $b_t(x)$ VIA A PINN OBJECTIVE

Equation (24) can be used to derive an objective for both $b_t(x)$ and F_t . The reason is that in this equation the unknown $\partial_t F_t$ can be viewed as factor that guarantees solvability: indeed, integrating both sides of (14) gives $0 = -\partial_t \int_{\mathbb{R}^d} U_t(x) \rho_t(x) dx + \partial_t F_t$, which, by (4), is satisfied if and only if F_t is (up to a constant fixed by $F_0 = -\log Z_0$) the exact free energy (3). This offers the possibility to learn both $b_t(x)$ and F_t variationally using an objective fitting the framework of physics informed neural networks (PINNs):

Proposition 4 (PINN objective). *Given any $T \in (0, 1]$ and any PDF $\hat{\rho}_t(x) > 0$ consider the objective for (\hat{b}, \hat{F}) given by:*

$$L_{PINN}^T[\hat{b}, \hat{F}] = \int_0^T \int_{\mathbb{R}^d} |\nabla \cdot \hat{b}_t(x) - \nabla U_t(x) \cdot \hat{b}_t(x) - \partial_t U_t(x) + \partial_t \hat{F}_t|^2 \hat{\rho}_t(x) dx dt. \quad (26)$$

Then $\min_{\hat{b}, \hat{F}} L_{PINN}^T[\hat{b}, \hat{F}] = 0$, and all minimizers (b, F) are such that $b_t(x)$ solves (24) and F_t is the free energy (3) for all $t \in [0, T]$.

This result is proven in Appendix 4.1: in practice, we will use $T \in (0, 1]$ for annealing but ultimately we are interested in the result when $T = 1$. Note that since the expectation over an arbitrary $\hat{\rho}_t(x)$ in (26), it can be used as an off-policy objective. It is however natural to use $\hat{\rho}_t(x) = \rho_t(x)$ since it allows us to put statistical weight in the objective precisely in the regions where we need $b_t(x)$ to transport probability mass. In either case, there is no need to backpropagate through simulation of the SDE used to produce data. We show how the expectation over $\rho_t(x)$ can be estimated without bias in Appendices 4.5 and 4.5.1 to arrive at an empirical estimator for (26) when $\hat{\rho}_t(x) = \rho_t(x)$. Note also that, while minimization of the objective (26) gives an estimate \hat{F}_t of the free energy, it is not needed at sampling time when solving (20). An objective similar to (26) was also recently posited in Máté & Fleuret (2023); Tian et al. (2024) for use with deterministic flows. Here, we devise it in the context of augmenting annealed Langevin dynamics.

One advantage of the PINN objective (26) is that we know that its minimum is zero, and hence we can track its value to monitor convergence when minimizing (26) by gradient descent as we do below. Another advantage of the loss (26) is that it controls the quality of the transport as measured by the Kullback-Leibler divergence:

Proposition 5 (KL control). *Let $\hat{\rho}_t$ be the solution to the transport equation*

$$\partial_t \hat{\rho}_t = -\nabla \cdot (\hat{b}_t \hat{\rho}_t), \quad \hat{\rho}_{t=0} = \rho_0 \quad (27)$$

where $\hat{b}_t(x)$ is some predefined velocity field. Then, we have

$$D_{KL}(\hat{\rho}_{t=1} || \rho_1) \leq \sqrt{L_{PINN}^{T=1}(\hat{b}, F)}. \quad (28)$$

where F_t is the free energy. In addition, given any estimate \hat{F}_t such that $\int_0^1 |\partial_t \hat{F}_t - \partial_t F_t|^2 dt \leq \delta$ for some $\delta \geq 0$, we have

$$D_{KL}(\hat{\rho}_{t=1} || \rho_1) \leq \sqrt{2L_{PINN}^{T=1}(\hat{b}, \hat{F})} + 2\delta. \quad (29)$$

This proposition is proven in Appendix 4.1. Notice that the bound (28) can be estimated by using $\partial_t F_t = \mathbb{E}[e^{A_t^b} \partial_t U_t(X_t^b)] / \mathbb{E}[e^{A_t^b}]$ in the PINN loss (26).

2.6 ESTIMATING THE DRIFT $b_t(x) = \nabla\phi_t(x)$ VIA ACTION MATCHING (AM)

In general (14) is solved by many $b_t(x)$. One way to get a unique (up to a constant in space and time) solution to this equation is to impose that the velocity be in gradient form, i.e. set $b_t(x) = \nabla\phi_t(x)$ for some scalar-valued potential $\phi_t(x)$. If we do so, (14) can be written as $\nabla \cdot (\nabla\phi_t(x)\rho_t) = -\partial_t\rho_t$, and it is easy to see that at all times $t \in [0, 1]$ the solution to this equation minimizes over $\hat{\phi}_t$ the objective

$$\begin{aligned} & \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla\hat{\phi}_t(x)|^2 \rho_t(x) - \hat{\phi}_t(x) \partial_t \rho_t(x) \right] dx \\ & = \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla\hat{\phi}_t(x)|^2 + (\partial_t U_t(x) - \partial_t F_t) \hat{\phi}_t(x) \right] \rho_t(x) dx. \end{aligned} \quad (30)$$

If we use (4) to set $\partial_t F_t = \int_{\mathbb{R}^d} \partial_t U_t(x) \rho_t(x) dx$ we can use the objective at the right hand-side of (30) to learn $\phi_t(x)$ locally in time (or globally if we integrate this objective on $t \in [0, 1]$). Alternatively, we can integrate the objective at the left hand-side of (30) over $t \in [0, T]$ and use integration by parts for the term involving $\partial_t \rho_t(x)$ to arrive at:

Proposition 6 (Action Matching objective). *Given any $T \in (0, 1]$ consider the objective for $\hat{\phi}_t(x)$:*

$$L_{AM}^T[\hat{\phi}] = \int_0^T \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla\hat{\phi}_t(x)|^2 + \partial_t \hat{\phi}_t(x) \right] \rho_t(x) dx dt + \int_{\mathbb{R}^d} [\hat{\phi}_0(x) \rho_0(x) - \hat{\phi}_T(x) \rho_T(x)] dx. \quad (31)$$

Then the minimizer $\phi_t(x)$ of (26) is unique (up to a constant) and $b_t(x) = \nabla\phi_t(x)$ satisfies (14) for all $t \in [0, T]$.

This proposition is proven in Appendix 4.1. This objective is analogous to the loss presented in Neklyudov et al. (2023), but adapted to the sampling problem. In practice, we will use again $T \in (0, 1]$ for annealing, but ultimately we are interested in the result at $T = 1$. Note that, unlike with the PINN objective (26), it is crucial that we use the correct $\rho_t(x)$ in the AM objective (31): that is, unlike (26), (31) cannot be turned into an off-policy objective.

Remark 2. *If we use $\hat{b}_t(x) = \nabla\hat{\phi}_t(x)$ in the SDEs in (20) and (21), we need to compute $\nabla \cdot \hat{b}_t(x) = \Delta\hat{\phi}_t(x)$, which is computationally costly. Fortunately, when $\varepsilon_t > 0$, the calculation of this Laplacian can be avoided by using the following alternative equation for $A_t^{\hat{b}}$:*

$$A_t^{\hat{b}} = \frac{1}{\varepsilon_t} [\hat{\phi}_t(X_t^{\hat{b}}) - \hat{\phi}_0(X_0^{\hat{b}})] - B_t, \quad (32)$$

where

$$dB_t = \partial_t U_t(X_t^{\hat{b}}) dt + \frac{1}{\varepsilon_t} \partial_t \hat{\phi}_t(X_t^{\hat{b}}) + \frac{1}{\varepsilon_t} |\nabla\hat{\phi}_t(X_t^{\hat{b}})|^2 dt + \sqrt{\frac{2}{\varepsilon_t}} \nabla\hat{\phi}_t(X_t^{\hat{b}}) \cdot dW_t. \quad (33)$$

This equation is derived in Appendix 4.1.

3 NUMERICAL EXPERIMENTS

In what follows, we test the NETS method, for both the PINN objective (26) and the action matching objective (31), on standard challenging sampling benchmarks. We then study how the method scales in comparison to baselines, particularly AIS on its own, by testing it on an increasingly high dimensional Gaussian Mixture Models (GMM). Following that, we show that it has orders of magnitude better statistical efficiency as compared to AIS on its own when applied to the study of lattice field theories, even past the phase transition of these theories and in 400 dimensions (an $L = 20 \times L = 20$ lattice).

3.1 40-MODE GAUSSIAN MIXTURE

A common benchmark for machine learning augmented samplers that originally appeared in the paper introducing Flow Annealed Importance Sampling Bootstrap (FAB) Midgley et al. (2023) is a 40-mode GMM in 2-dimensions for which the means of the mixture components span from -40 to 40 . The high variance and many wells make this problem challenging for re-weighting or locally updating MCMC processes. We choose as a time dependent potential $U_t(x)$ the linear interpolation (1) with U_0 the potential for a standard multivariate Gaussian with standard deviation scale $\sigma = 2$.




Figure 1: Comparison of the performance of annealed Langevin dynamics alone, transport alone, and annealed Langevin coupled with transport when sampling the 40-mode GMM from Midgley et al. (2023). **Left:** Annealed Langevin run for 250 steps with $\varepsilon_t = 4.0$, failing to capture the modes with 0% ESS. **Center:** Learning using the PINN loss and sampling with 100 steps and $\varepsilon_t = 0$ achieves an ESS of 95%. **Right:** Same learning and now sampling with $\varepsilon_t = 4.0$ achieves an ESS of 98%.

GMM ($d = 2$)		
Algorithm	ESS \uparrow	$\mathcal{W}_2 \downarrow$
FAB	0.653 ± 0.017	12.0 ± 5.73
PIS	0.295 ± 0.018	7.64 ± 0.92
DDS	0.687 ± 0.208	9.31 ± 0.82
pDEM	0.634 ± 0.084	12.20 ± 0.14
iDEM	0.734 ± 0.092	7.42 ± 3.44
CMCD-KL	0.268 ± 0.069	9.32 ± 0.71
CMCD-LV	0.655 ± 0.023	4.01 ± 0.25
NETS-AM $\varepsilon_t = 5$ (ours)	0.808 ± 0.031	3.89 ± 0.22
NETS-PINN $\varepsilon_t = 0$ (ours)	0.954 ± 0.003	3.55 ± 0.57
NETS-PINN $\varepsilon_t = 4$ (ours)	0.979 ± 0.002	3.14 ± 0.46
NETS-PINN-resample (ours)	0.993 ± 0.004	3.27 ± 0.31

Table 1: Performance of NETS in terms of ESS and \mathcal{W}_2 metrics for 40-mode GMM ($d = 2$) with comparative results quoted from Akhound-Sadegh et al. (2024) for reproducibility.

the 2-Wasserstein (\mathcal{W}_2) distance between the model and the target. As noted in Table 1, all proposed variants of NETS outperform existing methods. In addition, because our method can be turned into an SMC method by including resampling during the generation, we can push the acceptance rate of the same learned PINN model to nearly 100% by using a single resampling step when the ESS of the walkers dropped below 98%. NETS uses 100 sampling steps and an $\varepsilon_t = 0.0, 4.0$ in the SDE.

3.2 FUNNEL AND STUDENT-T MIXTURE

We next test NETS on Neal’s funnel, a challenging synthetic target distribution which exhibits correlations at different scales across its 10 dimensions, as well as the 50-dimensional Mixture of Student-T (MoS) distribution used in Blessing et al. (2024). The definitions of the target densities and the interpolating potentials are given in Appendix 4.6. Heuristically, the first dimension is Gaussian with variance $\sigma^2 = 9$, and the other 9 dimensions are conditionally Gaussian with variance $\exp(x_0)$, creating the funnel.

We again parameterize (\hat{b}, \hat{F}) or $\hat{\phi}$ using simple feed forward neural networks, this time of hidden size 512. We use 100 sampling steps for both, with diffusion coefficients given in the caption of Table 2. Following Blessing et al. (2024), we compute the maximum mean discrepancy (MMD) and \mathcal{W}_2 distance between 2000 samples from the model and 2000 samples from the target and compare to related methods in Table 2. NETS outperforms other methods with both losses on the high dimensional MoS target in both metrics. In addition this can be improved using SMC-style resampling in the interpolation when the ESS drops below 70%. NETS matches the best performance in MMD for the Funnel distribution, but it is slightly worse in \mathcal{W}_2 .

3.3 SCALING ON HIGH-DIMENSIONAL GMMs

In order to demonstrate that the method generalizes to high dimension, we study sampling from multimodal GMMs in higher and higher dimensions and observe how the performance scales. In

Algorithm	Funnel ($d = 10$)		MoS ($d = 50$)	
	MMD ↓	\mathcal{W}_2 ↓	MMD ↓	\mathcal{W}_2 ↓
FAB (Midgley et al., 2023)	0.032 ± 0.000	153.894 ± 3.916	0.093 ± 0.014	1204.160 ± 147.7
GMMVI (Arenz et al., 2023)	0.031 ± 0.000	105.620 ± 3.472	0.135 ± 0.017	1255.216 ± 296.9
PIS (Zhang & Chen, 2022)	--	--	0.218 ± 0.007	2113.172 ± 31.17
DDS (Vargas et al., 2023)	0.172 ± 0.031	142.890 ± 9.552	0.131 ± 0.001	2154.884 ± 3.861
AFT (Arbel et al., 2021)	0.159 ± 0.010	145.138 ± 6.061	0.395 ± 0.082	2648.410 ± 301.3
CRAFT (Arbel et al., 2021)	0.115 ± 0.003	134.335 ± 0.663	0.257 ± 0.024	1893.926 ± 117.3
CMCD-KL (Vargas et al., 2024)	0.095 ± 0.003	513.339 ± 192.4	--	--
NETS-AM (ours)	0.041 ± 0.001	435.793 ± 96.17	0.0396 ± 0.001	407.827 ± 69.64
NETS-PINN (ours)	0.033 ± 0.002	388.91 ± 141.5	0.032 ± 0.001	482.393 ± 174.6
NETS-PINN-resample (ours)	0.027 ± 0.003	343.78 ± 65.25	0.030 ± 0.000	400.076 ± 59.31

Table 2: Performance of NETS on Neal’s Funnel and Mixture of Student-T distributions, measured in MMD and \mathcal{W}_2 distances from the true distribution. Benchmarking is in accordance with the setup of Blessing et al. (2024). Diffusion coefficient $\epsilon_t = 5, 4$ was used for NETS-AM on the Funnel and MoS, respectively. Equivalently, $\epsilon_t = 5, 5$ were used by NETS-PINN. Bold numbers are within standard deviation the best performing. Note that NETS still has perfect sample in the $\epsilon_t \rightarrow \infty$ limit, but would require finer time discretization than the 100 sampling steps used here (see Figure 4).

addition we are curious to understand how the factor in the sampling SDE coming from annealed Langevin dynamics, ∇U , interacts with the learned drift \hat{b} or $\nabla \hat{\phi}$ as we change the diffusivity. We construct 8-mode target GMMs in $d = 36, 64, 128, 200$ dimensions and learn \hat{b} with the PINN loss in each scenario. We use the same feed forward neural network of width 512 and depth 4 to parameterize




Figure 2: Demonstration of high-dimensional sampling with our method using the PINN loss in (26) and a study of how diffusivity impacts performance, with and without transport. **Left:** NETS can achieve high ESS through transport alone, and the effect of increased diffusivity has more of a positive effect on performance with sampling than without. AIS cannot achieve ESS above ≈ 0 in high dimension. **Right:** Kernel density estimates of 2- d cross sections of the high-dimensional, multimodal distribution arising from the model and ground truth.

both \hat{b} and \hat{F} for all dimensions tested and train for 4000 training iterations. Figure 2 summarizes the results. On the left plot, we note that AIS on its own cannot produce any effective samples, while even in 200 dimensions, NETS works with transport alone with 60% ESS. As we increase the diffusivity ϵ_t and therefore the effect of the Langevin term coming from the gradient of the potential, we note all the methods converge to nearly independent sampling, and the discrepancy in performance across dimensions is diminished. Note that the caveat to achieve this is that the step size in the SDE integrator must be taken smaller to accommodate the increased diffusivity, especially for the $\epsilon_t = 80$ data point. The number of sampling steps used to discretize the SDEs in these experiments ranged from $K = 100$ for $\epsilon_t = 0$ up to $K = 2000$ for $\epsilon_t = 80$. Nonetheless, it suggests that diffusion can be more helpful when there is already some successful transport than without.

3.4 LATTICE φ^4 THEORY

We next apply NETS to the simulation of a statistical lattice field theory at and past the phase transition from which the lattice goes from disordered, to semi-ordered, to fully ordered (neighboring




Figure 3: Comparison of the performance of NETS to AIS on two different settings for the study of φ^4 theory. **Top row, left:** 10 example generative lattice configurations with parameters $L = 20$, $m^2 = -1.0$, $\lambda = 0.9$, which demarcates the phase transition to the antiferromagnetic phase. **Top row, right:** Performance of AIS (purple curve) vs. NETS (red curve) in terms of effective sample size over time of integration t , and a histogram of the average magnetization of 4000 lattice configurations, sampled with AIS, NETS, and HMC (superposed in this order). Note that NETS is closer to the HMC target and re-weights correctly. Re-weighted AIS was not plotted because the weights were too high variance. **Bottom row:** Equivalent setup for $L = 16$, $m^2 = -1.0$, $\lambda = 0.8$, past the phase transition and into the ordered phase. Note that the field configurations generated by NETS are either all positive across lattice sites or all negative. AIS fails to sample the correct distribution, and its weights are too high variance to be used on the histogram.

sites are highly correlated to be of the same sign and magnitude). We study the lattice φ^4 theory in $D = 2$ spacetime dimensions. The random variables in this circumstance are field configurations $\varphi \in \mathbb{R}^{L \times L}$, where L is the extent of space and time. The interpolating energy function under which we seek to sample is defined as:

$$U_t(\varphi) = \sum_x \left[-2 \sum_{\mu} \varphi_x \varphi_{x+\mu} \right] + (2D + m_t^2) \varphi_x^2 + \lambda_t \varphi_x^4, \quad (34)$$

where summation over x indicates summation over the lattices sites, and m_t^2 and λ_t are time dependent parameters of the theory that define the phase of the lattice (ranging from disordered to ordered, otherwise known as magnetized). A derivation of this energy function is given in Appendix 4.7. Importantly, sampling the lattice configurations becomes challenging when approaching the phase transition between the disordered and ordered phases. As an example, we identify the phase transition on $L = 16$ ($d = 256$) and $L = 20$ ($d = 400$) lattices and run NETS with the action matching loss, with $\hat{\phi}_t$ a simple feed forward neural network. We use the free theory $\lambda_0 = 0$ as the base distribution under which we initially draw samples. The definition of the target parameter values m_1^2 , λ_1 both at the phase transition and in the ordered phase are given in the Appendix 4.7. In Figure 3, the top row shows samples from NETS for $L = 20$ at the phase transition, where correlations begin to appear in the lattice configurations. NETS is almost 2 orders of magnitude more statistically efficient than AIS (the same setup without the transport) in sampling at the critical point, as seen in the plot showing ESS over time. Note also that NETS can produce unbiased estimates of the magnetization as compared to a Hybrid Monte Carlo (HMC) ground truth. The bottom row shows samples past the phase transition and into the ordered phase, where the lattices begin to take on either all positive or all negative values. Again in this regime, NETS is nearly 2 orders of magnitude more statistically efficient.

While NETS performs significantly better than conventional annealed samplers on the challenging field theory problem, algorithms built out of dynamical transport still experience slowdowns near phase transitions because of the difficulty of resolving the dynamics of the integrators near these critical points. As such, we need to use 1500-2000 steps in the integrator to properly resolve the dynamics of the SDE.

REFERENCES

- 540
541
542 Tara Akhound-Sadegh, Jarrid Rector-Brooks, Joey Bose, Sarthak Mittal, Pablo Lemos, Cheng-Hao
543 Liu, Marcin Sendera, Siamak Ravanbakhsh, Gauthier Gidel, Yoshua Bengio, Nikolay Malkin, and
544 Alexander Tong. Iterated denoising energy matching for sampling from boltzmann densities. In
545 Ruslan Salakhutdinov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett,
546 and Felix Berkenkamp (eds.), *Proceedings of the 41st International Conference on Machine
547 Learning*, volume 235 of *Proceedings of Machine Learning Research*, pp. 760–786. PMLR, 21–
548 27 Jul 2024. URL [https://proceedings.mlr.press/v235/akhound-sadegh24a.
549 html](https://proceedings.mlr.press/v235/akhound-sadegh24a.html).
- 550 M. S. Albergo, G. Kanwar, and P. E. Shanahan. Flow-based generative models for markov chain
551 monte carlo in lattice field theory. *Phys. Rev. D*, 100:034515, Aug 2019. doi: 10.1103/PhysRevD.
552 100.034515. URL <https://link.aps.org/doi/10.1103/PhysRevD.100.034515>.
- 553 Michael S Albergo and Eric Vanden-Eijnden. Building normalizing flows with stochastic interpolants.
554 In *The Eleventh International Conference on Learning Representations*, 2022.
- 555 Michael S Albergo, Nicholas M Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying
556 framework for flows and diffusions. *arXiv preprint arXiv:2303.08797*, 2023.
- 557 Michael Arbel, Alexander G. D. G. Matthews, and Arnaud Doucet. Annealed flow transport monte
558 carlo. In *Proceedings of the 38th International Conference on Machine Learning*, Proceedings of
559 Machine Learning Research, 18–24 Jul 2021.
- 560
561 Oleg Arenz, Philipp Dahlinger, Zihan Ye, Michael Volpp, and Gerhard Neumann. A unified per-
562 spective on natural gradient variational inference with gaussian mixture models. *Transactions
563 on Machine Learning Research*, 2023. ISSN 2835-8856. URL [https://openreview.net/
564 forum?id=tLBjsX4tjs](https://openreview.net/forum?id=tLBjsX4tjs).
- 565 Hamidreza Behjoo and Michael Chertkov. Harmonic path integral diffusion, 2024. URL [https:
566 //arxiv.org/abs/2409.15166](https://arxiv.org/abs/2409.15166).
- 567 Julius Berner, Lorenz Richter, and Karen Ullrich. An optimal control perspective on diffusion-based
568 generative modeling, 2024. URL <https://arxiv.org/abs/2211.01364>.
- 569
570 Denis Blessing, Xiaogang Jia, Johannes Esslinger, Francisco Vargas, and Gerhard Neumann. Beyond
571 ELBOs: A large-scale evaluation of variational methods for sampling. In Ruslan Salakhutdi-
572 nov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett, and Felix
573 Berkenkamp (eds.), *Proceedings of the 41st International Conference on Machine Learning*, vol-
574 ume 235 of *Proceedings of Machine Learning Research*, pp. 4205–4229. PMLR, 21–27 Jul 2024.
575 URL <https://proceedings.mlr.press/v235/blessing24a.html>.
- 576
577 Miodrag Bolić, Petar M. Djurić, and Sangjin Hong. Resampling algorithms for particle filters: A
578 computational complexity perspective. *EURASIP Journal on Advances in Signal Processing*, 2004
579 (15):403686, 2004. doi: 10.1155/S1110865704405149. URL [https://doi.org/10.1155/
580 S1110865704405149](https://doi.org/10.1155/S1110865704405149).
- 581 Claudio Bonanno, Alessandro Nada, and Davide Vadicchino. Mitigating topological freezing
582 using out-of-equilibrium simulations. *Journal of High Energy Physics*, 2024(4):126, 2024. doi:
583 10.1007/JHEP04(2024)126. URL [https://doi.org/10.1007/JHEP04\(2024\)126](https://doi.org/10.1007/JHEP04(2024)126).
- 584 Joan Bruna and Jiequn Han. Posterior sampling with denoising oracles via tilted transport, 2024.
585 URL <https://arxiv.org/abs/2407.00745>.
- 586
587 Michele Caselle, Elia Cellini, Alessandro Nada, and Marco Panero. Stochastic normalizing flows
588 as non-equilibrium transformations. *Journal of High Energy Physics*, 2022(7):15, 2022. doi:
589 10.1007/JHEP07(2022)015. URL [https://doi.org/10.1007/JHEP07\(2022\)015](https://doi.org/10.1007/JHEP07(2022)015).
- 590 Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David K Duvenaud. Neural ordinary
591 differential equations. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and
592 R. Garnett (eds.), *Advances in Neural Information Processing Systems*, volume 31. Curran Asso-
593 ciates, Inc., 2018. URL [https://proceedings.neurips.cc/paper_files/paper/
2018/file/69386f6bb1dfed68692a24c8686939b9-Paper.pdf](https://proceedings.neurips.cc/paper_files/paper/2018/file/69386f6bb1dfed68692a24c8686939b9-Paper.pdf).

- 594 Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. Diffusion schrödinger
595 bridge with applications to score-based generative modeling. In *Advances in Neural Information*
596 *Processing Systems*, volume 34, pp. 17695–17709, 2021.
- 597 Pierre Del Moral. Nonlinear filtering: Interacting particle resolution. *Comptes Rendus de l'Académie*
598 *des Sciences - Series I - Mathematics*, 325(6):653–658, 1997. ISSN 0764-4442. doi: [https://doi.org/10.1016/S0764-4442\(97\)84778-7](https://doi.org/10.1016/S0764-4442(97)84778-7). URL <https://www.sciencedirect.com/science/article/pii/S0764444297847787>.
- 602 Arnaud Doucet, Nando de Freitas, and Neil J. Gordon (eds.). *Sequential Monte Carlo Methods*
603 *in Practice*. Statistics for Engineering and Information Science. Springer, 2001. ISBN 978-
604 1-4419-2887-0. doi: 10.1007/978-1-4757-3437-9. URL [https://doi.org/10.1007/](https://doi.org/10.1007/978-1-4757-3437-9)
605 [978-1-4757-3437-9](https://doi.org/10.1007/978-1-4757-3437-9).
- 606 Mingzhou Fan, Ruida Zhou, Chao Tian, and Xiaoning Qian. Path-guided particle-based sampling. In
607 *Forty-first International Conference on Machine Learning*, 2024. URL [https://openreview.](https://openreview.net/forum?id=Kt4fwiuKqf)
608 [net/forum?id=Kt4fwiuKqf](https://openreview.net/forum?id=Kt4fwiuKqf).
- 609 Michael F. Faulkner and Samuel Livingstone. Sampling algorithms in statistical physics: a guide for
610 statistics and machine learning, 2023. URL <https://arxiv.org/abs/2208.04751>.
- 612 Marylou Gabrié, Grant M. Rotskoff, and Eric Vanden-Eijnden. Adaptive monte carlo augmented with
613 normalizing flows. *Proceedings of the National Academy of Sciences*, 119(10):e2109420119, 2022.
614 doi: 10.1073/pnas.2109420119. URL [https://www.pnas.org/doi/abs/10.1073/](https://www.pnas.org/doi/abs/10.1073/pnas.2109420119)
615 [pnas.2109420119](https://www.pnas.org/doi/abs/10.1073/pnas.2109420119).
- 616 Will Grathwohl, Ricky T. Q. Chen, Jesse Bettencourt, and David Duvenaud. Scalable reversible
617 generative models with free-form continuous dynamics. In *International Conference on Learning*
618 *Representations*, 2019. URL <https://openreview.net/forum?id=rJxgknCcK7>.
- 620 Carsten Hartmann, Christof Schütte, and Wei Zhang. Jarzynski’s equality, fluctuation theorems,
621 and variance reduction: Mathematical analysis and numerical algorithms. *Journal of Sta-*
622 *tistical Physics*, 175:1214–1261, 2018. URL [https://api.semanticscholar.org/](https://api.semanticscholar.org/CorpusID:59394118)
623 [CorpusID:59394118](https://api.semanticscholar.org/CorpusID:59394118).
- 624 W. K. Hastings. Monte carlo sampling methods using markov chains and their applications.
625 *Biometrika*, 57(1):97–109, 1970. ISSN 00063444, 14643510. URL [http://www.jstor.](http://www.jstor.org/stable/2334940)
626 [org/stable/2334940](http://www.jstor.org/stable/2334940).
- 627 Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. In *Advances in*
628 *neural information processing systems*, volume 33, pp. 6840–6851, 2020.
- 629 Jérôme Hénin, Tony Lelièvre, Michael R. Shirts, Omar Valsson, and Lucie Delemotte. Enhanced
630 sampling methods for molecular dynamics simulations [article v1.0]. *Living Journal of Computa-*
631 *tional Molecular Science*, 4(1):1583, Dec. 2022. doi: 10.33011/livecoms.4.1.1583. URL <https://livecomsjournal.org/index.php/livecoms/article/view/v4i1e1583>.
- 632 C. Jarzynski. Nonequilibrium equality for free energy differences. *Phys. Rev. Lett.*, 78:2690–2693,
633 Apr 1997. doi: 10.1103/PhysRevLett.78.2690. URL [https://link.aps.org/doi/10.](https://link.aps.org/doi/10.1103/PhysRevLett.78.2690)
634 [1103/PhysRevLett.78.2690](https://link.aps.org/doi/10.1103/PhysRevLett.78.2690).
- 635 Yaron Lipman, Ricky TQ Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching
636 for generative modeling. In *The Eleventh International Conference on Learning Representations*,
637 2022.
- 638 Jun S. Liu. Metropolized independent sampling with comparisons to rejection sampling and impor-
639 tance sampling. *Statistics and Computing*, 6(2):113–119, 1996. doi: 10.1007/BF00162521. URL
640 <https://doi.org/10.1007/BF00162521>.
- 641 Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate
642 and transfer data with rectified flow. In *The Eleventh International Conference on Learning*
643 *Representations*, 2022.

- 648 Nikolay Malkin, Salem Lahlou, Tristan Deleu, Xu Ji, Edward Hu, Katie Everett, Dinghuai Zhang,
649 and Yoshua Bengio. Gflownets and variational inference, 2023. URL [https://arxiv.org/
650 abs/2210.00580](https://arxiv.org/abs/2210.00580).
- 651 Alexander G. D. G. Matthews, Michael Arbel, Danilo J. Rezende, and Arnaud Doucet. Continual
652 repeated annealed flow transport monte carlo. In *Proceedings of the 39th International Conference
653 on Machine Learning*, Proceedings of Machine Learning Research, Jul 2022.
- 654 Laurence Illing Midgley, Vincent Stimper, Gregor N. C. Simm, Bernhard Schölkopf, and José Miguel
655 Hernández-Lobato. Flow annealed importance sampling bootstrap. In *The Eleventh International
656 Conference on Learning Representations*, 2023. URL [https://openreview.net/forum?
657 id=XCTVFJwS9LJ](https://openreview.net/forum?id=XCTVFJwS9LJ).
- 658 Bálint Máté and François Fleuret. Learning interpolations between boltzmann densities, 2023. URL
659 <https://arxiv.org/abs/2301.07388>.
- 660 Radford M. Neal. Probabilistic inference using markov chain monte carlo methods. Technical Report
661 CRG-TR-93-1, Department of Computer Science, University of Toronto, September 1993.
- 662 Radford M. Neal. Annealed importance sampling. *Statistics and Computing*, 11(2):125–
663 139, 2001. doi: 10.1023/A:1008923215028. URL [https://doi.org/10.1023/A:
664 1008923215028](https://doi.org/10.1023/A:1008923215028).
- 665 Kirill Neklyudov, Rob Brekelmans, Daniel Severo, and Alireza Makhzani. Action matching: Learning
666 stochastic dynamics from samples. In Andreas Krause, Emma Brunskill, Kyunghyun Cho, Barbara
667 Engelhardt, Sivan Sabato, and Jonathan Scarlett (eds.), *Proceedings of the 40th International
668 Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pp.
669 25858–25889. PMLR, 23–29 Jul 2023. URL [https://proceedings.mlr.press/v202/
670 neklyudov23a.html](https://proceedings.mlr.press/v202/neklyudov23a.html).
- 671 Frank Noé, Simon Olsson, Jonas Köhler, and Hao Wu. Boltzmann generators: Sampling equi-
672 librium states of many-body systems with deep learning. *Science*, 365(6457):eaaw1147, 2019.
673 doi: 10.1126/science.aaw1147. URL [https://www.science.org/doi/abs/10.1126/
674 science.aaw1147](https://www.science.org/doi/abs/10.1126/science.aaw1147).
- 675 Matthew D. Parno and Youssef M. Marzouk. Transport map accelerated markov chain monte carlo.
676 *SIAM/ASA Journal on Uncertainty Quantification*, 6(2):645–682, 2018. doi: 10.1137/17M1134640.
677 URL <https://doi.org/10.1137/17M1134640>.
- 678 Marcin Sendera, Minsu Kim, Sarthak Mittal, Pablo Lemos, Luca Scimeca, Jarrid Rector-Brooks,
679 Alexandre Adam, Yoshua Bengio, and Nikolay Malkin. Improved off-policy training of diffusion
680 samplers, 2024. URL <https://arxiv.org/abs/2402.05098>.
- 681 Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben
682 Poole. Score-based generative modeling through stochastic differential equations. *arXiv preprint
683 arXiv:2011.13456*, 2020.
- 684 Yifeng Tian, Nishant Panda, and Yen Ting Lin. Liouville flow importance sampler. In Ruslan
685 Salakhutdinov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett, and
686 Felix Berkenkamp (eds.), *Proceedings of the 41st International Conference on Machine Learning*,
687 volume 235 of *Proceedings of Machine Learning Research*, pp. 48186–48210. PMLR, 21–27 Jul
688 2024. URL <https://proceedings.mlr.press/v235/tian24c.html>.
- 689 Suriyanarayanan Vaikuntanathan and Christopher Jarzynski. Escorted free energy simulations:
690 Improving convergence by reducing dissipation. *Phys. Rev. Lett.*, 100:190601, May 2008.
691 doi: 10.1103/PhysRevLett.100.190601. URL [https://link.aps.org/doi/10.1103/
692 PhysRevLett.100.190601](https://link.aps.org/doi/10.1103/PhysRevLett.100.190601).
- 693 Francisco Vargas, Will Sussman Grathwohl, and Arnaud Doucet. Denoising diffusion samplers.
694 In *The Eleventh International Conference on Learning Representations*, 2023. URL <https://openreview.net/forum?id=8pvnfTAbulF>.

Francisco Vargas, Shreyas Padhy, Denis Blessing, and Nikolas Nüsken. Transport meets variational inference: Controlled monte carlo diffusions. In *The Twelfth International Conference on Learning Representations*, 2024. URL <https://openreview.net/forum?id=PP1rudnxiW>.

Kenneth G. Wilson. Confinement of quarks. *Phys. Rev. D*, 10:2445–2459, Oct 1974. doi: 10.1103/PhysRevD.10.2445. URL <https://link.aps.org/doi/10.1103/PhysRevD.10.2445>.

Qinsheng Zhang and Yongxin Chen. Path integral sampler: A stochastic control approach for sampling. In *International Conference on Learning Representations*, 2021.

Qinsheng Zhang and Yongxin Chen. Path integral sampler: A stochastic control approach for sampling. In *International Conference on Learning Representations*, 2022. URL https://openreview.net/forum?id=_uCb2ynRu7Y.

4 APPENDIX

4.1 PROOFS OF SEC. 2

Here we provide the proofs of the statements made in Sec. 2 which, for the reader convenience, we recall.

Proposition 1 (Jarzynski equality). *Let (X_t, A_t) solve the coupled system of SDE/ODE*

$$\begin{aligned} dX_t &= -\varepsilon_t \nabla U_t(X_t) dt + \sqrt{2\varepsilon_t} dW_t, & X_0 &\sim \rho_0, \\ dA_t &= -\partial_t U_t(X_t) dt, & A_0 &= 0, \end{aligned} \quad (10)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \frac{\mathbb{E}[e^{A_t} h(X_t)]}{\mathbb{E}[e^{A_t}]}, \quad Z_t/Z_0 = e^{-F_t+F_0} = \mathbb{E}[e^{A_t}], \quad (12)$$

where the expectations are taken over the law of (X_t, A_t) .

Proof. Let $f_t(x, a)$ with $(x, a) \in \mathbb{R}^{d+1}$ be the PDF of the joint process (X_t, A_t) defined by the SDE (10) and (11). This PDF solves the FPE

$$\partial_t f_t = \varepsilon_t \nabla \cdot (\nabla U_t f_t + \nabla f_t) + \partial_t U_t \partial_a f_t, \quad f_{t=0}(x, a) = \delta(a) \rho_0(x). \quad (35)$$

Define

$$g_t(x) = \int_{\mathbb{R}} e^a f_t(x, a) da. \quad (36)$$

We can derive an equation for $g_t(x)$ by multiplying both sides of the FPE (35) by e^a and integrating over $a \in \mathbb{R}$. Using

$$\begin{aligned} \int_{\mathbb{R}} e^a \partial_t f_t(x, a) da &= \partial_t \int_{\mathbb{R}} e^a f_t(x, a) da = \partial_t g_t, \\ \int_{\mathbb{R}} e^a \varepsilon_t \nabla \cdot (\nabla U_t f_t + \nabla f_t) da &= \varepsilon_t \nabla \cdot \left(\nabla U_t \int_{\mathbb{R}} e^a f_t(x, a) da + \nabla \int_{\mathbb{R}} e^a f_t(x, a) da \right) \\ &= \varepsilon_t \nabla \cdot (\nabla U_t g_t + \nabla g_t), \\ \int_{\mathbb{R}} e^a \partial_t U_t \partial_a f_t da &= \partial_t U_t \int_{\mathbb{R}} e^a \partial_a f_t da \\ &= -\partial_t U_t \int_{\mathbb{R}} e^a f_t da = -\partial_t U_t g_t, \end{aligned} \quad (37)$$

where we arrived at the second equality in the third equation by integration by parts, we deduce that

$$\partial_t g_t = \varepsilon_t \nabla \cdot (\nabla U_t g_t + \nabla g_t) - \partial_t U_t g_t, \quad g_{t=0}(x) = \rho_0(x) = e^{-U_0(x)+F_0}. \quad (38)$$

The solution to this parabolic PDE is unique and it can be checked by direct substitution that it is given by

$$g_t(x) = e^{-U_t(x)+F_0}. \quad (39)$$

Note that this solution is not normalized since it contains F_0 rather than F_t . In fact it is easy to see that

$$\int_{\mathbb{R}^d} g_t(x) dx = \int_{\mathbb{R}^{d+1}} e^a f_t(x, a) dx da = e^{-F_t+F_0}, \quad (40)$$

where the first equality follows from the definition of g_t and the second from its explicit expression and the definition of the free energy that implies $\int_{\mathbb{R}^d} e^{-U_t(x)} dx = e^{-F_t}$. Equation (39) is the second equation in (12). From (39) we also deduce that, given any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\begin{aligned} \frac{\int_{\mathbb{R}^{d+1}} e^a h(x) f_t(x, a) dx da}{\int_{\mathbb{R}^{d+1}} e^a f_t(x, a) dx da} &= \frac{\int_{\mathbb{R}^d} h(x) g_t(x) dx}{\int_{\mathbb{R}^d} g_t(x) dx} \\ &= \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x)+F_0} dx}{\int_{\mathbb{R}^d} e^{-U_t(x)+F_0} dx} \\ &= \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx}{\int_{\mathbb{R}^d} e^{-U_t(x)} dx} \\ &= e^{F_t} \int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx \\ &= \int_{\mathbb{R}^d} h(x) \rho_t(x) dx. \end{aligned} \quad (41)$$

Since by definition of $f_t(x, a)$ the left hand-side of this equation can be expressed as the ratio of expectations over (X_t, A_t) in the first equation in (12) we are done. \square

Proposition 2 (Sampling with perfect additional transport.). *Let $b_t(x)$ be a solution to (14) and let X_t^b satisfy the SDE*

$$dX_t^b = -\varepsilon_t \nabla U_t(X_t^b) dt + b_t(X_t^b) dt + \sqrt{2\varepsilon_t} dW_t, \quad X_0^b \sim \rho_0, \quad (16)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then $\rho_t(x)$ is the PDF of X_T^b , i.e. for all $t \in [0, 1]$ and, given any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \mathbb{E}[h(X_t^b)], \quad (17)$$

where the expectation at the right-hand side is taken over the law of (X_t^b) .

Proof. If b_t satisfies (14), then ρ_t satisfies the FPE (15). Since (16) is the SDE associated with this FPE, (17) holds. \square

Proposition 3 (Nonequilibrium Transport Sampler (NETS)). *Let $(X_t^{\hat{b}}, A_t^{\hat{b}})$ solve the coupled system of SDE/ODE*

$$dX_t^{\hat{b}} = -\varepsilon_t \nabla U_t(X_t^{\hat{b}}) dt + \hat{b}_t(X_t^{\hat{b}}) dt + \sqrt{2\varepsilon_t} dW_t, \quad X_0^{\hat{b}} \sim \rho_0, \quad (20)$$

$$dA_t^{\hat{b}} = \nabla \cdot \hat{b}_t(X_t^{\hat{b}}) dt - \nabla U_t(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}}) dt - \partial_t U_t(X_t^{\hat{b}}) dt, \quad A_0^{\hat{b}} = 0, \quad (21)$$

where $\varepsilon_t \geq 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \frac{\mathbb{E}[e^{A_t^{\hat{b}}} h(X_t^{\hat{b}})]}{\mathbb{E}[e^{A_t^{\hat{b}}}]}, \quad Z_t/Z_0 = e^{-F_t+F_0} = \mathbb{E}[e^{A_t^{\hat{b}}}], \quad (22)$$

where the expectations are taken over the law of $(X_t^{\hat{b}}, A_t^{\hat{b}})$.

810 *Proof.* We can follow the same steps as in the proof of Proposition 1 by considering the PDF $f_t^{\hat{b}}(x, a)$
 811 of $(X_t^{\hat{b}}, A_t^{\hat{b}})$. This PDF solves the FPE

$$812 \quad \partial_t f_t^{\hat{b}} = \varepsilon_t \nabla \cdot (\nabla U_t f_t^{\hat{b}} + \nabla f_t^{\hat{b}}) - \nabla \cdot (\hat{b}_t f_t^{\hat{b}}) - (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) \partial_a f_t^{\hat{b}},$$

$$813 \quad f_{t=0}^{\hat{b}}(x, a) = \delta(a) \rho_0(x). \quad (42)$$

814 Define

$$815 \quad g_t^{\hat{b}}(x) = \int_{\mathbb{R}} e^a f_t^{\hat{b}}(x, a) da. \quad (43)$$

816 We can derive an equation for $g_t^{\hat{b}}(x)$ by multiplying both sides of the FPE (42) by e^a and integrating
 817 over $a \in \mathbb{R}$. Using

$$818 \quad \int_{\mathbb{R}} e^a \partial_t f_t^{\hat{b}} da = \partial_t \int_{\mathbb{R}} e^a f_t^{\hat{b}} da = \partial_t g_t^{\hat{b}},$$

$$819 \quad \int_{\mathbb{R}} e^a \varepsilon_t \nabla \cdot (\nabla U_t f_t^{\hat{b}} + \nabla f_t^{\hat{b}}) da = \varepsilon_t \nabla \cdot \left(\nabla U_t \int_{\mathbb{R}} e^a f_t^{\hat{b}} da + \nabla \int_{\mathbb{R}} e^a f_t^{\hat{b}} da \right),$$

$$820 \quad - \int_{\mathbb{R}} e^a \nabla \cdot (\hat{b}_t f_t^{\hat{b}}) da = -\nabla \cdot \left(\hat{b}_t \int_{\mathbb{R}} e^a f_t^{\hat{b}} da \right)$$

$$821 \quad = -\nabla \cdot (\hat{b}_t g_t^{\hat{b}}), \quad (44)$$

$$822 \quad - \int_{\mathbb{R}} e^a (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) \partial_a f_t^{\hat{b}} da = -(\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) \int_{\mathbb{R}} e^a \partial_a f_t^{\hat{b}} da$$

$$823 \quad = (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) \int_{\mathbb{R}} e^a f_t^{\hat{b}} da$$

$$824 \quad = (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) g_t^{\hat{b}},$$

825 where we arrived at the second equality in the fourth equation by integration by parts, we deduce that

$$826 \quad \partial_t g_t^{\hat{b}} = \varepsilon_t \nabla \cdot (\nabla U_t g_t^{\hat{b}} + \nabla g_t^{\hat{b}}) - \nabla \cdot (\hat{b}_t g_t^{\hat{b}}) + (\nabla \cdot \hat{b}_t - \nabla U_t - \partial_t U_t) g_t^{\hat{b}},$$

$$827 \quad g_{t=0}^{\hat{b}}(x) = \rho_0(x) = e^{-U_0(x) + F_0}. \quad (45)$$

828 The solution to this parabolic PDE is unique and it can be checked by direct substitution that it is
 829 given by

$$830 \quad g_t^{\hat{b}}(x) = e^{-U_t(x) + F_0}. \quad (46)$$

831 This solution is not normalized since it contains F_0 rather than F_t , and it is easy to see that

$$832 \quad \int_{\mathbb{R}^d} g_t^{\hat{b}}(x) dx = \int_{\mathbb{R}^{d+1}} e^a f_t^{\hat{b}}(x, a) dx da = e^{-F_t + F_0}. \quad (47)$$

833 where the first equality follows from the definition of $g_t^{\hat{b}}$ and the second from its explicit expression
 834 and the definition of the free energy that implies $\int_{\mathbb{R}^d} e^{-U_t(x)} dx = e^{-F_t}$. Equation (47) is the second
 835 equation in (22). From (46) we also deduce that, given any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$836 \quad \frac{\int_{\mathbb{R}^{d+1}} e^a h(x) f_t^{\hat{b}}(x, a) dx da}{\int_{\mathbb{R}^{d+1}} e^a f_t^{\hat{b}}(x, a) dx da} = \frac{\int_{\mathbb{R}^d} h(x) g_t^{\hat{b}}(x) dx}{\int_{\mathbb{R}^d} g_t^{\hat{b}}(x) dx}$$

$$837 \quad = \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x) + F_0} dx}{\int_{\mathbb{R}^d} e^{-U_t(x) + F_0} dx}$$

$$838 \quad = \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx}{\int_{\mathbb{R}^d} e^{-U_t(x)} dx} \quad (48)$$

$$839 \quad = e^{F_t} \int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx$$

$$840 \quad = \int_{\mathbb{R}^d} h(x) \rho_t(x) dx.$$

841 Since by definition of $f_t^{\hat{b}}(x, a)$ the left hand-side of this equation can be expressed as the ratio of
 842 expectations over $(X_t^{\hat{b}}, A_t^{\hat{b}})$ in the first equation in (22) we are done. \square

Proposition 4 (PINN objective). *Given any $T \in (0, 1]$ and any PDF $\hat{\rho}_t(x) > 0$ consider the objective for (\hat{b}, \hat{F}) given by:*

$$L_{PINN}^T[\hat{b}, \hat{F}] = \int_0^T \int_{\mathbb{R}^d} |\nabla \cdot \hat{b}_t(x) - \nabla U_t(x) \cdot \hat{b}_t(x) - \partial_t U_t(x) + \partial_t \hat{F}_t|^2 \hat{\rho}_t(x) dx dt. \quad (26)$$

Then $\min_{\hat{b}, \hat{F}} L_{PINN}^T[\hat{b}, \hat{F}] = 0$, and all minimizers (b, F) are such that $b_t(x)$ solves (24) and F_t is the free energy (3) for all $t \in [0, T]$.

Proof. Clearly the minimum value of (26) is zero and the minimizing pair (\hat{b}, \hat{F}) must satisfy

$$\nabla \cdot \hat{b}_t - \nabla U_t \cdot \hat{b}_t - \partial_t U_t + \partial_t \hat{F}_t = 0 \quad (49)$$

By multiplying both sides of this equation by ρ_t it can be written as

$$\nabla \cdot (\hat{b}_t \rho_t) - \partial_t U_t \rho_t + \partial_t \hat{F}_t \rho_t = 0 \quad (50)$$

This equation requires a solvability condition obtained by integrating it over \mathbb{R}^d . This gives

$$-\int_{\mathbb{R}^d} \partial_t U_t(x) \rho_t(x) dx + \partial_t \hat{F}_t = 0, \quad (51)$$

which, by (4), implies that $\partial_t \hat{F}_t = \partial_t F_t$. In turn, this implies that (50) is equivalent to (14), i.e. \hat{b}_t solves (14). \square

Proposition 5 (KL control). *Let $\hat{\rho}_t$ be the solution to the transport equation*

$$\partial_t \hat{\rho}_t = -\nabla \cdot (\hat{b}_t \hat{\rho}_t), \quad \hat{\rho}_{t=0} = \rho_0 \quad (27)$$

where $\hat{b}_t(x)$ is some predefined velocity field. Then, we have

$$D_{KL}(\hat{\rho}_{t=1} || \rho_1) \leq \sqrt{L_{PINN}^{T=1}(\hat{b}, F)}. \quad (28)$$

where F_t is the free energy. In addition, given any estimate \hat{F}_t such that $\int_0^1 |\partial_t \hat{F}_t - \partial_t F_t|^2 dt \leq \delta$ for some $\delta \geq 0$, we have

$$D_{KL}(\hat{\rho}_{t=1} || \rho_1) \leq \sqrt{2L_{PINN}^{T=1}(\hat{b}, \hat{F})} + 2\delta. \quad (29)$$

Proof. Consider

$$D_{KL}(\hat{\rho}_t || \rho_t) = \int_{\mathbb{R}^d} \log \left(\frac{\hat{\rho}_t(x)}{\rho_t(x)} \right) \hat{\rho}_t(x) dx \quad (52)$$

where $\hat{\rho}_t$ satisfies (27). Taking the time-derivative of this expression we deduce that (using (27), $\rho_t(x) = e^{-U_t(x)+F_t}$, and multiple integrations by parts)

$$\begin{aligned} \partial_t D_{KL}(\hat{\rho}_t || \rho_t) &= \int_{\mathbb{R}^d} \left[\log \left(\frac{\hat{\rho}_t(x)}{\rho_t(x)} \right) \partial_t \hat{\rho}_t(x) - \frac{\partial_t \rho_t(x)}{\rho_t(x)} \hat{\rho}_t(x) \right] dx \\ &= \int_{\mathbb{R}^d} \left[-\log \left(\frac{\hat{\rho}_t(x)}{\rho_t(x)} \right) \nabla \cdot (\hat{b}_t(x) \hat{\rho}_t(x)) + (\partial_t U_t(x) - \partial_t F_t) \hat{\rho}_t(x) \right] dx \\ &= \int_{\mathbb{R}^d} \left[\hat{b}_t(x) \cdot \nabla \log \left(\frac{\hat{\rho}_t(x)}{\rho_t(x)} \right) + \partial_t U_t - \partial_t F_t \right] \hat{\rho}_t(x) dx \\ &= \int_{\mathbb{R}^d} \left[\hat{b}_t(x) \cdot \nabla \hat{\rho}_t(x) + (\hat{b}_t(x) \cdot \nabla U_t(x) + \partial_t U_t - \partial_t F_t) \hat{\rho}_t(x) \right] dx \\ &= \int_{\mathbb{R}^d} \left[-\nabla \cdot \hat{b}_t(x) + \hat{b}_t(x) \cdot \nabla U_t(x) + \partial_t U_t - \partial_t F_t \right] \hat{\rho}_t(x) dx \end{aligned} \quad (53)$$

Therefore

$$\begin{aligned} D_{KL}(\hat{\rho}_{t=1} || \rho_1) &= \int_0^1 \int_{\mathbb{R}^d} \left[-\nabla \cdot \hat{b}_t(x) + \hat{b}_t(x) \cdot \nabla U_t(x) + \partial_t U_t - \partial_t F_t \right] \hat{\rho}_t(x) dx dt \\ &\leq \left[\int_0^1 \int_{\mathbb{R}^d} \left| -\nabla \cdot \hat{b}_t(x) + \hat{b}_t(x) \cdot \nabla U_t(x) + \partial_t U_t - \partial_t F_t \right|^2 \hat{\rho}_t(x) dx dt \right]^{1/2} \\ &= \sqrt{L_{PINN}^{T=1}(\hat{b}, F)} \end{aligned} \quad (54)$$

918 which gives (28). To establish (29) observe that

$$\begin{aligned}
919 & L_{\text{PINN}}^1(\hat{b}, F) \\
920 & = \int_0^1 \int_{\mathbb{R}^d} \left| \nabla \cdot \hat{b}_t(x) - \hat{b}_t(x) \cdot \nabla U_t(x) - \partial_t U_t + \partial_t F_t \right|^2 \hat{\rho}_t(x) dx dt \\
921 & \leq 2 \int_0^1 \int_{\mathbb{R}^d} \left[\left| \nabla \cdot \hat{b}_t(x) - \hat{b}_t(x) \cdot \nabla U_t(x) - \partial_t U_t + \partial_t \hat{F}_t \right|^2 + \left| \partial_t F_t - \partial_t \hat{F}_t \right|^2 \right] \hat{\rho}_t(x) dx dt \quad (55) \\
922 & = 2L_{\text{PINN}}^1(\hat{b}, \hat{F}) + 2 \int_0^1 |\partial_t F_t - \partial_t \hat{F}_t|^2 dt
\end{aligned}$$

923 Therefore, if $\int_0^1 |\partial_t \hat{F}_t - \partial_t F_t|^2 dt \leq \delta$, we have

$$924 \quad L_{\text{PINN}}^1(\hat{b}, F) \leq 2L_{\text{PINN}}^1(\hat{b}, \hat{F}) + 2\delta \quad (56)$$

925 Combining this bound with (28) gives (29). \square

926 **Proposition 6** (Action Matching objective). *Given any $T \in (0, 1]$ consider the objective for $\hat{\phi}_t(x)$:*

$$927 \quad L_{\text{AM}}^T[\hat{\phi}] = \int_0^T \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla \hat{\phi}_t(x)|^2 + \partial_t \hat{\phi}_t(x) \right] \rho_t(x) dx dt + \int_{\mathbb{R}^d} [\hat{\phi}_0(x) \rho_0(x) - \hat{\phi}_T(x) \rho_T(x)] dx. \quad (31)$$

928 Then the minimizer $\phi_t(x)$ of (26) is unique (up to a constant) and $b_t(x) = \nabla \phi_t(x)$ satisfies (14) for

929 all $t \in [0, T]$.

930 *Proof.* By integrating by parts in time the term involving $\partial_t \phi_t$ in the AM objective (57), we can

931 express is as

$$932 \quad L_{\text{AM}}^T[\hat{\phi}] = \int_0^T \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla \hat{\phi}_t(x)|^2 \rho_t(x) - \phi_t(x) \partial_t \rho_t(x) \right] dx dt. \quad (57)$$

933 This is a convex objective in $\hat{\phi}$ whose minimizers satisfy

$$934 \quad \nabla \cdot (\nabla \hat{\phi}_t \rho_t) = -\partial_t \rho_t. \quad (58)$$

935 This is (14) written in terms of $b_t(x) = \nabla \phi_t(x)$. The solution of this equation is unique up

936 to a constant by the Fredholm alternative since its right hand-side satisfies the solvability condition $\int_{\mathbb{R}^d} \partial_t \rho_t(x) dx = 0$. \square

937 **Derivation of (32).** If $\hat{b}_t(x) = \nabla \hat{\phi}_t(x)$, the SDEs (20) and (21) reduce to

$$938 \quad dX_t^{\hat{b}} = -\varepsilon_t \nabla U_t(X_t^{\hat{b}}) dt + \hat{\nabla} \hat{\phi}_t(X_t^{\hat{b}}) dt + \sqrt{2\varepsilon_t} dW_t, \quad \hat{X}_0^{\hat{b}} \sim \rho_0, \quad (59)$$

$$939 \quad dA_t^{\hat{b}} = \Delta \hat{\phi}_t(X_t^{\hat{b}}) dt - \nabla U_t(X_t^{\hat{b}}) \cdot \nabla \hat{\phi}_t(X_t^{\hat{b}}) dt - \partial_t U_t(X_t^{\hat{b}}) dt, \quad A_0^{\hat{b}} = 0, \quad (60)$$

940 Since by Itô formula we have

$$\begin{aligned}
941 \quad d\hat{\phi}_t(X_t^{\hat{b}}) & = \partial_t \hat{\phi}_t(X_t^{\hat{b}}) dt - \varepsilon_t \nabla \hat{\phi}_t(X_t^{\hat{b}}) \cdot \nabla U_t(X_t^{\hat{b}}) dt + |\nabla \hat{\phi}_t(X_t^{\hat{b}})|^2 dt \\
942 & \quad + \sqrt{2\varepsilon_t} \nabla \hat{\phi}_t(X_t^{\hat{b}}) \cdot dW_t + \varepsilon_t \Delta \hat{\phi}_t(X_t^{\hat{b}}) dt,
\end{aligned} \quad (61)$$

943 we can express

$$\begin{aligned}
944 \quad \Delta \hat{\phi}_t(X_t^{\hat{b}}) dt & = \frac{1}{\varepsilon_t} d\hat{\phi}_t(X_t^{\hat{b}}) dt - \frac{1}{\varepsilon_t} \partial_t \hat{\phi}_t(X_t^{\hat{b}}) dt + \nabla \hat{\phi}_t(X_t^{\hat{b}}) \cdot \nabla U_t(X_t^{\hat{b}}) dt \\
945 & \quad - \frac{1}{\varepsilon_t} |\nabla \hat{\phi}_t(X_t^{\hat{b}})|^2 dt - \sqrt{\frac{2}{\varepsilon_t}} \nabla \hat{\phi}_t(X_t^{\hat{b}}) \cdot dW_t.
\end{aligned} \quad (62)$$

946 If we insert this expression in the SDE (60), we can write it as

$$947 \quad dA_t^{\hat{b}} = \frac{1}{\varepsilon_t} d\hat{\phi}_t(X_t^{\hat{b}}) dt + dB_t. \quad (63)$$

948 where dB_t is given by (33). Integrating (63) gives (32).

4.2 TIME-DISCRETIZED VERSION OF PROPOSITION 3

Here we show how to generalize the result in Proposition 3 if we time discretize the SDE in (20) using Euler-Maruyama scheme and use some suitable time-discretized version of the ODE (21).

Proposition 7. *Let $0 = t_0 < t_1 < \dots < t_K = 1$ be a time grid on $[0, 1]$, denote $\Delta t_k = t_{k+1} - t_k$ for $k = 0, \dots, K-1$, set $\tilde{X}_0^{\hat{b}} \sim \rho_0$ and $\tilde{A}_0^{\hat{b}} = 0$, and for $k = 0, \dots, K-1$ define $\tilde{X}_{t_{k+1}}^{\hat{b}}, \tilde{A}_{t_{k+1}}^{\hat{b}}$ recursively via*

$$\tilde{X}_{t_{k+1}}^{\hat{b}} = \tilde{X}_{t_k}^{\hat{b}} - \varepsilon_{t_k} \nabla U_{t_k}(\tilde{X}_{t_k}^{\hat{b}}) \Delta t_k + \hat{b}_{t_k}(\tilde{X}_{t_k}^{\hat{b}}) \Delta t_k + \sqrt{2\varepsilon_{t_k}} (W_{t_{k+1}} - W_{t_k}), \quad (64)$$

$$\tilde{A}_{t_{k+1}}^{\hat{b}} = \tilde{A}_{t_k}^{\hat{b}} + U_{t_k}(\hat{X}_{t_k}^{\hat{b}}) - U_{t_{k+1}}(\hat{X}_{t_{k+1}}^{\hat{b}}) + R_k^+(\tilde{X}_{t_k}^{\hat{b}}, \tilde{X}_{t_{k+1}}^{\hat{b}}) - R_k^-(\tilde{X}_{t_{k+1}}^{\hat{b}}, \tilde{X}_{t_k}^{\hat{b}}), \quad (65)$$

where we defined

$$R_k^\pm(x, y) = \frac{1}{4\varepsilon_{t_k} \Delta t_k} |y - x + \Delta t_k (\varepsilon_{t_k} \nabla U_{t_k}(x) \mp b_{t_k}(x))|^2 \quad (66)$$

Then for all $k = 0, \dots, K$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_{t_k}(x) dx = \frac{\mathbb{E}[e^{\tilde{A}_{t_k}^{\hat{b}}} h(\tilde{X}_{t_k}^{\hat{b}})]}{\mathbb{E}[e^{\tilde{A}_{t_k}^{\hat{b}}}]}, \quad Z_{t_k} = e^{-F_{t_k}} = \mathbb{E}[e^{\tilde{A}_{t_k}^{\hat{b}}}], \quad (67)$$

where the expectations are taken over the law of $(\tilde{X}_{t_k}^{\hat{b}}, \tilde{A}_{t_k}^{\hat{b}})$

Note that the weights in (67) correct for the bias coming from both the time evolution of $U_t(x)$ and the fact that the Euler-Maruyama update in (64) does not satisfy the detailed-balance condition locally. It can be checked by direct calculation that (65) is a consistent time-discretization of the ODE (21).

Proof. For simplicity of notations we will prove (67) for $k = K$: the argument for all the other $k = 1, \dots, K-1$ is similar. The update rule in (65) implies that

$$\begin{aligned} \tilde{A}_{t_K}^{\hat{b}} &= \sum_{k=0}^{K-1} \left(U_{t_k}(\hat{X}_{t_k}^{\hat{b}}) - U_{t_{k+1}}(\hat{X}_{t_{k+1}}^{\hat{b}}) + R_k^+(\tilde{X}_{t_k}^{\hat{b}}, \tilde{X}_{t_{k+1}}^{\hat{b}}) - R_k^-(\tilde{X}_{t_{k+1}}^{\hat{b}}, \tilde{X}_{t_k}^{\hat{b}}) \right) \\ &= U_0(\tilde{X}_{t_0}^{\hat{b}}) - U_K(\tilde{X}_{t_K}^{\hat{b}}) + \sum_{k=0}^{K-1} \left(R_k^+(\tilde{X}_{t_k}^{\hat{b}}, \tilde{X}_{t_{k+1}}^{\hat{b}}) - R_k^-(\tilde{X}_{t_{k+1}}^{\hat{b}}, \tilde{X}_{t_k}^{\hat{b}}) \right), \end{aligned} \quad (68)$$

Now, given the test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, consider

$$I[h] \equiv \mathbb{E}[e^{\tilde{A}_{t_K}^{\hat{b}}} h(\tilde{X}_{t_K}^{\hat{b}})] \quad (69)$$

Since the transition probability density function of the Euler-Maruyama update in (64) reads

$$\rho_{t_k}^+(x_{k+1}|x_k) = (4\pi\varepsilon_{t_k} \Delta t_k)^{-d/2} \exp(-R_k^+(x_k, x_{k+1})), \quad (70)$$

the joint probability density function of the path $(\tilde{X}_{t_0}^{\hat{b}}, \tilde{X}_{t_1}^{\hat{b}}, \dots, \tilde{X}_{t_K}^{\hat{b}})$ is given by

$$\begin{aligned} \rho(x_0, \dots, x_K) &= \exp(-U_0(x_0) + F_0) \prod_{k=0}^{K-1} \rho_{t_k}^+, \Delta t_k(x_{k+1}|x_k) \\ &= C \exp\left(-U_0(x_0) + F_0 - \sum_{k=0}^{K-1} R_k^+(x_k, x_{k+1})\right) \end{aligned} \quad (71)$$

where $C = \prod_{k=0}^{K-1} (4\pi\varepsilon_{t_k} \Delta t_k)^{-d/2}$. We can use this density along with the explicit expression for $\tilde{A}_{t_K}^{\hat{b}}$ in (68) to express the expectation (69) as an integral over $\rho(x_0, x_1, \dots, x_K)$

$$\begin{aligned} I[h] &= C \int_{\mathbb{R}^{d(K+1)}} dx_0 \cdots dx_K \exp\left(-U_0(x_0) + F_0 - \sum_{k=0}^{K-1} R_k^+(x_k, x_{k+1})\right) \\ &\quad \times \exp\left(U_0(x_0) - U_K(x_K) + \sum_{k=0}^{K-1} \left(R_k^+(x_k, x_{k+1}) - R_k^-(x_{k+1}, x_k)\right)\right) h(x_K) \end{aligned} \quad (72)$$

where the second exponential comes from the factor $e^{\tilde{A}_K^{\hat{b}}}$. (72) simplifies into

$$I[h] = C \int_{\mathbb{R}^{d(K+1)}} dx_0 \cdots dx_K \exp \left(-U_K(x_K) + F_0 - \sum_{k=0}^{K-1} R_k^-(x_{k+1}, x_k) \right) h(x_K) \quad (73)$$

In this expression we recognize a product of factors involving

$$\rho_{t_k}^-(x_k | x_{k+1}) = (4\pi\varepsilon_{t_k})^{-d/2} \exp \left(-R_k^-(x_{k+1}, x_k) \right), \quad (74)$$

which is the transition probability density function of the time-reversed update

$$\tilde{Y}_{t_k}^{\hat{b}} = \tilde{Y}_{t_{k+1}}^{\hat{b}} - \varepsilon_{t_k} \nabla U_{t_k}(\tilde{Y}_{t_{k+1}}^{\hat{b}}) \Delta t_k - \hat{b}_{t_k}(\tilde{Y}_{t_{k+1}}^{\hat{b}}) \Delta t_k + \sqrt{2\varepsilon_{t_k}} (W_{t_{k+1}} - W_{t_k}). \quad (75)$$

This implies in particular that we can perform the integrals in (73) sequentially over x_0, x_1, \dots, x_{K-1} to be left with

$$I[h] = \int_{\mathbb{R}^d} \exp(-U_K(x_K) + F_0) h(x_K) dx_K \quad (76)$$

Therefore

$$I[1] = \int_{\mathbb{R}^d} \exp(-U_K(x_K) + F_0) dx_K = e^{-F_K + F_0}, \quad (77)$$

which is the second equation in (67), and

$$\frac{I[h]}{I[1]} = e^{F_K - F_0} \int_{\mathbb{R}^d} \exp(-U_K(x_K) + F_0) h(x_K) dx_K = \int_{\mathbb{R}^d} h(x) \rho_{t_K}(x) dx \quad (78)$$

which is the first equation in (67). \square

4.3 SOLVING FOR THE OPTIMAL DRIFT VIA FEYNMAN-KAC FORMULA

Without loss of generality, we can always look for a solution to (24) in the form of $b_t(x) = \nabla \phi_t(x)$, so that this equation becomes the Poisson equation

$$\Delta \phi_t - \nabla U_t \cdot \nabla \phi_t = \partial_t U_t - \partial_t F_t. \quad (79)$$

The solution to this equation can be expressed via Feynman-Kac formula:

Proposition 8. *Let $X_\tau^{t,x}$ satisfy the following SDE*

$$dX_\tau^{t,x} = -\nabla U_t(X_\tau^{t,x}) d\tau + \sqrt{2} dW_\tau, \quad X_{\tau=0}^{t,x} = x \quad (80)$$

where U_t is evaluated fixed at $t \in [0, 1]$ fixed. Assume geometric ergodicity of the semi-group associated with (80), i.e. the probability distribution of the solutions to this SDE converges exponentially fast towards their unique equilibrium distribution with density $\rho_t(x)$. Then for all $(t, x) \in [0, 1] \times \mathbb{R}^d$ we have

$$\phi_t(x) = \int_0^\infty \mathbb{E}[\partial_t F_t - \partial_t U_t(X_\tau^{t,x})] d\tau \quad (81)$$

where the expectation is taken over the law of $X_\tau^{t,x}$.

Proof. By Ito formula,

$$\begin{aligned} d\phi_t(X_\tau^{t,x}) &= (\Delta \phi_t(X_\tau^{t,x}) - \nabla U_t(X_\tau^{t,x}) \cdot \nabla \phi_t(X_\tau^{t,x})) d\tau + \sqrt{2} \nabla \phi_t(X_\tau^{t,x}) \cdot dW_\tau \\ &= (\partial_t U_t(X_\tau^{t,x}) - \partial_t F_t) d\tau + \sqrt{2} \nabla \phi_t(X_\tau^{t,x}) \cdot dW_\tau \end{aligned} \quad (82)$$

where the differential is taken with respect to τ at t fixed, and we used (79) to get the second equality. If we integrate this relation on $\tau \in [0, T]$ and take expectation, we deduce that

$$\mathbb{E}[\phi_t(X_T^{t,x})] - \phi_t(x) = \int_0^T \mathbb{E}[\partial_t U_t(X_\tau^{t,x}) - \partial_t F_t] d\tau \quad (83)$$

where we use Ito isometry to zero the expectation of the martingale term involving $\sqrt{2} \nabla \phi_t(X_\tau^{t,x}) \cdot dW_\tau$. If we let $T \rightarrow \infty$, by ergodicity the first term at the left hand side converges towards a constant independent of (t, x) which we can neglect – this fixes the gauge of the solution to (79) which is unique only up to a constant. What remains in this limit is the expression (81). Note that the integral in this expression converges since $\mathbb{E}[\partial_t U_t(X_\tau^{t,x})] \rightarrow \partial_t F_t$ exponentially fast as $\tau \rightarrow \infty$ by assumption of geometric ergodicity. \square

Example: moving Gaussian distribution. Let us consider the case where

$$U_t(x) = \frac{1}{2}(x - b_t)^T A_t(x - b_t), \quad (84)$$

where $b_t \in \mathbb{R}^d$ is a time-dependent vector field and $A_t = A_t^T \in \mathbb{R}^d \times \mathbb{R}^d$ is a time-dependent positive-definite matrix: we assume that both b_t and A_t are C^1 in time, and also that $\dot{A}_t A_t = A_t \dot{A}_t$. The free energy in this example is

$$F_t = -\log Z_t, \quad Z_t = (2\pi)^{d/2} |\det A_t|^{-1/2}, \quad (85)$$

so that

$$\partial_t U_t(x) = -\dot{b}_t^T A_t(x - b_t) + \frac{1}{2}(x - b_t)^T \dot{A}_t(x - b_t), \quad \partial_t F_t = \frac{1}{2} \text{tr}(A_t^{-1} \dot{A}_t). \quad (86)$$

In this case, the SDE (80) reads

$$dX_\tau^{t,x} = -A_t(X_\tau^{t,x} - b_t)d\tau + \sqrt{2}dW_\tau, \quad X_{\tau=0}^{t,x} = x, \quad (87)$$

and its solution is

$$X_\tau^{t,x} = e^{-A_t \tau} x + (1 - e^{-A_t \tau}) b_t + \sqrt{2} \int_0^\tau e^{-A_t(\tau-\tau')} dW_{\tau'}. \quad (88)$$

This implies that (using Ito isometry)

$$\begin{aligned} \mathbb{E}[\partial_t U_t(X_\tau^{t,x})] &= -\dot{b}_t^T A_t e^{-A_t \tau} (x - b_t) + \frac{1}{2}(x - b_t)^T e^{-A_t \tau} \dot{A}_t e^{-A_t \tau} (x - b_t) \\ &\quad + \int_0^\tau \text{tr}(e^{-A_t \tau} \dot{A}_t e^{-A_t \tau}) d\tau \\ &= -\dot{b}_t^T A_t e^{-A_t \tau} (x - b_t) + \frac{1}{2}(x - b_t)^T e^{-A_t \tau} \dot{A}_t e^{-A_t \tau} (x - b_t) \\ &\quad + \frac{1}{2} \text{tr}(A_t^{-1} \dot{A}_t) - \frac{1}{2} \text{tr}(A_t^{-1} \dot{A}_t e^{-2A_t \tau}). \end{aligned} \quad (89)$$

Therefore, from (81), we have (using also (85))

$$\begin{aligned} \phi_t(x) &= \int_0^\infty (\dot{b}_t^T A_t e^{-A_t \tau} (x - b_t) - \frac{1}{2}(x - b_t)^T e^{-A_t \tau} \dot{A}_t e^{-A_t \tau} (x - b_t) \\ &\quad + \frac{1}{2} \text{tr}(A_t^{-1} \dot{A}_t e^{-2A_t \tau})) d\tau \\ &= \dot{b}_t \cdot (x - b_t) - \frac{1}{4}(x - b_t)^T \dot{A}_t A_t^{-1} (x - b_t) + \frac{1}{4} \text{tr}(A_t^{-1} \dot{A}_t A_t^{-1}). \end{aligned} \quad (90)$$

This solution checks out since it implies that

$$-\nabla U_t(x) \cdot \nabla \phi_t(x) + \Delta \phi_t(x) = -\dot{b}_t^T A_t (x - b_t) + \frac{1}{2}(x - b_t)^T \dot{A}_t (x - b_t) - \frac{1}{2} \text{tr}(A_t^{-1} \dot{A}_t), \quad (91)$$

which is $\partial_t U_t(x) - \partial_t F_t$ as it should.

4.4 EXTENSIONS AND GENERALIZATIONS

4.4.1 INERTIAL NETS

It is straightforward to generalize Proposition 9 so that the stochastic dynamics involves some memory/inertia:

Proposition 9. Let $(X_t^{\hat{b},\mu}, R_t^{\hat{b},\mu}, A_t^{\hat{b},\mu})$ solve the coupled system of SDE/ODE

$$dX_t^{\hat{b},\mu} = \hat{b}_t(X_t^{\hat{b},\mu}) dt + R_t^{\hat{b},\mu} dt, \quad X_0^{\hat{b},\mu} \sim \rho_0, \quad (92)$$

$$dR_t^{\hat{b},\mu} = -\mu \nabla U_t(X_t^{\hat{b},\mu}) dt - \mu \varepsilon_t^{-1} R_t^{\hat{b},\mu} dt + \mu \sqrt{2\varepsilon_t^{-1}} dW_t, \quad R_0^{\hat{b},\mu} \sim N(0, \mu Id), \quad (93)$$

$$dA_t^{\hat{b},\mu} = \nabla \cdot \hat{b}_t(X_t^{\hat{b},\mu}) dt - \nabla U_t(X_t^{\hat{b},\mu}) \cdot \hat{b}_t(X_t^{\hat{b},\mu}) dt - \partial_t U_t(X_t^{\hat{b},\mu}) dt, \quad A_0^{\hat{b},\mu} = 0, \quad (94)$$

where $\varepsilon_t > 0$ is a time-dependent diffusion coefficient, $\mu \geq 0$ is a mobility coefficient, and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t(x) dx = \frac{\mathbb{E}[e^{A_t^{\hat{b},\mu}} h(X_t^{\hat{b},\mu})]}{\mathbb{E}[e^{A_t^{\hat{b},\mu}}]}, \quad Z_t/Z_0 = e^{-F_t+F_0} = \mathbb{E}[e^{A_t^{\hat{b},\mu}}], \quad (95)$$

where the expectations are taken over the law of $(X_t^{\hat{b},\mu}, A_t^{\hat{b},\mu})$.

The proof of this proposition can be found at the end of this subsection. Note that when $\hat{b} = b$, the solution to (24), (94) is simply

$$A_t^{b,\gamma} = -F_t + F_0, \quad (96)$$

i.e. the weights are again deterministic with zero variance. In general, \hat{b} will not be the optimal one, in which case using the SDE in (92)-(94) gives us the extra parameter μ to play with post-training to improve the ESS. Below we show that (92)-(94) reduce to (20)-(21) in the limit as $\mu \rightarrow \infty$. It is also easy to see that, if we set $\mu = 0$ in (92)-(94), we simply get that $R_t^{\hat{b},\mu} = 0$ and hence (97) reduces to the ODE $dX_t^{\hat{b},\mu} = \hat{b}_t(X_t^{\hat{b},\mu})dt$. Finally it is worth noting that (92)-(93) can be cast into Langevin equations with some extra forces. Indeed, if we introduce the velocity $V_t^{\hat{b},\mu} = \hat{b}_t(X_t^{\hat{b},\mu}) + R_t^{\hat{b},\mu}$, (92)-(93) can be written as

$$\begin{aligned} dX_t^{\hat{b},\mu} &= V_t^{\hat{b},\mu} dt & X_0^{\hat{b},\mu} &\sim \rho_0, \\ dV_t^{\hat{b},\mu} &= -\mu \nabla U_t(X_t^{\hat{b},\mu}) dt + \mu \varepsilon_t^{-1} \hat{b}_t(X_t^{\hat{b},\mu}) dt - \partial_t \hat{b}_t(X_t^{\hat{b},\mu}) dt \\ &\quad + \nabla b_t(X_t^{\hat{b},\mu}) V_t^{\hat{b},\mu} dt - \mu \varepsilon_t^{-1} V_t^{\hat{b},\mu} dt + \mu \sqrt{2\varepsilon_t^{-1}} dW_t, & V_0^{\hat{b},\mu} &\sim N(\hat{b}_0(X_0^{\hat{b},\mu}), \mu \text{Id}) \end{aligned} \quad (98)$$

In these equations, the terms $\mu \varepsilon_t^{-1} \hat{b}_t - \partial_t \hat{b}_t$ can be interpreted as non-conservative forces added to $-\mu \nabla U_t$, and the term $\nabla b_t V_t^{\hat{b},\mu}$ as an extra friction term added to $-\mu \varepsilon_t^{-1} V_t^{\hat{b},\mu}$.

Proof of Proposition 9. Denote by $f_t^{\hat{b},\mu}(x, r, a)$ the joint PDF of $(X_t^{\hat{b},\mu}, R_t^{\hat{b},\mu}, A_t^{\hat{b},\mu})$. This PDF satisfies the FPE

$$\begin{aligned} \partial_t f_t^{\hat{b},\mu} &= -\nabla_x \cdot ([\hat{b}_t + r] f_t^{\hat{b},\mu}) + \mu \nabla U_t \cdot \nabla_r f + \mu \varepsilon_t^{-1} \nabla_r \cdot (r f_t^{\hat{b},\mu} + \mu \nabla_r f_t^{\hat{b},\mu}) \\ &\quad - (\nabla \cdot \hat{b}_t - \nabla U_t \cdot \hat{b}_t - \partial_t U_t) \partial_a f_t^{\hat{b},\mu}, \\ f_0^{\hat{b},\mu}(x, r, a) &= \rho_0(x) (2\pi\mu)^{-d/2} e^{-|r|^2/(2\mu)} \delta(a). \end{aligned} \quad (99)$$

Let

$$g_t^{\hat{b},\mu}(x, r) = \int_{\mathbb{R}} e^a f_t^{\hat{b},\mu}(x, r, a) da. \quad (100)$$

We can derive an equation for $g_t^{\hat{b},\mu}(x)$ by multiplying both sides of the FPE (99) by e^a and integrating over $a \in \mathbb{R}$. Using equations similar to (44), we arrive at

$$\begin{aligned} \partial_t g_t^{\hat{b},\mu} &= -\nabla_x \cdot ([\hat{b}_t + r] g_t^{\hat{b},\mu}) + \mu \nabla U_t \cdot \nabla_r f + \mu \varepsilon_t^{-1} \nabla_r \cdot (r g_t^{\hat{b},\mu} + \mu \nabla_r g_t^{\hat{b},\mu}) \\ &\quad + (\nabla \cdot \hat{b}_t - \nabla U_t \cdot \hat{b}_t - \partial_t U_t) g_t^{\hat{b},\mu}, \\ g_0^{\hat{b},\mu}(x, r) &= \rho_0(x) (2\pi\mu)^{-d/2} e^{-|r|^2/(2\mu)}. \end{aligned} \quad (101)$$

Since $\rho_0(x) = e^{-U_0(x)+F_0}$, it can be checked by direct substitution that the solution to this equation is

$$g_t^{\hat{b},\mu}(x, r) = e^{-U_t(x)+F_0} (2\pi\mu)^{-d/2} e^{-|r|^2/(2\mu)}. \quad (102)$$

Therefore

$$\int_{\mathbb{R}^{2d}} g_t^{\hat{b},\mu}(x, r) dx dr = \int_{\mathbb{R}^{2d+1}} e^a f_t^{\hat{b},\mu}(x, r, a) dx dr da = e^{-F_t+F_0}, \quad (103)$$

where the first equality follows from the definition of $g_t^{\hat{b},\mu}$ and the second from its explicit expression and the definition of the free energy that implies $\int_{\mathbb{R}^d} e^{-U_t(x)} dx = e^{-F_t}$. Equation (103) is the second

equation in (95). From (102) we also deduce that, given any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\begin{aligned}
\frac{\int_{\mathbb{R}^{2d+1}} e^a h(x) f_t^{\hat{b},\mu}(x, r, a) dx dr da}{\int_{\mathbb{R}^{2d+1}} e^a f_t^{\hat{b},\mu}(x, r, a) dx dr da} &= \frac{\int_{\mathbb{R}^{2d}} h(x) g_t^{\hat{b},\mu}(x, r) dx dr}{\int_{\mathbb{R}^{2d}} g_t^{\hat{b},\mu}(x, r) dx dr} \\
&= \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x)+F_0} dx}{\int_{\mathbb{R}^d} e^{-U_t(x)+F_0} dx} \\
&= \frac{\int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx}{\int_{\mathbb{R}^d} e^{-U_t(x)} dx} \tag{104} \\
&= e^{F_t} \int_{\mathbb{R}^d} h(x) e^{-U_t(x)} dx \\
&= \int_{\mathbb{R}^d} h(x) \rho_t(x) dx.
\end{aligned}$$

Since by definition of $f_t^{\hat{b},\mu}(x, r, a)$ the left hand-side of this equation can be expressed as the ratio of expectations over $(X_t^{\hat{b},\mu}, A_t^{\hat{b},\mu})$ in the first equation in (95) we are done. \square

To see what happens when $\mu \rightarrow \infty$, let us assume that $\varepsilon_t = \varepsilon$ (time-independent) and integrate (93) using Duhamel principle as

$$R_t^{\hat{b},\mu} = e^{-\mu\varepsilon^{-1}t} R_0^{\hat{b},\mu} - \mu \int_0^t e^{-\mu\varepsilon^{-1}(t-s)} \nabla U_s(X_s^{\hat{b},\mu}) ds + \mu \sqrt{2\varepsilon^{-1}} \int_0^t e^{-\mu\varepsilon^{-1}(t-s)} dW_s, \tag{105}$$

Letting $\mu \rightarrow \infty$, we see that the first term at the right hand side of (105) tends to zero, whereas the second one gives

$$\lim_{\mu \rightarrow \infty} \mu \int_0^t e^{-\mu\varepsilon^{-1}(t-s)} \nabla U_s(X_s^{\hat{b},\mu}) ds = \varepsilon \nabla U_t(X_t^{\hat{b},\mu}) \tag{106}$$

Finally, the third term at the right hand side of (105) is a Gaussian process with covariance

$$C_{t,t'}^\mu = 2\mu^2 \varepsilon^{-1} \int_0^{\min(t,t')} e^{-\mu\varepsilon^{-1}(t-s) - \mu\varepsilon^{-1}(t'-s)} ds = 2\mu \left(e^{-\mu\varepsilon^{-1}|t-t'|} - e^{-\mu\varepsilon^{-1}(t+t')} \right) \tag{107}$$

As a result, given any test function ϕ_t , we have

$$\lim_{\mu \rightarrow \infty} \int_{[0,1]^2} \phi_t C_{t,t'}^\mu \phi_{t'} dt dt' = 2\varepsilon \int_0^1 \phi_t^2 dt \tag{108}$$

which indicates that $C_{t,t'}^\mu$ converges weakly towards the Dirac distribution $\varepsilon \delta(t - t')$. Putting these results together shows that in the limit as $\mu \rightarrow \infty$, $R_t^{\hat{b},\mu} dt$ converges weakly towards $-\varepsilon \nabla U_t(X_t^{\hat{b},\mu}) dt + \sqrt{2\varepsilon} dW_t$, which, if inserted in (92), reduces this equation to (20). The case where ε_t depends on time can be treated similarly.

4.4.2 MULTIMARGINAL NETS

Let $\mathcal{U}(\alpha, x)$ be a potential depending on $\alpha \in D \subset \mathbb{R}^N$ with $N \in \mathbb{N}$ as well as $x \in \mathbb{R}^d$, and assumed to be continuously differentiable in both arguments. Assume that $e^{-\mathcal{U}(\alpha, x)}$ is integrable in x for all $\alpha \in D$, and define the family of PDF

$$\varrho(\alpha, x) = e^{-\mathcal{U}(\alpha, x) + \mathcal{F}(\alpha)}, \quad \mathcal{F}(\alpha) = -\log \int_{\mathbb{R}^d} e^{-\mathcal{U}(\alpha, x)} dx. \tag{109}$$

Finally, define the family of matrix-valued $\hat{B}(\alpha, x) : D \times \mathbb{R}^d \rightarrow \mathbb{R}^N \times \mathbb{R}^d$, assumed to be continuously differentiable in both arguments. These quantities allow us to give a generalization of Proposition 3 in which we can sample the PDF $\varrho(\alpha, x)$ along any differential path $\alpha_t \in D$:

Proposition 10. Let $\alpha : [0, 1] \rightarrow D$ be a differentiable path in D and define the vector field $b : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ as

$$\hat{b}_t^\alpha(x) = \dot{\alpha}_t^T \hat{\mathcal{B}}(\alpha_t, x) \quad (110)$$

as well as

$$U_t^\alpha(x) = \mathcal{U}(\alpha_t, x), \quad F_t^\alpha = \mathcal{F}(\alpha_t), \quad \rho_t^\alpha = \varrho(\alpha, x) = e^{-U_t^\alpha(x) + F_t^\alpha} \quad (111)$$

Let $(X_t^{\hat{b}, \alpha}, A_t^{\hat{b}, \alpha})$ solve the coupled system of SDE/ODE

$$dX_t^{\hat{b}, \alpha} = \hat{b}_t^\alpha(X_t^{\hat{b}, \alpha})dt - \varepsilon_t \nabla U_t^\alpha(X_t^{\hat{b}, \alpha})dt + \sqrt{2\varepsilon_t}dW_t \quad X_0^{\hat{b}, \alpha} \sim \rho_0^\alpha, \quad (112)$$

$$dA_t^{\hat{b}, \alpha} = \nabla \cdot \hat{b}_t^\alpha(X_t^{\hat{b}, \alpha})dt - \nabla U_t^\alpha(X_t^{\hat{b}, \alpha}) \cdot \hat{b}_t^\alpha(X_t^{\hat{b}, \alpha})dt - \partial_t U_t^\alpha(X_t^{\hat{b}, \alpha})dt, \quad A_0^{\hat{b}, \alpha} = 0, \quad (113)$$

where $\varepsilon_t > 0$ is a time-dependent diffusion coefficient and $W_t \in \mathbb{R}^d$ is the Wiener process. Then for all $t \in [0, 1]$ and any test function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\int_{\mathbb{R}^d} h(x) \rho_t^\alpha(x) dx = \frac{\mathbb{E}[e^{A_t^{\hat{b}, \alpha}} h(X_t^{\hat{b}, \alpha})]}{\mathbb{E}[e^{A_t^{\hat{b}, \alpha}}]}, \quad e^{-F_t^\alpha + F_0^\alpha} = \mathbb{E}[e^{A_t^{\hat{b}, \alpha}}], \quad (114)$$

where the expectations are taken over the law of $(X_t^{\hat{b}, \alpha}, A_t^{\hat{b}, \alpha})$.

We will omit to give the proof of this proposition since it is a simple consequence of Proposition 3. The interest in formulating the problem in this new way is that it is easy to see that the right hand side of (113) (with $\partial_t F_t^\alpha$ added for convenience) can be written as

$$\begin{aligned} & \nabla \cdot \hat{b}_t^\alpha(x) - \nabla U_t^\alpha(x) \cdot \hat{b}_t^\alpha(x) - \partial_t U_t^\alpha(x) + \partial_t F_t^\alpha \\ &= \dot{\alpha}_t^T \left(\nabla_x \cdot \hat{\mathcal{B}}(\alpha_t, x) - \hat{\mathcal{B}}(\alpha_t, x) \nabla_x \mathcal{U}(\alpha_t, x) - \nabla_\alpha \mathcal{U}(\alpha_t, x) + \nabla_\alpha \mathcal{F}(\alpha_t) \right). \end{aligned} \quad (115)$$

Therefore if we zero this term for all $(\alpha, x) \in D \times \mathbb{R}^d$ by picking the right $\hat{\mathcal{B}}(\alpha, x)$ we will obtain that (113) reduces to $dA_t^{\hat{b}, \alpha} = -F_t^\alpha dt$, i.e. $A_t^{\hat{b}, \alpha} = F_t^\alpha - F_0^\alpha$. Finding this optimal $\mathcal{B}(\alpha, x)$ can be obtained using the following result:

Proposition 11 (Multimarginal PINN objective). Consider the objective for $(\hat{\mathcal{B}}, \hat{\mathcal{F}})$ given by:

$$\begin{aligned} & L_{PINN}^\alpha[\hat{\mathcal{B}}, \hat{\mathcal{F}}] \\ &= \int_D \int_{\mathbb{R}^d} \left| \nabla_x \cdot \hat{\mathcal{B}}(\alpha, x) - \hat{\mathcal{B}}(\alpha, x) \nabla_x \mathcal{U}(\alpha, x) - \nabla_\alpha \mathcal{U}(\alpha, x) + \nabla_\alpha \hat{\mathcal{F}}(\alpha) \right|^2 \hat{\varrho}(\alpha, x) f(\alpha) dx d\alpha \end{aligned} \quad (116)$$

where $\hat{\varrho}(\alpha, x) > 0$ is a PDF in x for all $\alpha \in D$, and $f(\alpha)$ is a PDF in α . Then $\min_{\mathcal{B}, \mathcal{F}} L_{PINN}^\alpha[\hat{\mathcal{B}}, \hat{\mathcal{F}}] = 0$, and all minimizers $(\mathcal{B}, \mathcal{F})$ are such that and $\mathcal{B}(\alpha, x)$ solves

$$\forall (\alpha, x) \in D \times \mathbb{R}^d : \quad 0 = \nabla_x \cdot \hat{\mathcal{B}}(\alpha, x) - \hat{\mathcal{B}}(\alpha, x) \nabla_x \mathcal{U}(\alpha, x) - \nabla_\alpha \mathcal{U}(\alpha, x) + \nabla_\alpha \mathcal{F}(\alpha), \quad (117)$$

and $\mathcal{F}(\alpha)$ is the free energy (109) for all $\alpha \in D$.

We will omit to give the proof of this proposition since it is a simple generalization of the proof of Proposition 4.

4.5 IMPLEMENTATION

The computation of the divergence $\nabla \cdot b_t(x)$ in the PINN objective given in (26) can be avoided by using Hutchinson's trace estimator, see Appendix 4.5.1. If we minimize (26) *off-policy*, i.e. with samples from some $\hat{\rho}_t \neq \rho_t$, this is perfectly valid, but may be inefficient for learning \hat{b}_t over the support necessary for the problem. If we decide instead to set $\hat{\rho}_t(x) = \rho_t(x)$, since the SDEs in (20) and (21) can be used with any $\hat{b}_t(x)$ to estimate expectation over $\rho_t(x)$ via (22), we can write the PINN objective on-policy as

$$L_{PINN}^T[\hat{b}, \hat{F}] = \int_0^T \frac{1}{\mathbb{E}[e^{A_t^{\hat{b}}}] } \mathbb{E} \left[e^{A_t^{\hat{b}}} \left| \nabla \cdot \hat{b}_t(X_t^{\hat{b}}) - \nabla U_t(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}}) - \partial_t U_t(X_t^{\hat{b}}) + \partial_t \hat{F}_t \right|^2 \right] dt \quad (118)$$

1296 These expectations can be estimated empirically over a population of solutions to (20) and (21).
 1297 Crucially, since we can switch from off-policy to on-policy after taking the gradient of the PINN
 1298 objective, *when computing the gradient of (118) over $\hat{b}_t(x)$, $(X_t^{\hat{b}}, A_t^{\hat{b}})$ can be considered independent*
 1299 *of $\hat{b}_t(x)$ and do not need to be differentiated over.* In other words, the method does not require
 1300 backpropagation through the simulation even if used on-policy, i.e. even though it uses the current
 1301 value of \hat{b}_t to estimate the loss and its gradient. Finally note that we can use the ODE (21) for $A_t^{\hat{b}}$ to
 1302 write (118) as

$$1303 L_{\text{PINN}}^T[\hat{b}, \hat{F}] = \int_0^T \frac{1}{\mathbb{E}[e^{A_t^{\hat{b}}}] \mathbb{E}[e^{A_t^{\hat{b}}}] \mathbb{E}[e^{A_t^{\hat{b}}} | \partial_t A_t^{\hat{b}} + \partial_t \hat{F}_t]^2} dt \quad (119)$$

1304 Since $\mathbb{E}[e^{A_t^{\hat{b}}} \partial_t A_t^{\hat{b}}] / \mathbb{E}[e^{A_t^{\hat{b}}}] = \partial_t \log \mathbb{E}[e^{A_t^{\hat{b}}}] = -\partial_t F_t$, (119) clearly shows that this loss controls the
 1305 variance of $\partial_t A_t^{\hat{b}}$, which directly connects the Jarzynski weights to the PINN objective.

1306 Learning $b_t(x)$ and F_t for $t \in [0, 1]$ from the start can be challenging if the initial $\hat{b}_t(x)$ is far from
 1307 exact and the weights gets large variance as t increases. This problem can be alleviated by estimating
 1308 $b_t(x)$ sequentially. In practice, this amounts to annealing T from a small initial value to $T = 1$, in
 1309 such a way that $b_t(x)$ is learned sufficiently accurately so that variance of the weights remains small.
 1310 This variance can be estimated on the fly, which also give us an estimate of the effective sample size
 1311 (ESS) of the population at all times $t \in [0, 1]$.

1312 Note that we can also employ resampling strategies of the type used in SMC to keep the variance of
 1313 the weights low Doucet et al. (2001); Bolić et al. (2004).

1314 We can proceed similarly with the AM loss (31) by rewriting it as

$$1315 L_{\text{AM}}^T[\hat{\phi}] = \int_0^T \frac{\mathbb{E}[e^{A_t^{\hat{\phi}}} [\frac{1}{2} |\nabla \hat{\phi}_t(X_t^{\hat{\phi}})|^2 + \partial_t \phi_t(X_t^{\hat{\phi}})]]}{\mathbb{E}[e^{A_t^{\hat{\phi}}}]^2} dt + \frac{\mathbb{E}[e^{A_0^{\hat{\phi}}} \phi_0(X_0^{\hat{\phi}})]}{\mathbb{E}[e^{A_0^{\hat{\phi}}}]^2} - \frac{\mathbb{E}[e^{A_T^{\hat{\phi}}} \phi_T(X_T^{\hat{\phi}})]}{\mathbb{E}[e^{A_T^{\hat{\phi}}}]^2}. \quad (120)$$

1316 These expectations can be estimated empirically over solutions to (20) and (21) with $\hat{b}_t(x) = \nabla \hat{\phi}_t(x)$.
 1317 The above implementation is detailed in Algorithm 1.

1325 4.5.1 HUTCHINSON’S TRACE ESTIMATOR FOR THE EVALUATION OF $\nabla_t \hat{b}_t(x)$

1326 It is well-known that, if $\nabla \nabla b_t(x)$ is bounded,

$$1327 \nabla \cdot \hat{b}_t(x) = \frac{1}{2\delta} \mathbb{E}[\eta \cdot (\hat{b}_t(x + \delta\eta) - \hat{b}_t(x - \delta\eta))] + O(\delta^2), \quad (121)$$

1328 where $0 < \delta \ll 1$ is an adjustable parameter and $\eta \sim N(0, \text{Id})$. Indeed we have

$$1329 \frac{1}{2\delta} \eta \cdot (\hat{b}_t(x + \delta\eta) - \hat{b}_t(x - \delta\eta)) = \eta^T \nabla b_t(x) \eta + O(\delta^2), \quad (122)$$

1330 which implies (121) after taking the expectation over η .

1331 We can use this formula to estimate the PINN loss via

$$1332 L_{\text{PINN}}^{T, \delta}[\hat{b}, \hat{F}] = \int_0^T \mathbb{E}[R_t^\delta(x_t, \eta) R_t^\delta(x_t, \eta')] dt \quad (123)$$

1333 where the expectation is now taken independent over $x_t \sim \hat{\rho}_t$, $\eta \sim N(0, \text{Id})$, and $\eta' \sim N(0, \text{Id})$, and
 1334 we defined

$$1335 R_t^\delta(x, \eta) = \frac{1}{2\delta} \eta \cdot (\hat{b}_t(x + \delta\eta) - \hat{b}_t(x - \delta\eta)) - \nabla U_t(x) \cdot \hat{b}_t(x) - \partial_t U_t(x) + \partial_t \hat{F}_t \quad (124)$$

1336 The expectation in (123) is unbiased since $\eta \perp \eta'$, and its accuracy can be controlled by lowering δ .

1346 4.6 DETAILS ON NUMERICAL EXPERIMENTS

1347 In the following we include details for reproducing the experiments presented in Section 3. An
 1348 overview of the training procedure is given in Algorithm 1. Note that the SDE for the weights can
 1349 be replaced with (32) when learning with $\hat{\phi}_t$, as one would do with the action matching loss (31).

Algorithm 1 Training: Note that for both objectives the resultant set of walkers across time slices $\{x_k^i\}$ are detached from the computational graph when taking a gradient step (*off-policy learning*).

```

1: Initialize:  $n$  walkers,  $x_0 \sim \rho_0$ ,  $A_0 = 0$ ,  $K$  time steps, model parameters for  $\{\hat{b}_t, \hat{F}_t\}$  or  $\hat{\phi}_t$ 
   respectively, diffusion coefficient  $\varepsilon_t$ , learning rate  $\eta$ 
2: repeat
3:   Randomize time grid:  $t_0, t_1, \dots, t_K \sim \text{Uniform}(0, T)$ , sort such that  $t_0 < t_1 < \dots < t_K$ 
4:   for  $k = 0, \dots, K$  do
5:      $\Delta t_k = t_{k+1} - t_k$ ,
6:     for each walker  $i = 1, \dots, n$  do
7:        $x_{t_{k+1}}^i = x_{t_k}^i - \varepsilon_{t_k} \nabla U_{t_k}(x_{t_k}^i) \Delta t_k + \hat{b}_{t_k}(x_{t_k}^i) \Delta t_k + \sqrt{2\varepsilon_{t_k}}(W_{t_{k+1}}^i - W_{t_k}^i)$ 
8:        $A_{t_{k+1}}^i = A_{t_k}^i - \partial_t U_{t_k}(x_{t_k}^i) \Delta t_k - \hat{b}_{t_k}(x_{t_k}^i) \cdot \nabla U_{t_k}(x_{t_k}^i) \Delta t_k + \nabla \cdot \hat{b}_{t_k}(x_{t_k}^i) \Delta t_k$ 
9:     end for
10:  end for
11:  Estimate (118) or (120), respectively, by replacing the expectation by an empirical average
   over the  $n$  walkers and the time integral by an empirical average over  $t_0, \dots, t_K$ .
12:  Take gradient descent step to update the model parameters.
13: until converged

```

4.6.1 PERFORMANCE METRICS

Effective sample size. We can compute the self-normalized ESS as

$$\text{ESS}_t = \frac{\left(N^{-1} \sum_{i=1}^N \exp(A_t^i)\right)^2}{N^{-1} \sum_{i=1}^N \exp(2A_t^i)} \quad (125)$$

at time t along the SDE trajectory. We can use the ESS both as a quality metric and as a trigger for when to perform resampling of the walkers based on the weights, using, e.g. systematic resampling (Doucet et al., 2001; Bolić et al., 2004). Systematic resampling is one of many resampling techniques from particle filtering wherein some walkers are killed and some are duplicated based on their importance weights.

2-Wasserstein distance. The 2-Wasserstein distance reported in Table 1 were computed with 2000 samples from the model and the target density using the Python Optimal Transport library.

Maximum Mean Discrepancy (MMD). We use the MMD code from Blessing et al. (2024) to benchmark the performance of NETS on Neal’s funnel. We use the definition of the MMD as

$$\text{MMD}^2(\hat{\rho}, \rho) \approx \frac{1}{n(n-1)} \sum_{i,j}^n k(\hat{x}_i, \hat{x}_j) + \frac{1}{m(m-1)} \sum_{i,j}^m k(x_i, x_j) - \frac{2}{nm} \sum_i^n \sum_j^m k(\hat{x}_i, x_j) \quad (126)$$

where $\hat{x} \sim \hat{\rho}$ is from the model distribution and $x \sim \rho$ is from the target and $k : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is chosen to be the radial basis kernel with unit bandwidth.

4.6.2 40-MODE GMM

The 40-mode GMM is defined with the mean vectors given as:

1404
 1405
 1406
 1407
 1408
 1409
 1410
 1411
 1412
 1413
 1414
 1415
 1416
 1417
 1418
 1419
 1420
 1421
 1422
 1423
 1424
 1425
 1426
 1427
 1428
 1429
 1430
 1431
 1432
 1433
 1434
 1435
 1436
 1437
 1438
 1439
 1440
 1441
 1442
 1443
 1444
 1445
 1446
 1447
 1448
 1449
 1450
 1451
 1452
 1453
 1454
 1455
 1456
 1457

$$\begin{aligned}
 \mu_1 &= (-0.2995, 21.4577), & \mu_2 &= (-32.9218, -29.4376), \\
 \mu_3 &= (-15.4062, 10.7263), & \mu_4 &= (-0.7925, 31.7156), \\
 \mu_5 &= (-3.5498, 10.5845), & \mu_6 &= (-12.0885, -7.8626), \\
 \mu_7 &= (-38.2139, -26.4913), & \mu_8 &= (-16.4889, 1.4817), \\
 \mu_9 &= (15.8134, 24.0009), & \mu_{10} &= (-27.1176, -17.4185), \\
 \mu_{11} &= (14.5287, 33.2155), & \mu_{12} &= (-8.2320, 29.9325), \\
 \mu_{13} &= (-6.4473, 4.2326), & \mu_{14} &= (36.2190, -37.1068), \\
 \mu_{15} &= (-25.1815, -10.1266), & \mu_{16} &= (-15.5920, 34.5600), \\
 \mu_{17} &= (-25.9272, -18.4133), & \mu_{18} &= (-27.9456, -37.4624), \\
 \mu_{19} &= (-23.3496, 34.3839), & \mu_{20} &= (17.8487, 19.3869), \\
 \mu_{21} &= (2.1037, -20.5073), & \mu_{22} &= (6.7674, -37.3478), \\
 \mu_{23} &= (-28.9026, -20.6212), & \mu_{24} &= (25.2375, 23.4529), \\
 \mu_{25} &= (-17.7398, -1.4433), & \mu_{26} &= (25.5824, 39.7653), \\
 \mu_{27} &= (15.8753, 5.4037), & \mu_{28} &= (26.8195, -23.5521), \\
 \mu_{29} &= (7.4538, -31.0122), & \mu_{30} &= (-27.7234, -20.6633), \\
 \mu_{31} &= (18.0989, 16.0864), & \mu_{32} &= (-23.6941, 12.0843), \\
 \mu_{33} &= (21.9589, -5.0487), & \mu_{34} &= (1.5273, 9.2682), \\
 \mu_{35} &= (24.8151, 38.4078), & \mu_{36} &= (-30.8249, -14.6588), \\
 \mu_{37} &= (15.7204, 33.1420), & \mu_{38} &= (34.8083, 35.2943), \\
 \mu_{39} &= (7.9606, -34.7833), & \mu_{40} &= (3.6797, -25.0242)
 \end{aligned}$$

These means follow the definition given in the FAB (Midgley et al., 2023) code base that has been subsequently used in recent papers. The time-dependent potential $U_t(x)$ is given by the interpolation of means.

4.6.3 NEAL’S 10- d FUNNEL

The Neal’s Funnel distribution is a 10- d probability distribution defined as

$$x_0 \sim \mathcal{N}(0, \sigma^2), \quad x_{1:9} \sim \mathcal{N}(0, e^{x_0}) \quad (127)$$

where $\sigma = 3$ and we use subscripts here as a dimensional index and not as a time index like in the rest of the paper. Following this, we use as a definition of the interpolating potential:

$$U_t(x) = \frac{1}{2}x_0^2(1-t + \frac{t}{\sigma^2}) + \frac{1}{2} \sum_{i=1}^{d-1} e^{-tx_0} x_i^2 + (d-1)tx_0 \quad (128)$$

so that at $t = 0$, we have $U_0(x) = \frac{1}{2}x_0^2 + \frac{1}{2} \sum_{i=1}^{d-1} x_i^2$ and at time $t = 1$ we have the funnel potential given as $U_1(x) = \frac{1}{2\sigma^2}x_0 + \frac{1}{2} \sum_{i=1}^{d-1} e^{-x_0} x_i^2 + (d-1)x_0$.

4.6.4 50- d MIXTURE OF STUDENT-T DISTRIBUTIONS

Following Blessing et al. (2024), we use their mixture of 10 student-T distributions in 50 dimensions. We construct U_t via interpolation of means from a single standard student-T distribution (mean 0). We use the same neural network as used in the GMM experiments.

To further drive home the fact that our annealed Langevin dynamics with transport can be taken post-training to the $\epsilon \rightarrow \infty$ limit to approach perfect sampling, we provide the following ablation from our model learned with the action matching loss given in Figure 4.

4.7 DETAILS OF THE φ^4 MODEL

We consider the Euclidean scalar ϕ^4 theory given by the action

$$S_{\text{Euc}}[\varphi] = \int [\partial_\mu \varphi(x) \partial^\mu \varphi(x) + m^2 \varphi^2(x) + \lambda \varphi^4(x)] d^D x \quad (129)$$

1458
 1459
 1460
 1461
 1462
 1463
 1464
 1465
 1466
 1467
 1468
 1469
 1470
 1471
 1472
 1473
 1474
 1475
 1476
 1477
 1478
 1479
 1480
 1481
 1482
 1483
 1484
 1485
 1486
 1487
 1488
 1489
 1490
 1491
 1492
 1493
 1494
 1495
 1496
 1497
 1498
 1499
 1500
 1501
 1502
 1503
 1504
 1505
 1506
 1507
 1508
 1509
 1510
 1511




Figure 4: Reduction in W_2 distance from taking the $\epsilon \rightarrow \infty$ limit in sampling with NETS. Note that the resolution of the SDE integration must increase to accommodate the higher stochasticity. Average taken over 3 sampling runs of 2000 walkers each.

where we use Einstein summation to denote the dot product with respect to the Euclidean metric and D is the spacetime dimension. We are interested in acquiring a variant of this expression that provides a fast computational realization when put onto the lattice. Using Green’s identity (integrating by parts) we note that

$$\int (\partial_\mu \varphi(x) \partial^\mu \varphi(x) d^d x = \int \partial_\mu \varphi \cdot \partial_\mu \varphi d^d x = - \int \varphi(x) \partial_\mu \partial^\mu \varphi(x) d^d x + \text{vanishing surface term} \quad (130)$$

so that

$$S_{\text{Euc}}[\varphi] = \int -\varphi(x) \partial_\mu \partial^\mu \varphi(x) + m^2 \varphi^2(x) + \lambda \varphi^4(x) d^d x. \quad (131)$$

Discretizing S_{Euc} onto the lattice

$$\Lambda = \{a(n_0, \dots, n_{d-1}) \mid n_i \in \{0, 1, 2, \dots, L\}, i = 0, 1, \dots, d, a \in \mathbb{R}_+\},$$

where a is the lattice spacing used to define the physical point $x = an$, we use the forward difference operator to define

$$\partial_\mu \varphi(x) \rightarrow \frac{1}{a} [\varphi(x + \mu) - \varphi(x)] \quad \partial_\mu \partial^\mu \varphi(x) \rightarrow \frac{1}{a^2} [\varphi(x + \mu) - 2\varphi(x) + \varphi(x - \mu)]. \quad (132)$$

Using these expressions, we write the discretized lattice action as

$$S_{\text{Lat}} = \sum_{x \in \Lambda} a^D \left[\sum_{\mu=1}^D -\frac{1}{a^2} [\varphi_{x+\mu} \varphi_x - 2\varphi_x^2 + \varphi_{x-\mu} \varphi_x] + m^2 \varphi_x^2 + \lambda \varphi_x^4 \right] \quad (133)$$

$$= \sum_x a^D \left[2Da^{-2} \varphi_x^2 - a^{-2} \sum_\mu [\varphi_{x+\mu} \varphi_x + \varphi_{x-\mu} \varphi_x] + m^2 \varphi_x^2 + \lambda \varphi_x^4 \right] \quad (134)$$

$$= \sum_x a^d \left[2Da^{-2} \varphi_x^2 - 2a^{-2} \sum_\mu [\varphi_x \varphi_{x+\mu}] + m^2 \varphi_x^2 + \lambda \varphi_x^4 \right] \quad (135)$$

$$= \sum_x a^D \left[-2a^{-2} \sum_\mu \varphi_x \varphi_{x+\mu} + (2a^{-2} D + m^2) \varphi_x^2 + \lambda \varphi_x^4 \right] \quad (136)$$

where we have used the fact that on the lattice $\sum_x \varphi_x \varphi_{x+\hat{\mu}} = \sum_x \varphi_{x-\hat{\mu}} \varphi_x$ to get the third equality. It is useful to put the action in a form that is independent of the lattice spacing a . To do so, we introduce the re-scaled lattice field as

$$\varphi_x \rightarrow a^{D/2-1} \varphi_x, \quad m^2 \rightarrow a^2 m^2, \quad \text{and} \quad \lambda \rightarrow a^{4-D} \lambda. \quad (137)$$

1512 Plugging these rescalings into (136) gives us the final expression

$$1513 \quad S_{\text{Lat}} = \sum_x \left[-2 \sum_{\mu} \varphi_x \varphi_{x+\mu} \right] + (2D + m^2) \varphi_x^2 + \lambda \varphi_x^4, \quad (138)$$

1514 which we are to use in simulation.

1515 4.7.1 FREE THEORY $\lambda = 0$

1516 Turning off the interaction makes it possible to analytically solve the theory. To do this, introduce the discrete Fourier transform relations

$$1517 \quad \varphi_k = \frac{1}{\sqrt{L^D}} \sum_x \varphi_x e^{-ik \cdot x} \quad (139)$$

$$1518 \quad \varphi_x = \frac{1}{\sqrt{L^D}} \sum_k \varphi_k e^{ik \cdot x} \quad (140)$$

1519 for discrete wavenumbers $k = \frac{2l\pi}{L}$ with $l = 0, \dots, L-1$. Plugging in (140) into the first part of (136), we get the expanded sum

$$1520 \quad \sum_x \left[-2 \sum_{\mu} \hat{\varphi}_x \hat{\varphi}_{x+\mu} \right] \rightarrow -\frac{2}{L^d} \sum_x \sum_{\mu} \sum_k \sum_{k'} \varphi_k \varphi_{k'} e^{i(k+k') \cdot x} e^{ik' \cdot \mu} \quad (141)$$

$$1521 \quad = -2 \sum_{\mu} \sum_k \sum_{k'} \delta_{k,-k'} \varphi_k \varphi_{k'} e^{ik' \cdot \mu} \quad (142)$$

$$1522 \quad = -2 \sum_{\mu} \sum_k \varphi_k \varphi_{-k} e^{-ik \cdot \mu} \quad (143)$$

$$1523 \quad = -2 \sum_{\mu} \sum_k \varphi_k \varphi_{k^*} e^{-ik \cdot \mu} \quad (144)$$

$$1524 \quad = -2 \sum_{\mu} \sum_k |\varphi_k|^2 [\cos k_{\mu} + i \sin k_{\mu}] = - \sum_{\mu} \sum_k |\varphi_k|^2 \cos k_{\mu} \quad (145)$$

1525 where ϕ^* indicates conjugation, and we got the first equality by the orthogonality of the Fourier modes, the second by the Kronecker delta, and the third by the reality of the scalar field. Proceeding similarly for the terms proportional to φ^2 gives us the expression

$$1526 \quad S_k = \sum_k \left[m^2 + 2D - 2 \sum_{\mu} \cos k_{\mu} \right] |\varphi|^2 \quad (146)$$

1527 The above equation can be written in quadratic form to highlight that the field may be sampled analytically

$$1528 \quad S_k = \frac{1}{L^d} \sum_k \varphi_k M_{k,-k} \varphi_{-k} \quad (147)$$

$$1529 \quad \text{where } M_{k,-k} = \left[m^2 + 2D - 2 \sum_{\mu} \cos k_{\mu} \right] \delta_{k,-k} \quad (148)$$

1530 Note that this free theory can be sampled for any $m^2 > 0$.

1531 4.7.2 φ^4 NUMERICAL DETAILS

1532 We numerically realize the above lattice theory in D=2 spacetime dimensions. We use an interpolating potential with time dependent $m_t^2 = (1-t)m_0^2 + tm_1^2$, $\lambda_t = (1-t)\lambda_0 + t\lambda_1$ where λ_0 is always chosen to be 0 (though we note that you could run this sampler for any U_0 that you could sample from easily, not just analytically but also with existing MCMC methods). For the $L = 20$ ($d = L \times L = 400$ dimensional) experiments, we identify the critical point of the theory (where the lattices go from ordered to disordered) using HMC by studying the distribution of the magnetization of the

field configurations as $M[\varphi^i(x)] = \sum_x \varphi^i(x)$, where summation is taken over all lattice sites on the i^{th} lattice configuration. We identify this at $m_1^2 = -1.0$, $\lambda_1 = 0.9$ and use these as the target theory parameters on which to perform the sampling. For the $L = 16$ test ($d = 256$), we go past this phase transition into the ordered phase of the theory, which we identify via HMC simulations at $m_1^2 = -1.0$, $\lambda_1 = 0.8$.

4.8 LINK WITH VARGAS ET AL. (2024)

Consider the process $Y_t^{\hat{b}}$ solution to the SDE

$$dY_t^{\hat{b}} = \varepsilon_t \nabla U_t(Y_t^{\hat{b}}) dt + 2\varepsilon_t \nabla \log \rho_t^{\hat{b}}(Y_t^{\hat{b}}) dt + \hat{b}_t(Y_t^{\hat{b}}) dt + \sqrt{2\varepsilon_t} dW_t, \quad Y_0^{\hat{b}} \sim \rho_0. \quad (149)$$

where $\rho_t^{\hat{b}}$ denotes the PDF of the process $X_t^{\hat{b}}$ defined by the SDE (20), i.e. the solution to the FPE

$$\partial_t \rho_t^{\hat{b}} = \varepsilon_t \nabla \cdot (\nabla U_t \rho_t^{\hat{b}} + \nabla \rho_t^{\hat{b}}) - \nabla \cdot (\hat{b}_t \rho_t^{\hat{b}}), \quad \rho_0^{\hat{b}} = \rho_0 \quad (150)$$

The process $Y_t^{\hat{b}}$ has a simple interpretation: it is the time-reversed of the process run using the time-reversed potential U_{1-t} and $-\hat{b}_{1-t}$: that is, if the additional drift \hat{b}_t was the perfect one solution to (24), the law of $X^{\hat{b}} = (X_t^{\hat{b}})_{t \in [0,1]}$ and $Y^{\hat{b}} = (Y_t^{\hat{b}})_{t \in [0,1]}$ should coincide. This suggests to learn b using as objective a divergence of the path measure of $X^{\hat{b}}$ from that of $Y^{\hat{b}}$. This is essentially what is suggested in Vargas et al. (2024), and for the reader convenience let us re-derive some of their results in our notations.

The Kullback-Leibler divergence (or relative entropy) of the path measure of $X^{\hat{b}}$ from that of $Y^{\hat{b}}$ reads

$$\text{KL}(X^{\hat{b}} \| Y^{\hat{b}}) = \frac{1}{4} \varepsilon_t \int_0^1 \mathbb{E}[|\nabla U_t(X_t^{\hat{b}}) + \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})|^2] dt \quad (151)$$

This objective is akin to the one used in score-based diffusion modeling (SBDM) and simply says that one way to adjust \hat{b} is by matching the score of $\rho_t^{\hat{b}}$ to that of ρ_t . As written (151) is not explicit since we do not know $\nabla \log \rho_t^{\hat{b}}$. We can however make it explicit after a few manipulations similar to those used in SBDM. To this end, notice first that, by Ito formula, we have

$$\begin{aligned} d \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) &= \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot (-\varepsilon_t \nabla U_t(X_t^{\hat{b}}) + \hat{b}_t(X_t^{\hat{b}})) dt + \varepsilon_t \Delta \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) dt \\ &\quad + \sqrt{2\varepsilon_t} \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot dW_t \end{aligned} \quad (152)$$

which implies that

$$\begin{aligned} \varepsilon_t \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot \nabla U_t(X_t^{\hat{b}}) dt &= -d \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) + \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}}) dt \\ &\quad + \varepsilon_t \Delta \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) dt + \sqrt{2\varepsilon_t} \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot dW_t \end{aligned} \quad (153)$$

Inserting this expression in (151) after expanding the square, and noticing that the martingale term involving dW_t disappears by Ito isometry and that the term $d \log \rho_t^{\hat{b}}(X_t^{\hat{b}})$ can be integrated in time we arrive at

$$\begin{aligned} \text{KL}(X^{\hat{b}} \| Y^{\hat{b}}) &= \frac{1}{4} \varepsilon_t \int_0^1 \mathbb{E}[|\nabla U_t(X_t^{\hat{b}})|^2 + |\nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})|^2 + 2\nabla U_t(X_t^{\hat{b}}) \cdot \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})] dt \\ &= \frac{1}{4} \int_0^1 \mathbb{E}[\varepsilon_t |\nabla U_t(X_t^{\hat{b}})|^2 + \varepsilon_t |\nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})|^2 + \varepsilon_t \nabla U_t(X_t^{\hat{b}}) \cdot \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})] dt \\ &\quad + \frac{1}{4} \int_0^1 \mathbb{E}[\nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}}) + \varepsilon_t \Delta \log \rho_t^{\hat{b}}(X_t^{\hat{b}})] dt \\ &\quad + \frac{1}{4} \mathbb{E}[\log \rho_0(X_0)] - \frac{1}{4} \mathbb{E}[\log \rho_1^{\hat{b}}(X_1)] \end{aligned} \quad (154)$$

where we used $\rho_0^{\hat{b}} = \rho_0$. We can now use the following identities, each obtained using $\rho_t^{\hat{b}} \nabla \log \rho_t^{\hat{b}} = \nabla \rho_t^{\hat{b}}$ and one integration by parts:

$$\begin{aligned} \mathbb{E}[|\nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})|^2] &= \int_{\mathbb{R}^d} |\nabla \log \rho_t^{\hat{b}}(x)|^2 \rho_t^{\hat{b}}(x) dx \\ &= \int_{\mathbb{R}^d} \nabla \log \rho_t^{\hat{b}}(x) \cdot \nabla \rho_t^{\hat{b}}(x) dx \\ &= - \int_{\mathbb{R}^d} \Delta \log \rho_t^{\hat{b}}(x) \rho_t^{\hat{b}}(x) dx \\ &= -\mathbb{E}[\Delta \log \rho_t^{\hat{b}}(X_t^{\hat{b}})], \end{aligned} \quad (155)$$

$$\begin{aligned} \mathbb{E}[\nabla U_t(X_t^{\hat{b}}) \cdot \nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}})] &= \int_{\mathbb{R}^d} \nabla U_t(x) \cdot \nabla \log \rho_t^{\hat{b}}(x) \rho_t^{\hat{b}}(x) dx \\ &= \int_{\mathbb{R}^d} \nabla U_t(x) \cdot \nabla \rho_t^{\hat{b}}(x) dx \\ &= - \int_{\mathbb{R}^d} \Delta U_t(x) \rho_t^{\hat{b}}(x) dx \\ &= -\mathbb{E}[\Delta U_t(X_t^{\hat{b}})] \end{aligned} \quad (156)$$

and

$$\begin{aligned} \mathbb{E}[\nabla \log \rho_t^{\hat{b}}(X_t^{\hat{b}}) \cdot \hat{b}_t(X_t^{\hat{b}})] &= \int_{\mathbb{R}^d} \nabla \log \rho_t^{\hat{b}}(x) \cdot \hat{b}_t(x) \rho_t^{\hat{b}}(x) dx \\ &= - \int_{\mathbb{R}^d} \nabla \cdot \hat{b}_t(x) \rho_t^{\hat{b}}(x) dx \\ &= -\mathbb{E}[\nabla \cdot \hat{b}_t(X_t^{\hat{b}})] \end{aligned} \quad (157)$$

Inserting these expressions in (154), it reduces to

$$\begin{aligned} \text{KL}(X^{\hat{b}} \| Y^{\hat{b}}) &= \frac{1}{4} \int_0^1 \mathbb{E}[\varepsilon_t |\nabla U_t(X_t^{\hat{b}})|^2 - \varepsilon_t \Delta U_t(X_t^{\hat{b}}) - \nabla \cdot b_t(X_t^{\hat{b}})] dt \\ &\quad + \frac{1}{4} \mathbb{E}[\log \rho_0(X_0)] - \frac{1}{4} \mathbb{E}[\log \rho_1^{\hat{b}}(X_1)] \end{aligned} \quad (158)$$

This objective is still not practical because it involves $\log \rho_1^{\hat{b}}$, which is unknown. There is however a simple way to fix this, by adding a term in the Kullback-Leibler divergence (151)

$$\text{KL}'(X^{\hat{b}} \| Y^{\hat{b}}) = \text{KL}(X^{\hat{b}} \| Y^{\hat{b}}) + \frac{1}{4} \mathbb{E}[\log(\rho_1^{\hat{b}}(X_1^{\hat{b}})/\rho_1(X_1^{\hat{b}}))] \quad (159)$$

This additional term is proportional to the Kullback-Leibler divergence of $\rho_1^{\hat{b}}$ from the target PDF ρ_1 . Using (158) as well as $\rho_1(x) = e^{-U_1(x)+F_1}$, we can now express (159) as

$$\begin{aligned} \text{KL}'(X^{\hat{b}} \| Y^{\hat{b}}) &= \frac{1}{4} \int_0^1 \mathbb{E}[\varepsilon_t |\nabla U_t(X_t^{\hat{b}})|^2 - \varepsilon_t \Delta U_t(X_t^{\hat{b}}) - \nabla \cdot b_t(X_t^{\hat{b}})] dt \\ &\quad + \frac{1}{4} \mathbb{E}[\log \rho_0(X_0)] + \frac{1}{4} \mathbb{E}[U_1(X_1)] - \frac{1}{4} F_1. \end{aligned} \quad (160)$$

This is Equation (24) in Vargas et al. (2024) in which we set $\nabla \hat{\phi}_t(x) = \hat{b}_t(x)$ and we used that, for any $c_t : \mathbb{R}^d \rightarrow \mathbb{R}^d$, we have

$$\mathbb{E} \int_0^1 c_t(X_t^{\hat{b}}) \cdot \overleftarrow{d} W_t = \sqrt{2\varepsilon_t} \int_0^1 \mathbb{E}[\nabla \cdot c_t(X_t^{\hat{b}})] dt. \quad (161)$$

Note that we can neglect the term $\frac{1}{4} \mathbb{E}[\log \rho_0(X_0)]$ in (158) since it does not depend on \hat{b} , so that the minimization of (158) can be cast into the minimization of (after multiplication by 4)

$$\begin{aligned} &\int_0^1 \mathbb{E}[\varepsilon_t |\nabla U_t(X_t^{\hat{b}})|^2 - \varepsilon_t \Delta U_t(X_t^{\hat{b}}) - \nabla \cdot b_t(X_t^{\hat{b}})] + \mathbb{E}[U_1(X_1) - F_1] \\ &= \int_0^1 \int_{\mathbb{R}^d} [\varepsilon_t |\nabla U_t(x)|^2 - \varepsilon_t \Delta U_t(x) - \nabla \cdot b_t(x)] \rho_t^{\hat{b}}(x) dx + \int_{\mathbb{R}^d} (U_1(x) - F_1) \rho_1^{\hat{b}}(x) dx \end{aligned} \quad (162)$$

1674 where $\rho_t^{\hat{b}}$ solves (150).
1675

1676 Let us check that the minimizer of (162) is $\hat{b}_t = b_t$, the solution to (24), so that we also have
1677 $\rho_t^{\hat{b}} = \rho_t^b = \rho_t$. To this end, notice that the minimization of (162) can be performed with the method
1678 Lagrange multiplier, using the extended objective

$$1679 \int_0^1 \int_{\mathbb{R}^d} [\varepsilon_t |\nabla U_t(x)|^2 - \varepsilon_t \Delta U_t(x) - \nabla \cdot \hat{b}_t(x)] \rho_t^{\hat{b}}(x) dx dt + \int_{\mathbb{R}^d} (U_1(x) - F_1) \rho_1^{\hat{b}}(x) dx$$

$$1680 + \int_0^1 \int_{\mathbb{R}^d} \lambda_t(x) (\partial_t \rho_t^{\hat{b}} - \varepsilon_t \nabla \cdot (\nabla U_t \rho_t^{\hat{b}} + \nabla \rho_t^{\hat{b}}) + \nabla \cdot (\hat{b}_t \rho_t^{\hat{b}})) dx dt$$
(163)

1684 where $\lambda_t(x)$ is the Lagrange multiplier to be determined. Taking the first variation of this objective
1685 over λ , $\rho^{\hat{b}}$, and \hat{b} , we arrive at the Euler-Lagrange equations
1686

$$1687 0 = \partial_t \rho_t^{\hat{b}} - \varepsilon_t \nabla \cdot (\nabla U_t \rho_t^{\hat{b}} + \nabla \rho_t^{\hat{b}}) + \nabla \cdot (\hat{b}_t \rho_t^{\hat{b}}), \quad \rho_0^{\hat{b}} = \rho_0,$$

$$1688 0 = \varepsilon_t |\nabla U_t|^2 - \varepsilon_t \Delta U_t - \nabla \cdot \hat{b}_t$$

$$1689 - \partial_t \lambda_t + \varepsilon_t \nabla U_t \cdot \nabla \lambda_t - \varepsilon_t \Delta \lambda_t - \hat{b}_t \cdot \nabla \lambda_t \quad \lambda_1 = -U_1 + F_1$$
(164)

$$1690 0 = \nabla \rho_t^{\hat{b}} - \rho_t^{\hat{b}} \nabla \lambda_t$$

1693 We can check that $\hat{b}_t(x) = b_t(x)$, $\rho_t^{\hat{b}}(x) = \rho_t(x) = e^{-U_t(x)+F_t}$, and $\lambda_t(x) = -U_t(x) + F_t$ is a
1694 solution: indeed this solves the first and the last equations in (164) and reduces the second to
1695

$$1696 0 = [\varepsilon_t |\nabla U_t|^2 - \varepsilon_t \Delta U_t - \nabla \cdot b_t]$$

$$1697 + \partial_t U_t - \partial_t F_t - \varepsilon_t |\nabla U_t|^2 + \Delta U_t + \nabla b_t \cdot \nabla U_t$$
(165)

$$1698 = -\nabla \cdot b_t + \partial_t U_t - \partial_t F_t + \nabla b_t \cdot \nabla U_t$$

1700 which is satisfied since b_t solves (24).
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727