

TRANSFORMERS PROVABLY LEARN TWO-MIXTURE OF LINEAR CLASSIFICATION VIA GRADIENT FLOW

Anonymous authors

Paper under double-blind review

ABSTRACT

Understanding how transformers learn and utilize hidden connections between words is crucial to understand the behavior of large language models. To study this mechanism, we consider the task of two-mixture of linear classification which features a hidden correspondence structure between words, and study the training dynamics of a symmetric two-headed transformer with ReLU neurons. Motivated by the stage-wise learning phenomenon observed in our experiments, we design and theoretically analyze a three-stage training algorithm, which can effectively characterize the actual gradient descent dynamics when we simultaneously train the neuron weights and the softmax attention. The first stage is a neuron learning stage, where the neurons align with the underlying signals. The second stage is an attention feature learning stage, where we analyze the feature learning process of how the attention learns to utilize the relationship between the tokens to solve certain hard samples. In the meantime, the attention features evolve from a nearly non-separable state (at the initialization) to a well-separated state. The third stage is a convergence stage, where the population loss is driven towards zero. The key technique in our analysis of softmax attention is to identify a critical sub-system inside a large dynamical system and bound the growth of the non-linear sub-system by a linear system. Finally, we discuss the setting with more than two mixtures. We empirically show the difficulty of generalizing our analysis of the gradient flow dynamics to the case even when the number of mixtures equals three, although the transformer can still successfully learn such distribution. On the other hand, we show by construction that there exists a transformer that can solve mixture of linear classification given any arbitrary number of mixtures.

1 INTRODUCTION

Since the invention of self-attention (Vaswani et al., 2017), transformers have become the dominating backbone architecture in many machine learning applications such as computer vision (Dosovitskiy et al., 2020; Liu et al., 2021), natural language processing (Devlin et al., 2018) and protein structure prediction (Jumper et al., 2021). Within the past two years, ChatGPT and GPT-4 (OpenAI, 2023; Bubeck et al., 2023) along with other large language models (LLMs) (Touvron et al., 2023; Manyika & Hsiao, 2023; Gemini, 2023; Anthropic, 2024) have demonstrated astonishing abilities in language understanding, math Olympiad (AlphaProof & AlphaGeometry, 2024) and coding, etc.

Despite the wide range of success of transformers, how those models can achieve such impressive performance still remains largely unknown. One mystery lies in how transformers are trained to utilize the connections between words to solve various tasks. For example, consider the prompt “Bob is watching television in the living room. Where is Bob?”. To answer the question, the model must utilize the (hidden) correspondence between “Where” and “living room”.

There are many previous works studying the mechanism of self-attention under different settings through training dynamics such as (Li et al., 2024b; Huang et al., 2024; Li et al., 2022; Tarzanagh et al., 2023b;a; Vasudeva et al., 2024). However, many of those previous works were either focusing on attentions that are position-based or studying a setting where the words in the dictionary don’t have explicit relationships between each other. The goal of this work is to study some simple yet informative setting such that we can get a clear understanding how the softmax attention learns and utilizes certain connections between words to solve tasks through training. To achieve this goal, we

054 consider the task of mixtures of linear classification. In this task, we have a dictionary consisting
 055 of group identifiers and classification features, and there is a hidden **correspondence structure**:
 056 each group identifier is associated with one classification feature. Each input contains a group
 057 identifier and a corresponding signed classification feature. The rest of tokens are sampled from
 058 signed classification features from other groups. The label of the sequence is determined by the sign
 059 of the classification feature corresponding to the group identifier. Thus, a model needs to capture the
 060 correspondence structure between group identifiers and classification features to solve this task. [To
 061 see how attention can solve this task, later in our training dynamics analysis, we are going to show
 062 that the softmax attention builds such correspondence structure by allocating more weights on the
 063 group identifier when it sees classification feature queries.](#)

064 Although the intuition suggests that we can construct a transformer to solve this task, it is still
 065 unknown that how the transformer models can be *trained* to learn such a hidden correspondence
 066 structure. This inspires the following intriguing question:

067 *How do transformers learn and utilize the hidden correspondence structure to
 068 solve mixture of linear classification via gradient descent?*
 069

070 **Our contributions.** In this work, we study the training dynamics of a two-headed transformer given
 071 two mixtures. Our contributions are summarized as follows:

- 072 1. As a guidance for our theory, we first conduct experiments to observe the training dynamics
 073 of the transformer where we train all the weights simultaneously. Our experiment results
 074 show a clear stage-wise learning phenomenon where the neuron weights learn before the
 075 attention modules.
- 076 2. Motivated by our experimental observations, we design a three-stage layer-wise training al-
 077 gorithm and further analyze its gradient flow dynamics. We characterize the feature learn-
 078 ing process where the attention features evolve from a nearly non-separable state to a well-
 079 separated state. In particular, our analysis captures how the self-attention associates the
 080 group signals and classification signals to solve this task. Our analysis closely reflects the
 081 actual behavior of the gradient descent dynamics when we train all weights simultaneously.
- 082 3. In order to analyze the change in softmax attention, we formulate a set of relevant variables
 083 whose evolution can be characterized by a large non-linear dynamical system of ordinary
 084 differential equations. To make the analysis of the dynamical system tractable, we reduce
 085 the dimension of the system by identifying some key variables and forming a smaller sub-
 086 system. We are able to relate the evolution of the smaller system with a linear dynamical
 087 system to track its dynamics. Our proof techniques can be of independent interest.
- 088 4. We explore the training dynamics of the transformers when the number of mixtures goes
 089 beyond two. We empirically show the difficulty of analyzing training dynamics even when
 090 the number of mixture equals three, although the two-headed transformer model can still
 091 learn this distribution. On the other hand, we give a general two-headed transformer con-
 092 struction that can solve mixture of linear classification given arbitrary number of mixtures.

093 1.1 RELATED WORK

095 **Mixture of linear regression/classification.** Mixture of linear regression/classification is a classi-
 096 cal model in statistics and machine learning (De Veaux, 1989; Jordan & Jacobs, 1994) which was
 097 applied in areas such as object recognition (Quattoni et al., 2004) and machine translation (Liang
 098 et al., 2006). This problem can be solved by tensor-based methods (Anandkumar et al., 2014; 2012;
 099 Hsu et al., 2012; Chaganty & Liang, 2013) and the expectation-maximization algorithm (Khalili &
 100 Chen, 2007; Yi et al., 2014; Balakrishnan et al., 2017; Wang et al., 2015). Previously, mixture of
 101 linear classification has been used in studying mixtures of experts (Chen et al., 2022).

102 **Training dynamics of transformers.** The training dynamics of transformers have been studied
 103 under various settings. One particular category is in-context learning. For example, in-context linear
 104 regression has been studied under linear attention (Zhang et al., 2024), softmax attention (Huang
 105 et al., 2023), non-linear embedding (Yang et al.), and multi-task with multiple heads (Chen et al.,
 106 2024). Other in-context learning settings include classification (Li et al., 2024b), causal structure
 107 (Nichani et al., 2024), Markov chain or n-gram (Edelman et al., 2024; Makkuva et al., 2024; Chen
 et al.), nearest neighbor (Li et al., 2024c) and chain-of-thought (Li et al., 2024a).

For non-in-context settings, it has been shown that trained transformer can learn spatial structures (Jelassi et al., 2022), topic models (Li et al., 2023b), and feature-position correlation (Huang et al., 2024). In addition, (Tian et al., 2023) showed that self-attention behaves like a discriminative scan algorithm. There is one line of works studying a setting where the attention weight can converge to a SVM solution (Tarzanagh et al., 2023b;a; Vasudeva et al., 2024). In addition, (Li et al., 2022) studied a classification task with label-relevant and label-irrelevant tokens. (Li et al., 2023a) provided analysis of training graph transformers for node classification tasks. (Wang et al.) showed transformers can learn a sparse token selection task which lead to an optimization-based separation between transformers and MLPs. There are also works trying to prove convergence of transformer training via NTK (Wu et al., 2023; Deora et al.) and mean-field (Kim & Suzuki, 2024; Gao et al.).

Our work also study a non-in-context learning setting. However, many of the previous works were either considering attentions that are position-based or studying a setting where the words don't have explicit relationships with each other. Our study complements the above work by considering a setting where the dictionary possesses a correspondence structure.

2 SETTINGS

Notations. For a vector $v \in \mathbb{R}^d$, we use $\text{diag}(v)$ to denote a diagonal matrix with v being the diagonal entries. When we subtract the vector v by a scalar a , we subtract each entry of v by a , i.e., $v - a \in \mathbb{R}^d$ and $(v - a)_i = v_i - a$. For a set \mathcal{S} with elements in \mathbb{R} and $|\mathcal{S}| = n$, we can vectorize the set \mathcal{S} in any arbitrary but fixed order and denote the vector as $\text{vec}(\mathcal{S}) \in \mathbb{R}^n$. We use $\tilde{\Omega}, \tilde{\Theta}, \tilde{O}$ to hide polylogarithmic factors.

2.1 DATA DISTRIBUTION

Definition 2.1 (Mixture of linear classification). *Define the K -mixture of linear classification data distribution \mathcal{D} as follows: Assume $d \geq 2K$. Let $\mathcal{C} = \{c_k\}_{k=1}^K \subset \mathbb{R}^d$ denote the group signals and $\mathcal{V} = \{v_k\}_{k=1}^K \subset \mathbb{R}^d$ denote the classification signals, where $\mathcal{C} \cup \mathcal{V}$ is orthonormal. Let L be the number of tokens per sample. Each data entry $(X, y) \in \mathbb{R}^{d \times L} \times \{\pm 1\}$ is created as follows:*

1. Sample $y \sim \text{Uniform}(\{\pm 1\})$.
2. Sample $k \sim \text{Uniform}([K])$ and $l_0 \sim \text{Uniform}([L])$. Set $X_{l_0} = c_k$.
3. Sample $l_1 \sim \text{Uniform}([L] \setminus \{l_0\})$ and set $X_{l_1} = yv_k$.
4. For each $l_2 \in [L] \setminus \{l_0, l_1\}$, sample $k' \stackrel{i.i.d.}{\sim} \text{Uniform}([K] \setminus \{k\})$ and set $X_{l_2} = \epsilon v_{k'}$, where $\epsilon \stackrel{i.i.d.}{\sim} \text{Uniform}(\{\pm 1\})$.

Based on our data distribution, the model needs to utilize the correspondence between the group signals and classification signals to solve this task. We further elaborate our data model as follows. We take the sequence $X = [c_1, y_2 v_2, y_1 v_1, y_3 v_3]$ with $y_1, y_2, y_3 \in \{\pm 1\}$ as an example. Based on the form of X , the token c_1 indicates that this instance is in group 1. Then the label of this sequence depends on the sign associated with v_1 (that shares the same index with c_1), and hence is given by y_1 .

For the training dynamics part of this work, we consider $K = 2$ and $L = 3$, and we discuss the setting with arbitrary number of mixtures in Section 6. Since the samples are invariant of permutation between tokens, we can ignore the order of the tokens and define $X_{k,s,s'} := (c_k, sv_k, s'v_{k'})$ for $k \neq k' \in [2], s, s' \in \{\pm 1\}$. We refer to the samples $X_{k,s,s}$ as **consistent samples** and samples $X_{k,s,-s}$ as **conflicting samples**.

2.2 TRANSFORMER ARCHITECTURE AND LOSS FUNCTION

Given inputs $X \in \mathbb{R}^{d \times L}$, we first define a single softmax attention head $H : \mathbb{R}^{d \times L} \rightarrow \mathbb{R}$ as

$$H(X) = \sum_{l=1}^L w^\top X \text{softmax}(X^\top W_K^\top W_Q x_l)$$

where $w \in \mathbb{R}^d$, $b \in \mathbb{R}$, $W_K, W_Q \in \mathbb{R}^{m \times d}$, and x_l denotes the l -th token in X . Then, we define our one-layer symmetric two-headed transformer with ReLU neurons $f: \mathbb{R}^{d \times L} \rightarrow \mathbb{R}$ as

$$f(X) = \sigma(H_+(X) + b) - \sigma(H_-(X) + b) \quad (1)$$

where $b \in \mathbb{R}$ and σ is the ReLU activation function. We denote the weights in H_+ as w_+, W_{K+}, W_{Q+} and similarly for H_- . We introduce a shorthand notation $K_+ = W_{K+}X$ and $Q_+ = W_{Q+}X$. For $\mu, \nu \in X$, we define the output of attention, or softmax probability, as $p_{q \leftarrow \mu, k \leftarrow \nu}^{(+)}(X) := \text{softmax}\left(X^\top W_{K_+}^\top W_{Q_+} \mu\right)_{l(\nu)}$, where $l(\nu)$ denotes the index such that $X_{l(\nu)} = \nu$.

Loss Function. We train $w_\pm, W_{K_\pm}, W_{Q_\pm}$ to minimize the population loss $\mathbb{E}_{(X,y)}[\ell(yf(X))]$ where $\ell(x) = \log(1 + \exp(-x))$ is the binary cross-entropy loss. The gradient of the cross-entropy loss is given as $\ell'(x) = -\frac{1}{1+e^x}$ and we denote $g(x) := 1/(1+e^x)$.

3 GUIDING EXPERIMENTS AND ALGORITHM

3.1 EXPERIMENT RESULTS

To understand the dynamics of the training process, we train the two-headed transformer in Equation (1) on the data distribution in Definition 2.1. In our experiments, all the weights in the transformer are trained simultaneously via gradient descent with learning rate 0.1.




Figure 1: Experiment results. The attention scores and neuron alignment are of the positive head.

We make the following observations from our experimental results in Figure 1, which will guide us to further develop our three-stage training algorithm and our analysis in Section 4:

- Learning for different types of samples follows different orders. From Figure 1a, the loss for the samples with consistent signs drops in the early stage of training, while the loss for the samples with conflicting signs only start to drop after sufficient amount of training.
- There is also a learning order of the weights. From Figure 1b, the neuron weights start to align with the underlying signals since the beginning of the training. On the other hand, by Figure 1c, only after sufficient amount of training, the attention scores start to exhibit noticeable change. We further conduct experiments by fixing the neuron weights as Gaussian and only training the attention. Our results show that the model is not trainable in this case.
- The time when the attention scores exhibit noticeable change coincides with the time when the loss for samples with conflicting signs drops. This suggests that attentions play a central role in learning those samples.
- The attention weights are positively associating the group signals and classification signals from the same group and negatively associating them from different groups. This can be seen from Figure 1c that the score $c_1^\top W_{K_+}^{(t)\top} W_{Q_+}^{(t)} v_1$ is increasing to a large positive magnitude and the score $c_1^\top W_{K_+}^{(t)\top} W_{Q_+}^{(t)} v_2$ is decreasing to a large negative magnitude.

In order to understand the mechanism of neurons and attentions separately, the second observation suggests to decompose the training process into stages: neurons are trained first, followed by training the attention. The process can be finished by training neurons again to drive the loss towards zero.

3.2 THREE-STAGE TRAINING ALGORITHM

Motivated by our experiment results, we introduce our three-stage layer-wise training algorithm in Algorithm 1. In our layer-wise training algorithm, in the first stage we train the neuron weights w_+, w_- . In the second stage, we train the attention weights $W_{K+}, W_{Q+}, W_{K-}, W_{Q-}$ on the samples with conflicting signs. In the third stage, we train the neuron weights again. Our algorithm design closely reflects our experimental observation in Section 3.

Algorithm 1 Three-stage Training

Initialize $w_{\pm}^{(0)} = 0$, $W_{K\pm}^{(0)}, W_{Q\pm}^{(0)} \sim \mathcal{N}(0, \omega^2 \frac{1}{m})$ and b to be a sufficiently small positive constant such as $1/2$.

while $t \leq T_1$ **do** ▷ Stage 1: training the neuron weights with all samples

$$\frac{dw_{\pm}^{(t)}}{dt} = -\mathbb{E}_{(X,y)}[\nabla_{w_{\pm}} \ell(yf^{(t)}(X))].$$

$$[W_{K\pm}^{(t)}, W_{Q\pm}^{(t)}] = [W_{K\pm}^{(0)}, W_{Q\pm}^{(0)}].$$

end while

while $t \in [T_1, T_2]$ **do** ▷ Stage 2: training the softmax attention with the conflicting samples

$$\frac{d[W_{K\pm}^{(t)}, W_{Q\pm}^{(t)}]}{dt} = -\mathbb{E}_{(X,y)}[\nabla_{[W_{K\pm}, W_{Q\pm}]} \ell(yf^{(t)}(X)) | X = X_{k,s,-s}].$$

$$w_{\pm}^{(t)} = w_{\pm}^{(T_1)}.$$

end while

while $t \in [T_2, T_3]$ **do** ▷ Stage 3: training the neuron weights to minimize the population loss

$$\frac{dw_{\pm}^{(t)}}{dt} = -\mathbb{E}_{(X,y)}[\nabla_{w_{\pm}} \ell(yf^{(t)}(X))].$$

$$[W_{K\pm}^{(t)}, W_{Q\pm}^{(t)}] = [W_{K\pm}^{(T_2)}, W_{Q\pm}^{(T_2)}].$$

end while

4 MAIN RESULTS

Before we state the main results, we first introduce the parameter conditions used in our analysis.

Condition 1. *We use the following parameter conditions:*

- *The attention initialization scale satisfies $\omega < C < 1$ for some sufficiently small constant C .*
- *The attention weight embedding dimension $m \geq \text{poly}(1/\omega)$.*

We now state the guarantee for Algorithm 1 under an idealized infinite-width transformer model which we will introduce later in Definition 5.1. Our experiment results show that this idealized model is indeed an accurate model that captures the real training dynamics.

Theorem 4.1 (Main theorem). *Under Condition 1 and the infinite-width transformer model in Definition 5.1, there exists $T_1 = \Theta(1)$, $T_2 - T_1 = O(\log(1/\omega))$, $T_3 - T_2 = O(1/\epsilon)$ such that after running Algorithm 1, we have $\mathbb{E}[\ell(yf^{(T_3)}(X))] \leq \epsilon$. The neuron weights of the transformer satisfies*

- *Positive alignment with the signed classification signals $w_+^{(T_3)\top} v_k = \tilde{\Theta}(1)$ and $w_-^{(T_3)\top} (-v_k) = \tilde{\Theta}(1)$.*
- *Positive alignment with the group signals $w_{\pm}^{(T_3)\top} c_k = \tilde{\Theta}(1)$.*
- *Symmetry between heads and groups: $w_+^{(T_3)\top} v_k = w_-^{(T_3)\top} (-v_k)$, $w_+^{(T_3)\top} v_1 = w_+^{(T_3)\top} v_2$, $w_+^{(T_3)\top} c_k = w_-^{(T_3)\top} c_k$, $w_+^{(T_3)\top} c_1 = w_+^{(T_3)\top} c_2$.*

In particular, the output of the softmax attention for sample $X_{1,+,-}$ satisfies the following properties:

- *Concentration on the group signal key:*

$$p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, T_3)}(X_{1,+,-}) - p_{q \leftarrow v_1, k \leftarrow c_1}^{(-, T_3)}(X_{1,+,-}) = \Theta(1)$$

$$\begin{aligned}
& p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+, T_3)}(X_{1,+, -}) - p_{q \leftarrow -v_2, k \leftarrow c_1}^{(-, T_3)}(X_{1,+, -}) = \Theta(1) \\
& p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_3)}(X_{1,+, -}) - p_{q \leftarrow c_1, k \leftarrow c_1}^{(-, T_3)}(X_{1,+, -}) \geq 0
\end{aligned}$$

• *Near symmetry within a single attention head:*

$$\begin{aligned}
& \left| p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) - p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) \right| \leq \tilde{O}(\omega) \\
& \left| p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) - p_{q \leftarrow -v_2, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) \right| \leq \tilde{O}(\omega) \\
& \left| p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) - p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) \right| \leq \tilde{O}(\omega).
\end{aligned}$$

We now describe how the transformer can correctly classify the sample $X_{1,+, -}$. Based on Theorem 4.1, the output of the transformer $f^{(T_3)}(X_{1,+, -})$ can be analyzed as follows: by the attention’s concentration on the group signal key, we have

$$\begin{aligned}
& w_+^{(T_3)\top} c_1 \left(p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_3)}(X_{1,+, -}) \right) \\
& - w_-^{(T_3)\top} c_1 \left(p_{q \leftarrow v_1, k \leftarrow c_1}^{(-, T_3)}(X_{1,+, -}) + p_{q \leftarrow -v_2, k \leftarrow c_1}^{(-, T_3)}(X_{1,+, -}) + p_{q \leftarrow c_1, k \leftarrow c_1}^{(-, T_3)}(X_{1,+, -}) \right) \geq \Omega(1).
\end{aligned}$$

On the other hand, due to attention’s near symmetry property, regardless of queries, we have $w_+^{(T_3)\top} v_1$ and $w_+^{(T_3)\top} (-v_2)$ approximately cancelling each other:

$$\begin{aligned}
& w_+^{(T_3)\top} (-v_2) \left(p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow -v_2, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+, T_3)}(X_{1,+, -}) \right) \\
& + w_+^{(T_3)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) + p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_3)}(X_{1,+, -}) \right) \approx 0.
\end{aligned}$$

Therefore, the positive head can output a value larger than the negative head and thus the sample can be correctly classified. Our results on the behavior of the softmax attention in Theorem 4.1 agrees with our experiment results in Figure 1c. By symmetry, the trained transformer will behave similarly on other samples with conflicting signs.

We now introduce another interpretation of Theorem 4.1 that relates to logistic linear regression. Notice that the transformer architecture in Equation (1) can be rewritten as

$$f(X) = [w_+^\top, w_-^\top, 1] \underbrace{\begin{bmatrix} \dot{\sigma}(H_+(X) + b) \sum_{l=1}^L X \text{softmax}(X^\top W_{K^+}^\top W_{Q^+} x_l) \\ -\dot{\sigma}(H_-(X) + b) \sum_{l=1}^L X \text{softmax}(X^\top W_{K^-}^\top W_{Q^-} x_l) \\ \dot{\sigma}(H_+(X) + b)b - \dot{\sigma}(H_-(X) + b)b \end{bmatrix}}_{\mathcal{F}(X)} \quad (2)$$

We define the vector $\mathcal{F}(X) \in \mathbb{R}^{2d+1}$ to be the **attention feature** of input X . Thus, if the transformer model can successfully learn the data distribution \mathcal{D} , we must have the set of attention features $\{\mathcal{F}(X)\}_{X \in \mathcal{D}}$ linearly separable. We show in Lemma 5.4 that the attention features under uniform attention is not linearly separable and we describe how the attention features evolve and become well-separated after training in Section 5.2.

Finally, for samples with consistent signs, although their attention features also change after stage 2, we are able to show that those changes do not affect the *separability* of their attention features.

5 PROOF OUTLINE

In order to analyze the training dynamics of the softmax attention, first of all, we need to keep track of the dynamics of the score variables $\mu^\top W_K^\top W_Q \nu$ for $\mu, \nu \in \mathcal{C} \cup \mathcal{V}$. However, upon writing down the gradient flow updates for $\mu^\top W_K^\top W_Q \nu$, it can be seen that their updates are governed by the variables in the form of $\delta \mu^\top W_K^\top W_K \nu$ and $\mu^\top W_Q^\top W_Q \nu$ which we call **key and query self-score variables**. By keeping track of the gradient flow updates for those three types of variables all at once, we now have a complete dynamical system where all the terms appearing in the updates are tracked. The gradient flow dynamical system can be found in Appendix B.2.

Before we analyze the training dynamics, we observe one complication brought by random initialization: from our data distribution in Definition 2.1, the two mixtures are symmetric to each other and thus, ideally, the transformer’s behavior should be the same for each class, e.g., $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 = c_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2$. However, due to random initialization, $c_1^\top W_{K+}^{(0)\top} W_{Q+}^{(0)} v_1 \neq c_2^\top W_{K+}^{(0)\top} W_{Q+}^{(0)} v_2$. In order to solve the above complication, we consider the infinite-width transformer where the embedding dimension m of the softmax attention goes to infinity. Within this model, we can derive nice symmetry properties between two groups and between two heads (Appendix C), that are convenient for our analysis.

Definition 5.1 (Infinite-width transformer). *The infinite-width transformer is defined as the architecture in Equation (1) with $m \rightarrow \infty$.*

Our experiment results show that as long as the initialization scale ω is small enough, and the embedding width is large enough, the infinite-width transformer can indeed serve as an accurate model reflecting the true dynamics. For clarity, we present all of our results and analysis under the infinite-width transformer model. The finite-width transformer is a discretized version of the infinite-width transformer. Thus, the actual gradient flow dynamics can be seen as the infinite-width dynamics with perturbation introduced by finite-width discretization. We discuss how to take the discretization into account in Appendix G.

5.1 STAGE 1: NEURONS ALIGN WITH THE SIGNALS

In stage 1 of Algorithm 1, we train the neuron weights and keep the attention weights fixed at their random initialization. Since the neuron weights are zero-initialized, the outputs of the two attention heads are both zero at initialization. Recall that we fix the bias term b in our transformer (Equation (1)) to be some small positive constant. Thus, there exists a period of time from initialization that the outputs of ReLU will be positive for *all* samples. We first characterize how the neuron weights align with the signals in such a case.

Lemma 5.2 (Neuron alignment, stage 1.1). *Under Definition 5.1, for $t \in [0, T_1]$, if $H_+^{(t)}(X), H_-^{(t)}(X) > -b$ for all $X \in \mathcal{D}$, then for $k \in \{1, 2\}$,*

$$\frac{dw_+^{(t)\top} v_k}{dt} = \frac{1}{8} \left(\sum_{k' \in [2], s=s'} g^{(t)}(X_{k',s,s'}) \right), \quad \frac{dw_-^{(t)\top} v_k}{dt} = -\frac{1}{8} \left(\sum_{k' \in [2], s=s'} g^{(t)}(X_{k',s,s'}) \right).$$

At the same time, for $k \in \{1, 2\}$,

$$\frac{dw_+^{(t)\top} c_k}{dt} = 0, \quad \frac{dw_-^{(t)\top} c_k}{dt} = 0.$$

From the above lemma, we can see that when the output of ReLUs are positive for all samples, the neuron weights will align with the signed classification signals while not aligning with the group signals at all. A direct consequence is that the attention head output $H_+^{(t)}(X_{1,-,-})$ will keep decreasing until it becomes smaller than $-b$, as long as b is not too large. Then the neuron weight w_+ is no longer able to receive gradient from sample $X_{1,-,-}$. We can show that when this happens, the neurons start to positively align with the group signals.

Lemma 5.3 (Neuron alignment with group signals, stage 1.2). *Under Definition 5.1, for $t \in [0, T_1]$, if $H_+^{(t)}(X_{k,-,-}) < -b$ for $k \in \{1, 2\}$, then*

$$\frac{dw_+^{(t)\top} c_k}{dt} = \frac{1}{8} g^{(t)}(X_{k,+,+}).$$

At the same time, the neurons still keep aligning with the signed classification signals. Thus, if $T_1 = \Theta(1)$, then we have $w_+^{(t)\top} v_k = \Theta(1)$ and $w_+^{(t)\top} c_k = \Theta(1)$ after stage 1.

Although the neuron weights can learn the signals in stage 1, it is not hard to show that the transformer will output zero for samples with conflicting signs like $X_{1,+,-}$ and thus those samples are not correctly classified. In fact, we can show that under the averaged initial value model, our data distribution \mathcal{D} is not learnable by merely training the neuron weights.

Lemma 5.4 (Non-separability of uniform attention feature). *There exist no neuron weights w_{\pm} such that $yf(X) > 0$ for all $X \in \{X_{k,s,-s}\}_{k \in [2], s \in \{\pm 1\}}$ when the softmax attention is uniform.*

Recall our definition of attention feature in Equation (2). Lemma 5.4 can be interpreted as that the uniform attention feature is not linearly separable. Due to randomness, the actual attention feature at initialization is not exactly uniform. However, by a simple continuity argument, Lemma 5.4 can be translated as the maximum margin of the initial attention feature is at most $\tilde{O}(\omega^2/\sqrt{m})$.¹

5.2 STAGE 2: ATTENTION LEARNS FROM THE SAMPLES WITH CONFLICTING SIGNS

Since by Lemma 5.4, the samples with conflicting signs are hard to learn by merely training the neuron weights, in stage 2, we directly train the softmax attention weights $W_{K\pm}, W_{Q\pm}$ on those hard samples. As we mentioned in the beginning of Section 5, in order to keep track of how the softmax attention changes during training, we need to analyze a dynamical system with all the score variables $\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu$, key self-score variables $\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu$ and query self-score variables $\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu$. From the gradient flow update in Appendix B.2, observe that we can write the dynamical system in the following form:

$$\frac{d}{dt} \begin{bmatrix} \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \end{bmatrix} = A^{(t)} \begin{bmatrix} \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \end{bmatrix}$$

where $A^{(t)}$ is a square matrix with dimension $|\{\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu}| + |\{\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu\}_{\mu, \nu}| + |\{\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu}| = 16 + 10 + 10 = 36$. Thus, directly analyzing the dynamical system can easily become intractable. We need to come up with ways to utilize the structure of the problem to reduce the dimension we need to analyze.

One observation from Figure 1c is that the score variables $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ and $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2$ exhibit the maximum magnitude change compared with all other score variables. In fact, these two quantities play a key role in making $H_+(X_{1,+,-}) > H_-(X_{1,+,-})$. Thus, if we are able to prove that $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ will become sufficiently large at the end of stage 2, then we can make good progress on proving that the attention features become separable.

To make our presentation simpler, for a moment, assume we have the near symmetry properties:

$$\begin{aligned} c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 &\approx -c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 \\ c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 &\approx -c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \\ v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 &\approx -v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \end{aligned}$$

which can indeed be made rigorous in Lemma E.4. Applying this near symmetry property to the gradient of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ at $t = T_1$ (see Appendix E.5.1), we have

$$\begin{aligned} \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \right|_{t=T_1} &\approx \frac{1}{3} g^{(T_1)} 4 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_1 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\ &\quad + \frac{1}{3} g^{(T_1)} 4 \left(c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right). \end{aligned}$$

Due to symmetry and cancellation at initialization, it only depends on 2 self-score variables: $c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1$ and $v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1$. At the same time, the gradient of $c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1$ and $v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1$ at $t = T_1$ also depends only on $c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1$. Thus, if this simpler dependence can hold through the rest of stage 2, then we get a much smaller 3×3 system to analyze. Unfortunately, this is not true. Take the gradient of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ as an example. Training the

¹Recall that the maximum margin of a set \mathcal{S} is defined as $\gamma := \max_{\|w\|_2=1} \min_{x \in \mathcal{S}} w^\top x$.

attention weights can break the symmetry that holds at T_1 , and the gradient of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ will start to depend on other self-score variables. The way to overcome this issue is to realize that as long as the maximum change in the output of the softmax is not too large, we can bound the effects from other self-score variables and the gradient of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ is still mainly contributed by $c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1$ and $v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1$.

Now, we have successfully reduce the dimension of the system that we need to analyze to 3×3 . However, there is still a problem we need to handle: the system has a matrix with *non-linear* dependence on those 3 variables. Generally, it is impossible to derive a close-form solution. Fortunately, we can show that (1) those 3 variables are increasing throughout the training and (2) when the change of the softmax attention (from initialization) is within a certain range, the growth of those 3 variables can be lower bounded by a linear dynamical system.

Lemma 5.5 (Informal version of Lemma E.2). *For $t \geq T_1$, when all the score variables are bounded within some range, there exists a constant $b > 0$ such that*

$$\frac{d}{dt} \begin{bmatrix} c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \\ c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \\ v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \end{bmatrix} \geq b \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \\ c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \\ v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \end{bmatrix}.$$

With Lemma 5.5, we can prove that the score $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ will become sufficiently large at the end of stage 2.

On the other hand, when the query token is the group signal, we can prove

$$v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 < 0, \quad v_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 > 0.$$

By symmetry, this implies $p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,-}) > p_{q \leftarrow c_1, k \leftarrow c_1}^{(-,t)}(X_{1,+,-})$. This finishes the proof of the trained attention’s property on the concentration on the group signal token in Theorem 4.1.

Finally, when v_1 and v_2 are the keys, we can prove the near symmetry property between scores:

$$\begin{aligned} v_1^\top W_{K+}^{(t)\top} W_{Q+} v_1 &\approx -v_2^\top W_{K+}^{(t)\top} W_{Q+} v_1, \\ v_1^\top W_{K+}^{(t)\top} W_{Q+} v_2 &\approx -v_2^\top W_{K+}^{(t)\top} W_{Q+} v_2 \\ v_1^\top W_{K+}^{(t)\top} W_{Q+} c_1 &\approx -v_2^\top W_{K+}^{(t)\top} W_{Q+} c_1 \end{aligned}$$

which can be translated to the near symmetry property within a single head in Theorem 4.1.

5.3 STAGE 3: CONVERGENCE

In stage 3, we train the neuron weights again. At the end of stage 2, the attention features for samples with conflicting signs becomes well-separated. In addition, we can show that the attention features for the samples with consistent signs are still well-separated. Thus, the training in stage 3 behaves like logistic linear regression on linearly separable data and training the neuron weight can drive the population loss towards zero. This proves the part in Theorem 4.1 that the population loss can become sufficiently small after training.

6 GENERAL NUMBER OF MIXTURES

In this section, we study the case when the mixtures have more than two groups. Our experiment results can be found in Appendix A.4. We summarize our experimental findings below.

Observation 1. *The transformer architecture in Equation (1) can learn the data distribution for $K \geq 3$. However, the near symmetry property in the attention no longer holds. In addition, the training takes longer with more groups.*

On the other hand, we observe that for $K = 3$, the training dynamics is still stage-wise and there are score variables exhibiting major growth compared with others. Thus, the techniques developed in our work can be used to study the training dynamics for $K \geq 3$.

Although the training dynamics for $K \geq 3$ has some differences from the $K = 2$ case, we are able to provide a construction of transformers that can solve any arbitrary K mixtures.

Lemma 6.1. *There exists a transformer architecture solving K -mixture of linear classification.*

Proof. The construction is given as follows. Let

$$W_{K+}^\top W_{Q+} = 2C \cdot \sum_{k=1}^K v_k c_k^\top + C \cdot \sum_{k=1}^K c_k c_k^\top + C \cdot \sum_{k=1}^K v_k v_k^\top,$$

$$W_{K-}^\top W_{Q-} = -2C \cdot \sum_{k=1}^K v_k c_k^\top + C \cdot \sum_{k=1}^K c_k c_k^\top + C \cdot \sum_{k=1}^K v_k v_k^\top$$

The transformer is given as

$$f(X) = \sum_{l_1} \sum_{k=1}^K v_k^\top X \text{softmax}(X^\top W_{K+}^\top W_{Q+} X_{l_1}) + \sum_{l_1} \sum_{k=1}^K -v_k^\top X \text{softmax}(X^\top W_{K-}^\top W_{Q-} X_{l_1})$$

Let $C \rightarrow \infty$. First of all, consider classification signal query v_k . Then,

$$\sum_{k=1}^K v_k^\top X \text{softmax}(X^\top W_{K+}^\top W_{Q+} v_k) + \sum_{k=1}^K -v_k^\top X \text{softmax}(X^\top W_{K+}^\top W_{Q+} v_k)$$

$$= v_k^\top v_k - v_k^\top v_k = 0$$

Similar analysis for query $-v_k$. On the other hand, for group signal query c_k , suppose the corresponding classification signal in X is v_k , we have

$$\sum_{k=1}^K v_k^\top X \text{softmax}(X^\top W_{K+}^\top W_{Q+} c_k) + \sum_{k=1}^K -v_k^\top X \text{softmax}(X^\top W_{K+}^\top W_{Q+} c_k)$$

$$= v_k^\top v_k - v_k^\top c_k = 1$$

Similar analysis if the corresponding classification signal is $-v_k$. Thus, we have $yf(X) > 0$ for all $(X, y) \in \mathcal{D}$. \square

7 DISCUSSION

In this work, we studied the training dynamics of a two-headed transformer solving two-mixture of linear classification. We characterized the feature learning process of how the softmax attention utilizes the relationship between the class signals and their corresponding classification signals in this task. One open problem is that the current proof techniques critically utilize the fact that both the transformer architecture and the data distribution are symmetric. Can we discover some hidden structures of the problem such that we can analyze the training dynamics for $K \geq 3$? More interestingly, we demonstrated that during training, the attention features evolved from a near non-separable state to a well-separated state, i.e, the maximum margin of the attention features increases during training. What properties does the data distribution need to have such that the maximum margin can increase in training? We leave those interesting questions as future works.

REFERENCES

- AlphaProof and teams AlphaGeometry. Ai achieves silver-medal standard solving international mathematical olympiad problems. 2024. URL <https://deepmind.google/discover/blog/ai-solves-imo-problems-at-silver-medal-level/>.
- Animashree Anandkumar, Daniel Hsu, and Sham M Kakade. A method of moments for mixture models and hidden markov models. In *Conference on learning theory*, pp. 33–1. JMLR Workshop and Conference Proceedings, 2012.
- Animashree Anandkumar, Rong Ge, Daniel J Hsu, Sham M Kakade, Matus Telgarsky, et al. Tensor decompositions for learning latent variable models. *J. Mach. Learn. Res.*, 15(1):2773–2832, 2014.

- 540 Anthropic. The claude 3 model family: Opus, sonnet, haiku. 2024.
541
- 542 Sivaraman Balakrishnan, Martin J Wainwright, and Bin Yu. Statistical guarantees for the em algo-
543 rithm: From population to sample-based analysis. 2017.
- 544 Sébastien Bubeck, Varun Chandrasekaran, Ronen Eldan, Johannes Gehrke, Eric Horvitz, Ece Ka-
545 mar, Peter Lee, Yin Tat Lee, Yuanzhi Li, Scott Lundberg, et al. Sparks of artificial general
546 intelligence: Early experiments with gpt-4. *arXiv preprint arXiv:2303.12712*, 2023.
547
- 548 Arun Tejasvi Chaganty and Percy Liang. Spectral experts for estimating mixtures of linear regres-
549 sions. In *International Conference on Machine Learning*, pp. 1040–1048. PMLR, 2013.
550
- 551 Siyu Chen, Heejune Sheen, Tianhao Wang, and Zhuoran Yang. Unveiling induction heads: Provable
552 training dynamics and feature learning in transformers. In *ICML 2024 Workshop on Theoretical
553 Foundations of Foundation Models*.
- 554 Siyu Chen, Heejune Sheen, Tianhao Wang, and Zhuoran Yang. Training dynamics of multi-head
555 softmax attention for in-context learning: Emergence, convergence, and optimality. *arXiv preprint
556 arXiv:2402.19442*, 2024.
557
- 558 Zixiang Chen, Yihe Deng, Yue Wu, Quanquan Gu, and Yuanzhi Li. Towards understanding the
559 mixture-of-experts layer in deep learning. *Advances in neural information processing systems*,
560 35:23049–23062, 2022.
- 561 Richard D De Veaux. Mixtures of linear regressions. *Computational Statistics & Data Analysis*, 8
562 (3):227–245, 1989.
563
- 564 Puneesh Deora, Rouzbeh Ghaderi, Hossein Taheri, and Christos Thrampoulidis. On the optimization
565 and generalization of multi-head attention. *Transactions on Machine Learning Research*.
- 566 Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. Bert: Pre-training of deep
567 bidirectional transformers for language understanding. *arXiv preprint arXiv:1810.04805*, 2018.
568
- 569 Alexey Dosovitskiy, Lucas Beyer, Alexander Kolesnikov, Dirk Weissenborn, Xiaohua Zhai, Thomas
570 Unterthiner, Mostafa Dehghani, Matthias Minderer, Georg Heigold, Sylvain Gelly, et al. An im-
571 age is worth 16x16 words: Transformers for image recognition at scale. In *International Confer-
572 ence on Learning Representations*, 2020.
- 573 Benjamin L Edelman, Ezra Edelman, Surbhi Goel, Eran Malach, and Nikolaos Tsilivis. The
574 evolution of statistical induction heads: In-context learning markov chains. *arXiv preprint
575 arXiv:2402.11004*, 2024.
576
- 577 Cheng Gao, Yuan Cao, Zihao Li, Yihan He, Mengdi Wang, Han Liu, Jason M Klusowski, and
578 Jianqing Fan. Global convergence in training large-scale transformers.
579
- 580 Team Gemini. Gemini: a family of highly capable multimodal models. *arXiv preprint
581 arXiv:2312.11805*, 2023.
- 582 Daniel J Hsu, Sham M Kakade, and Percy S Liang. Identifiability and unmixing of latent parse trees.
583 *Advances in neural information processing systems*, 25, 2012.
584
- 585 Yu Huang, Yuan Cheng, and Yingbin Liang. In-context convergence of transformers. *arXiv preprint
586 arXiv:2310.05249*, 2023.
- 587 Yu Huang, Zixin Wen, Yuejie Chi, and Yingbin Liang. Transformers provably learn feature-position
588 correlations in masked image modeling. *arXiv preprint arXiv:2403.02233*, 2024.
589
- 590 Samy Jelassi, Michael Sander, and Yuanzhi Li. Vision transformers provably learn spatial structure.
591 *Advances in Neural Information Processing Systems*, 35:37822–37836, 2022.
592
- 593 Michael I Jordan and Robert A Jacobs. Hierarchical mixtures of experts and the em algorithm.
Neural computation, 6(2):181–214, 1994.

- 594 John Jumper, Richard Evans, Alexander Pritzel, Tim Green, Michael Figurnov, Olaf Ronneberger,
595 Kathryn Tunyasuvunakool, Russ Bates, Augustin Židek, Anna Potapenko, et al. Highly accurate
596 protein structure prediction with alphafold. *nature*, 596(7873):583–589, 2021.
- 597 Abbas Khalili and Jiahua Chen. Variable selection in finite mixture of regression models. *Journal*
598 *of the American Statistical Association*, 102(479):1025–1038, 2007.
- 600 Juno Kim and Taiji Suzuki. Transformers learn nonlinear features in context: Nonconvex mean-field
601 dynamics on the attention landscape. *arXiv preprint arXiv:2402.01258*, 2024.
- 602 Hongkang Li, Meng Wang, Sijia Liu, and Pin-Yu Chen. A theoretical understanding of shallow
603 vision transformers: Learning, generalization, and sample complexity. In *The Eleventh Interna-*
604 *tional Conference on Learning Representations*, 2022.
- 606 Hongkang Li, Meng Wang, Tengfei Ma, Sijia Liu, Zaixi Zhang, and Pin-Yu Chen. What improves
607 the generalization of graph transformer? a theoretical dive into self-attention and positional en-
608 coding. *NeurIPS 2023 Workshop: New Frontiers in Graph Learning*, 2023, 2023a.
- 609 Hongkang Li, Meng Wang, Songtao Lu, Xiaodong Cui, and Pin-Yu Chen. How do nonlinear trans-
610 formers acquire generalization-guaranteed cot ability? In *ICML 2024 Workshop on Theoretical*
611 *Foundations of Foundation Models*, 2024a. URL [https://openreview.net/forum?id=](https://openreview.net/forum?id=HiZDrDvxLP)
612 [HiZDrDvxLP](https://openreview.net/forum?id=HiZDrDvxLP).
- 613 Hongkang Li, Meng Wang, Songtao Lu, Xiaodong Cui, and Pin-Yu Chen. Training nonlinear trans-
614 formers for efficient in-context learning: A theoretical learning and generalization analysis. *arXiv*
615 *preprint arXiv:2402.15607*, 2024b.
- 617 Yuchen Li, Yuanzhi Li, and Andrej Risteski. How do transformers learn topic structure: Towards a
618 mechanistic understanding. In *International Conference on Machine Learning*, pp. 19689–19729.
619 PMLR, 2023b.
- 620 Zihao Li, Yuan Cao, Cheng Gao, Yihan He, Han Liu, Jason M Klusowski, Jianqing Fan, and Mengdi
621 Wang. One-layer transformer provably learns one-nearest neighbor in context. 2024c.
- 622 Percy Liang, Alexandre Bouchard-Côté, Dan Klein, and Ben Taskar. An end-to-end discriminative
623 approach to machine translation. In *Proceedings of the 21st International Conference on Compu-*
624 *tational Linguistics and 44th Annual Meeting of the Association for Computational Linguistics*,
625 pp. 761–768, 2006.
- 627 Ze Liu, Yutong Lin, Yue Cao, Han Hu, Yixuan Wei, Zheng Zhang, Stephen Lin, and Baining Guo.
628 Swin transformer: Hierarchical vision transformer using shifted windows. In *Proceedings of the*
629 *IEEE/CVF international conference on computer vision*, pp. 10012–10022, 2021.
- 630 Ashok Vardhan Makkuva, Marco Bondaschi, Chanakya Ekbote, Adway Girish, Alliot Nagle, Hyeji
631 Kim, and Michael Gastpar. Local to global: Learning dynamics and effect of initialization for
632 transformers. *arXiv preprint arXiv:2406.03072*, 2024.
- 633 James Manyika and Sissie Hsiao. An overview of bard: an early experiment with generative ai. *AI*
634 *Google Static Documents*, 2, 2023.
- 636 Eshaan Nichani, Alex Damian, and Jason D Lee. How transformers learn causal structure with
637 gradient descent. *arXiv preprint arXiv:2402.14735*, 2024.
- 638 OpenAI. Gpt-4 technical report, 2023.
- 640 Ariadna Quattoni, Michael Collins, and Trevor Darrell. Conditional random fields for object recog-
641 nition. *Advances in neural information processing systems*, 17, 2004.
- 642 Davoud Ataee Tarzanagh, Yingcong Li, Christos Thrampoulidis, and Samet Oymak. Transformers
643 as support vector machines. In *NeurIPS 2023 Workshop on Mathematics of Modern Machine*
644 *Learning*, 2023a.
- 645 Davoud Ataee Tarzanagh, Yingcong Li, Xuechen Zhang, and Samet Oymak. Max-margin token
646 selection in attention mechanism. In *Thirty-seventh Conference on Neural Information Processing*
647 *Systems*, 2023b.

- 648 Yuandong Tian, Yiping Wang, Beidi Chen, and Simon Du. Scan and snap: Understanding training
649 dynamics and token composition in 1-layer transformer. *arXiv preprint arXiv:2305.16380*, 2023.
650
- 651 Hugo Touvron, Thibaut Lavril, Gautier Izacard, Xavier Martinet, Marie-Anne Lachaux, Timothée
652 Lacroix, Baptiste Rozière, Naman Goyal, Eric Hambro, Faisal Azhar, et al. Llama: Open and
653 efficient foundation language models. *arXiv preprint arXiv:2302.13971*, 2023.
- 654 Bhavya Vasudeva, Puneesh Deora, and Christos Thrampoulidis. Implicit bias and fast convergence
655 rates for self-attention. *arXiv preprint arXiv:2402.05738*, 2024.
656
- 657 Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez,
658 Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural informa-
659 tion processing systems*, 30, 2017.
- 660 Zhaoran Wang, Quanquan Gu, Yang Ning, and Han Liu. High dimensional em algorithm: Statistical
661 optimization and asymptotic normality. *Advances in neural information processing systems*, 28,
662 2015.
- 663 Zixuan Wang, Stanley Wei, Daniel Hsu, and Jason D Lee. Transformers provably learn sparse token
664 selection while fully-connected nets cannot. In *Forty-first International Conference on Machine
665 Learning*.
666
- 667 Yongtao Wu, Fanghui Liu, Grigorios Chrysos, and Volkan Cevher. On the convergence of encoder-
668 only shallow transformers. In *Thirty-seventh Conference on Neural Information Processing Sys-
669 tems*, 2023.
- 670 Tong Yang, Yu Huang, Yingbin Liang, and Yuejie Chi. In-context learning with representations:
671 Contextual generalization of trained transformers. In *ICML 2024 Workshop on Theoretical Foun-
672 dations of Foundation Models*.
673
- 674 Xinyang Yi, Constantine Caramanis, and Sujay Sanghavi. Alternating minimization for mixed linear
675 regression. In *International Conference on Machine Learning*, pp. 613–621. PMLR, 2014.
- 676 Ruiqi Zhang, Spencer Frei, and Peter L Bartlett. Trained transformers learn linear models in-context.
677 *Journal of Machine Learning Research*, 25(49):1–55, 2024.
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701

702	CONTENTS	
703		
704	1 Introduction	1
705		
706	1.1 Related Work	2
707		
708	2 Settings	3
709		
710	2.1 Data distribution	3
711	2.2 Transformer architecture and loss function	3
712		
713	3 Guiding Experiments and Algorithm	4
714		
715	3.1 Experiment results	4
716	3.2 Three-stage Training Algorithm	5
717		
718	4 Main Results	5
719		
720	5 Proof Outline	6
721		
722	5.1 Stage 1: Neurons align with the signals	7
723	5.2 Stage 2: Attention learns from the samples with conflicting signs	8
724	5.3 Stage 3: Convergence	9
725		
726		
727	6 General Number of Mixtures	9
728		
729	7 Discussion	10
730		
731	A Additional Experiment Results	16
732		
733	A.1 Softmax attention	16
734	A.2 Only train the attention	16
735	A.3 Full attention scores of two mixtures	17
736	A.4 Experiments on Multi-mixture Models	17
737		
738		
739	B Gradient Flow	18
740		
741	B.1 Gradients of each weight matrices	18
742	B.2 Gradient Flow Dynamical System	19
743		
744	C Symmetry	19
745		
746	D Stage 1: Training neurons to learn the signals	23
747		
748	D.1 Stage 1.1: Growth of neuron alignment with classification signals	23
749	D.2 Stage 1.2: Growth of neuron alignment with group signals	24
750	D.3 Properties at the end of stage 1	25
751		
752	E Stage 2: Self-attention learns from the conflicting (hard) samples	26
753		
754	E.1 Growth of scores	26
755		

756	E.2 Growth of asymmetry	29
757	E.3 When the query is the group signal	31
758	E.4 Properties at the end of stage 2	34
759	E.5 The updates of the softmax attention at $t = T_1$	35
760	E.5.1 The scores	35
761	E.5.2 The key self-scores	37
762	E.5.3 The query self-scores	39
763	E.6 Sum of partial gradients	40
764	E.7 Expanding the updates	42
765	E.7.1 The key self-score	42
766	E.7.2 The query self-score	45
767	E.7.3 The scores	47
768		
769	F Stage 3	52
770		
771	G Handling the finite-width discretization	55
772		
773	H Probability	57
774		
775		
776		
777		
778		
779		
780		
781		
782		
783		
784		
785		
786		
787		
788		
789		
790		
791		
792		
793		
794		
795		
796		
797		
798		
799		
800		
801		
802		
803		
804		
805		
806		
807		
808		
809		

A ADDITIONAL EXPERIMENT RESULTS

A.1 SOFTMAX ATTENTION

We provide the softmax probability of the positive head on different samples.




Figure 2: Softmax attention of the positive head on different samples. The vertical axis denotes the query tokens and the horizontal axis denotes the key tokens.


A.2 ONLY TRAIN THE ATTENTION

Recall that in Section 3, we observed there is a learning order when all the weights are trained simultaneously: the attention weights learn after the neuron weights. To further study whether this learning order is necessary, we conduct another experiment where we initialize the neurons as Gaussian and only train the attention. Our result suggests that in this case, the transformer is no longer trainable.



Figure 3: The loss for training only the attention while keeping the neurons fixed at initialization.

A.3 FULL ATTENTION SCORES OF TWO MIXTURES




(a) Score

Figure 4: Full attention scores for $K = 2$.

A.4 EXPERIMENTS ON MULTI-MIXTURE MODELS

In this section, we provide experiment results on synthetic data for multi-mixture models. Specifically, Figure 5, Figure 6 and Figure 7 respectively indicate that the training dynamics of 3-mixture, 4-mixture and 8-mixture models are all showing some similarities as that for 2-mixture model, demonstrating the broad applicability of our developed theoretical results.



(a) Loss

(b) Neuron alignment

(c) Scores with width 65

(d) Scores with width 260

Figure 5: The experiment results for the case with $K = 3$ mixtures.

918
919
920
921
922
923
924
925
926
927
928
929
930




Figure 6: The experiment results for the case with $K = 4$ mixtures.

931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949




Figure 7: The experiment results for the case with $K = 8$ mixtures.

B GRADIENT FLOW

B.1 GRADIENTS OF EACH WEIGHT MATRICES

958
959
960
961
962
963
964
965
966
967
968
969
970
971

$$\begin{aligned}
 \frac{dw_+^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L X p_l^{(+,t)} \right] \\
 \frac{dw_-^{(t)}}{dt} &= -\mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_-^{(t)} \sum_{l=1}^L X p_l^{(+,t)} \right] \\
 \frac{dW_{K+}^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \sum_{h=1}^L w_+^\top X_h \sum_{h'=1}^L \frac{\partial p_{l,h}^{(+,t)}}{\partial s_{l,h'}^{(+)}} q_{+l}^{(t)} x_{h'}^\top \right] \\
 &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \sum_{h=1}^L w_+^\top X_h \sum_{h'=1}^L p_{l,h}^{(+,t)} (\mathbb{I}(h = h') - p_{l,h'}^{(+,t)}) q_{+l}^{(t)} x_{h'}^\top \right] \\
 &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(-w_+^\top X p_l^{(+,t)} q_{+l}^{(t)} (X p_l^{(+,t)})^\top + q_{+l}^{(t)} w_+^\top X \text{diag}(p_l^{(+,t)}) X^\top \right) \right]
 \end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L q_{+l}^{(t)} \left(-w_+^\top X p_l^{(+,t)} p_l^{(+,t)\top} + w_+^\top X \text{diag}(p_l^{(+,t)}) \right) X^\top \right] \\
&= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top \right] \\
\frac{dW_{Q+}^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L w_+^\top \sum_{h=1}^L X_h \sum_{h'=1}^L p_{l,h}^{(+,t)} (\mathbb{I}(h=h') - p_{l,h'}^{(+,t)}) k_{+h}^{(t)} x_l^{(i)\top} \right] \\
&= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(-w_+^\top X p_l^{(+,t)} (K_+^{(t)} p_l^{(+,t)}) + K_+^{(t)} \text{diag}(p_l^{(+,t)}) X^\top w_+ \right) x_l^\top \right] \\
&= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L K_+^{(t)} \left(-w_+^\top X p_l^{(+,t)} p_l^{(+,t)} + \text{diag}(p_l^{(+,t)}) X^\top w_+ \right) x_l^\top \right] \\
&= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top \right]
\end{aligned}$$

B.2 GRADIENT FLOW DYNAMICAL SYSTEM

Let $\mu, \nu \in \mathcal{V} \cup \mathcal{C}$. We have

$$\begin{aligned}
\frac{dw_+^{(t)\top} \mu}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \mu \right] \\
\frac{d\nu^\top W_{K+}^{(t)\top} W_{Q+\mu}^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top \nu \right] \\
&\quad + \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top \mu \right] \\
\frac{d\nu^\top W_{K+}^{(t)\top} W_{K+\mu}^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top \mu \right] \\
&\quad + \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top \nu \right] \\
\frac{d\nu^\top W_{Q+}^{(t)\top} W_{Q+\mu}^{(t)}}{dt} &= \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top \mu \right] \\
&\quad + \mathbb{E} \left[g^{(t)}(X) y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top \nu \right]
\end{aligned}$$

C SYMMETRY

In this section, we derive symmetry results for the infinite-width transformer model. **We first characterize the initial values of the softmax attention weights:**

Lemma C.1. *The initial values of the attention weight in the infinite-width transformer satisfy:*

$$\nu^\top W_{K\pm}^{(0)\top} W_{Q\pm\mu}^{(0)} = \mathbb{E}_{\widehat{W}_K, \widehat{W}_Q} \left[\nu^\top \widehat{W}_K^\top \widehat{W}_Q \mu \right] = 0$$

$$\begin{aligned} \nu^\top W_{K\pm}^{(0)\top} W_{K\pm}^{(0)} \mu &= \begin{cases} \mathbb{E}_{\widehat{W}_K} \left[\nu^\top \widehat{W}_K^\top \widehat{W}_K \mu \right] = 0, & \nu \neq \mu \\ \mathbb{E}_{\widehat{W}_K} \left[\mu^\top \widehat{W}_K^\top \widehat{W}_K \mu \right] = \omega^2, & \nu = \mu \end{cases} \\ \nu^\top W_{Q\pm}^{(0)\top} W_{Q\pm}^{(0)} \mu &= \begin{cases} \mathbb{E}_{\widehat{W}_Q} \left[\nu^\top \widehat{W}_Q^\top \widehat{W}_Q \mu \right] = 0, & \nu \neq \mu \\ \mathbb{E}_{\widehat{W}_Q} \left[\mu^\top \widehat{W}_Q^\top \widehat{W}_Q \mu \right] = \omega^2, & \nu = \mu \end{cases} \end{aligned}$$

where $\widehat{W}_K, \widehat{W}_Q \in \mathbb{R}^{m \times d}$ and $(\widehat{W}_K)_{ij}, (\widehat{W}_Q)_{ij} \sim \mathcal{N}(0, \omega^2 \frac{1}{m})$.

Proof. Let $(W_{K+})_i, (W_{Q+})_i$ denote the i -th row of W_{K+}, W_{Q+} , respectively. Then, we have $\nu^\top W_{K\pm}^{(0)\top} W_{Q\pm}^{(0)} \mu = \sum_{i=1}^m (W_{K+}^{(0)})_i^\top \nu (W_{Q+}^{(0)})_i \mu$. Notice that $(W_{K+}^{(0)})_i^\top \nu \sim \mathcal{N}(0, \omega^2 \frac{1}{m})$, $(W_{Q+}^{(0)})_i \mu \sim \mathcal{N}(0, \omega^2 \frac{1}{m})$. Thus, $\nu^\top W_{K\pm}^{(0)\top} W_{Q\pm}^{(0)} \mu$ has the same distribution as $\frac{1}{m} \sum_{i=1}^m a_i b_i$ where $a_i, b_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \omega^2)$. By the Law of Large Number,

$$\lim_{m \rightarrow \infty} \nu^\top W_{K\pm}^{(0)\top} W_{Q\pm}^{(0)} \mu = \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{i=1}^m a_i b_i = \mathbb{E}[a_1 b_1] = \mathbb{E}[a_1] \mathbb{E}[b_1] = 0.$$

The rest can be proved similarly with an additional fact that for $w \sim \mathcal{N}(0, I)$, $w^\top \mu$ is independent of $w^\top \nu$ if $\mu \perp \nu$. \square

The symmetry results contain two parts: symmetry between two groups and symmetry between two attention heads.

Lemma C.2 (Symmetry between two groups). *Under the infinite-width transformer model in Definition 5.1, let $\mu_1, \nu_1 \in \{c_1, v_1\}$ and $\mu_2, \nu_2 \in \{c_2, v_2\}$ such that μ_1, μ_2 are the same type signals (i.e., either $\mu_1, \mu_2 \in \mathcal{C}$ or $\mu_1, \mu_2 \in \mathcal{V}$) and ν_1, ν_2 are the same type signals. For all t ,*

$$w_+^{(t)\top} \mu_1 = w_+^{(t)\top} \mu_2 \quad (3)$$

$$\nu_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_1 = \nu_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_2, \quad \nu_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_1 = \nu_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_2 \quad (4)$$

$$\nu_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu_1 = \nu_2^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu_2, \quad \nu_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu_2 = \nu_2^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu_1 \quad (5)$$

$$\nu_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu_1 = \nu_2^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu_2, \quad \nu_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu_2 = \nu_2^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu_1 \quad (6)$$

The symmetry between the two groups also holds for the negative head.

Corollary C.3. *Let $s, s' \in \{\pm 1\}$. For all t ,*

$$p^{(+,t)}(X_{1,s,s'}) = p^{(+,t)}(X_{2,s,s'}), \quad p^{(-,t)}(X_{1,s,s'}) = p^{(-,t)}(X_{2,s,s'})$$

$$H_+^{(t)}(X_{1,s,s'}) = H_+^{(t)}(X_{2,s,s'}), \quad H_-^{(t)}(X_{1,s,s'}) = H_-^{(t)}(X_{2,s,s'})$$

$$f^{(t)}(X_{1,s,s'}) = f^{(t)}(X_{2,s,s'})$$

Proof of Lemma C.2. First of all, by Definition 5.1, all the equalities hold at $t = 0$. Thus, we only need to show the gradient flow updates satisfy all the equalities if all the equalities hold at time t .

Assume all the equalities hold at time t . This implies that $p^{(+,t)}(X_{1,s,s'}) = p^{(+,t)}(X_{2,s,s'})$ (recall that since our model doesn't have any positional encoding, we can consider ordered inputs in our analysis), which further implies that $H_+^{(t)}(X_{1,s,s'}) = H_+^{(t)}(X_{2,s,s'})$, $H_-^{(t)}(X_{1,s,s'}) = H_-^{(t)}(X_{2,s,s'})$ and $f^{(t)}(X_{1,s,s'}) = f^{(t)}(X_{2,s,s'})$ for all $s, s' \in \{\pm 1\}$.

First, we prove Equation (3):

$$\begin{aligned} \frac{dw_+^{(t)\top} \mu_1}{dt} &= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] \mu_1 \\ &= \mathbb{E}_{s,s'} \left[\frac{1}{2} g^{(t)}(X_{1,s,s'}) s \dot{\sigma}_+^{(t)}(X_{1,s,s'}) \sum_{l=1}^L \left(X_{1,s,s'} p_l^{(+,t)}(X_{1,s,s'}) \right)^\top \right] \mu_1 \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} g^{(t)}(X_{2,s,s'}) s \dot{\sigma}_+^{(t)}(X_{2,s,s'}) \sum_{l=1}^L \left(X_{2,s,s'} p_l^{(+,t)}(X_{2,s,s'}) \right)^\top \mu_1 \Big] \\
& = \mathbb{E}_{s,s'} \left[\frac{1}{2} g^{(t)}(X_{2,s,s'}) s \dot{\sigma}_+^{(t)}(X_{2,s,s'}) \sum_{l=1}^L \left(X_{2,s,s'} p_l^{(+,t)}(X_{2,s,s'}) \right)^\top \mu_2 \right. \\
& \quad \left. + \frac{1}{2} g^{(t)}(X_{1,s,s'}) s \dot{\sigma}_+^{(t)}(X_{1,s,s'}) \sum_{l=1}^L \left(X_{1,s,s'} p_l^{(+,t)}(X_{1,s,s'}) \right)^\top \mu_2 \right] \\
& \hspace{15em} \text{(by } p^{(+,t)}(X_{1,s,s'}) = p^{(+,t)}(X_{2,s,s'}) \text{)} \\
& = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] \mu_2 = \frac{dw_+^{(t)\top} \mu_2}{dt}
\end{aligned}$$

Next, we prove the first equality in Equation (4):

$$\begin{aligned}
& \frac{d\nu_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_1}{dt} \\
& = \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X_{1,s,s'} - w_+^{(t)\top} X_{1,s,s'} p_l^{(+,t)} \right) X_{1,s,s'}^\top \nu_1 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X_{2,s,s'} - w_+^{(t)\top} X_{2,s,s'} p_l^{(+,t)} \right) X_{2,s,s'}^\top \nu_1 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X_{1,s,s'}^\top w_+ - w_+^\top X_{1,s,s'} p_l^{(+,t)} \right) p_l^{(+,t)} (X_{1,s,s'})_l^\top \mu_1 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X_{2,s,s'}^\top w_+ - w_+^\top X_{2,s,s'} p_l^{(+,t)} \right) p_l^{(+,t)} (X_{2,s,s'})_l^\top \mu_1 \right] \\
& = \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu_2^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X_{2,s,s'} - w_+^{(t)\top} X_{2,s,s'} p_l^{(+,t)} \right) X_{2,s,s'}^\top \nu_2 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu_2^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X_{1,s,s'} - w_+^{(t)\top} X_{1,s,s'} p_l^{(+,t)} \right) X_{1,s,s'}^\top \nu_2 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu_2^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X_{2,s,s'}^\top w_+ - w_+^\top X_{2,s,s'} p_l^{(+,t)} \right) p_l^{(+,t)} (X_{2,s,s'})_l^\top \mu_2 \right] \\
& \quad + \frac{1}{2} \mathbb{E}_{s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu_2^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X_{1,s,s'}^\top w_+ - w_+^\top X_{1,s,s'} p_l^{(+,t)} \right) p_l^{(+,t)} (X_{1,s,s'})_l^\top \mu_2 \right] \\
& = \frac{d\nu_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_2}{dt}
\end{aligned}$$

The rest equalities in Equation (4), Equation (5) and Equation (6) can be proved similarly. \square

Symmetry between two heads: notice that for the negative head, we can view it as training the model $-f(X)$ on the dataset where the label is $-y$ and the set of classification signals is $\{-v_1, -v_2\}$.

Lemma C.4 (Symmetry of two heads). *Suppose Definition 5.1 holds. Let $\mu_1, \mu_2 \in \mathcal{V} \cup \mathcal{C}$. For all t ,*

$$w_+^{(t)\top} \mu_1 = w_-^{(t)\top} \mu_1 \cdot (-1)^{\mathbb{I}(\mu_1 \in \mathcal{V})} \quad (7)$$

$$\mu_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu_1 = \mu_2^\top W_{K-}^{(t)\top} W_{Q-}^{(t)} \mu_1 \cdot (-1)^{\mathbb{I}(\mu_2 \in \mathcal{V})} (-1)^{\mathbb{I}(\mu_1 \in \mathcal{V})} \quad (8)$$

$$\mu_2^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu_1 = \mu_2^\top W_{K-}^{(t)\top} W_{K-}^{(t)} \mu_1 \cdot (-1)^{\mathbb{I}(\mu_2 \in \mathcal{V})} (-1)^{\mathbb{I}(\mu_1 \in \mathcal{V})} \quad (9)$$

$$\mu_2^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu_1 = \mu_2^\top W_{Q-}^{(t)\top} W_{Q-}^{(t)} \mu_1 \cdot (-1)^{\mathbb{I}(\mu_2 \in \mathcal{V})} (-1)^{\mathbb{I}(\mu_1 \in \mathcal{V})} \quad (10)$$

Corollary C.5. Let $s, s' \in \{\pm 1\}$. For all t ,

$$\begin{aligned} p^{(+,t)}(X_{k,s,s'}) &= p^{(-,t)}(X_{k,-s,-s'}) \\ H_+^{(t)}(X_{k,s,s'}) &= H_-^{(t)}(X_{k,-s,-s'}) \\ sf^{(t)}(X_{k,s,s'}) &= -sf^{(t)}(X_{k,-s,-s'}). \end{aligned}$$

Proof of Lemma C.4. First, all the equalities are satisfied when $t = 0$ by Definition 5.1. Assume all the equalities hold at time t and we are going to show that the gradients with respect to t also satisfy the equalities. Equation (8) implies that $p^{(+,t)}(X_{k,s,s'}) = p^{(-,t)}(X_{k,-s,-s'})$, which further implies $H_+^{(t)}(X_{k,s,s'}) = H_-^{(t)}(X_{k,-s,-s'})$ and $sf^{(t)}(X_{k,s,s'}) = -sf^{(t)}(X_{k,-s,-s'})$.

Define $\mu' := \mu \cdot (-1)^{\mathbb{I}(\mu \in \mathcal{V})}$ for $\mu \in \mathcal{V} \cup \mathcal{C}$ and $X'_{k,s,s'} := [c'_k, v'_k, v'_{k'}] = [c_k, -v_k, -v_{k'}] = X_{k,-s,-s'}$.

We first prove Equation (7):

$$\begin{aligned} \frac{dw_+^{(t)\top} \mu_1}{dt} &= \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X_{k,s,s'} p_l^{(+,t)} \right)^\top \right] \mu_1 \\ \frac{dw_-^{(t)\top} \mu'_1}{dt} &= -\mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L \left(X_{k,s,s'} p_l^{(-,t)} \right)^\top \right] \mu'_1 \\ &= \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L \left(X'_{k,s,s'} p_l^{(-,t)} \right)^\top \right] \mu'_1 \end{aligned}$$

which implies

$$\frac{dw_+^{(t)\top} \mu_1}{dt} = \frac{dw_-^{(t)\top} \mu'_1}{dt}.$$

We next prove Equation (8):

$$\begin{aligned} &\frac{d\nu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)}}{dt} \\ &= \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \mu^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X_{k,s,s'} - w_+^{(t)\top} X_{k,s,s'} p_l^{(+,t)} \right) X_{k,s,s'}^\top \nu \right] \\ &\quad + \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_+^{(t)} \sum_{l=1}^L \nu^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X_{k,s,s'}^\top w_+ - w_+^\top X_{k,s,s'} p_l^{(+,t)} \right) p_l^{(+,t)} (X_{k,s,s'})_l^\top \mu \right] \\ &\frac{d(\nu')^\top W_{K-}^{(t)\top} W_{Q-}^{(t)} \mu'}{dt} \\ &= -\mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L (\mu')^\top W_{Q-}^{(t)\top} q_{-l}^{(t)} p_l^{(-,t)\top} \text{diag} \left(w_-^{(t)\top} X_{k,s,s'} - w_-^{(t)\top} X_{k,s,s'} p_l^{(-,t)} \right) X_{k,s,s'}^\top \nu' \right] \\ &\quad - \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L (\nu')^\top W_{K-}^{(t)\top} K_-^{(t)} \text{diag} \left(X_{k,s,s'}^\top w_- - w_-^\top X_{k,s,s'} p_l^{(-,t)} \right) p_l^{(-,t)} (X_{k,s,s'})_l^\top \mu' \right] \\ &= \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L (\mu')^\top W_{Q-}^{(t)\top} q_{-l}^{(t)} p_l^{(-,t)\top} \text{diag} \left(w_-^{(t)\top} X'_{k,s,s'} - w_-^{(t)\top} X'_{k,s,s'} p_l^{(-,t)} \right) X_{k,s,s'}^\top \nu' \right] \\ &\quad + \mathbb{E}_{k,s,s'} \left[g^{(t)} s \dot{\sigma}_-^{(t)} \sum_{l=1}^L (\nu')^\top W_{K-}^{(t)\top} K_-^{(t)} \text{diag} \left(X_{k,s,s'}^\top w_- - w_-^\top X'_{k,s,s'} p_l^{(-,t)} \right) p_l^{(-,t)} (X'_{k,s,s'})_l^\top \mu' \right] \end{aligned}$$

1188
1189
1190 which implies

$$1191 \frac{d\nu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu}{dt} = \frac{d(\nu')^\top W_{K-}^{(t)\top} W_{Q-}^{(t)} \mu'}{dt}. \quad 1192$$

1193
1194 The rest of equalities Equation (9) and Equation (10) can be proved similarly. \square
1195

1196 D STAGE 1: TRAINING NEURONS TO LEARN THE SIGNALS

1197 D.1 STAGE 1.1: GROWTH OF NEURON ALIGNMENT WITH CLASSIFICATION SIGNALS

1198
1199 Due to our zero-initialization on the neuron weights, both attention heads are outputting zero. Since
1200 we add a small positive bias in the ReLU, for a short amount of time from the random initialization,
1201 we can ignore the ReLU activation.
1202

1203 **Lemma D.1** (Neuron alignment with the classification signal, stage 1.1). *Under Definition 5.1, for*
1204 *$t \in [0, T_1]$, if $H_+^{(t)}(X), H_-^{(t)}(X) > -b$ for all $X \in \mathcal{D}$, then for $k \in \{1, 2\}$,*
1205

$$1206 \frac{dw_+^{(t)\top} v_k}{dt} = \frac{1}{8} \left(\sum_{k' \in [2], s=s'} g^{(t)}(X_{k',s,s'}) \right)$$

$$1207 \frac{dw_-^{(t)\top} v_k}{dt} = -\frac{1}{8} \left(\sum_{k' \in [2], s=s'} g^{(t)}(X_{k',s,s'}) \right). \quad 1208$$

1209
1210 *Proof.* Take $k = 1$. By the gradient flow update, we have
1211

$$1212 \frac{dw_+^{(t)\top} v_1}{dt} = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] v_1$$

$$1213 = \mathbb{E} \left[g^{(t)} y \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] v_1$$

$$1214 = \frac{1}{8} \left(\sum_{s=s'} g^{(t)}(X_{1,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow v_1}^{(+,t)}(X_{1,s,s'}) \right) + \frac{1}{8} \left(\sum_{s \neq s'} s g^{(t)}(X_{1,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow s v_1}^{(+,t)}(X_{1,s,s'}) \right)$$

$$1215 + \frac{1}{8} \left(\sum_{s=s'} g^{(t)}(X_{2,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow v_1}^{(+,t)}(X_{2,s,s'}) \right) - \frac{1}{8} \left(\sum_{s \neq s'} s' g^{(t)}(X_{2,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow s' v_1}^{(+,t)}(X_{2,s,s'}) \right)$$

$$1216 = \frac{1}{8} \left(\sum_{k' \in [2], s=s'} g^{(t)}(X_{k',s,s'}) \right). \quad \text{(by Corollary C.3)}$$

1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232 \square

1233 **Lemma D.2** (No neuron alignment of group signals, stage 1.1). *Under Definition 5.1, as long as*
1234 *$H_+^{(t)}(X), H_-^{(t)}(X) > -b$ for all X , and for $k \in \{1, 2\}$,*

$$1235 \frac{dw_+^{(t)\top} c_k}{dt} = 0, \quad \frac{dw_-^{(t)\top} c_k}{dt} = 0.$$

1236
1237
1238 *Proof.* By the gradient flow update, we have
1239

$$1240 \frac{dw_+^{(t)\top} c_k}{dt} = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] c_k$$

$$\begin{aligned}
1242 &= \mathbb{E} \left[g^{(t)} y \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] c_k \\
1243 &= \frac{1}{8} \left(\sum_{s'} g^{(t)}(X_{k,+,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(+,t)} - \sum_{s'} g^{(t)}(X_{k,-,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(+,t)} \right) \\
1244 &= 0 \quad \text{(by and Corollary C.5)}
\end{aligned}$$

1249 \square

1251 **Corollary D.3** (Duration of stage 1.1). *There exists a time $T_{0.5} = \Theta(1)$ such that*
1252 $H_s^{(T_{0.5})}(X_{k,-s,-s}) \leq -b$ for $s \in \{\pm 1\}$.

1254 *Proof.* Lemma D.1 and Lemma D.2 imply that $g^{(t)}(X) = \Omega(1)$ for $t \leq O(1)$. Since b is chosen to
1255 be a sufficiently small constant, there exists a time $T_{0.5} = \Theta(1)$ such that $H_s^{(T_{0.5})}(X_{k,-s,-s}) \leq -b$
1256 for $s \in \{\pm 1\}$. \square

1258 D.2 STAGE 1.2: GROWTH OF NEURON ALIGNMENT WITH GROUP SIGNALS

1260 In this sub-stage, we are going to show that (1) the neuron weight will keep positively aligning with
1261 the signed classification signal and (2) the neuron weights start to positively align with the group
1262 signal.

1263 **Lemma D.4** (Growth of classification signals, stage 1.2). *Under Definition 5.1, for $t \in [0, T_1]$, if*
1264 $H_+^{(t)}(X_{k,-,-}) < -b$ for all $k \in [2]$, then

$$1266 \frac{dw_+^{(t)\top} v_k}{dt} = \frac{1}{8} \left(\sum_{k' \in [2]} g^{(t)}(X_{k',+,+}) \right).$$

1269 Similarly, if $H_-^{(t)}(X_{k,+,+}) < -b$ for all $k \in [2]$, then

$$1272 \frac{dw_-^{(t)\top} v_k}{dt} = -\frac{1}{8} \left(\sum_{k' \in [2]} g^{(t)}(X_{k',-,-}) \right).$$

1275 *Proof.* Take $k = 1$. If $H_+^{(t)}(X_{k',-,-}) < -b$ for all $k' \in [2]$, then

$$\begin{aligned}
1277 \frac{dw_+^{(t)\top} v_1}{dt} &= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] v_1 \\
1278 &= \frac{1}{8} \left(g^{(t)}(X_{1,+,+}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow v_1}^{(+,t)}(X_{1,+,+}) \right) + \frac{1}{8} \left(\sum_{s \neq s'} s g^{(t)}(X_{1,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow s v_1}^{(+,t)}(X_{1,s,s'}) \right) \\
1281 &+ \frac{1}{8} \left(g^{(t)}(X_{2,+,+}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow v_1}^{(+,t)}(X_{2,+,+}) \right) - \frac{1}{8} \left(\sum_{s \neq s'} s' g^{(t)}(X_{2,s,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow s' v_1}^{(+,t)}(X_{2,s,s'}) \right) \\
1284 &= \frac{1}{8} \left(\sum_{k' \in [2]} g^{(t)}(X_{k',+,+}) \right). \quad \text{(by Corollary C.3)}
\end{aligned}$$

1290 \square

1292 **Lemma D.5** (Growth of group signals, stage 1.2). *For $t \in [0, T_1]$, if $H_+^{(t)}(X_{k,-,-}) < -b$ for*
1293 $k \in \{1, 2\}$, then

$$1295 \frac{dw_+^{(t)\top} c_k}{dt} = \frac{1}{8} g^{(t)}(X_{k,+,+}).$$

1296 Similarly, if $H_-^{(t)}(X_{k,+,+}) < -b$ for $k \in \{1, 2\}$, then

$$1297 \frac{dw_-^{(t)\top} c_k}{dt} = \frac{1}{8} g^{(t)}(X_{k,-,-}).$$

1301 *Proof.* If $H_+^{(t)}(X_{k,-,-}) < -b$, then

$$1302 \frac{dw_+^{(t)\top} c_k}{dt} = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] c_k$$

$$1303 = \frac{1}{8} \left(\sum_{s'} g^{(t)}(X_{k,+,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(+,t)}(X_{k,+,s'}) - g^{(t)}(X_{k,-,+}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(+,t)}(X_{k,-,+}) \right)$$

$$1304 = \frac{1}{8} g^{(t)}(X_{k,+,+}). \quad (\text{by Corollary C.5})$$

1311 Similarly, if $H_-^{(t)}(X_{k,+,+}) < -b$ for $k \in [2]$, then

$$1312 \frac{dw_-^{(t)\top} c_k}{dt} = -\mathbb{E} \left[g^{(t)} y \dot{\sigma}_-^{(t)} \sum_{l=1}^L \left(X p_l^{(-,t)} \right)^\top \right] c_k$$

$$1313 = -\frac{1}{8} \left(-\sum_{s'} g^{(t)}(X_{k,-,s'}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(-,t)}(X_{k,-,s'}) + g^{(t)}(X_{k,+,-}) \sum_{l=1}^L p_{q \leftarrow l, k \leftarrow c_k}^{(-,t)}(X_{k,+,-}) \right)$$

$$1314 = \frac{1}{8} g^{(t)}(X_{k,-,-}). \quad (\text{by Corollary C.5})$$

□

1324 D.3 PROPERTIES AT THE END OF STAGE 1

1325 **Corollary D.6** (Activation of different types of samples). *For $t \in [T_{0.5}, T_1]$ and $s \in \{\pm 1\}$, $k \in [K]$, only $\dot{\sigma}_s^{(t)}(X_{k,-s,-s}) = 0$ and all the rest of cases equal 1.*

1329 *Proof.* First of all, by the neuron signal alignment lemma stage 1.2 (Lemma D.4 and Lemma D.5), we have

$$1330 \frac{dw_+^{(t)\top} (v_1 + v_2)}{dt} > \frac{dw_+^{(t)\top} c_1}{dt}.$$

1331 Since the attention is uniform during stage 1, by Corollary D.3, we continue to have $\dot{\sigma}_+(X_{k,-,-}) = 0$ in stage 1.2. □

1332 **Corollary D.7** (Neuron alignments at the end of stage 1). *There exists a time $T_1 = \Theta(1)$ such that $w_+^{(T_1)\top} \mu = \Theta(1)$ for all $\mu \in \mathcal{V} \cup \mathcal{C}$.*

1333 *Proof.* This is a direct consequence of Lemma D.4 and Lemma D.5. □

1342 **As a result of stage 1, we can show that the transformer model now can produce a positive margin on the consistent samples but it fails to classify the conflicting samples.**

1343 **Theorem D.8** (Stage 1). *Under Definition 5.1, we have $f^{(T_1)}(X_{k,s,s}) = s \cdot \Omega(1)$ and $f^{(T_1)}(X_{k,s,-s}) = 0$.*

1344 *Proof.* Since the attention during stage 1 is uniform, this is a direct consequence of Lemma D.1, Lemma D.4, Lemma D.5 and the symmetry between two heads in Lemma C.4. □

E STAGE 2: SELF-ATTENTION LEARNS FROM THE CONFLICTING (HARD) SAMPLES

E.1 GROWTH OF SCORES

The following result states that there exists a range where (1) the growth of $c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k$ and $-c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_{k'}$ dominates other score variables; (2) the growth of $v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_k, -v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_{k'}$ dominates other query self-score variables; and (3) the growth of $c_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_k, -c_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_{k'}$ dominates other key self-score variables.

Lemma E.1. *For $t \geq T_1$, there exists a constant $C > 0$ such that as long as $\max_{\mu, \nu \in \mathcal{V} \cup \mathcal{C}} |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \leq C$, then there exist a constant $0 < c < 1$ such that*

$$c \cdot \min_{k \neq k' \in [K]} \left(c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k, -c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_{k'} \right) \geq \max \left(\left(\bigcup_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}} \left\{ |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \right\} \right) \setminus \left(\bigcup_{k \neq k' \in [K]} \left\{ |c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k|, |c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_{k'}| \right\} \right) \right) \quad (11)$$

$$c \cdot \min_{k \neq k' \in [K]} \left(v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_k, -v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_{k'} \right) \geq \max \left(\bigcup_{k \neq k' \in [K]} \left\{ |v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_k|, |v_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_{k'}|, |c_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_{k'}| \right\} \right) \quad (12)$$

$$c \cdot \min_{k \neq k' \in [K]} \left(c_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_k, -c_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_{k'} \right) \geq \max \left(\bigcup_{k \neq k' \in [K]} \left\{ |v_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_k|, |v_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_{k'}|, |v_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_{k'}| \right\} \right) \quad (13)$$

and

$$\|W_{Q+}^{(t)} v_k\|_2^2 \geq \|W_{Q+}^{(t)} c_k\|_2^2, \quad \|W_{K+}^{(t)} c_k\|_2^2 \geq \|W_{K+}^{(t)} v_k\|_2^2. \quad (14)$$

Proof. The proof is by inspecting the gradient flow updates in Appendix E.5 and proving by induction. At $t = T_1$, all the scores equal 0 and further, we have

$$\begin{aligned} \left. \frac{dc_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k}{dt} \right|_{t=T_1}, \quad \left. -\frac{dc_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_{k'}}{dt} \right|_{t=T_1} &= \frac{4}{3} g^{(T_1)} \omega^2 m \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+, T_1)} \right) \\ &= \frac{4}{3} g^{(T_1)} \omega^2 \frac{2}{3} w_+^{(T_1)\top} c_1 \end{aligned}$$

On the other hand, we have

$$\begin{aligned} \arg \max \frac{d}{dt} \Big|_{t=T_1} & \left(\left(\bigcup_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}} \left\{ |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \right\} \right) \setminus \left(\bigcup_{k \neq k' \in [K]} \left\{ |c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k|, |c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_{k'}| \right\} \right) \right) \\ &= \left. \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} \right|_{t=T_1} \end{aligned}$$

and

$$\left. \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} \right|_{t=T_1} = \frac{4}{3} g^{(T_1)} \omega^2 w_+^{(T_1)\top} X p_l^{(+, T_1)} = \frac{4}{3} g^{(T_1)} \omega^2 \frac{1}{3} w_+^{(T_1)\top} c_1$$

Thus, Equation (11) is satisfied at $t = T_1$.

Next, we have $\|W_{Q+}^{(T_1)} v_k\|_2^2 = \omega^2$, $v_k^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_{k'} = 0$, $v_k^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_k = 0$, $v_k^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_{k'} = 0$ and $c_k^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_{k'} = 0$. Further,

$$\left. \frac{dv_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_{k'}}{dt} \right|_{t=T_1}, \left. \frac{dv_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_k}{dt} \right|_{t=T_1}, \left. \frac{dv_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_{k'}}{dt} \right|_{t=T_1}, \left. \frac{dc_k^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_{k'}}{dt} \right|_{t=T_1} = 0$$

Hence, Equation (12) is satisfied at $t = T_1$.

Similarly, we have $\|W_{K+}^{(T_1)} c_k\|_2^2 = \omega^2$, $c_k^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_{k'} = 0$, $v_k^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_k = 0$, $v_k^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_{k'} = 0$ and $v_k^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} v_{k'} = 0$. In addition,

$$\left. \frac{dc_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_{k'}}{dt} \right|_{t=T_1}, \left. \frac{dv_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_k}{dt} \right|_{t=T_1}, \left. \frac{dv_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_{k'}}{dt} \right|_{t=T_1}, \left. \frac{dv_k^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_{k'}}{dt} \right|_{t=T_1} = 0$$

Thus, Equation (13) is satisfied at $t = T_1$. Finally, Equation (14) is also satisfied at $t = T_1$.

Now, assume Equation (11), Equation (12) and Equation (13) hold at some $t > T_1$. We are going to show that the gradients at t also satisfy those inequalities. We are going to prove the inequality between $c_k^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_k$ and $v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1$. The proof for the rest of cases is similar by checking the gradient flow dynamical system in Appendix E.5 as they all follow similar structures.

First of all, notice that if we replace $p^{(+,t)}$ with $p^{(+,T_1)}$ in $\frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt}$ and $\frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt}$, it holds that there exists a constant $0 < c' < 1$ such that

$$\begin{aligned} & c' \cdot \frac{1}{3} g^{(t)} 2 \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \right) \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X p_l^{(+,T_1)} \right) \\ & + \frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 - c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \right) \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X p_l^{(+,T_1)} \right) \\ & \geq \left(\frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 \right) \left(-w_+^{(t)\top} X p_l^{(+,T_1)} \right) \right. \\ & \left. + \frac{1}{3} g^{(t)} 2 \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 - v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \right) \left(-w_+^{(t)\top} X p_l^{(+,T_1)} \right) \right) \end{aligned}$$

since $w_+^{(t)\top} c_1 - w_+^{(t)\top} X p_l^{(+,T_1)} = \frac{2}{3} w_+^{(t)\top} c_1$ and $w_+^{(t)\top} X p_l^{(+,T_1)} = \frac{1}{3} w_+^{(t)\top} c_1$. Next, we calculate the deviation of replacing $p^{(+,t)}$ with $p^{(+,T_1)}$ in the gradients

$$\begin{aligned} & \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \\ & - \left(\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,T_1)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,T_1)} \right) X^\top c_1 \right] \right. \\ & \left. + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,T_1)} \right) p_l^{(+,T_1)} x_l^\top v_1 \right] \right) \\ & = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} \left(p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) \right. \right. \\ & \quad \left. \left. - p_l^{(+,T_1)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,T_1)} \right) \right) X^\top c_1 \right] \\ & + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \left(\text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} \right. \right. \end{aligned}$$

$$- \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+, T_1)} \right) p_l^{(+, T_1)} x_l^\top v_1 \Big]$$

Thus, there exists a constant $C > 0$ not too large such that $\max_{\mu, \nu \in \mathcal{V} \cup \mathcal{C}} |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \leq C$, and thus $\|p^{(+, t)} - p^{(+, T_1)}\|_\infty = O(1)$ and then

$$\begin{aligned} & \left| \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} - \left(\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+, T_1)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+, T_1)} \right) X^\top c_1 \right] \right. \\ & \quad \left. + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+, T_1)} \right) p_l^{(+, T_1)} x_l^\top v_1 \right] \right) \Big| \\ & \leq c'' \left(\frac{1}{3} g^{(t)} 2 \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \right) \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X p_l^{(+, T_1)} \right) \right. \\ & \quad \left. + \frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 - c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \right) \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X p_l^{(+, T_1)} \right) \right) \end{aligned}$$

for some sufficiently small constant $c'' > 0$. This implies that if c satisfies $c' + \Theta(1) < c < 1$ then $c \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \geq \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt}$. \square

Based on Lemma E.1, we can show that there exists a time range where the dynamical system exhibits exponential growth.

Lemma E.2 (Score growth lemma). *For $t \geq T_1$, there exists a constant $C > 0$ such that as long as $\max_{\mu, \nu \in \mathcal{V} \cup \mathcal{C}} |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \leq C$, then there exists a constant $b > 0$ such that*

$$\frac{d}{dt} \begin{bmatrix} c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \\ c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \\ v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \end{bmatrix} \geq b \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \\ c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \\ v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \end{bmatrix}$$

Proof. First, as long as $\|p_l^{(+, t)}(X) - p_l^{(+, T_1)}(X)\|_\infty = O(1)$ for all X and l , we have $g^{(t)}(X) \geq \Omega(1)$. Now, applying Lemma E.1 to the update of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ in Appendix E.5.1, we can derive that there exists a constant $b_1 > 0$ such that

$$\frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \geq b_1 \left(c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \right)$$

and similarly, there exists b_2, b_3 such that

$$\begin{aligned} \frac{dc_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1}{dt} & \geq b_2 \cdot c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \\ \frac{dv_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} & \geq b_3 \cdot c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 \end{aligned}$$

Take $b = \min(b_1, b_2, b_3)$ we finish the proof. \square

Lemma E.3. *There exists a T_2 such that $T_2 - T_1 = O(\log \frac{1}{\omega^2})$ and*

$$c_1^\top W_{K+}^{(T_2)\top} W_{Q+}^{(T_2)} v_1, -c_1^\top W_{K+}^{(T_2)\top} W_{Q+}^{(T_2)} v_2 = \Theta(1).$$

Proof. We can define a process $x(t)$ by

$$x(0) = \begin{bmatrix} 0 \\ \omega^2 \\ \omega^2 \end{bmatrix}, \quad \frac{dx}{dt} = \frac{b}{\sqrt{m}} Ax, \quad A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

and by Lemma E.2, we have

$$x(t) \leq \begin{bmatrix} c_1^\top W_{K+}^{(T_1+t)\top} W_{Q+}^{(T_1+t)} v_1 \\ c_1^\top W_{K+}^{(T_1+t)\top} W_{K+}^{(T_1+t)} c_1 \\ v_1^\top W_{Q+}^{(T_1+t)\top} W_{Q+}^{(T_1+t)} v_1 \end{bmatrix}.$$

Now, we analyze $x(t)$. We can compute the eigen-decomposition of A as

$$\det(A - \lambda I) = \det \left(\begin{bmatrix} -\lambda & 1 & 1 \\ 1 & -\lambda & 0 \\ 1 & 0 & -\lambda \end{bmatrix} \right) = -\lambda^3 + 2\lambda = 0 \Rightarrow \lambda = 0, \pm\sqrt{2}.$$

The eigenvectors can be found by finding the null space of $A - \lambda I$:

$$\begin{bmatrix} -\lambda & 1 & 1 \\ 1 & -\lambda & 0 \\ 1 & 0 & -\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Thus, the eigenvector for $\lambda = 0$ is

$$C \cdot \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

and the eigenvectors for the non-zero eigenvalues are

$$C \cdot \begin{bmatrix} \lambda \\ 1 \\ 1 \end{bmatrix}$$

Therefore, plugging in the initial value $x(0)$, we have

$$x(t) = \frac{\omega^2}{2} \left(\exp(\sqrt{2}bt) \begin{bmatrix} \sqrt{2} \\ 1 \\ 1 \end{bmatrix} + \exp(-\sqrt{2}bt) \begin{bmatrix} -\sqrt{2} \\ 1 \\ 1 \end{bmatrix} \right)$$

Hence, there exists $T_2 - T_1 \leq O(\log \frac{1}{\omega^2})$ such that $c_1^\top W_{K+}^{(T_2)\top} W_{Q+}^{(T_2)} v_1 \geq x(T_2 - T_1) \geq \Omega(1)$. \square

E.2 GROWTH OF ASYMMETRY

The following result shows that there exists a near-symmetry between the score and self-score variables.

Lemma E.4. For $t \geq T_1$, there exists a constant $C > 0$ such that as long as $\max_{\mu, \nu \in \mathcal{V}_{UC}} |\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu| \leq C$, if we define

$\Delta^{(t)} :=$

$$\max_{\mu \in \mathcal{V}_{UC}} \left\{ \left| (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu \right|, \left| (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu \right|, \left| \mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (c_1 + c_2) \right|, \left| \mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2) \right|, \left| (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu \right|, \left| (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu \right|, \left| (c_1 + c_2)^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu \right|, \left| (v_1 + v_2)^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu \right| \right\}$$

then

$$\frac{\frac{d\Delta^{(t)}}{dt}}{\frac{dc_1^\top W_{K+}^{(T')}^\top W_{Q+}^{(T')} v_1}{dt}} \leq \tilde{O}(\omega).$$

and

$$\Delta^{(t)} \leq \tilde{O}(\omega)$$

1566 *Proof.* First of all, by the symmetry between two groups in Lemma C.2, we have

$$\begin{aligned}
1567 & (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 = c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2) \\
1568 & (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 = c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (c_1 + c_2) \\
1569 & (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 = v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2) \\
1570 & (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 = v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (c_1 + c_2)
\end{aligned}$$

1574 Thus,

$$\begin{aligned}
1575 & \Delta^{(t)} = \max_{\mu \in \mathcal{V} \cup \mathcal{C}} \left\{ (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu, (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu, \right. \\
1576 & \left. (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu, (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \mu, (c_1 + c_2)^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu, (v_1 + v_2)^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \mu \right\}
\end{aligned}$$

1581 At $t = T_1$, we have $\Delta^{(T_1)} = \omega^2$ and by the updates in Appendix E.5, we have

$$\left. \frac{d\Delta^{(t)}}{dt} \right|_{t=T_1} = 0$$

1585 Define $\Delta p^{(t)} = \max_{l_1, l_2 \in [L], X \in \mathcal{D}} |p_{l_1, l_2}^{(t)}(X) - \frac{1}{3}|$. For $t \geq T_1$, from the gradients of the score

1586 variables in Appendix E.5 and Appendix E.6, we have

$$\left| \frac{d\Delta^{(t)}}{dt} \right| \leq O\left(\left|\Delta^{(t)} \Delta p^{(t)}\right|\right) \quad (15)$$

1589 This can be seen from the following. Take $\frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt}$ as an example: Define

$$\begin{aligned}
1592 & \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \right|_{p^{(+,t)}=p^{(+,T_1)}} \\
1593 & := \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,T_1)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,T_1)} \right) X^\top c_1 \Big| X_{k,s,-s} \right] \\
1594 & + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,T_1)} \right) p_l^{(+,T_1)} x_l^\top v_1 \Big| X_{k,s,-s} \right]
\end{aligned}$$

1602 Then, following similar steps in deriving Appendix E.5, it is not hard to derive that if we replace

1603 $p^{(+,t)}$ by $p^{(+,T_1)}$ in $\frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt}$, we have

$$\begin{aligned}
1604 & \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \right|_{p^{(+,t)}=p^{(+,T_1)}} \\
1605 & = \frac{1}{3} g^{(t)} 2 \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
1606 & + \frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 - c_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right)
\end{aligned}$$

1612 and thus,

$$\left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2)}{dt} \right|_{p^{(+,t)}=p^{(+,T_1)}=\frac{1}{3}} = 0.$$

1617 Further, by the chain rule, we have

$$\frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2)}{dt} = c_1^\top W_{K+}^{(t)\top} \frac{dW_{Q+}^{(t)} (v_1 + v_2)}{dt} + \frac{dc_1^\top W_{K+}^{(t)\top}}{dt} W_{Q+}^{(t)} (v_1 + v_2)$$

Therefore, expanding $\frac{dW_{Q+}^{(t)}(v_1+v_2)}{dt}$ via Appendix E.6, we have

$$\left| \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)}(v_1+v_2)}{dt} \right| = \left| \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)}(v_1+v_2)}{dt} - \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)}(v_1+v_2)}{dt} \right|_{p^{(+,t)}=p^{(+,T_1)}=\frac{1}{3}}$$

$$\leq O\left(\left|\Delta^{(t)}\Delta p^{(t)}\right|\right)$$

Applying similar analysis to all other asymmetry variables, we get Equation (15).

Now, we analyze the growth of $\Delta^{(t)}$. Consider a specific time T' such that $c_1^\top W_{K+}^{(T')\top} W_{Q+}^{(T')} v_1 = \Theta(\omega)$ for the first time and by Lemma E.1, we have $|\Delta p^{(t)}| \leq O(\omega)$ for all $t \leq T'$. Further, by Lemma E.2 and similar analysis in Lemma E.3, we have $T' - T_1 \leq O(\log(1/\omega))$. Thus,

$$\left|\Delta^{(T')}\right| \leq \left|\Delta^{(T_1)}\right| + \int_{T_1}^{T'} \left|\frac{d\Delta^{(t)}}{dt}\right| dt \leq \omega^2 + \tilde{O}(\omega^2).$$

This implies that

$$\frac{\Delta^{(T')}}{c_1^\top W_{K+}^{(T')\top} W_{Q+}^{(T')} v_1} \leq \frac{\tilde{O}(\omega^2)}{\Theta(\omega)} \leq \tilde{O}(\omega),$$

Finally, comparing the gradient of $\Delta^{(t)}$ with the gradient of $c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1$ for $t \in [T', T_2]$ (following similar analysis in Lemma E.1), we have

$$\frac{\frac{d\Delta^{(t)}}{dt}}{\frac{dc_1^\top W_{K+}^{(T')\top} W_{Q+}^{(T')} v_1}{dt}} \leq \tilde{O}(\omega).$$

where the \tilde{O} is independent of t . Notice that this inequality on the ratio of gradients also holds for $t \leq T'$ since $\Delta p^{(t)} \leq O(\omega)$ for $t \leq T'$. This implies that $|\Delta^{(T_2)}| \leq |\Delta^{(T')}| + \tilde{O}(\omega) |c_1^\top W_{K+}^{(T_2)\top} W_{Q+}^{(T_2)} v_1 - c_1^\top W_{K+}^{(T')\top} W_{Q+}^{(T')} v_1| \leq \tilde{O}(\omega)$. \square

E.3 WHEN THE QUERY IS THE GROUP SIGNAL

The main goal of this section is to analyze the group signal queries. To do this, we need to know the sign of the group signal queries. This can be analyzed by induction-like argument over all the scores and self-scores below.

Lemma E.5 (Sign of all the score and self-score variables). *For $t \in [T_1, T_2]$,*

$$\begin{aligned} c_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 &\geq 0, & c_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 &\leq 0 \\ v_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 &\leq 0, & v_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 &\geq 0 \\ c_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 &\geq 0, & c_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 &\leq 0 \\ v_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 &\leq 0, & v_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 &\geq 0 \\ c_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 &\geq 0, & c_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 &\leq 0 \\ c_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 &\geq 0, & c_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 &\leq 0 \\ v_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 &\geq 0, & v_1 W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 &\leq 0 \\ c_1 W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 &\leq 0, & c_1 W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 &\geq 0 \\ c_1 W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 &\geq 0, & c_1 W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 &\leq 0 \\ v_1 W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 &\geq 0, & v_1 W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 &\leq 0 \end{aligned}$$

Proof. First of all, at $t = T_1$, all the variables are zero and thus the results are satisfied. Now, assume all the inequalities hold for some $t \geq T_1$, we show that the gradients of those score variables

also satisfy those inequalities. We first prove the inequalities in the first column. First of all, by Lemma E.2, we have

$$\frac{dc_1 W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt}, \frac{dc_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1}{dt}, \frac{dv_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \geq 0.$$

We now prove $v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 \leq 0$ as an example and the signs of all the other score variables can be analyzed similarly. By Appendix E.7, we have

$$\begin{aligned} \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} &= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\ &\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,-,+} \right] \\ &\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \end{aligned}$$

We first analyze the sign of the sum of the first two terms.

$$\begin{aligned} &\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\ &\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,-,+} \right] \\ &= g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) \left(p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 w_+^{(t)\top} v_1 \left(-p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\ &\quad \left. + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,+,-}) w_+^{(t)\top} v_1 + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) w_+^{(t)\top} v_1 \right) \\ &+ g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\ &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \cdot w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\ &\quad \left. + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) w_+^{(t)\top} (-v_1) - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) w_+^{(t)\top} v_1 \right) \end{aligned}$$

Define

$$\Delta_s^{(t)} := \max_{\mu \in \mathcal{VUC}} \left\{ \left| (c_1 + c_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu \right|, \left| (v_1 + v_2)^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \mu \right|, \right.$$

$$\left| \mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (c_1 + c_2) \right|, \left| \mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (v_1 + v_2) \right| \Big\}$$

Notice that

$$\begin{aligned} & p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)}(X_{1, +, -}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{2, +, -}) = O(\Delta s^{(t)}) \\ & p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)} w_+^{(t)\top} X_{1, +, -} - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)} w_+^{(t)\top} X_{1, +, -} = p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, t)}(X_{1, +, -})(1 - p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, t)}(X_{1, +, -}))/2 + O(\Delta s^{(t)}) \\ & p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)} w_+^{(t)\top} X_{1, -, +} - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)} w_+^{(t)\top} X_{1, -, +} = p_{q \leftarrow -v_1, k \leftarrow c_1}^{(+, t)}(X_{1, -, +})(1 - p_{q \leftarrow -v_1, k \leftarrow c_1}^{(+, t)}(X_{1, -, +}))/2 + O(\Delta s^{(t)}) \end{aligned}$$

Now consider the function $y(x) = (1-x)x/2$. Notice that if $x_1 < \frac{1}{3}$ and $\frac{1}{3} < x_2 < \frac{2}{3}$, then $y(x_2) > y(x_1)$. Therefore, there exists a constant $C > 0$ such that if $|c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1| \leq C$, we have

$$p_{q \leftarrow v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, +, -})(1 - p_{q \leftarrow v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, +, -}))/2 > p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +})(1 - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +}))/2$$

Next,

$$\begin{aligned} & p_{q \leftarrow v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, +, -})(1 - p_{q \leftarrow v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, +, -}))/2 - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +})(1 - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +}))/2 \\ & = \Theta(c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1) > O(\Delta s^{(t)}) \end{aligned}$$

where the inequality is by Lemma E.4. Thus,

$$\begin{aligned} & c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{1, +, -} - p_{q \leftarrow v_1}^{(+, t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{2, +, -} - p_{q \leftarrow -v_1}^{(+, t)} \right) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)}(X_{1, +, -}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{2, +, -}) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{1, -, +} - p_{q \leftarrow -v_1}^{(+, t)} \right) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{2, -, +} - p_{q \leftarrow v_1}^{(+, t)} \right) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)}(X_{2, -, +}) \right) \\ & \leq 0 \end{aligned}$$

Similarly,

$$\begin{aligned} & c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{1, +, -} - p_{q \leftarrow -v_2}^{(+, t)} \right) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{2, +, -} - p_{q \leftarrow v_2}^{(+, t)} \right) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 w_+^{(t)\top} v_1 \left(-p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+, t)}(X_{1, +, -}) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+, t)}(X_{2, +, -}) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{1, -, +} - p_{q \leftarrow v_2}^{(+, t)} \right) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+, t)} \left(-w_+^{(t)\top} X_{2, -, +} - p_{q \leftarrow -v_2}^{(+, t)} \right) \right) \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \cdot w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +}) - p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+, t)}(X_{2, -, +}) \right) \\ & \leq 0 \end{aligned}$$

Finally, since $v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 \leq 0$, $v_2^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 \geq 0$,

$$\begin{aligned} & c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, t)}(X_{1, +, -}) w_+^{(t)\top} v_1 + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+, t)}(X_{2, +, -}) w_+^{(t)\top} v_1 \\ & + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+, t)}(X_{1, -, +}) w_+^{(t)\top} (-v_1) - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+, t)}(X_{2, -, +}) w_+^{(t)\top} v_1 \\ & \leq 0 \end{aligned}$$

Thus,

$$\frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} \leq 0.$$

The inequalities in the second column can be proved by noticing the symmetric structure of the gradients of score variables in Appendix E.6. Thus, the gradient of the variables in the second column has the exact opposite sign of the corresponding variables in the first column. \square

E.4 PROPERTIES AT THE END OF STAGE 2

Lemma E.6 (Activation in stage 2). For $t \in [T_1, T_2]$,

$$\forall k \in [2], s \in \{\pm 1\} : \sigma_+^{(t)}(X_{k,s,-s}) > 0$$

Proof. This is a direct consequences of Corollary D.7 and Lemma E.4. \square

Theorem E.7 (Attention at the end of stage 2). There exists a $T_2 = O(\log(1/\omega))$ such that

- For samples $X_{1,+,+}$ and $X_{1,-,-}$,
 - when the query token is the group signal c_1 ,

$$\left| p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow c_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow c_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

- when the query token is the classification signal,

$$\left| p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_2, k \leftarrow c_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_2, k \leftarrow v_2}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_2, k \leftarrow v_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_2, k \leftarrow c_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_2, k \leftarrow v_2}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_2, k \leftarrow v_1}^{(+, T_2)}(X_{1,+,+}) - p_{q \leftarrow v_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,-,-}) \right| \leq \tilde{O}(\omega)$$

- For samples $X_{1,+,-}$ and $X_{1,-,+}$,

- when the query is the group signal c_1 ,

$$p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,+,-}) > p_{q \leftarrow c_1, k \leftarrow c_1}^{(+, T_2)}(X_{1,-,+})$$

$$\left| p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,+,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow c_1, k \leftarrow v_2}^{(+, T_2)}(X_{1,-,+}) - p_{q \leftarrow c_1, k \leftarrow v_1}^{(+, T_2)}(X_{1,-,+}) \right| \leq \tilde{O}(\omega)$$

- when the query is the classification signal,

$$p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, t)}(X_{1,+,-}) - p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, t)}(X_{1,-,+}) = \Theta(1)$$

$$p_{q \leftarrow v_2, k \leftarrow c_1}^{(+, t)}(X_{1,+,-}) - p_{q \leftarrow v_2, k \leftarrow c_1}^{(+, t)}(X_{1,-,+}) = \Theta(1)$$

$$\left| p_{q \leftarrow v_1, k \leftarrow v_1}^{(+, t)}(X_{1,+,-}) - p_{q \leftarrow v_1, k \leftarrow v_2}^{(+, t)}(X_{1,+,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_2, k \leftarrow v_1}^{(+, t)}(X_{1,+,-}) - p_{q \leftarrow v_2, k \leftarrow v_2}^{(+, t)}(X_{1,+,-}) \right| \leq \tilde{O}(\omega)$$

$$\left| p_{q \leftarrow v_2, k \leftarrow c_1}^{(+, t)}(X_{1,+,-}) - p_{q \leftarrow v_1, k \leftarrow c_1}^{(+, t)}(X_{1,+,-}) \right| \leq \tilde{O}(\omega)$$

Proof. The proof is a direct consequence of Lemma E.3, Lemma E.4 and Lemma E.5. \square

E.5 THE UPDATES OF THE SOFTMAX ATTENTION AT $t = T_1$

In this section, we give out the explicit updates for the scores and self-scores at $t = T_1$.

The equations in this section are derived by the following:

1. Notice that $p_{l_1, l_2}^{(+, T_1)} = \frac{1}{3}$ for $l_1, l_2 \in [L]$ and thus $w_+^{(T_1)} X_{k, s, -s} p_l^{(+, T_1)} = \frac{1}{3} w_+^{(T_1)\top} c_k$ where $s \in \{\pm 1\}$.
2. Therefore, by Corollary D.7, $w_+^{(T_1)\top} c_k = \Omega(1)$ and thus $\dot{\sigma}_+^{(T_1)}(X_{1, +, -}) = \dot{\sigma}_+^{(T_1)}(X_{1, -, +}) = \dot{\sigma}_+^{(T_1)}(X_{2, +, -}) = \dot{\sigma}_+^{(T_1)}(X_{2, -, +}) = 1$.
3. By symmetry between two groups and two heads in Lemma C.2 and Lemma C.4, we have $w_+^{(T_1)\top} c_1 = w_+^{(T_1)\top} c_2$ and $g^{(T_1)}(X_{1, +, -}) = g^{(T_1)}(X_{1, -, +}) = g^{(T_1)}(X_{2, +, -}) = g^{(T_1)}(X_{2, -, +})$. Thus, we denote $g^{(T_1)} = g^{(T_1)}(X_{1, +, -})$.

E.5.1 THE SCORES

$$\begin{aligned}
& \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \right|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+, t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+, t)} \right) X^\top c_1 \left| X_{k, s, -s} \right. \right] \Big|_{t=T_1} \\
&+ \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+, t)} \right) p_l^{(+, t)} x_l^\top v_1 \left| X_{k, s, -s} \right. \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+, T_1)} \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+, T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
& \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2}{dt} \right|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+, T_1)} \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 + c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+, T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 + c_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
& \left. \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} \right|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+, t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+, t)} \right) X^\top c_1 \left| X_{k, s, -s} \right. \right] \Big|_{t=T_1}
\end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K_+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \left| X_{k,s,-s} \right. \right] \Bigg|_{t=T_1} \\
& = \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_1 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& = \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_1 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \frac{dc_1^\top W_{K_+}^{(t)\top} W_{Q_+}^{(t)} c_2}{dt} \Bigg|_{t=T_1} \\
& = \frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_2 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_1 \right) \left(w_+^{(T_1)} c_2 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 - c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& = \frac{1}{3} g^{(t)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_2 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} v_1 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_2 \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 - c_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \frac{dv_1^\top W_{K_+}^{(t)\top} W_{Q_+}^{(t)} c_1}{dt} \Bigg|_{t=T_1} \\
& = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q_+}^{(t)\top} q_{+l} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \left| X_{k,s,-s} \right. \right] \Bigg|_{t=T_1} \\
& \quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K_+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \left| X_{k,s,-s} \right. \right] \Bigg|_{t=T_1} \\
& = \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_1 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - v_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& = \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_1 - c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - v_1^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \frac{dv_2^\top W_{K_+}^{(t)\top} W_{Q_+}^{(t)} c_1}{dt} \Bigg|_{t=T_1} \\
& = \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_1 + c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - v_2^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
& = \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_1 + c_1^\top W_{Q_+}^{(T_1)\top} W_{Q_+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \quad + \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_1 - v_2^\top W_{K_+}^{(T_1)\top} W_{K_+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right)
\end{aligned}$$

$$\begin{aligned}
& \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \Big|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&+ \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
& \frac{dv_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2}{dt} \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 + v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 + v_1^\top W_{K+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right)
\end{aligned}$$

E.5.2 THE KEY SELF-SCORES

$$\begin{aligned}
& \frac{dv_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2}{dt} \Big|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_2 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&+ \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_2^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) w_+^{(T_1)\top} X p_l^{(+,T_1)} \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-v_2^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 + v_2^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) w_+^{(T_1)\top} X p_l^{(+,T_1)} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \frac{1}{3} w_+^{(T_1)\top} c_1 \\
&+ \frac{1}{3} g^{(T_1)} 2 \left(-v_2^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 + v_2^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \frac{1}{3} w_+^{(T_1)\top} c_1 \\
& \frac{dv_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1}{dt} \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 4 \left(v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_1 - v_1^\top W_{K+}^{(T_1)\top} W_{Q+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{3} g^{(T_1)} 4 \left(v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_1 - v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
&\frac{dc_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_2}{dt} \Big|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top c_2 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_2^\top W_{K^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top c_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 \right) \left(w_+^{(T_1)\top} c_2 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_2 \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&\frac{dc_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_1}{dt} \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 4 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 4 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&\frac{dc_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_1}{dt} \Big|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top c_1 \Big| X_{k,s,-s} \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_1 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_1 - c_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} X c_1 \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 - v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&\frac{dc_2^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_1}{dt} \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_1 - c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 - v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 \right) \left(w_+^{(T_1)\top} c_2 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{3}g^{(T_1)}2 \left(c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_1 - c_2^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} c_2 \right) \left(-\frac{1}{3}w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_2 - v_1^\top W_{K^+}^{(T_1)\top} W_{Q^+}^{(T_1)} v_1 \right) \left(\frac{2}{3}w_+^{(T_1)\top} c_2 \right)
\end{aligned}$$

E.5.3 THE QUERY SELF-SCORES

$$\begin{aligned}
&\left. \frac{dv_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} v_2}{dt} \right|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_2 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_2^\top W_{Q^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&= \frac{1}{3}g^{(T_1)}2 \left(-v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 + v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(v_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3}g^{(T_1)}2 \left(-v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 + v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(\frac{2}{3}w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(v_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(\frac{2}{3}w_+^{(T_1)\top} c_1 \right) \\
&\left. \frac{dv_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} v_1}{dt} \right|_{t=T_1} \\
&= \frac{1}{3}g^{(T_1)}2 \left(v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3}g^{(T_1)}2 \left(v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(\frac{2}{3}w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_1 - v_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} c_2 \right) \left(\frac{2}{3}w_+^{(T_1)\top} c_1 \right) \\
&\left. \frac{dc_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} c_2}{dt} \right|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_2 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_2^\top W_{Q^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&= \frac{1}{3}g^{(T_1)}2 \left(c_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_2 - c_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_1 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(c_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_1 - c_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3}g^{(T_1)}2 \left(c_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_2 - c_1^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_1 \right) \left(-\frac{1}{3}w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3}g^{(T_1)}2 \left(c_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_1 - c_2^\top W_{Q^+}^{(T_1)\top} W_{K^+}^{(T_1)} v_2 \right) \left(-\frac{1}{3}w_+^{(T_1)\top} c_1 \right)
\end{aligned}$$

$$\begin{aligned}
& \left. \frac{dc_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} \right|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 + c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(-c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 + c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \left. \frac{dc_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2}{dt} \right|_{t=T_1} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_2 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_2^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \left| X_{k,s,-s} \right. \right] \Big|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T-1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_2^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - v_2^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right) \\
& \left. \frac{dc_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \right|_{t=T_1} \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(w_+^{(T_1)\top} c_1 - w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-w_+^{(T_1)\top} X p_l^{(+,T_1)} \right) \\
&= \frac{1}{3} g^{(T_1)} 2 \left(c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_1 - c_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} c_2 \right) \left(\frac{2}{3} w_+^{(T_1)\top} c_1 \right) \\
&\quad + \frac{1}{3} g^{(T_1)} 2 \left(v_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_1 - v_1^\top W_{Q+}^{(T_1)\top} W_{K+}^{(T_1)} v_2 \right) \left(-\frac{1}{3} w_+^{(T_1)\top} c_1 \right)
\end{aligned}$$

E.6 SUM OF PARTIAL GRADIENTS

$$\begin{aligned}
& \frac{dW_{Q+}^{(t)} v_1}{dt} + \frac{dW_{Q+}^{(t)} v_2}{dt} : \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L K_+^{(t)} \text{diag} \left(X^\top w_+^{(t)} - w_+^{(t)\top} X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \left| X_{1,+,-} \right. \right] \\
& \quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L K_+^{(t)} \text{diag} \left(X^\top w_+^{(t)} - w_+^{(t)\top} X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_2 \left| X_{2,+,-} \right. \right]
\end{aligned}$$

2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213

$$\begin{aligned}
&= g^{(t)} \left(W_{K^+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \right. \\
&\quad + W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) p_{q \leftarrow v_2, k \leftarrow v_2}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) p_{q \leftarrow v_2, k \leftarrow c_2}^{(+,t)} \\
&\quad \left. + W_{K^+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \right)
\end{aligned}$$

$$\frac{dW_{Q^+}^{(t)}(c_1 + c_2)}{dt} :$$

$$\begin{aligned}
&\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top (c_1 + c_2) \middle| X_{k,+,-} \right] \\
&= g^{(t)} \left(W_{K^+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \right. \\
&\quad + W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) p_{q \leftarrow c_2, k \leftarrow v_2}^{(+,t)} \\
&\quad + W_{K^+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) p_{q \leftarrow c_2, k \leftarrow c_2}^{(+,t)} \\
&\quad \left. + W_{K^+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \right)
\end{aligned}$$

$$\frac{dW_{K^+}^{(t)} v_1}{dt} + \frac{dW_{K^+}^{(t)} v_2}{dt} :$$

$$\begin{aligned}
&\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{1,+,-} \right] \\
&\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_2 \middle| X_{2,+,-} \right] \\
&= g^{(t)} \left(W_{Q^+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
&\quad + W_{Q^+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
&\quad + W_{Q^+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
&\quad + W_{Q^+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow v_2}^{(+,t)} \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \\
&\quad + W_{Q^+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_2}^{(+,t)} \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
&\quad \left. + W_{Q^+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow v_2}^{(+,t)} \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right)
\end{aligned}$$

2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267

$$\begin{aligned}
& \frac{dW_{K+}^{(t)}(c_1 + c_2)}{dt} : \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top (c_1 + c_2) \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(W_{Q+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
& \quad + W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad + W_{Q+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow c_2}^{(+,t)} \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad + W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow c_2}^{(+,t)} \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. + W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow c_2}^{(+,t)} \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right)
\end{aligned}$$

E.7 EXPANDING THE UPDATES

To help us analyze the sign of the scores and self-scores in Lemma E.5, we rearrange the gradient flow updates for those variables into forms that are convenient for such analysis.

E.7.1 THE KEY SELF-SCORE

$$\begin{aligned}
& \frac{dc_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1}{dt} = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,s,-s} \right] \\
& \quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top c_1 \middle| X_{k,s,-s} \right] \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
& \quad + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\
& \quad \left. + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+} - p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+} - p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\
& + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) w_+^{(t)\top} v_1 \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,+}, -) - p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+}, -) \right) \\
& + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+} - p_{q \leftarrow -c_1}^{(+,t)} \right) - c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow -c_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+} - p_{q \leftarrow -c_2}^{(+,t)} \right) \\
& + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,+}, -) w_+^{(t)\top} v_1 + c_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow -c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+}, -) w_+^{(t)\top} v_1 \Big) \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \Big| X_{k,-,+} \right] \\
= & g^{(t)} \left(c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_1}^{(+,t)} \right) \right. \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -c_1}^{(+,t)} \right) \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& - c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& - c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow -c_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -c_2}^{(+,t)} \right) \\
& \left. - c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \\
= & g^{(t)} \left(c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-}, +) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,-}, +) \right) \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-}, +) - p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-}, +) \right) \\
& + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -c_1}^{(+,t)} \right) - c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow -c_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-} + p_{q \leftarrow -c_2}^{(+,t)} \right) \\
& \left. + c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 w_+^{(t)\top} (-v_1) p_{q \leftarrow -c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-}, +) - c_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 w_+^{(t)\top} v_1 p_{q \leftarrow -c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-}, +) \right) \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top c_1 \Big| X_{k,s,-s} \right] \\
= & g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 p_{q \leftarrow -v_1, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+} - p_{q \leftarrow -v_1}^{(+,t)} \right) \right. \\
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+} - p_{q \leftarrow -c_1}^{(+,t)} \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+} - p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_1}^{(+,t)} \right) \\
& - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow -c_1, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -c_1}^{(+,t)} \right) \\
& \left. - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow -v_2, k \leftarrow -c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-} + p_{q \leftarrow -v_2}^{(+,t)} \right) \right)
\end{aligned}$$

2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334
2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375

$$\begin{aligned}
& \frac{dv_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1}{dt} = 2\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,s,-s} \right] \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 \left(p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) \left(p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} (-v_2) w_+^{(t)\top} v_1 \left(p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) - v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) w_+^{(t)\top} v_1 + v_1^\top W_{K+}^{(t)} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) w_+^{(t)\top} v_1 \right) \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^\top X - w_+^\top X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,-,+} \right] \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) \right. \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\
& \quad \left. + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} v_2 w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1 w_+^{(t)\top} (-v_1) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - v_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_2 w_+^{(t)\top} v_1 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+})
\end{aligned}$$

E.7.2 THE QUERY SELF-SCORE

$$\begin{aligned}
& \frac{dc_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1}{dt} \\
& = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \middle| X_{k,s,-s} \right] \\
& \quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \right. \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+,t)} \\
& \quad - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow v_2}^{(+,t)} \\
& \quad - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow c_2}^{(+,t)} \\
& \quad \left. - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \right) \\
& = g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \right) \right. \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow v_2}^{(+,t)} \right) \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow v_2}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \\
& \quad \left. - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow c_2}^{(+,t)} \right) \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \middle| X_{k,-,+} \right] \\
& = g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \right)
\end{aligned}$$

$$\begin{aligned}
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_1}^{(+,t)} \\
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} \\
& - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} \\
& - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \\
& - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} \\
& = g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(\left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} \right) \right. \\
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 \left(\left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} \right) \\
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} (X_{1,-,+}) - p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} (X_{2,-,+}) \right) \\
& + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_1}^{(+,t)} \\
& \left. - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \right) \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^\top \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
& = g^{(t)} \left(v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow v_1}^{(+,t)} \right. \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow c_1}^{(+,t)} \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow -v_2}^{(+,t)} \\
& - v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow -v_1}^{(+,t)} \\
& - v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow c_1}^{(+,t)} \\
& \left. - v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow v_2}^{(+,t)} \right) \\
& = g^{(t)} \left(v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow -v_1}^{(+,t)} \right) \right. \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q\leftarrow c_1, k\leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q\leftarrow c_1, k\leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \right) \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow v_2}^{(+,t)} \right) \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q\leftarrow c_1, k\leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q\leftarrow c_1, k\leftarrow v_2}^{(+,t)} (X_{1,-,+}) \right) \\
& + v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow c_1}^{(+,t)} \\
& \left. - v_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow c_1}^{(+,t)} \right) p_{q\leftarrow c_1, k\leftarrow c_1}^{(+,t)} \right) \\
& \frac{dc_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} = 2\mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^\top \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right]
\end{aligned}$$

2484
 2485
 2486
 2487
 2488
 2489
 2490
 2491
 2492
 2493
 2494
 2495
 2496
 2497
 2498
 2499
 2500
 2501
 2502
 2503
 2504
 2505
 2506
 2507
 2508
 2509
 2510
 2511
 2512
 2513
 2514
 2515
 2516
 2517
 2518
 2519
 2520
 2521
 2522
 2523
 2524
 2525
 2526
 2527
 2528
 2529
 2530
 2531
 2532
 2533
 2534
 2535
 2536
 2537

$$\begin{aligned}
 & \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
 &= g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \right. \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} \\
 &\quad - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \\
 &\quad - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
 &\quad \left. - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
 &= g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \right) \right. \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \right) \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(\left(-w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} (X_{1,-,+}) \right) \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
 &\quad \left. - c_1^\top W_{Q+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \right)
 \end{aligned}$$

E.7.3 THE SCORES

$$\begin{aligned}
 \frac{dc_1^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} c_1}{dt} &= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top c_1 \middle| X_{k,s,-s} \right] \\
 &\quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
 &= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top c_1 \middle| X_{k,s,-s} \right] \\
 &= g^{(t)} \left(c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow v_1}^{(+,t)}} \right) \right. \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) \\
 &\quad + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow -v_2}^{(+,t)}} \right) \\
 &\quad - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow (-v_1), k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow -v_1}^{(+,t)}} \right) \\
 &\quad - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) \\
 &\quad \left. - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)} \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow v_2}^{(+,t)}} \right) \right)
 \end{aligned}$$

$$\begin{aligned}
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{K^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
& = g^{(t)} \left(c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \right. \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} \\
& \quad - c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \\
& \quad - c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
& \quad \left. - c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
& = g^{(t)} \left(c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \right) \right. \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \right) \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} (-v_2) \left(\left(-w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} (X_{1,-,+}) \right) \\
& \quad + c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
& \quad \left. - c_1^\top W_{K^+}^{(t)\top} W_{K^+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+p_{q \leftarrow c_1}^{(+,t)}} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \right) \\
& \frac{dv_1^\top W_{K^+}^{(t)\top} W_{Q^+}^{(t)} v_1}{dt} = \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,s,-s} \right] \\
& \quad + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K^+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \middle| X_{k,s,-s} \right] \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{Q^+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow v_1}^{(+,t)}} \right) \right. \\
& \quad + v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow c_1}^{(+,t)}} \right) \\
& \quad + v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-p_{q \leftarrow -v_2}^{(+,t)}} \right) \\
& \quad - v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-p_{q \leftarrow v_2}^{(+,t)}} \right) \\
& \quad - v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-p_{q \leftarrow c_2}^{(+,t)}} \right) \\
& \quad \left. - v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-p_{q \leftarrow -v_1}^{(+,t)}} \right) \right) \\
& = g^{(t)} \left(v_1^\top W_{Q^+}^{(t)\top} W_{Q^+}^{(t)} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-p_{q \leftarrow v_1}^{(+,t)}} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-p_{q \leftarrow -v_1}^{(+,t)}} \right) \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) \left(p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \right) \\
& + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& + \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \right) w_+^{(t)\top} v_1 \\
& + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+ \sum_{l=1}^L v_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,-,+} \right] \\
& = g^{(t)} \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_2) p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} (-v_1) \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) \right) \right. \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) \right) \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& \quad + \left(v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \right) w_+^{(t)\top} (-v_1) \\
& \quad + v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad \left. - v_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \right) \\
& \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+ \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \right. \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} \\
& \quad \left. + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) p_{q \leftarrow v_1, k \leftarrow v_2}^{(+,t)} \right)
\end{aligned}$$

$$\begin{aligned}
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)} v_2 - w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} \\
= & g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} \right) \right. \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)} v_1 \left(p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \left(\left(-w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)} v_2 \left(p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} (X_{2,+,-}) \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)} c_1 - w_+^{(t)\top} X_{1,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_1}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)} c_2 - w_+^{(t)\top} X_{2,+,-} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top v_1 \left| X_{k,-,+} \right. \right] \\
= & g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} \right. \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_1}^{(+,t)} \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} \\
= & g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(\left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} \right) \right. \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q\leftarrow v_1, k\leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q\leftarrow v_1, k\leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \left(\left(-w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q\leftarrow v_1, k\leftarrow v_2}^{(+,t)} (X_{1,-,+}) - p_{q\leftarrow v_1, k\leftarrow -v_2}^{(+,t)} (X_{2,-,+}) \right) \\
& + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_1}^{(+,t)} \\
& -v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_2 \left(w_+^{(t)\top} c_2 - w_+^{(t)\top} X_{2,-,+} p_{q\leftarrow v_1}^{(+,t)} \right) p_{q\leftarrow v_1, k\leftarrow c_2}^{(+,t)} \\
\frac{dv_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1}{dt} = & \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \left| X_{k,s,-} \right. \right]
\end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L v_1^\top W_{K_+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \middle| X_{k,s,-s} \right] \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q_+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,+,-} \right] \\
& = g^{(t)} \left(c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) \right. \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow v_1}^{(+,t)} \right) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow -v_1}^{(+,t)} \right) \right) \right. \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow c_1}^{(+,t)} \right) - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_2) \left(p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,+,-} p_{q \leftarrow -v_2}^{(+,t)} \right) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,+,-} p_{q \leftarrow v_2}^{(+,t)} \right) \right) \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_2 w_+^{(t)\top} v_1 \left(-p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \right) \\
& \quad \left. + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) w_+^{(t)\top} v_1 + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) w_+^{(t)\top} v_1 \right) \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L c_1^\top W_{Q_+}^{(t)\top} q_{+l}^{(t)} p_l^{(+,t)\top} \text{diag} \left(w_+^{(t)\top} X - w_+^{(t)\top} X p_l^{(+,t)} \right) X^\top v_1 \middle| X_{k,-,+} \right] \\
& = g^{(t)} \left(c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_1) p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) \right. \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_2 p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_2) p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \\
& \quad - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \\
& \quad \left. - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} v_1 p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) \right) \\
& = g^{(t)} \left(c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow -v_1}^{(+,t)} \right) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow v_1}^{(+,t)} \right) \right) \right. \\
& \quad + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} (-v_1) w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& \quad \left. + c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow c_1}^{(+,t)} \right) - c_1^\top W_{Q_+}^{(t)\top} W_{Q_+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow c_2}^{(+,t)} \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \left(p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{1,-,+} p_{q \leftarrow v_2}^{(+,t)} \right) + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \left(-w_+^{(t)\top} X_{2,-,+} p_{q \leftarrow -v_2}^{(+,t)} \right) \right) \\
& + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} v_2 \cdot w_+^{(t)\top} (-v_1) \left(p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) - p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right) \\
& + c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_1 p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) w_+^{(t)\top} (-v_1) - c_1^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} c_2 p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) w_+^{(t)\top} v_1 \Big) \\
\mathbb{E} & \left[g^{(t)} y \dot{\sigma}_+ \sum_{l=1}^L v_1^\top W_{K+}^{(t)\top} K_+^{(t)} \text{diag} \left(X^\top w_+ - w_+^\top X p_l^{(+,t)} \right) p_l^{(+,t)} x_l^\top c_1 \Big| X_{k,s,-s} \right] \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 \left(w_+^{(t)\top} v_1 - w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} \right. \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(w_+^{(t)\top} (-v_2) - w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} \\
& \quad - v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_1) \left(w_+^{(t)\top} (-v_1) - w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \\
& \quad - v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} \\
& \quad \left. - v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 \left(w_+^{(t)\top} v_2 - w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
& = g^{(t)} \left(v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 \left(\left(-w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} \right) \right. \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_1 w_+^{(t)\top} v_1 \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \right) \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} (-v_2) \left(\left(-w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} + \left(-w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} \right) \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} v_2 w_+^{(t)\top} v_2 \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} (X_{1,+,-}) - p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} (X_{1,-,+}) \right) \\
& \quad + v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,+} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} (X_{1,+,-}) \\
& \quad \left. - v_1^\top W_{K+}^{(t)\top} W_{K+}^{(t)} c_1 \left(w_+^{(t)\top} c_1 - w_+^{(t)\top} X_{1,-} p_{q \leftarrow c_1}^{(+,t)} \right) p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} (X_{1,-,+}) \right)
\end{aligned}$$

F STAGE 3

In stage 3, we show that training the neuron weights can drive the training loss to be arbitrarily small. This is possible since the attention features now are linearly separable with at least a constant margin. This can be seen by considering the neuron weight $w_+ = c_1 + c_2 + v_1 + v_2$ and $w_- = c_1 + c_2 - v_1 - v_2$. Simple calculation can show that this will make $yf(X) \geq \Omega(1)$ for all $(X, y) \in \mathcal{D}$.

Lemma F.1 (Growth of classification token alignment). *Let $g_+^{(t)} := g^{(t)}(X_{1,+,+})$ and $g_-^{(t)} := g^{(t)}(X_{1,+,-})$. For $t \in [T_2, T_3]$, $k \in [2]$,*

$$\frac{dw_+^{(t)\top} v_k}{dt} = \Theta(g_+^{(t)}) + \tilde{O}(\omega g_-^{(t)}).$$

Proof. First of all, by symmetry, we have $g_+^{(t)} = g^{(t)}(X_{2,+,+}) = g^{(t)}(X_{1,-,-}) = g^{(t)}(X_{2,-,-})$ and $g_-^{(t)} = g^{(t)}(X_{2,+,-}) = g^{(t)}(X_{1,-,+}) = g^{(t)}(X_{2,-,+})$. In stage 3, we have

$$\frac{dw_+^{(t)\top} v_1}{dt}$$

$$\begin{aligned}
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] v_1 \\
&= \frac{1}{8} g_+^{(t)} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) \right. \\
&\quad + \dot{\sigma}_+^{(t)} (X_{1,-,-}) \left(p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) \right) \\
&\quad + p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) \\
&\quad \left. + \dot{\sigma}_+^{(t)} (X_{2,-,-}) \left(p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) \right) \right) \\
&+ \frac{1}{8} g_-^{(t)} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) \right. \\
&\quad + p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \\
&\quad - p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) - p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \\
&\quad \left. - p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) - p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \right)
\end{aligned}$$

By Theorem E.7, we have

$$\begin{aligned}
&p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,+}) \\
&\quad + \dot{\sigma}_+^{(t)} (X_{1,-,-}) \left(p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,-}) \right) \\
&\quad + p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,+,+}) \\
&\quad + \dot{\sigma}_+^{(t)} (X_{2,-,-}) \left(p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,-,-}) \right) \\
&= \Theta(1)
\end{aligned}$$

and by symmetry between two groups Lemma C.2

$$\begin{aligned}
&p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{1,+,-}) \\
&\quad + p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) + p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{1,-,+}) \\
&\quad - p_{q \leftarrow c_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) - p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) - p_{q \leftarrow v_2, k \leftarrow -v_1}^{(+,t)} (X_{2,+,-}) \\
&\quad - p_{q \leftarrow c_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) - p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) - p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} (X_{2,-,+}) \\
&= \tilde{O}(\omega)
\end{aligned}$$

Thus,

$$\frac{dw_+^{(t)\top} v_1}{dt} = \Theta(g_+^{(t)}) + \tilde{O}(\omega g_-^{(t)}).$$

□

Lemma F.2 (Growth of group token alignment). *Let $g_+^{(t)} := g^{(t)}(X_{1,+,+})$ and $g_-^{(t)} := g^{(t)}(X_{1,+,-})$. For $t \in [T_2, T_3]$, $k \in [2]$,*

$$\frac{dw_+^{(t)\top} c_k}{dt} \geq \Omega(g_-^{(t)}) + \tilde{O}(\omega g_+^{(t)})$$

Proof. In stage 3, by Theorem E.7, we have

$$\begin{aligned}
&\frac{dw_+^{(t)\top} c_1}{dt} \\
&= \mathbb{E} \left[g^{(t)} y \dot{\sigma}_+^{(t)} \sum_{l=1}^L \left(X p_l^{(+,t)} \right)^\top \right] c_1
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{8}g_+^{(+,t)} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) \right. \\
&\quad \left. - \dot{\sigma}_+^{(+,t)}(X_{1,-,-}) \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) \right) \right) \\
&+ \frac{1}{8}g_-^{(+,t)} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,-}) + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,-}) + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)}(X_{1,+,-}) \right. \\
&\quad \left. - p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,+}) - p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,+}) - p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)}(X_{1,-,+}) \right) \\
&\geq \Theta(g_-^{(+,t)}) + \tilde{O}(\omega g_+^{(+,t)})
\end{aligned}$$

□

Theorem F.3 (Convergence). *There exists T_3 such that $T_3 - T_2 = O(1/\epsilon)$ and $\mathbb{E}[\ell(yf^{(t)}(X))] \leq \epsilon$.*

Proof. Let $g_+^{(+,t)} := g^{(+,t)}(X_{1,+,+})$ and $g_-^{(+,t)} := g^{(+,t)}(X_{1,+,-})$. For consistent samples $X_{k,s,s}$, we have

$$\begin{aligned}
\frac{df^{(+,t)}(X_{1,+,+})}{dt} &= \frac{dw_+^{(+,t)\top} v_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \right) \\
&\quad + \frac{dw_+^{(+,t)\top} v_2}{dt} \left(p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow v_2}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow v_2}^{(+,t)} \right) \\
&\quad + \frac{dw_+^{(+,t)\top} c_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)} \right) \\
&\quad - \frac{dw_-^{(+,t)\top} v_1}{dt} \dot{\sigma}_-^{(+,t)} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)} \right) \\
&\quad - \frac{dw_-^{(+,t)\top} v_2}{dt} \dot{\sigma}_-^{(+,t)} \left(p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow v_2}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow v_2}^{(+,t)} \right) \\
&\quad - \frac{dw_-^{(+,t)\top} c_1}{dt} \dot{\sigma}_-^{(+,t)} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)} \right)
\end{aligned}$$

By symmetry, we have

$$\begin{aligned}
&- \frac{dw_-^{(+,t)\top} v_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_1}^{(+,t)}(X_{1,+,+}) \right) \\
&- \frac{dw_-^{(+,t)\top} v_2}{dt} \left(p_{q \leftarrow c_1, k \leftarrow v_2}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow v_2}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow v_2}^{(+,t)}(X_{1,+,+}) \right) \\
&- \frac{dw_-^{(+,t)\top} c_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) + p_{q \leftarrow v_2, k \leftarrow c_1}^{(+,t)}(X_{1,+,+}) \right) \\
&= \frac{dw_+^{(+,t)\top} v_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow -v_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_1}^{(+,t)}(X_{1,-,-}) \right) \\
&+ \frac{dw_+^{(+,t)\top} v_2}{dt} \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_1, k \leftarrow -v_2}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_2, k \leftarrow -v_2}^{(+,t)}(X_{1,-,-}) \right) \\
&- \frac{dw_+^{(+,t)\top} c_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_1, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) + p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+,t)}(X_{1,-,-}) \right)
\end{aligned}$$

Thus,

$$\begin{aligned}
\frac{df^{(+,t)}(X_{1,+,+})}{dt} &= \Omega \left(\frac{dw_+^{(+,t)\top} v_1}{dt} + \frac{dw_+^{(+,t)\top} v_2}{dt} \right) + \tilde{O} \left(\omega \frac{dw_-^{(+,t)\top} c_1}{dt} \right) \\
&= \Omega(g_+^{(+,t)}) + \tilde{O}(\omega g_-^{(+,t)}) \quad (\text{by Lemma F.1})
\end{aligned}$$

For conflicting samples $X_{k,s,-s}$, we have

$$\begin{aligned}
\frac{df^{(t)}(X_{1,+,-})}{dt} &= \frac{dw_+^{(t)\top} v_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow v_1}^{(+,t)} + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(+,t)} \right) \\
&\quad - \frac{dw_+^{(t)\top} v_2}{dt} \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow -v_2}^{(+,t)} + p_{q \leftarrow -v_2, k \leftarrow -v_2}^{(+,t)} \right) \\
&\quad + \frac{dw_+^{(t)\top} c_1}{dt} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow v_1, k \leftarrow c_1}^{(+,t)} + p_{q \leftarrow -v_2, k \leftarrow c_1}^{(+,t)} \right) \\
&\quad + \frac{dw_-^{(t)\top} v_1}{dt} \dot{\sigma}_-^{(t)} \left(p_{q \leftarrow c_1, k \leftarrow v_1}^{(-,t)} + p_{q \leftarrow v_1, k \leftarrow v_1}^{(-,t)} + p_{q \leftarrow -v_2, k \leftarrow v_1}^{(-,t)} \right) \\
&\quad - \frac{dw_-^{(t)\top} v_2}{dt} \dot{\sigma}_-^{(t)} \left(p_{q \leftarrow c_1, k \leftarrow -v_2}^{(-,t)} + p_{q \leftarrow v_1, k \leftarrow -v_2}^{(-,t)} + p_{q \leftarrow -v_2, k \leftarrow -v_2}^{(-,t)} \right) \\
&\quad - \frac{dw_-^{(t)\top} c_1}{dt} \dot{\sigma}_-^{(t)} \left(p_{q \leftarrow c_1, k \leftarrow c_1}^{(-,t)} + p_{q \leftarrow v_1, k \leftarrow c_1}^{(-,t)} + p_{q \leftarrow -v_2, k \leftarrow c_1}^{(-,t)} \right) \\
&= \Omega \left(\frac{dw_-^{(t)\top} c_1}{dt} \right) + \tilde{O} \left(\omega \frac{dw_+^{(t)\top} v_1}{dt} \right) \\
&= \Omega(g_-^{(t)}) + \tilde{O}(\omega g_+^{(t)}) \tag{by Lemma F.2}
\end{aligned}$$

Thus,

$$\begin{aligned}
\frac{d\mathbb{E}[\ell(yf^{(t)}(X))]}{dt} &= \frac{1}{2} \frac{d\ell(f^{(t)}(X_{1,+,+}))}{dt} + \frac{1}{2} \frac{d\ell(f^{(t)}(X_{1,+,-}))}{dt} \\
&= \frac{1}{2} \ell'(f^{(t)}(X_{1,+,+})) \frac{df^{(t)}(X_{1,+,+})}{dt} + \frac{1}{2} \ell'(f^{(t)}(X_{1,+,-})) \frac{df^{(t)}(X_{1,+,-})}{dt} \\
&= -\Omega \left((g_+^{(t)})^2 + (g_-^{(t)})^2 \right) \\
&= -\Omega \left(\ell^2(f^{(t)}(X_{1,+,+})) + \ell^2(f^{(t)}(X_{1,+,-})) \right)
\end{aligned}$$

which implies that

$$\mathbb{E}[\ell(yf^{(t)}(X))] \leq O \left(\frac{1}{t - T_2 + 1} \right).$$

□

G HANDLING THE FINITE-WIDTH DISCRETIZATION

In this section, we describe how to incorporate the finite-width discretization. As we mentioned in the beginning of Section 5, the gradient flow for the finite width transformer can be seen as the gradient flow for infinite-width transformer model with perturbation. Thus, we need to calculate the deviation of the finite-width gradient flow from the infinite-width gradient flow. Since we only train the neuron weights in stage 1 and stage 3, we only need to consider how the finite-width discretization will affect stage 2.

First, at initialization, by the standard concentration bound, we have the following:

Lemma G.1. *With probability at least $1 - \delta$ over the randomness of the initialization of W_K, W_Q , for any $\mu, \nu \in \mathcal{C} \cup \mathcal{V}$,*

$$\begin{aligned}
\left| \left\langle W_K^{(0)} \mu, W_Q^{(0)} \nu \right\rangle \right| &\leq \omega^2 \left(\sqrt{\frac{4}{m} \log \frac{8}{\delta}} + \frac{4}{m} \log \frac{8}{\delta} \right) \\
\left| \left\langle W_K^{(0)} \mu, W_K^{(0)} \nu \right\rangle \right| &\leq \omega^2 \left(\sqrt{\frac{4}{m} \log \frac{8}{\delta}} + \frac{4}{m} \log \frac{8}{\delta} \right),
\end{aligned}$$

$$\left| \langle W_Q^{(0)} \mu, W_Q^{(0)} \nu \rangle \right| \leq \omega^2 \left(\sqrt{\frac{4}{m} \log \frac{8}{\delta}} + \frac{4}{m} \log \frac{8}{\delta} \right),$$

and for any $\mu \in \mathcal{C} \cup \mathcal{V}$,

$$\begin{aligned} \|W_K^{(0)} \mu\|_2^2 &= \omega^2 \left(1 \pm \left(\sqrt{\frac{4}{m} \log \frac{8}{\delta}} + \frac{4}{m} \log \frac{8}{\delta} \right) \right) \\ \|W_Q^{(0)} \mu\|_2^2 &= \omega^2 \left(1 \pm \left(\sqrt{\frac{4}{m} \log \frac{8}{\delta}} + \frac{4}{m} \log \frac{8}{\delta} \right) \right) \end{aligned}$$

Proof. Since $W_K^{(0)} \mu, W_Q^{(0)} \nu \sim \mathcal{N}(0, \omega^2 I)$, we apply Lemma H.2 to get the corresponding bound. \square

Next, recall that from Section 5.2 the score and self-score variables evolve according to the dynamical system

$$\frac{d}{dt} \begin{bmatrix} \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \end{bmatrix} = A^{(t)} \begin{bmatrix} \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{K+}^{(t)\top} W_{K+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \\ \text{vec}(\{\mu^\top W_{Q+}^{(t)\top} W_{Q+}^{(t)} \nu\}_{\mu, \nu \in \mathcal{C} \cup \mathcal{V}}) \end{bmatrix}$$

for some matrix $A^{(t)}$ with $\|A^{(t)}\|_\infty = O(1)$. By Lemma E.3, stage 2 lasts $O(\log(1/\omega))$ time. Then, we can bound the effect of the perturbation to the dynamical system as follows. Consider the dynamical system

$$\frac{dx(t)}{dt} = A(t)x(t), \quad x(0) = x_0$$

and the perturbed dynamical system

$$\frac{dx_\epsilon(t)}{dt} = (A(t) + \Delta A(t))x_\epsilon(t), \quad x_\epsilon(0) = x_0 + \epsilon$$

where ϵ and $\Delta A(t)$ are the perturbation introduced by the finite-width discretization. Notice that we have $\|\Delta A(s)\| \leq c\|\delta x(s)\|$ for some constant c for all $s \in [T_1, T_2]$. Define $\delta x(t) = x_\epsilon(t) - x(t)$. We have

$$\frac{d\delta x(t)}{dt} = A(t)\delta x(t) + \Delta A(t)x_\epsilon(t)$$

This implies that

$$\begin{aligned} \|\delta x(t)\| &\leq \int_0^t \|A(s)\delta x(s) + \Delta A(s)x_\epsilon(s)\| ds \\ &\leq \int_0^t \|A(s)\| \|\delta x(s)\| + \|\Delta A(s)\| \|x_\epsilon(s)\| ds \\ &\leq \int_0^t \|\delta x(s)\| (\|A(s)\| + c\|x_\epsilon(s)\|) ds \end{aligned}$$

Applying Grönwall's inequality, we have

$$\|\delta x(t)\| \leq \|\delta x(0)\| \exp \left(\int_0^t \|A(s)\| + c\|x_\epsilon(s)\| \right) = \text{poly}(1/\omega) / \sqrt{m}$$

since $\|A(t)\| \leq O(1)$ and $\|x_\epsilon(s)\| \leq O(1)$ and $t \leq O(\log(1/\omega))$. Thus, picking $m \geq \text{poly}(1/\omega)$ can make the perturbation to the gradient flow system at most $O(\omega)$.

3024 H PROBABILITY

3025
3026 **Lemma H.1** (Bernstein's inequality for bounded random variables). *Assume Z_1, \dots, Z_n are n i.i.d.*
3027 *random variables with $\mathbb{E}[Z_i] = 0$ and $|Z_i| \leq M$ for all $i \in [n]$ almost surely. Let $Z = \sum_{i=1}^n Z_i$.*
3028 *Then, for all $t > 0$,*

$$3029 \mathbb{P}[Z > t] \leq \exp\left(-\frac{t^2/2}{\sum_{j=1}^n \mathbb{E}[Z_j^2] + Mt/3}\right) \leq \exp\left(-\min\left\{\frac{t^2}{2\sum_{j=1}^n \mathbb{E}[Z_j^2]}, \frac{t}{2M}\right\}\right)$$

3030
3031 which implies with probability at least $1 - \delta$,

$$3032 Z \leq \sqrt{2\sum_{j=1}^n \mathbb{E}[Z_j^2] \log \frac{1}{\delta}} + 2M \log \frac{1}{\delta}.$$

3033
3034 **Lemma H.2.** *For $w_1, w_2 \in \mathbb{R}^m$ with $w_1, w_2 \stackrel{i.i.d.}{\sim} \mathcal{N}(0, I_m/m)$,*

$$3035 \mathbb{P}\left[\left|\|w_1\|_2^2 - 1\right| \geq \sqrt{\frac{4}{m} \log \frac{2}{\delta}} + \frac{4}{m} \log \frac{2}{\delta}\right] \leq \delta$$

$$3036 \mathbb{P}\left[|\langle w_1, w_2 \rangle| \geq \sqrt{\frac{4}{m} \log \frac{2}{\delta}} + \frac{4}{m} \log \frac{2}{\delta}\right] \leq \delta$$

3037
3038 *Proof.* We have

$$3039 \mathbb{E}[\|w_1\|_2^2] = \mathbb{E}\left[\sum_{i=1}^m w_{1,i}^2\right] = 1$$

3040 Notice that $w_{1,i}^2$ is a sub-Gamma random variable with parameters $(\frac{4}{m^2}, \frac{4}{m})$. Thus, by Bernstein's
3041 inequality,

$$3042 \mathbb{P}\left[\left|\|w_1\|_2^2 - \mathbb{E}[\|w_1\|_2^2]\right| \geq \sqrt{\frac{4}{m} \log \frac{2}{\delta}} + \frac{4}{m} \log \frac{2}{\delta}\right] \leq \delta$$

3043
3044 Next,

$$3045 \mathbb{E}[\langle w_1, w_2 \rangle] = \mathbb{E}\left[\sum_{i=1}^m w_{1,i} w_{2,i}\right] = 0$$

3046 By Bernstein's inequality,

$$3047 \mathbb{P}\left[|\langle w_1, w_2 \rangle| \geq \sqrt{\frac{4}{m} \log \frac{2}{\delta}} + \frac{4}{m} \log \frac{2}{\delta}\right] \leq \delta$$

3048 □