

REVISITING CONVOLUTION ARCHITECTURE IN THE REALM OF DNA FOUNDATION MODELS

Anonymous authors

Paper under double-blind review

ABSTRACT

In recent years, A variety of methods based on Transformer and state space model (SSM) architectures have been proposed, advancing foundational DNA language models. However, there is a lack of comparison between these recent approaches and the classical architecture—convolutional networks (CNNs)—on foundation model benchmarks. This raises the question: *are CNNs truly being surpassed by these recent approaches based on transformer and SSM architectures?* In this paper, we develop a simple yet well-designed CNN-based method, named ConvNova. ConvNova identifies and proposes three effective designs: 1) dilated convolutions, 2) gated convolutions, and 3) a dual-branch framework for gating mechanisms. Through extensive empirical experiments, we demonstrate that ConvNova significantly outperforms recent methods on more than half of the tasks across several foundation model benchmarks. For example, in histone-related tasks, ConvNova surpasses the second-best method by an average of 5.8%, while generally utilizing fewer parameters and enabling faster computation. Additionally, the experiments observed findings that may be related to biological characteristics. This indicates that CNNs are still a strong competitor compared to Transformers and SSMs. We anticipate that this work will spark renewed interest in CNN-based methods for DNA foundation models.




Figure 1: A) The trade-off between model size and accuracy (AUC score) of different methods. ConvNova achieves the current SoTA performance. B) The trade-off between input sequence length and runtime of different methods. ConvNova exhibits clear superiority over HyenaDNA and Caduceus. All models are around 7M parameters, tested on A100 80GB with batch size 1.

1 INTRODUCTION

DNA, being consistently at the heart of scientific research, has brought about significant advancements in various fields, including aging research, synthetic biology, and disease treatment (Chucair-Elliott et al., 2024; Moe-Behrens et al., 2013). Conversely, unlike NLP, comprehending DNA poses a formidable challenge, owing to the myriad undiscovered biological principles underlying its structure and function. Fortunately, the rapid progress of genome language modeling has showcased its

054 dominance in numerous subsequent applications. These include the prediction of promoters (Zhang
055 et al., 2022), gene expression (Avsec et al., 2021), DNA methylation (Jin et al., 2022), analysis of
056 chromatin state (Lee et al., 2022), prediction of promoter-enhancer interactions (Chen et al., 2022),
057 TF-DNA binding prediction (Wang et al., 2022), variant effect prediction (Rozowsky et al., 2023),
058 and gene network prediction (Theodoris et al., 2023).

059 Versatile sequence models have been established as foundational models for genetics, typically falling
060 into two categories: Transformer-based methods (e.g., DNABERT (Ji et al., 2021), DNABERT-
061 2 (Zhou et al.), NucleotideTransformer (Dalla-Torre et al., 2023)) and SSM-inspired methods (e.g.,
062 HyenaDNA (Nguyen et al., 2024), Caduceus (Schiff et al., 2024)). These approaches have made
063 significant progress in genomic language modeling tasks and are now widely adopted. Prior to these
064 developments, however, convolutional neural networks (CNNs) were the dominant technique for
065 DNA modeling in bioinformatics, giving rise to notable models such as Basset (Kelley et al., 2016),
066 Bassenji (Kelley et al., 2018), and LegNet (Penzar et al., 2023). Despite their past success, CNN
067 architectures have not been systematically compared to the newer DNA foundation models in recent
068 research. Comparisons between CNNs and modern approaches have been limited to specific domains,
069 such as NTV2 (Dalla-Torre et al., 2023) and Caduceus (Schiff et al., 2024). In these cases, only a few
070 domain-specific CNN models, like Spliceator (Scalzitti et al., 2021) and some benchmark baselines,
071 have been evaluated.

072 In this landscape, we rethink and reevaluate whether CNN methods are indeed less effective than the
073 current leading paradigms, including Transformer-based methods and SSM-inspired methods, which
074 have been the primary focus of recent research in DNA foundation models. The motivation behind
075 this is straightforward, as CNNs still hold advantages over existing methods: 1) The input lengths
076 for downstream tasks on DNA vary greatly (ranging from tens to thousands). Transformers are not
077 robust to sequence length variations (Press et al., 2021), and their performance can be affected. 2)
078 Transformers have a computational complexity of $O(n^2)$, higher than CNN methods. Additionally,
079 to avoid overly long sequences, transformers also require tokenizers, which affect translational
080 invariance (Zhou et al.). 3) Mamba (Gu and Dao, 2023) is better suited for tasks involving long
081 sequences with autoregressive properties (Yu and Wang, 2024), while DNA sequences typically do
082 not exhibit autoregressive characteristics. 4) CNNs possess a local inductive bias, which can provide
083 better data utilization efficiency and are more friendly towards tasks with small training data volumes.

084 Based on this motivation, we propose a simple yet well-designed CNN method, named ConvNova.
085 Extensive empirical experiments demonstrate that ConvNova outperforms recent Transformer or
086 SSM-inspired methods on more than half of the tasks, indicating that CNNs continue to demon-
087 strate superiority. Through an analysis of the design of ConvNova, we identify and propose three
088 key designs that enable CNNs to achieve superior performance: 1) dilated convolution (Yu and
089 Koltun, 2015), 2) gated convolution (Yu et al., 2019), and 3) a dual-branch framework for the gated
090 convolution. Specifically, we found that for DNA tasks, increasing the receptive field of the CNN
091 through downsampling (U-Net (Ronneberger et al., 2015) style) severely degrades CNN performance.
092 However, dilated convolution can circumvent this issue by enlarging the receptive field without the
093 need for downsampling. Additionally, gated convolution can significantly enhance CNN performance;
094 we hypothesize that this is because DNA may contain a substantial amount of irrelevant segments,
095 and the gating mechanism can suppress this information. Lastly, we discovered that splitting the
096 network pathways into two branches, with one branch exclusively providing gating signals to the
097 other, can improve network performance. For detailed designs, please refer to Section 3 and Section
098 4.

099 Comprehensive experiments demonstrate the superiority of ConvNova. On the NT benchmarks,
100 ConvNova achieves SoTA performance on 12 out of 18 datasets. Notably, in the H3K4me3 task,
101 it outperforms the second-best method by 10.5%, and in the histone-related tasks, it surpasses the
102 second-best method by an average of 5.8%. In the gene finding benchmark (long sequences modeling),
103 ConvNova significantly surpasses HyenaDNA (Nguyen et al., 2024) (55% vs. 35%). Compared to
104 the transformer model with much larger parameters (7.4M vs. 336M), ConvNova still achieves better
105 performance (55% vs. 52%).

106 By adjusting the receptive field size of ConvNova, specifically through controlling the dilation stride,
107 we reveal that the performance of the vast majority of tasks improves with the increase of the receptive
field of the CNN. However, we find that some specific tasks, including H3K4me2, H3K4me3, and
H3K14ac, exhibit the opposite phenomenon. Existing wet lab experimental data indicate that some of

these tasks exhibit pronounced localized characteristics; therefore, this phenomenon may be caused by biological characteristics. For detailed findings and analysis, please refer to Section 5.

In brief, the main contributions of our work can be summarized as follows:

- We reexamine the convolutional paradigm for DNA foundation models. Analyses and extensive experimentation demonstrate the continued competitiveness of convolutional neural networks (CNNs) for downstream tasks. This may prompt the community to reconsider CNNs for such tasks.
- We propose ConvNova and identify three key designs for genomic language modeling: dilated convolution, gated convolution, and a dual-branch design. With these designs, ConvNova achieves superior performance on numerous benchmarks while maintaining fewer parameters.
- ConvNova outperforms recent methods on over half of the tasks across various DNA foundation model benchmarks, typically using fewer parameters and achieving faster performance.

2 PRELIMINARIES AND RELATED WORK

2.1 TRANSFORMER-BASED DNA MODELS

Transformers have become a dominant force across various fields in the deep learning community, and recent research has begun to explore their potential within the realm of genomics. DNABERT (Ji et al., 2021) was the first to establish the transformer as a foundational model for DNA, utilizing k-mer tokenization. Nucleotide Transformer (NT) (Dalla-Torre et al., 2023) scaled up the transformer model (with sizes of 500M and 2.5B) and investigated the impact of different pretraining datasets (HG38¹ and Multispecies²) on 18 downstream tasks. Their experiments demonstrate that multispecies data and larger models generally yield superior performance. DNABERT-2 (Zhou et al.), also pretrained on multispecies data, tackled several problems encountered with DNABERT. They further explored the Byte Pair Encoding (BPE) tokenization and ALiBi positional embedding as effective strategies for modeling DNA with transformer models. Additionally, they utilized FlashAttention to accelerate the processing. Meanwhile, BigBird (Zaheer et al., 2020), despite not being specifically designed, had also been applied in genomics.

While adept at capturing long-range dependencies, transformer-based models in genomics face challenges compared to CNNs. CNNs possess an inductive bias towards local, translation-invariant features (Battaglia et al., 2018), aligning well with many genomic patterns. In contrast, transformers lack this inherent bias for local structure modeling. Additionally, the attention layer’s $O(n^2)$ complexity poses computational challenges, especially for long DNA sequences.

2.2 SSM-INSPIRED DNA MODELS

Another approach that has gained traction involves the use of State Space Models (SSMs). HyenaDNA (Nguyen et al., 2024) is a decoder-only, sequence-to-sequence model that incorporates a modified convolution operator (Hyena (Poli et al., 2023)). This innovative approach enables the expansion of input sequences of the model to 1M-size nucleotides and significantly outperforms transformer-based models in a variety of tasks with a model size that is 300 times smaller. Additionally, Mamba (Gu and Dao, 2023), a state-space-model inspired architecture, has demonstrated promising, albeit preliminary, experiments on DNA sequences. Caduceus (Schiff et al., 2024) refines the Mamba Block into an RC-equivariant architecture. By taking into account the reverse complementary strand, Caduceus surpasses both HyenaDNA and transformer-based models in downstream tasks, marking a significant advancement in the field.

¹https://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.26

²Based on genomes available on https://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.26, for more details can refer to NT’s original paper

2.3 CNN-BASED DNA MODELS

Convolutional Neural Network (CNN)-based DNA models are powerful deep learning tools designed to analyze and interpret DNA sequences by automatically extracting significant motifs and patterns. Early models like DeepBind (Alipanahi et al., 2015) predicted the sequence specificities of DNA and RNA-binding proteins, while DeepSEA (Zhou and Troyanskaya, 2015) used CNNs to predict the effects of noncoding variants. Basset (Kelley et al., 2016) advanced the field by learning the regulatory code of the accessible genome using deep CNNs.

Recent advancements have further enhanced the capabilities of CNN-based genomic models. Enformer (Avsec et al., 2021) integrates convolutional layers with attention mechanisms to predict gene expression from DNA sequences, effectively capturing short- and long-range regulatory interactions. Borzoi (Linder et al., 2023) builds upon the Basenji (Kelley et al., 2018) family by incorporating dilated convolutions and skip connections, improving performance on various genomic prediction tasks. LegNet (Penzar et al., 2023) employs a convolutional architecture for predicting gene expression and single-nucleotide variant effects in regulatory regions, achieving first place in the DREAM 2022 challenge.

These models demonstrate the CNNs’ strengths, capturing local sequence features and motifs crucial for understanding genetic information. However, despite their success, they have not been incorporated into the broader consideration of DNA foundation models.

3 METHOD

In this section, we first present our overall network architecture (§ 3.1), followed by an introduction to the core component of this architecture - the Gated Convolution Block (§ 3.2). Subsequently, we delve into the selection of convolutional network methods, specifically whether to use dilation or down-sampling (§ 4.5). Finally, we discuss our pretraining approach and downstream usage (§ 3.3).

3.1 CONVNOVA

Figure 2 shows the overall illustration of ConvNova. DNA sequences are mapped to the hidden space through a convolution operation, followed by N gated convolution blocks (GCBs), and finally processed through a multilayer perceptron (MLP). Different output heads are used in pretraining and downstream tasks. Each GCB utilizes dilated convolution to increase the receptive field and aggregate the features through a dual-branch structure.

3.2 GATED CONVOLUTION BLOCK

Yu et al. (2019) introduces gated convolution as a solution to a problem inherent to standard convolution, which indiscriminately treats all input pixels as valid. This method enhances partial convolution by offering a learnable dynamic feature selection mechanism for each channel at every spatial location throughout all layers. This mechanism allows for more nuanced and effective feature selection, improving the performance of convolutional neural networks.

With the motivation to effectively retain and forget information, we propose Gated Convolutional Blocks (GCBs). **The dual-branch structure of these blocks is designed to facilitate independent feature extraction, thereby promoting complementary representation learning.** Let’s define $\mathbf{A} \in \mathbb{R}^{l \times d}$ as the feature extracted by the left branch and $\mathbf{B} \in \mathbb{R}^{l \times d}$ as the right branch feature.

We first process the left branch feature \mathbf{A} through a LayerNorm layer and then pass it through a convolution layer, followed by the GELU activation function to obtain the intermediate features \mathbf{h} of the current layer. In the other path, the right branch features \mathbf{B} are processed through other LayerNorm and convolution layers that do not share weight with the left branch, followed by the sigmoid activation function to obtain the intermediate features \mathbf{g} .

Next, we update the left branch features \mathbf{A} by multiplying the intermediate features \mathbf{g} with the intermediate features \mathbf{h} . Simultaneously, the right branch features \mathbf{g} are used to update the right branch features \mathbf{B} themselves. Additionally, the kernel size for all convolutional layers is consistently set to 9. This entire process, which integrates both branches, is described by Eq.1 and is illustrated in

216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269




Figure 2: **ConvNova architecture.** The ConvNova model processes double stands as inputs. The sequences are initially subjected to one-hot encoding and subsequently pass through a convolution layer. The processed data then enters a series of Gated Convolution Blocks (GCBs), the specifics of which are elaborated in section § 3.2. The output from the GCBs is then fed into an MLP. The final stage of the framework bifurcates into two distinct heads: the downstream head and the pretraining head.

the Gated Convolution Block of Figure 2:

$$\begin{aligned}
 \mathbf{h} &= \text{GELU}(\text{convolution}(\mathbf{A})) \\
 \mathbf{g} &= \text{sigmoid}(\text{convolution}(\mathbf{B})) \\
 \mathbf{A} &= \mathbf{A} + \mathbf{h} \odot \mathbf{g} \\
 \mathbf{B} &= \mathbf{B} + \mathbf{g}
 \end{aligned}
 \tag{1}$$

A stage is composed of several GCBs, with the dilation rate for each Block set to $[1, 1, d, d^2, d^3, \dots]$. In different experiment, we use different stages. By default, 1 stage uses 5 GCBs. You can find the information in Table 9.

3.3 MLM PRETRAINING AND DOWNSTREAM USAGE

MLM Pretraining We use the bidirectional masked language model (MLM) pretraining method and have observed significant performance improvement in downstream tasks.

In this work, 10% of the nucleotides in a primary DNA strand are masked, and the model is trained to predict the type of the masked tokens. Therefore no extra labels are required. The objective of this pretraining is to infer the type of nucleotides that have been masked by utilizing the surrounding nucleotides. The optimization objective is to minimize the cross-entropy loss. [The pretraining data is HG38, same as HyenaDNA \(Nguyen et al., 2024\).](#)

Downstream Usage Upon completing the pretraining phase for 400 epochs, we proceed to finetune ConvNova on the downstream tasks, keeping the entire model adjustable. We employ the pooling method of HyenaDNA to get an embedding for each DNA sequence and utilize a linear mapping to the output label dimension to serve as the finetuned head. In cases where it’s not significantly indicated, the setting for our downstream tasks remains consistent with that of HyenaDNA [\(Nguyen et al., 2024\).](#)

Table 1: **NT Benchmark results.** MCC/F1-score is reported for pretrained NTv2, HyenaDNA, DNABERT-2, Caduceus-Ph, and ConvNova. The best values per task are bold, and the second-best are underlined. \pm indicates the error range across five random seeds.

	NTv2 (500M)	HyenaDNA (1.6M)	DNABERT-2 (117M)	Caduceus-Ph (1.9M)	ConvNova (1.7M)
<i>Histone</i>					
H3	78.17 \pm 2.54	78.14 \pm 1.70	79.31 \pm 0.68	<u>80.48</u> \pm 1.04	81.50 \pm 0.80
H3K4me1	51.64 \pm 1.12	44.52 \pm 2.59	48.34 \pm 4.63	<u>52.83</u> \pm 0.96	56.60 \pm 1.01
H3K4me2	37.24 \pm 2.25	42.68 \pm 2.66	43.02 \pm 2.92	<u>49.88</u> \pm 2.65	57.45 \pm 2.27
H3K4me3	50.30 \pm 1.77	50.41 \pm 3.15	45.43 \pm 3.33	<u>56.72</u> \pm 2.58	67.15 \pm 0.93
H3K9ac	61.05 \pm 1.40	58.50 \pm 1.75	60.04 \pm 1.27	<u>63.27</u> \pm 2.29	68.10 \pm 1.91
H3K14ac	57.22 \pm 2.19	56.71 \pm 2.40	54.49 \pm 4.99	<u>60.84</u> \pm 2.94	70.71 \pm 2.32
H3K36me3	60.50 \pm 1.75	59.92 \pm 1.06	57.58 \pm 2.38	<u>61.12</u> \pm 1.44	68.31 \pm 1.19
H3K79me3	65.78 \pm 2.34	66.25 \pm 3.65	64.38 \pm 0.48	<u>67.17</u> \pm 2.03	72.08 \pm 1.23
H4	79.87 \pm 1.34	78.15 \pm 1.58	78.18 \pm 0.98	<u>80.10</u> \pm 1.00	81.12 \pm 0.93
H4ac	55.22 \pm 2.20	54.15 \pm 2.96	51.80 \pm 0.10	<u>59.26</u> \pm 3.67	66.10 \pm 1.20
<i>Regulatory</i>					
Enhancer	54.51 \pm 1.94	53.13 \pm 4.52	52.50 \pm 1.44	<u>55.20</u> \pm 2.56	57.60 \pm 2.52
Enhancer Types	43.36 \pm 1.75	<u>48.16</u> \pm 2.48	44.32 \pm 1.18	47.17 \pm 2.85	49.75 \pm 2.82
Promoter All	96.82 \pm 0.47	95.57 \pm 0.18	96.23 \pm 0.17	<u>96.65</u> \pm 0.16	96.82 \pm 0.22
Promoter non-TATA	97.45 \pm 0.69	95.86 \pm 0.37	<u>97.17</u> \pm 0.17	96.31 \pm 0.50	96.76 \pm 0.21
Promoter TATA	<u>96.53</u> \pm 0.81	95.88 \pm 0.53	96.99 \pm 0.49	96.21 \pm 0.81	96.34 \pm 0.38
<i>Splice sites</i>					
Splice Acceptor	97.99 \pm 0.66	96.98 \pm 0.49	<u>97.49</u> \pm 0.36	94.21 \pm 0.37	96.23 \pm 0.41
Splice Donor	98.50 \pm 0.43	95.27 \pm 1.07	94.33 \pm 0.27	94.69 \pm 0.67	<u>96.62</u> \pm 0.61
Splice All	98.15 \pm 1.01	94.05 \pm 1.08	93.75 \pm 1.25	92.87 \pm 1.73	<u>96.33</u> \pm 0.31

4 EXPERIMENTS

Some other studies have demonstrated that adopting multispecies data for pretraining can enhance downstream task performance. However, in this work, all pretraining tasks are performed on the human reference genome³ to align with the previous work (HyenaDNA and Caduceus). In addition, we employ a tokenization scheme at the character or base pair level, which has been proven to function well in SSM-inspired models. We also show the speed comparison between ConvNova and other models along sequence length (10^3 to 10^6). At a sequence length of 10^6 , ConvNova is 1.35 times faster than its fastest counterpart—HyenaDNA, as shown in Figure 1. Please refer to A.1 for additional details on the pretraining dataset and recipes.

4.1 SHORT RANGE

4.1.1 NUCLEOTIDE TRANSFORMER BENCHMARK

We start the evaluation with the recently proposed Nucleotide Benchmark (Dalla-Torre et al., 2023). These datasets encompass three types of tasks: histone marker prediction, regulatory annotation prediction, and splice site annotation prediction.

To assess performance, we follow the methodology outlined in (Nguyen et al., 2024), applying different metrics depending on the task: Matthews Correlation Coefficient (MCC) for histone marker tasks and various enhancer-related tasks, F1 score for various promoter-related tasks and splice site annotation tasks. The baselines consist of NT-v2(500M), DNABERT-2, similar-sized HyenaDNA, and Caduceus-ph. Because in (Schiff et al., 2024), Caduceus-ph outperforms Caduceus-ps in almost all functions except promoter TATA and H3K36me3, we only choose Caduceus-ph as our baseline.

³https://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.26

Table 2: **Gene Finding results.** MCC-score is reported for ConvNova and baseline models. The best value is in bold.

GeneFinding	NT-H (500M)	DNABERT-2 (117M)	GENA-LM (336M)	HyenaDNA (6.5M)	ConvNova (7.4M)
	0.41	0.43	0.52	0.35	0.55

For HyenaDNA and ConvNova, we adopt the pooling methodology from HyenaDNA (Nguyen et al., 2024) and implement a linear layer to derive the logits.

Furthermore, we recognize the instability inherent in the datasets, where different random seeds may result in variations of up to 5 points. Therefore, we provide the mean and standard deviation (\pm) calculated from 10 random seeds.

The results of this benchmark suite are showcased in Table 1, demonstrating the competitive performance of ConvNova. Notably, ConvNova outperforms Caduceus-ph, the second-best model, by 10 points despite having fewer parameters. Across nearly all tasks, ConvNova exhibits superior performance compared to a similarly sized HyenaDNA model, except for the splice sites acceptor. Additionally, ConvNova surpasses the transformer-based DNABERT-2 in 15 out of 18 tasks and NT-v2 in 12 out of 18 tasks despite having significantly fewer parameters. For more details, see A.2.1.

Furthermore, we have also included previously popular supervised CNNs, LegNet (Penzar et al., 2023) and Basenji (Kelley et al., 2018), in this benchmark. For more details, please refer to A.4.

4.1.2 GENOMIC BENCHMARK

Next, we conduct a comprehensive evaluation using eight datasets introduced by GenomicsBenchmarks (Grešová et al., 2023) with eight regulatory element classification tasks with sequence lengths spanning from 200 to approximately 2000bps. Our baselines comprise HyenaDNA, Caduceus-ph, and the original CNN model described in (Grešová et al., 2023). We conduct experiments using five distinct random seeds and report the mean and the disparity between the maximum/minimum and the mean results.

As shown in Table 3, ConvNova models outperform baselines in 5 tasks. In other tasks, ConvNova is competitive with the best models. Additionally, we also compared the performance of supervised CNNs, LegNet and Basenji, in this benchmark. See A.2.4 and A.4 for additional experiment details.

Table 3: **Genomic Benchmark results.** Top-1 accuracy (\uparrow) is reported for pretrained HyenaDNA, Caduceus-Ph, ConvNova, and the original CNN baseline (trained from scratch). The best values are in bold, and the second-best is underlined. \pm indicates the error range across five random seeds.

Task	CNN (264K)	HyenaDNA (436K)	Caduceus-Ph (470K)	ConvNova (386K)
<i>Enhancers</i>				
Mouse Enhancers	0.730 \pm 0.032	<u>0.779</u> \pm 0.013	0.754 \pm 0.074	0.784 \pm 0.009
Human Enhancers Cohn	0.702 \pm 0.021	0.718 \pm 0.008	0.747 \pm 0.004	<u>0.743</u> \pm 0.005
Human Enhancer Ensembl	0.744 \pm 0.122	0.832 \pm 0.006	<u>0.893</u> \pm 0.008	0.900 \pm 0.004
<i>Species Classification</i>				
Coding vs. Intergenic	0.892 \pm 0.008	0.904 \pm 0.008	<u>0.915</u> \pm 0.003	0.943 \pm 0.001
Human vs. Worm	0.942 \pm 0.002	0.961 \pm 0.002	0.973 \pm 0.001	<u>0.967</u> \pm 0.002
<i>Regulatory Elements</i>				
Human Regulatory	0.872 \pm 0.005	0.862 \pm 0.004	<u>0.872</u> \pm 0.011	0.873 \pm 0.002
Human Non-TATA Promoters	0.861 \pm 0.009	0.887 \pm 0.005	<u>0.946</u> \pm 0.007	0.951 \pm 0.003
Human OCR Ensembl	0.698 \pm 0.013	0.744 \pm 0.019	0.828 \pm 0.006	<u>0.793</u> \pm 0.004

4.2 LONG RANGE

4.2.1 BEND GENE FINDING

Gene finding (Marin et al., 2023) is a multiclass classification task aimed at predicting the nucleotide types within DNA sequences (exons, introns, donors, acceptors, noncoding regions). This task aids in gene annotation and coding sequence identification. The dataset, derived from GENCODE annotations, includes labels $\{E_F, D_F, I_F, A_F, E_R, D_R, I_R, A_R, NC\}$.

The task employs the GENCODE dataset (Harrow et al., 2012), comprising 5,976 DNA sequences spanning 1,433 to 14,000 bps. The objective is to classify each nucleotide based on its gene structure context, using local signals to predict exon-intron boundaries and long-range dependencies for accurate annotation. As shown in Table 2, our ConvNova model achieved the highest MCC of 0.55 on the gene-finding task, outperforming NT-H (0.41), DNABERT-2 (0.43), GENA-LM (0.52), and HyenaDNA (0.35). This demonstrates ConvNova’s ability to leverage local and long-range dependencies for improved gene annotation.




Figure 3: **Chromatin Profile Prediction results.** AUC Score (\uparrow) is reported for ConvNova, HyenaDNA, NT v2, DNABERT-2, and DeepATT performance on transcription factors (TF), DNase I hypersensitive sites (DHS), histone modifications (HM), and average score.

4.2.2 CHROMATIN PROFILE PREDICTION

This classification task (Zhou and Troyanskaya, 2015) focuses on predicting chromatin profiles and epigenetic markers from DNA sequences, which is crucial for quantifying the functional effects of non-coding variants. The dataset comprises 919 chromatin features, including transcription factor (TF) binding profiles, DNase I hypersensitive sites (DHS), and histone mark (HM) profiles. For a given DNA sequence of length 1,000 bps, the task involves jointly predicting 919 binary classes corresponding to the chromatin profile of a central region of the sequence. We fine-tune our pretrained ConvNova models for 10 epochs, along with all baseline models, which range from 436K (tiny HyenaDNA) to 500M (NT-v2). Notably, ConvNova outperforms all baseline models, including DeepATT (Li et al., 2021), which is a previously state-of-the-art supervised model for this task, as illustrated in Figure 3.

Moreover, we provide compelling evidence of ConvNova’s scalability when ample data is available. As we augment the model’s parameter count from 6.8M to 51.1M, ConvNova shows a discernible improvement in the performance capabilities exhibited. See Appendix A.2.2 for experiment details and comparing different model sizes and training methods, whether pretrained or from scratch. Remarkably, without pretraining, ConvNova outperforms all Hyena-DNA models and DeepATT under the same epoch hyperparameter settings, even with a smaller parameter size.

Table 4: **Comparison of dilation and downsampling in ConvNova.** Performance (MCC-score or F1-score) is reported on selected NT benchmark tasks. The best values are in bold.

Task	Downsampling (1.7M)	Dilation (1.7M)
H3	65.20	81.62
H3K4me1	34.88	56.37
H3K9ac	48.81	68.98
Promoter All	93.69	96.99
Splice Sites All	45.65	96.45

4.3 ABLATION

4.4 DILATION OR DOWNSAMPLING

In image processing, the U-Net (Ronneberger et al., 2015) architecture has been proven highly effective through its use of downsampling and upsampling. Downsampling reduces image size while increasing the receptive field, enabling the capture of broader contextual information. Additionally, dilation, another method to increase the receptive field, expands the convolutional operation’s receptive field by introducing gaps within the convolution kernel.

In DNA sequence modeling tasks, both the upsampling-downsampling structure and dilation convolution represent potentially viable approaches. For the dilation mechanism, we use the design shown in Figure 2. For the downsampling architecture, we implement a variant of ConvNova with U-Net style, maintaining the gating mechanism. We retain the same pretraining settings and parameter size (1.7M) and conduct comparative experiments on randomly selected tasks within the NT benchmark.

Due to the significant performance gap observed as shown in Table 4, we opt not to conduct multiple experiments using ten different random seeds. Instead, we only utilize a single random seed. It is evident that while the downsampling (U-Net style) architecture can approximate the performance achieved using dilation in the ‘promoter all’ task, it significantly lags behind in other tasks, even decreasing by 50 points in ‘splice sites all’.

4.5 KEY DESIGNS

We conduct ablation experiments on the gate mechanism and the double-branch design in ConvNova. We implement an ordinary gate convolution model with the same parameter count (1.7M) named "Single Branch" in Table 5 to compare with the double-branch structure, and an additive aggregation model (1.7M) named "w/o Gate" in Table 5 to assess the role of the gate mechanism. We select homologous protein tasks from the NT benchmark and maintain the same method of selecting ten random seeds as described in Section 4.1.1 for comparison. For the details, refer to A.3.1.

The results indicate that both the feature aggregation method and the gate mechanism in ConvNova are crucial design components, effectively supporting the overall model capability.

For detailed ablation on kernel size, dilation rate, please refer to A.3.2.

5 DISCUSSION

We utilize dilation to control the receptive field in histone tasks from the NT-benchmark. H3K4me2, H3K4me3, and H3K14ac demonstrate significantly improved classification accuracy when the receptive field covers approximately 15% of the input length, compared to a global receptive field.

The enhancement effects of H3K14ac (Figure 6A) and H3K4me3 (Figure 6C) align with previous biological research (Ramakrishnan et al.,

Table 5: **Ablation study results.** MCC-score (\uparrow) is reported for performance comparison across different models. ‘w/o Gate’ represents the ablation of the Gate mechanism, while ‘Single Branch’ is the ablation of the double-branch structure in ConvNova. The best values are in bold, and the second-best is underlined. \pm indicates the error range across experiments

Task	w/o Gate	Single Branch	ConvNova
H3	78.96 \pm 1.46	<u>80.95</u> \pm 1.61	81.50 \pm 0.80
H3K4me1	57.83 \pm 1.87	<u>56.45</u> \pm 1.58	56.60 \pm 1.01
H3K4me2	52.52 \pm 2.59	50.20 \pm 1.92	53.72 \pm 2.42
H3K4me3	54.45 \pm 1.31	<u>58.66</u> \pm 0.97	60.20 \pm 1.91
H3K9ac	63.54 \pm 2.20	<u>66.87</u> \pm 0.69	68.10 \pm 1.91
H3K14ac	63.83 \pm 1.03	<u>65.34</u> \pm 2.12	66.19 \pm 1.84
H3K36me3	64.12 \pm 1.46	<u>66.77</u> \pm 1.42	68.31 \pm 1.19
H3K79me3	69.49 \pm 1.75	<u>71.23</u> \pm 1.37	72.08 \pm 1.23
H4	79.27 \pm 1.09	<u>80.55</u> \pm 0.61	81.12 \pm 0.93
H4ac	62.02 \pm 2.28	<u>63.07</u> \pm 1.10	64.75 \pm 1.90

Table 6: **ConvNova performance on Histone tasks in NT Benchmark.** Results are reported for models with 15% and 100% full sequence receptive fields. The best values are in bold.

Task	15% Receptive Field	100% Receptive Field
H3K4me2	57.45 \pm 2.27	53.72 \pm 2.42
H3K4me3	67.15 \pm 0.93	60.20 \pm 1.91
H3K14ac	70.71 \pm 2.32	66.19 \pm 1.84
H3K9ac	65.49 \pm 1.83	68.10 \pm 1.91

2016; Regadas et al., 2021), which has shown that these histone marks are highly enriched around transcription start sites (TSS), indicating their strong localized characteristics. As illustrated in Figure 6A, the pronounced enrichment of H3K14ac and H3K4me3 within this localized region supports the idea that a smaller receptive field is sufficient for effective classification in these tasks.

In contrast, H3K9ac (Figure 6D) exhibits a more uniform distribution across the gene body, which explains the suboptimal performance of a small receptive field in this case.

Interestingly, H3K4me2 (Figure 6B) also shows an enhancement effect with a small receptive field despite its enrichment being predominantly located in the middle of genes.

This observation raises two intriguing possibilities:

Gene Position Context: The enrichment of H3K4me2 in the middle of genes may indicate its role in transcriptional regulation during elongation. This localization could facilitate the binding of transcriptional machinery, enhancing transcriptional efficiency. Additionally, since our data were collected through ChIP-chip experiments on cells before and after oxidative stress, this enrichment might reflect interactions with genes associated with the histone methyltransferase Set1, suggesting a complex relationship that requires further exploration.

Contextual Locality: Alternatively, the improved performance of H3K4me2 in small receptive fields may not be tied to its position within genes but rather could stem from its involvement in other biological processes not captured during the experimental conditions. The sequences associated with H3K4me2 might exhibit strong local characteristics in contexts outside the immediate transcriptional environment, potentially involving regulatory mechanisms that are yet to be fully understood.

These possibilities highlight the need for further exploration to clarify the specific roles of H3K4me2 and its implications in various biological contexts.

6 CONCLUSION

In this study, we’ve revisited the convolutional paradigm for DNA modeling. Through analysis and extensive experimentation, we’ve shown that convolutional neural networks maintain a competitive edge in DNA modeling tasks. This reevaluation prompts a reconsideration of CNNs, a relatively early method, for such functions within the community. We’ve introduced ConvNova, featuring three key design elements: dilated convolution, gated convolution, and a dual-branch structure. With these innovations, ConvNova achieves state-of-the-art performance across multiple benchmarks with small model sizes. Additionally, we’ve investigated the varying demands for receptive field sizes across different tasks. While some tasks exhibit expected behaviors, others may suggest the presence of undiscovered biological phenomena. The **limitation** is that this work does not consider the multi-species pretraining dataset. Additionally, we have not considered the specific genomic regions used in pretraining; it is possible that training exclusively on known functional regions could lead to improved performance. Furthermore, the current foundation model benchmarks are primarily limited to classification tasks, which results in a lack of diversity in the types of tasks evaluated. The **positive societal impact** is that this work can accelerate the research of DNA. No **negative societal impact** is perceived.

REFERENCES

- Babak Alipanahi, Andrew Delong, Matthew T Weirauch, and Brendan J Frey. Predicting the sequence specificities of DNA- and RNA-binding proteins by deep learning. *Nature biotechnology*, 33(8): 831–838, 2015.
- Žiga Avsec, Vikram Agarwal, Daniel Visentin, Joseph R Ledsam, Agnieszka Grabska-Barwinska, Kyle R Taylor, Yannis Assael, John Jumper, Pushmeet Kohli, and David R Kelley. Effective gene expression prediction from sequence by integrating long-range interactions. *Nature methods*, 18(10):1196–1203, 2021.
- Peter W Battaglia, Jessica B Hamrick, Victor Bapst, Alvaro Sanchez-Gonzalez, Vinicius Zambaldi, Mateusz Malinowski, Andrea Tacchetti, David Raposo, Adam Santoro, Ryan Faulkner, et al. Relational inductive biases, deep learning, and graph networks. *arXiv preprint arXiv:1806.01261*, 2018.

- 540 Ken Chen, Huiying Zhao, and Yuedong Yang. Capturing large genomic contexts for accurately
541 predicting enhancer-promoter interactions. *Briefings in Bioinformatics*, 23(2):bbab577, 2022.
542
- 543 Ana J Chucair-Elliott, Sarah R Ocanas, Kevin Pham, Adeline Machalinski, Scott Plafker, Michael B
544 Stout, Michael H Elliott, and Willard M Freeman. Age-and sex-divergent translatomic responses
545 of the mouse retinal pigmented epithelium. *Neurobiology of Aging*, 2024.
- 546 Hugo Dalla-Torre, Liam Gonzalez, Javier Mendoza-Revilla, Nicolas Lopez Carranza, Adam Henryk
547 Grzywaczewski, Francesco Oteri, Christian Dallago, Evan Trop, Bernardo P de Almeida, Hassan
548 Sirelkhatim, et al. The nucleotide transformer: Building and evaluating robust foundation models
549 for human genomics. *bioRxiv*, pages 2023–01, 2023.
- 550 Katarína Grešová, Vlastimil Martinek, David Čechák, Petr Šimeček, and Panagiotis Alexiou. Ge-
551 nomic benchmarks: a collection of datasets for genomic sequence classification. *BMC Genomic
552 Data*, 24(1):25, 2023.
553
- 554 Albert Gu and Tri Dao. Mamba: Linear-time sequence modeling with selective state spaces. *arXiv
555 preprint arXiv:2312.00752*, 2023.
556
- 557 Jennifer Harrow, Adam Frankish, Jose M Gonzalez, Electra Tapanari, Mark Diekhans, Felix Kokocin-
558 ski, Bronwen L Aken, Daniel Barrell, Amonida Zadissa, Stephen Searle, et al. GENCODE:
559 the reference human genome annotation for The ENCODE Project. *Genome research*, 22(9):
560 1760–1774, 2012.
- 561 Yanrong Ji, Zhihan Zhou, Han Liu, and Ramana V Davuluri. DNABERT: Pre-trained bidirectional
562 encoder representations from transformers model for DNA-language in genome. *Bioinformatics*,
563 37(15):2112–2120, 2021.
- 564 Junru Jin, Yingying Yu, Ruheng Wang, Xin Zeng, Chao Pang, Yi Jiang, Zhongshen Li, Yutong Dai,
565 Ran Su, Quan Zou, et al. iDNA-ABF: multi-scale deep biological language learning model for the
566 interpretable prediction of DNA methylations. *Genome biology*, 23(1):219, 2022.
567
- 568 David R Kelley, Jasper Snoek, and John L Rinn. Basset: Learning the regulatory code of the
569 accessible genome with deep convolutional neural networks. *Genome research*, 26(7):990–999,
570 2016.
- 571 David R Kelley, Yakir A Reshef, Maxwell Bileschi, David Belanger, Cory Y McLean, and Jasper
572 Snoek. Sequential regulatory activity prediction across chromosomes with convolutional neural
573 networks. *Genome research*, 28(5):739–750, 2018.
574
- 575 Dohoon Lee, Jeewon Yang, and Sun Kim. Learning the histone codes with large genomic windows
576 and three-dimensional chromatin interactions using transformer. *Nature Communications*, 13(1):
577 6678, 2022.
- 578 Jiawei Li, Yuqian Pu, Jijun Tang, Quan Zou, and Fei Guo. Deepatt: a hybrid category attention neural
579 network for identifying functional effects of dna sequences. *Briefings in bioinformatics*, 22(3):
580 bbaa159, 2021.
- 581 Johannes Linder, Divyanshi Srivastava, Han Yuan, Vikram Agarwal, and David R. Kelley. Predicting
582 RNA-seq coverage from DNA sequence as a unifying model of gene regulation. *bioRxiv*, 2023.
583 doi: 10.1101/2023.08.30.555582.
584
- 585 Frederikke Isa Marin, Felix Teufel, Marc Horlacher, Dennis Madsen, Dennis Pultz, Ole Winther,
586 and Wouter Boomsma. BEND: Benchmarking DNA language models on biologically meaningful
587 tasks. In *The Twelfth International Conference on Learning Representations*, 2023.
- 588 Gerd HG Moe-Behrens, Rene Davis, and Karmella A Haynes. Preparing synthetic biology for the
589 world. *Frontiers in microbiology*, 4:5, 2013.
590
- 591 Eric Nguyen, Michael Poli, Marjan Faizi, Armin Thomas, Michael Wornow, Callum Birch-Sykes,
592 Stefano Massaroli, Aman Patel, Clayton Rabideau, Yoshua Bengio, et al. HyenaDNA: Long-range
593 genomic sequence modeling at single nucleotide resolution. *Advances in neural information
processing systems*, 36, 2024.

- 594 Dmitry Penzar, Daria Nogina, Elizaveta Noskova, Arsenii Zinkevich, Georgy Meshcheryakov, Andrey
595 Lando, Abdul Muntakim Rafi, Carl De Boer, and Ivan V Kulakovskiy. Legnet: a best-in-class deep
596 learning model for short dna regulatory regions. *Bioinformatics*, 39(8):btad457, 2023.
597
- 598 Michael Poli, Stefano Massaroli, Eric Nguyen, Daniel Y Fu, Tri Dao, Stephen Baccus, Yoshua
599 Bengio, Stefano Ermon, and Christopher Ré. Hyena hierarchy: Towards larger convolutional
600 language models. In *International Conference on Machine Learning*, pages 28043–28078. PMLR,
601 2023.
- 602 Ofir Press, Noah A Smith, and Mike Lewis. Train short, test long: Attention with linear biases
603 enables input length extrapolation. *arXiv preprint arXiv:2108.12409*, 2021.
604
- 605 Saravanan Ramakrishnan, Srijana Pokhrel, Sowmiya Palani, Christian Pflueger, Timothy J Parnell,
606 Bradley R Cairns, Srividya Bhaskara, and Mahesh B Chandrasekharan. Counteracting H3K4
607 methylation modulators set1 and jhd2 co-regulate chromatin dynamics and gene transcription.
608 *Nature communications*, 7(1):11949, 2016.
- 609 Isabel Regadas, Olle Dahlberg, Roshan Vaid, Oanh Ho, Sergey Belikov, Gunjan Dixit, Sebastian
610 Deindl, Jiayu Wen, and Mattias Mannervik. A unique histone 3 lysine 14 chromatin signature
611 underlies tissue-specific gene regulation. *Molecular Cell*, 81(8):1766–1780, 2021.
612
- 613 Olaf Ronneberger, Philipp Fischer, and Thomas Brox. U-Net: Convolutional networks for biomedical
614 image segmentation. In *Medical image computing and computer-assisted intervention–MICCAI
615 2015: 18th international conference, Munich, Germany, October 5-9, 2015, proceedings, part III
616 18*, pages 234–241. Springer, 2015.
- 617 Joel Rozowsky, Jiahao Gao, Beatrice Borsari, Yucheng T Yang, Timur Galeev, Gamze Gürsoy,
618 Charles B Epstein, Kun Xiong, Jinrui Xu, Tianxiao Li, et al. The EN-TE_x resource of multi-tissue
619 personal epigenomes & variant-impact models. *Cell*, 186(7):1493–1511, 2023.
620
- 621 Nicolas Scalzitti, Arnaud Kress, Romain Orhand, Thomas Weber, Luc Moulinier, Anne Jeannin-
622 Girardon, Pierre Collet, Olivier Poch, and Julie D Thompson. Spliceator: multi-species splice site
623 prediction using convolutional neural networks. *BMC bioinformatics*, 22:1–26, 2021.
- 624 Yair Schiff, Chia-Hsiang Kao, Aaron Gokaslan, Tri Dao, Albert Gu, and Volodymyr Kuleshov.
625 Caduceus: Bi-directional equivariant long-range dna sequence modeling. *arXiv preprint
626 arXiv:2403.03234*, 2024.
627
- 628 Christina V Theodoris, Ling Xiao, Anant Chopra, Mark D Chaffin, Zeina R Al Sayed, Matthew C
629 Hill, Helene Mantineo, Elizabeth M Brydon, Zexian Zeng, X Shirley Liu, et al. Transfer learning
630 enables predictions in network biology. *Nature*, 618(7965):616–624, 2023.
- 631 Zixuan Wang, Meiqin Gong, Yuhang Liu, Shuwen Xiong, Maocheng Wang, Jiliu Zhou, and Yongqing
632 Zhang. Towards a better understanding of TF-DNA binding prediction from genomic features.
633 *Computers in Biology and Medicine*, 149:105993, 2022.
634
- 635 Fisher Yu and Vladlen Koltun. Multi-scale context aggregation by dilated convolutions. *arXiv
636 preprint arXiv:1511.07122*, 2015.
- 637 Jiahui Yu, Zhe Lin, Jimei Yang, Xiaohui Shen, Xin Lu, and Thomas S Huang. Free-form image
638 inpainting with gated convolution. In *Proceedings of the IEEE/CVF international conference on
639 computer vision*, pages 4471–4480, 2019.
640
- 641 Weihao Yu and Xinchao Wang. MambaOut: Do we really need mamba for vision? *arXiv preprint
642 arXiv:2405.07992*, 2024.
- 643 Manzil Zaheer, Guru Guruganesh, Kumar Avinava Dubey, Joshua Ainslie, Chris Alberti, Santiago
644 Ontanon, Philip Pham, Anirudh Ravula, Qifan Wang, Li Yang, et al. Big bird: Transformers for
645 longer sequences. *Advances in neural information processing systems*, 33:17283–17297, 2020.
646
- 647 Pengyu Zhang, Hongming Zhang, and Hao Wu. iPro-WAEL: a comprehensive and robust framework
for identifying promoters in multiple species. *Nucleic Acids Research*, 50(18):10278–10289, 2022.

648 Jian Zhou and Olga G Troyanskaya. Predicting effects of noncoding variants with deep learning–based
649 sequence model. *Nature methods*, 12(10):931–934, 2015.

650
651 Zhihan Zhou, Yanrong Ji, Weijian Li, Pratik Dutta, Ramana V Davuluri, and Han Liu. DNABERT-2:
652 Efficient foundation model and benchmark for multi-species genomes. In *The Twelfth International
653 Conference on Learning Representations*.

654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701

A APPENDIX

All the experiments can be conducted with 4 RTX-4090 GPUs.

A.1 PRETRAINING

We adhere to most of the pretraining hyperparameters and settings used for HyenaDNA, including the dataset selection. However, we implemented some small differences. We pretrained on sequences with a maximum length of 16K, while other pretraining sequence lengths were 1K or 2K, aligning with HyenaDNA’s recommendation to use sequences 2-4 times longer than those required for downstream tasks. We set the global batch size to 512 and trained for 400 epochs with a learning rate of $1e-3$. For MLM pretraining, selected nucleotides were "masked" by replacing them with "N" to predict the original nucleotide type. Table 7 and Table 8 illustrate the parameter settings for models of various sizes and the time needed to pretrain. [The pretraining dataset is HG38⁴](#).

Table 7: **Pretraining hyperparameters.** Values used during pretraining are reported.

Hyperparameter	Value
Learning Rate	1×10^{-3}
Batch Size	256
Weight Decay	0.1
Dropout	0.0
Optimizer	AdamW
Optimizer Momentum	$\beta_1 = 0.9, \beta_2 = 0.999$
Learning Rate Scheduler	Cosine Decay
Training Epochs	200

Table 8: **Pretraining time.** The time required for pretraining different model sizes with varying sequence lengths on 4 RTX-4090 GPUs is reported.

Parameters	386K	1.7M	7.4M	27.4M
Sequence Length	1K	1K	16K	2K
Time	50 mins	80 mins	14 hours	4 hours

A.2 DOWNSTREAM TASKS

All hyperparameters for the ConvNova model on downstream tasks can be found in Table 9.

Table 9: **Hyperparameters for ConvNova model on all downstream tasks.** Specific settings for each task include sequence length, dilation rate, hidden dimension, number of GCBs, and model size.

Tasks	Sequence Length	Dilation Rate	Hidden Dim	Num GCB	Size
NT	1K	4/1	128	5	1.7M
Deepstarr	1K	4	128	5	1.7M
Genomic Benchmark	1K	4	64	5	386K
Chromatin Profile Prediction	2K	4	512/256	5	27.4M/6.8M
Gene Finding	16K	4	196	10	7.4M

A.2.1 NUCLEOTIDE TRANSFORMER BENCHMARK

Objective: Sequence classification.

⁴https://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.26

Models and Setup: We follow the HyenaDNA setup (Nguyen et al., 2024), splitting the dataset into 90/10 training and test sets and fine-tuning DNABERT-2, HyenaDNA, NT-v2, and Caduceus-Ph for 20 epochs using pre-trained weights from Hugging Face.

Experiment Configuration: All models are tested with 32-bit floating point precision, and ConvNova uses consistent hyperparameters across tasks as detailed in Table 9, with ten random seeds to report mean values, shown in Tables 16 and 17. Optimizer we use AdamW and share the same setting as Table 7. Learning rate we use 1e-3 for ConvNova and official learning rate for the baselines.

A.2.2 CHROMATIN PROFILE PREDICTION

Objective: Perform 919 binary classification tasks on DNA fragments using 4.4 million training data points. Even small improvements (<1%) are significant for this challenging task.

Models and Setup: The baseline models include HyenaDNA (0.4M, 3.3M, 6.5M), NT-v2 (50M, 100M, 250M, 500M), DNABERT-2, and DeepATT (previous state-of-the-art CNN for this task), trained for 10 epochs. All models are tested using 32-bit floating point precision.

Experiment Configuration: The train/test split follows the DeepSEA paper methodology. The optimizer used is AdamW with settings from Table 7, and a learning rate of 1e-3 for ConvNova, with official learning rates used for the baselines.

We provide more detailed comparisons of different models in Table 10. Besides the pretrained ConvNova model, we also train ConvNova from scratch. Remarkably, even without pretraining, the model outperformed all versions of HyenaDNA at similar parameter sizes, further demonstrating ConvNova’s superior ability to encode long-sequence information. Hyperparameters can be found in Table 9.

Table 10: **Chromatin Profile Prediction detailed comparison.** AUC Score (\uparrow) is reported for ConvNova, HyenaDNA, NT v2, DNABERT-2, DeepATT performance on transcription factors (TF), DNase I hypersensitive sites (DHS), histone modifications (HM), and average score. The best values per task are bolded, and the second best are underlined.

Tasks	Pretrain										From Scratch			
	ConvNova		Hyena-DNA			NTv2				DNABERT-2	ConvNova			DeepATT
	6.8M	27.4M	0.4M	3.3M	6.5M	50M	100M	250M	500M	117M	1.7M	6.8M	27.4M	7.8M
TF	96.77	96.99	94.42	94.72	95.91	96.17	96.23	96.64	<u>96.76</u>	96.16	96.62	96.63	96.77	96.35
HM	86.54	86.70	84.64	84.96	85.46	86.23	86.20	86.81	<u>86.76</u>	86.18	85.74	85.74	85.76	86.16
DHS	93.42	93.71	90.58	90.90	92.33	92.50	92.65	<u>93.13</u>	93.39	92.50	92.93	92.94	93.03	92.52
Avg.	92.25	92.47	89.89	90.19	91.23	91.63	91.69	<u>92.19</u>	92.30	91.61	91.76	91.77	91.85	91.68

A.2.3 GENE FINDING

Objective: Classify each nucleotide based on its gene structure context, predicting exon-intron boundaries and capturing long-range dependencies for accurate annotation.

Models and Setup: The baseline models include NT-H, DNABERT2, HyenaDNA, and GENA_LM. ConvNova is tuned for 10 epochs, while the results of the other baselines are taken from BEND.


Experiment Configuration: The optimizer used is AdamW with settings from Table 7, and a learning rate of 1e-3 for ConvNova.

Dilation and Layer impact on Gene Finding We find that the gene-finding task requires long-range dependency. Both decreasing dilation rate and the number of layers can result in performance decline in Figure 4. Models are trained from scratch in all experiments. For Dilation impact, the model dimension is 256 and contains 5 GCBs. For GCB counts impact, the model dimension is set to 256, and the dilation rate is set to 4. For additional hyperparameter settings in this task, readers can refer to Table 9.

A.2.4 GENOMIC BENCHMARK

Objective: Sequence classification task.

Figure 4: The impact of dilation rate and GCB counts on the MCC in ConvNova model.



Models and Setup: All models are fine-tuned for ten epochs. ConvNova is used as the backbone with a width of 64, 5 GCBs, and a parameter size of 386K, pretrained on sequences of length 1000. The baselines used in this task are CaduceusPh, HyenaDNA, and the CNN model provided in the benchmark. We follow the train-valid split (90/10) as provided by HyenaDNA (Nguyen et al., 2024).

Experiment Configuration: The models are trained with 16-precision float points. We use five random seeds and report the mean result across all models. Hyperparameters for this task can be found in Table 9. The optimizer used is AdamW with settings from Table 7, and a learning rate of 1e-3 for ConvNova, with official learning rates used for the baselines.

A.3 ABLATIONS EXPERIMENTS

A.3.1 GATING MECHANISM AND DUAL BRANCH

In this section, we provide more details about the ablation experiments. There are two variants of the ConvNova model: "Single Gate" and "Dual Branch", and one variant without any gating mechanism.

Single Gate Our "Single Gate" design follows a sequential process as shown in Eq.2. First, a single Conv1D layer processes the input feature $\mathbf{A} \in \mathbb{R}^{l \times d}$ with weight matrix $\mathbf{W}_d \in \mathbb{R}^{k \times d \times d}$ and bias term $\mathbf{b}_d \in \mathbb{R}^d$ to produce $\mathbf{z} \in \mathbb{R}^{l \times d}$. This output is then processed through two parallel activation paths - GELU activation and sigmoid activation (σ). The final output is obtained through residual addition of element-wise multiplication (\odot) of activated \mathbf{h} and gated \mathbf{g} . To maintain parameter count parity with ConvNova, we increase the dimension of the Conv1D layer.

$$\begin{aligned}
 \mathbf{z} &= \mathbf{W}_d * \mathbf{A} + \mathbf{b}_d \\
 \mathbf{h} &= \text{GELU}(\mathbf{z}) \\
 \mathbf{g} &= \sigma(\mathbf{z}) \\
 \mathbf{A} &= \mathbf{A} + \mathbf{h} \odot \mathbf{g}
 \end{aligned}
 \tag{2}$$

Dual Gate Our "Dual Branch" design processes input features through two parallel convolutional paths with gating mechanisms, as shown in Eq.3. The inputs $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{l \times d}$ are processed through two separate Conv1D layers with weights $\mathbf{W}_{1,d/\sqrt{2}}, \mathbf{W}_{2,d/\sqrt{2}} \in \mathbb{R}^{k \times d \times d/\sqrt{2}}$ and biases $\mathbf{b}_{1,d/\sqrt{2}}, \mathbf{b}_{2,d/\sqrt{2}} \in \mathbb{R}^{d/\sqrt{2}}$ to produce intermediate features $\mathbf{z}_1, \mathbf{z}_2 \in \mathbb{R}^{l \times d/\sqrt{2}}$. The first path applies GELU activation to \mathbf{z}_1 , while the second applies sigmoid activation to \mathbf{z}_2 . The final output \mathbf{A} is obtained through residual addition of element-wise multiplication (\odot) of activated \mathbf{h} and gated \mathbf{g} , and \mathbf{B} is obtained through residual addition of gated \mathbf{g} .

864
865
866
867
868
869
870
871
872

$$\begin{aligned}
\mathbf{z}_1 &= \mathbf{W}_{1,d/\sqrt{2}} * \mathbf{A} + \mathbf{b}_{1,d/\sqrt{2}} \\
\mathbf{z}_2 &= \mathbf{W}_{2,d/\sqrt{2}} * \mathbf{B} + \mathbf{b}_{2,d/\sqrt{2}} \\
\mathbf{h} &= \text{GELU}(\mathbf{z}_1) \\
\mathbf{g} &= \sigma(\mathbf{z}_2) \\
\mathbf{A} &= \mathbf{A} + \mathbf{h} \odot \mathbf{g} \\
\mathbf{B} &= \mathbf{B} + \mathbf{g}
\end{aligned}
\tag{3}$$

873
874
875
876
877

Addition w/o Gate This variant uses simple addition operations for feature aggregation without any gating mechanism. As shown in Eq.4, the input features $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{l \times d}$ are processed through convolutional layers and GELU activation to produce intermediate features $\mathbf{h}, \mathbf{g} \in \mathbb{R}^{l \times d/\sqrt{2}}$. These are then combined through addition to produce the final outputs \mathbf{A} and \mathbf{B} .

878
879
880
881
882
883

$$\begin{aligned}
\mathbf{h} &= \text{GELU}(\mathbf{W}_{1,d/\sqrt{2}} * \mathbf{A} + \mathbf{b}_{1,d/\sqrt{2}}) \\
\mathbf{g} &= \text{GELU}(\mathbf{W}_{2,d/\sqrt{2}} * \mathbf{B} + \mathbf{b}_{2,d/\sqrt{2}}) \\
\mathbf{A} &= \mathbf{A} + \mathbf{h} + \mathbf{g} \\
\mathbf{B} &= \mathbf{B} + \mathbf{g}
\end{aligned}
\tag{4}$$

884

A.3.2 DILATION MECHANISM, KERNEL SIZE AND LOCAL DEPENDENCY ANALYSIS

885
886
887
888

In this section, we present the results of the ablation experiments on the dilation rate and kernel size. We perform experiments on the NT benchmark, testing kernel sizes of 3, 5, 7, 9, 11 and dilation rates of 1, 2, 3, 4. We report the average results across 18 tasks from the entire benchmark.

889
890
891

As shown in Table 11, the performance generally improves with increasing dilation rate and kernel size. However, to balance compatibility with long-range tasks and maintain relatively small model parameters, we ultimately select a kernel size of 9 and a dilation rate of 4.

892
893
894

Table 11: Performance of different kernel sizes and dilation rates across 18 tasks in the NT benchmark. The values represent the average performance for all tasks.

895
896
897
898
899
900
901

Kernel Size \ Dilation Rate	1	2	3	4
3	63.710	67.440	69.630	71.705
5	63.939	70.324	73.049	73.688
7	68.353	71.141	74.049	74.598
9	68.959	72.257	74.241	74.530
11	69.408	73.437	74.835	74.549

902
903
904
905
906
907
908

Furthermore, we observe that even for short-range tasks (all having a sequence length of 500 bp), the intensity of local dependencies varies significantly. For instance, the tasks H3K4me2, H3K4me3, and H3K14ac, as discussed in Section 5, exhibit strong local dependencies, with better performance observed when the dilation rate is small. On the other hand, tasks like splice site prediction show a much stronger global dependency, which benefits from a larger dilation rate. See table 12 for more details.

909
910

A.4 SUPERVISED METHODS AGAINST FOUNDATION MODELS

911
912
913
914
915
916
917

We conduct a comparison between previous supervised models and DNA foundation models on both the NT benchmark and genomic benchmark. For our comparison, we select the popular Basenji and the recently established state-of-the-art model, LegNet, which excels in short DNA regulatory regions. Originally, these supervised models are designed for specific tasks. We apply them to the benchmarks used for foundation models. As observed in Table 13 and Table 14, although they show decent performance on specific tasks, they do not achieve the same level of consistency across various datasets as foundation models do. This inconsistency may arise from the enhanced modeling capabilities afforded by pretraining in foundation models.

Table 12: Performance of different dilation rates and kernel sizes on the NT benchmark. For example, k3_d1 represents a kernel size of 3 and a dilation rate of 1, with similar notation for other configurations.

Task	Histone Modifications							
	H3	H3K4me1	H3K4me2	H3K4me3	H3K9ac	H3K14ac	H3K36me3	H3K79me3
k3_d1	70.14	43.14	41.26	49.12	54.00	54.80	54.48	62.38
k3_d2	74.66	50.54	47.52	56.50	58.92	60.16	63.18	67.50
k3_d3	79.24	52.44	49.50	55.18	61.60	62.72	63.66	68.50
k3_d4	78.72	51.94	47.54	54.12	62.36	62.30	63.74	68.48
k5_d1	70.76	50.26	48.28	56.86	58.02	61.62	60.44	65.38
k5_d2	76.80	55.88	53.50	62.30	63.58	66.24	65.04	70.30
k5_d3	80.88	55.58	52.16	58.86	65.62	65.22	66.02	70.84
k5_d4	80.48	55.12	50.46	56.56	65.60	63.58	65.58	69.70
k7_d1	71.76	53.12	50.72	59.78	61.06	65.16	62.80	67.32
k7_d2	77.72	56.62	53.66	62.54	65.96	66.92	66.06	70.78
k7_d3	81.20	56.42	53.38	58.04	66.56	65.30	66.54	70.96
k7_d4	81.62	56.46	51.02	57.60	66.38	64.24	65.96	69.10
k9_d1	72.52	54.70	52.90	60.50	62.26	66.02	63.94	69.14
k9_d2	78.02	56.76	55.38	62.04	66.80	67.46	67.62	71.52
k9_d3	81.58	56.16	52.20	58.64	66.70	65.00	66.28	71.84
k9_d4	81.18	55.36	48.54	56.60	65.84	64.40	66.08	70.02
k11_d1	72.60	55.90	53.78	62.80	63.02	66.40	66.14	69.58
k11_d2	79.62	56.74	55.20	63.84	67.08	67.24	68.20	71.54
k11_d3	81.66	56.12	53.98	58.66	66.88	64.62	66.84	71.22
k11_d4	80.66	55.16	47.84	55.76	66.82	63.20	66.28	70.18

Task	promoter			splice_sites			enhancer	
	all	non_tata	tata	all	acceptor	donor	types	
k3_d1	94.68	94.82	94.54	47.08	79.72	78.16	54.04	48.96
k3_d2	96.06	96.04	95.74	49.26	80.62	79.84	52.36	49.76
k3_d3	96.50	96.44	96.20	60.36	84.10	83.10	53.84	50.66
k3_d4	96.62	96.64	96.38	91.42	87.54	87.12	56.46	48.96
k5_d1	95.10	95.30	94.88	48.20	79.84	78.20	52.44	51.30
k5_d2	96.36	96.48	96.32	53.84	82.22	81.28	52.96	50.50
k5_d3	96.68	96.80	96.54	88.70	86.96	85.74	55.68	49.62
k5_d4	96.76	96.74	96.18	96.38	96.08	95.62	56.14	51.84
k7_d1	95.36	95.46	95.46	48.50	79.72	78.92	55.70	51.48
k7_d2	96.54	96.48	96.32	59.54	83.76	83.12	53.26	49.00
k7_d3	96.76	96.70	96.28	95.38	88.86	88.82	57.42	49.88
k7_d4	96.76	96.68	96.48	96.24	97.02	96.40	58.02	50.02
k9_d1	95.56	95.66	95.28	49.16	80.36	79.40	53.58	49.60
k9_d2	96.54	96.54	96.34	70.96	84.78	84.16	54.06	48.12
k9_d3	96.70	96.66	96.40	95.94	91.98	90.98	56.82	48.94
k9_d4	96.74	96.82	96.92	96.54	97.08	96.48	58.60	50.00
k11_d1	95.68	95.84	95.06	49.08	80.78	79.04	54.56	49.18
k11_d2	96.40	96.46	96.40	84.60	85.62	84.84	55.92	48.20
k11_d3	96.64	96.58	96.22	95.84	96.86	95.62	57.16	48.58
k11_d4	96.82	96.66	96.60	96.76	97.16	86.46	61.68	50.80

Table 13: **Supervised Methods against Foundation Models on NT Benchmark.** MCC/F1-score is reported for pretrained NTv2, HyenaDNA, DNABERT-2, Caduceus-Ph, ConvNova, Basenji, LegNet ConvNova*(* means trained from scratch). The best values per task are bold, and the second-best are underlined. \pm indicates the error range across five random seeds.

	NTv2 (500M)	HyenaDNA (1.6M)	DNABERT-2 (117M)	Caduceus-Ph (1.9M)	ConvNova (1.7M)	Basenji (7.4M)	LegNet (2.1M)	ConvNova* (1.7M)
<i>Histone</i>								
H3	78.17 \pm 2.54	78.14 \pm 1.70	79.31 \pm 0.68	80.48 \pm 1.04	81.50 \pm 0.80	78.05 \pm 1.83	76.24 \pm 0.83	<u>81.18</u> \pm 1.68
H3K4me1	51.64 \pm 1.12	44.52 \pm 2.59	48.34 \pm 4.63	52.83 \pm 0.96	56.60 \pm 1.01	42.70 \pm 3.16	47.47 \pm 1.36	<u>55.36</u> \pm 2.26
H3K4me2	37.24 \pm 2.25	42.68 \pm 2.66	43.02 \pm 2.92	<u>49.88</u> \pm 2.65	57.45 \pm 2.27	34.73 \pm 1.66	45.75 \pm 0.56	<u>48.54</u> \pm 3.24
H3K4me3	50.30 \pm 1.77	50.41 \pm 3.15	45.43 \pm 3.33	<u>56.72</u> \pm 2.58	67.15 \pm 0.93	38.85 \pm 1.49	52.67 \pm 2.54	<u>56.60</u> \pm 2.70
H3K9ac	61.05 \pm 1.40	58.50 \pm 1.75	60.04 \pm 1.27	63.27 \pm 2.29	68.10 \pm 1.91	56.34 \pm 2.86	58.21 \pm 0.92	<u>65.84</u> \pm 1.14
H3K14ac	57.22 \pm 2.19	56.71 \pm 2.40	54.49 \pm 4.99	60.84 \pm 2.94	70.71 \pm 2.32	49.88 \pm 1.60	56.64 \pm 2.12	<u>64.40</u> \pm 0.90
H3K36me3	60.50 \pm 1.75	59.92 \pm 1.06	57.58 \pm 2.38	61.12 \pm 1.44	68.31 \pm 1.19	52.73 \pm 1.84	57.54 \pm 1.03	<u>66.08</u> \pm 1.88
H3K79me3	65.78 \pm 2.34	66.25 \pm 3.65	64.38 \pm 0.48	67.17 \pm 2.03	72.08 \pm 1.23	62.43 \pm 1.29	63.78 \pm 1.72	<u>70.02</u> \pm 0.72
H4	79.87 \pm 1.34	78.15 \pm 1.58	78.18 \pm 0.98	80.10 \pm 1.00	81.12 \pm 0.93	77.60 \pm 2.02	78.73 \pm 1.93	<u>80.78</u> \pm 0.82
H4ac	55.22 \pm 2.20	54.15 \pm 2.96	51.80 \pm 0.10	59.26 \pm 3.67	66.10 \pm 1.20	46.05 \pm 1.77	55.29 \pm 2.60	<u>63.56</u> \pm 1.14
<i>Regulatory</i>								
Enhancer	54.51 \pm 1.94	53.13 \pm 4.52	52.50 \pm 1.44	55.20 \pm 2.56	57.60 \pm 2.52	57.54 \pm 2.75	55.25 \pm 3.23	<u>58.60</u> \pm 3.80
Enhancer Types	43.36 \pm 1.75	48.16 \pm 2.48	44.32 \pm 1.18	47.17 \pm 2.85	49.75 \pm 2.82	52.67 \pm 2.94	44.72 \pm 2.04	<u>50.00</u> \pm 1.80
Promoter All	96.82 \pm 0.47	95.57 \pm 0.18	96.23 \pm 0.17	96.65 \pm 0.16	96.82 \pm 0.22	95.79 \pm 0.24	96.52 \pm 0.33	<u>96.74</u> \pm 0.06
Promoter non-TATA	97.45 \pm 0.69	95.86 \pm 0.37	<u>97.17</u> \pm 0.17	96.31 \pm 0.50	96.76 \pm 0.21	95.99 \pm 0.18	96.45 \pm 0.27	<u>96.82</u> \pm 0.28
Promoter TATA	96.53 \pm 0.81	95.88 \pm 0.53	96.99 \pm 0.49	96.21 \pm 0.81	96.34 \pm 0.38	95.89 \pm 0.78	96.25 \pm 0.45	<u>96.92</u> \pm 0.32
<i>Splice sites</i>								
Splice Acceptor	97.99 \pm 0.66	96.98 \pm 0.49	97.49 \pm 0.36	94.21 \pm 0.37	96.23 \pm 0.41	97.65 \pm 0.34	85.80 \pm 0.58	<u>97.08</u> \pm 0.32
Splice Donor	98.50 \pm 0.43	95.27 \pm 1.07	94.33 \pm 0.27	94.69 \pm 0.67	96.62 \pm 0.61	<u>97.82</u> \pm 0.66	84.72 \pm 0.95	<u>96.48</u> \pm 0.32
Splice All	98.15 \pm 1.01	94.05 \pm 1.08	93.75 \pm 1.25	92.87 \pm 1.73	96.33 \pm 0.31	<u>98.07</u> \pm 0.45	54.71 \pm 2.38	<u>96.54</u> \pm 0.36

Table 14: **Supervised Methods against Foundation Models on Genomic Benchmark.** Top-1 accuracy (\uparrow) is reported for pretrained HyenaDNA, Caduceus-Ph, ConvNova, Basenji, LegNet, and the original CNN baseline (trained from scratch). The best values are in bold, and the second-best is underlined. \pm indicates the error range across five random seeds.

Task	CNN (264K)	HyenaDNA (436K)	Caduceus-Ph (470K)	ConvNova (386K)	Basenji (7.4N)	LegNet (2.1M)
<i>Enhancers</i>						
Mouse Enhancers	0.730 \pm 0.032	<u>0.779</u> \pm 0.013	0.754 \pm 0.074	0.784 \pm 0.009	0.659 \pm 0.155	0.504 \pm 0.000
Human Enhancers Cohn	0.702 \pm 0.021	0.718 \pm 0.008	0.747 \pm 0.004	<u>0.743</u> \pm 0.005	0.712 \pm 0.030	0.739 \pm 0.004
Human Enhancer Ensembl	0.744 \pm 0.122	0.832 \pm 0.006	0.893 \pm 0.008	<u>0.900</u> \pm 0.004	0.905 \pm 0.007	0.879 \pm 0.002
<i>Species Classification</i>						
Coding vs. Intergenic	0.892 \pm 0.008	0.904 \pm 0.008	0.915 \pm 0.003	0.943 \pm 0.001	0.905 \pm 0.004	<u>0.939</u> \pm 0.002
Human vs. Worm	0.942 \pm 0.002	0.961 \pm 0.002	0.973 \pm 0.001	<u>0.967</u> \pm 0.002	0.957 \pm 0.003	0.965 \pm 0.001
<i>Regulatory Elements</i>						
Human Regulatory	0.872 \pm 0.005	0.862 \pm 0.004	<u>0.872</u> \pm 0.011	0.873 \pm 0.002	0.764 \pm 0.005	0.764 \pm 0.006
Human Non-TATA Promoters	0.861 \pm 0.009	0.887 \pm 0.005	<u>0.946</u> \pm 0.007	0.951 \pm 0.003	0.919 \pm 0.006	0.942 \pm 0.007
Human OCR Ensembl	0.698 \pm 0.013	0.744 \pm 0.019	0.828 \pm 0.006	<u>0.793</u> \pm 0.004	0.766 \pm 0.009	<u>0.802</u> \pm 0.004

A.5 NEIGHBORHOOD IMPORTANCE IN DNA MODELING

To support our hypothesis that the inductive bias of CNNs—emphasizing neighboring nucleotides—is beneficial for DNA modeling, we make modifications to the Rotary Position Embedding (RoPE) mechanism. Specifically, we adjust the original θ values, enabling each attention head to have distinct θ values.

Additionally, we modified the initialization of the bias term in the linear layers of the K and Q projections. The bias was initialized to $[0, 0, 0, 0, 0, \dots, 1]$, while all other parameters retained the initialization strategy of NTv2 (standard deviation $\sigma = 0.02$, mean $\mu = 0$). This adjustment ensures that the initialization process places greater emphasis on neighboring nucleotides. Refer to Fig A.4 for an illustration of the improved attention map, which places greater emphasis on neighboring tokens.

To validate our hypothesis, we train NTv2 from scratch on three tasks where ConvNova with small dilation values performs better (H3K4me2, H3K4me3, and H3K14ac). The results, presented in Table 15, show that the adjusted NTv2 consistently outperform its original version. However, despite these improvements, our naive modifications to the transformer architecture does not surpass ConvNova trained from scratch(See Table 13). To enhance transformer and state space model (SSM) designs for DNA modeling further dedicated efforts is needed.

Our primary goal here is to validate the hypothesis and provide an explanation for why CNNs might outperform transformers in this context.

Table 15: Performance comparison of NTv2 and NTv2*, where NTv2* represents our modified version. Results demonstrate that NTv2* achieves significant improvements.

Task	NTv2*	NTv2
H3K14ac	46.65	34.42
H3K4me2	33.10	25.79
H3K4me3	34.62	21.40




Figure 5: Visualization of the attention maps for NTv2*(Left) and NTv2(Right). The asterisk (*) denotes the modified version. The modified initialization places more emphasis on neighboring tokens in the attention map.

Table 16: Results for all models on NT benchmark across genomic regions. Ten random seeds were used for each model. The best values per task are bolded.

Seed	2222	42	43	44	45	46	47	48	49	50
Promoter All										
ConvNova (4 dilation)	96.99	96.84	96.81	96.82	96.75	96.82	96.64	96.60	96.96	96.96
Caduceus-Ph	96.66	96.61	96.64	96.61	96.51	96.79	96.77	96.49	96.73	96.68
HyenaDNA	95.55	95.62	95.44	95.58	95.62	95.58	95.56	95.65	95.75	95.39
DNABERT-2	96.38	96.27	96.15	96.21	96.05	96.19	96.31	96.22	96.29	96.11
NTv2	96.47	97.19	96.95	96.35	97.12	96.63	97.21	96.85	96.40	97.00
Promoter Non-TATA										
ConvNova (4 dilation)	96.55	96.81	96.73	96.83	96.88	96.61	96.77	96.83	96.87	96.72
Caduceus-Ph	96.14	95.81	96.14	96.52	96.79	95.98	96.47	96.36	96.39	96.50
HyenaDNA	95.81	96.23	95.66	96.00	95.96	95.60	95.63	96.04	95.87	95.83
DNABERT-2	97.18	97.02	97.31	97.22	97.04	97.29	97.15	97.03	97.26	97.12
NTv2	96.89	97.44	98.01	96.81	97.29	98.14	97.52	96.78	97.93	97.66
Promoter TATA										
ConvNova (4 dilation)	96.57	96.08	95.92	96.57	96.07	96.40	96.57	96.09	96.41	96.72
Caduceus-Ph	95.40	96.72	96.40	95.73	96.23	96.74	96.24	95.71	96.56	96.41
HyenaDNA	95.92	96.06	96.23	95.89	96.41	95.55	95.58	96.16	95.58	95.42
DNABERT-2	97.38	97.12	96.52	97.41	97.18	96.53	97.40	97.15	96.54	96.87
NTv2	95.75	96.89	97.13	95.92	96.48	96.12	97.05	96.73	95.88	97.34
Splice Sites All										
ConvNova (4 dilation)	96.46	96.18	96.6	96.41	96.28	96.50	95.82	96.30	95.93	95.85
Caduceus-Ph	96.29	96.33	96.21	96.36	96.58	96.04	96.19	95.84	96.03	96.27
HyenaDNA	96.97	96.99	96.93	96.74	96.96	97.22	97.05	96.89	97.47	96.61
DNABERT-2	94.32	94.18	93.71	94.29	94.21	93.72	94.34	94.23	93.74	94.03
NTv2	97.14	98.47	98.97	97.88	98.63	98.09	97.57	98.42	98.28	98.05
Splice Sites Acceptor										
ConvNova (4 dilation)	96.65	96.01	96.87	96.70	96.86	96.59	96.37	97.06	96.50	96.59
Caduceus-Ph	97.36	96.41	96.74	96.61	96.24	96.83	96.74	96.87	96.50	96.56
HyenaDNA	94.20	95.37	95.21	94.97	96.06	95.94	95.2	94.48	95.73	95.56
DNABERT-2	94.57	94.21	94.23	94.55	94.19	94.24	94.59	94.18	94.25	94.28
NTv2	98.57	97.89	98.14	97.45	98.64	97.72	98.09	97.33	98.38	97.69
Splice Sites Donor										
ConvNova (4 dilation)	96.57	96.34	96.52	96.51	96.44	96.02	96.04	96.38	96.12	96.33
Caduceus-Ph	92.47	93.21	94.74	92.61	92.78	93.56	91.14	94.36	92.57	91.22
HyenaDNA	94.38	94.59	95.04	93.09	94.63	93.38	94.52	93.67	92.97	94.22
DNABERT-2	92.51	94.76	93.72	92.48	94.81	93.74	92.53	94.79	93.71	94.52
NTv2	98.93	98.67	98.43	98.12	98.54	98.79	98.26	98.68	98.37	98.21
Enhancer										
ConvNova (4 dilation)	59.00	55.08	57.09	58.54	55.33	58.26	58.77	58.42	58.10	57.39
Caduceus-Ph	53.72	53.64	54.04	53.50	57.76	56.75	55.94	55.35	54.66	56.63
HyenaDNA	55.34	53.88	54.43	51.44	51.56	54.28	55.53	48.61	53.18	53.05
DNABERT-2	52.11	51.06	52.15	53.67	53.23	53.24	51.56	52.08	52.88	53.06
NTv2	53.25	55.80	54.10	52.99	56.45	54.75	53.40	55.50	54.99	53.85
Enhancer Types										
ConvNova (4 dilation)	49.08	48.32	49.24	52.56	48.72	48.90	51.24	49.45	49.96	49.98
Caduceus-Ph	47.78	44.58	45.93	46.01	48.04	50.02	47.98	47.09	46.93	47.35
HyenaDNA	47.71	47.77	48.06	48.49	50.34	49.67	49.45	48.43	45.68	46.09
DNABERT-2	43.68	44.17	43.29	45.50	43.23	45.29	44.22	45.18	45.40	43.28
NTv2	42.15	44.90	43.20	41.85	44.25	43.80	42.50	45.10	43.05	42.75

Table 17: (Cont.) Results for all models on NT benchmark across genomic regions.

Task/Model	2222	42	43	44	45	46	47	48	49	50
H3										
ConvNova (4 dilation)	81.62	81.78	81.04	82.24	82.12	82.06	81.36	81.12	80.91	80.69
ConvNova (1 dilation)	76.90	77.14	77.28	77.73	76.80	76.59	77.24	77.26	77.68	76.96
Caduceus-Ph	80.29	79.85	78.63	81.47	80.24	81.50	81.35	80.35	79.95	81.02
HyenaDNA	76.88	77.01	78.23	77.24	79.84	78.39	77.56	79.27	79.11	77.90
DNABERT-2	79.61	79.34	79.12	78.90	79.02	79.61	78.68	79.42	79.99	79.41
NTv2	76.55	80.32	78.30	77.52	80.71	79.25	76.89	77.82	76.94	77.38
H3K4me1										
ConvNova (4 dilation)	58.88	56.99	56.37	56.20	56.70	55.59	55.71	55.71	56.48	57.41
ConvNova (1 dilation)	55.67	57.8	56.62	57.64	55.28	55.65	55.80	56.32	57.44	55.82
Caduceus-Ph	53.31	53.00	54.54	51.87	53.40	53.42	52.34	51.91	52.01	52.45
HyenaDNA	43.04	44.54	44.04	44.55	46.02	44.9	41.93	45.26	46.13	44.77
DNABERT-2	50.32	47.21	48.09	49.38	49.22	43.71	46.91	50.22	47.82	50.51
NTv2	51.56	50.96	52.35	51.81	50.52	52.03	52.31	52.72	51.25	50.93
H3K4me2										
ConvNova (4 dilation)	55.80	54.68	51.81	52.52	54.51	52.20	51.30	55.01	54.91	54.45
ConvNova (1 dilation)	56.53	59.72	57.26	58.60	56.13	56.35	58.72	56.25	58.67	56.24
Caduceus-Ph	47.96	47.23	52.47	50.14	51.35	50.14	52.18	50.01	48.72	48.56
HyenaDNA	40.5	40.02	42.43	43.32	41.80	44.07	43.78	42.39	44.82	43.65
DNABERT-2	43.10	42.10	45.20	41.90	43.01	40.10	43.80	44.40	43.40	43.20
NTv2	37.76	36.32	37.59	37.57	35.56	37.87	39.49	37.04	36.35	36.86
H3K4me3										
ConvNova (4 dilation)	59.23	62.11	60.42	59.98	59.82	60.10	60.58	58.73	61.45	59.60
ConvNova (1 dilation)	67.13	66.8	67.97	67.79	66.45	67.01	66.75	68.00	67.42	66.22
Caduceus-Ph	56.93	56.09	58.64	54.56	57.95	57.84	58.54	56.49	54.14	56.03
HyenaDNA	47.65	52.13	50.12	52.10	52.32	51.43	49.46	49.72	47.26	51.90
DNABERT-2	46.30	46.40	46.60	46.00	45.80	45.60	46.40	44.80	44.30	42.10
NTv2	51.70	48.94	50.52	49.17	52.03	50.15	49.87	51.34	50.75	48.53
H3K9ac										
ConvNova (4 dilation)	68.98	69.98	66.18	67.54	68.39	66.18	67.54	68.39	68.67	69.10
ConvNova (1 dilation)	63.85	65.33	67.12	65.15	66.24	63.66	65.38	66.82	65.17	66.20
Caduceus-Ph	63.78	62.28	64.20	65.04	61.06	60.98	65.24	63.59	63.55	62.95
HyenaDNA	57.57	57.51	58.37	60.25	58.57	59.78	59.19	59.31	57.53	56.88
DNABERT-2	58.80	60.40	61.00	58.82	60.41	60.98	58.77	60.78	61.23	59.20
NTv2	60.45	61.90	62.22	59.98	60.85	61.57	62.10	60.02	61.78	59.65
H3K14ac										
ConvNova (4 dilation)	66.24	66.61	65.91	68.03	65.98	65.83	65.6	66.58	64.93	66.22
ConvNova (1 dilation)	69.76	73.03	69.26	72.57	69.95	69.88	71.03	69.46	72.23	69.92
Caduceus-Ph	61	59.36	62.4	60.7	62.14	57.9	62.83	61.31	59.65	61.08
HyenaDNA	56.08	57.59	54.5	54.31	57.31	55.19	58.26	57.09	57.95	58.85
DNABERT-2	56.31	55.72	53.42	55.14	55.61	55.36	49.52	54.54	53.81	55.72
NTv2	55.03	58.65	56.74	59.18	57.01	55.82	57.33	58.40	56.10	57.92
H3K36me3										
ConvNova (4 dilation)	67.12	67.58	68.44	69.43	68.07	68.54	69.13	67.75	67.85	69.22
ConvNova (1 dilation)	69.31	67.01	66.1	66.43	67.17	68.31	66.66	66.24	66.59	67.01
Caduceus-Ph	62.53	61.53	61.16	62.3	59.68	61.42	61.46	60.33	60.81	59.93
HyenaDNA	60.27	60.52	59.8	59.01	58.88	59.66	60.73	60.09	60.98	59.3
DNABERT-2	59.12	58.11	58.11	57.16	56.14	57.15	58.25	58.21	58.65	55.27
NTv2	58.75	61.42	60.15	62.10	59.81	61.68	60.25	58.90	60.77	61.12
H3K79me3										
ConvNova (4 dilation)	72.24	71.37	71.05	70.85	72.41	72.73	72.67	72.19	72.35	72.91
ConvNova (1 dilation)	70.99	71.83	69.34	72.57	71.17	70.66	71.85	70.23	71.24	71.67
Caduceus-Ph	68.78	68.28	67.77	68.02	65.14	66.89	66.37	66.49	66.58	67.42
HyenaDNA	62.6	65.71	67.33	66.82	66.97	65.52	65.23	67.68	67.63	66.99
DNABERT-2	64.10	64.10	65.10	64.10	64.12	65.13	64.32	64.21	64.82	63.91
NTv2	64.23	66.14	67.22	65.99	63.87	65.43	68.12	64.68	66.79	65.31

Table 18: (Cont.) Results for all models on NT benchmark across genomic regions.

Task/Model	2222	42	43	44	45	46	47	48	49	50
H4										
ConvNova (4 dilation)	80.44	80.97	82.36	80.76	81.37	81.17	81.62	80.19	80.74	81.54
ConvNova (1 dilation)	77.29	76.05	77.06	76.83	78.82	76.29	76.88	77.42	76.81	79.06
Caduceus-Ph	80.51	80.02	80.21	79.94	81.12	79.72	79.36	79.82	80.25	80.33
HyenaDNA	78.33	77.61	77.60	77.84	77.85	77.96	79.73	78.58	79.24	76.79
DNABERT-2	78.82	78.42	77.83	78.83	78.42	77.26	78.62	78.23	77.54	78.14
NTv2	78.53	80.21	79.65	79.11	81.20	78.92	79.76	81.00	80.45	79.83
H4ac										
ConvNova (4 dilation)	64.78	65.04	62.44	64.47	66.65	63.18	64.95	66.23	65.63	64.14
ConvNova (1 dilation)	64.9	66.89	66.77	66.9	65.68	65.32	66.14	65.77	66.82	65.77
Caduceus-Ph	58.21	60.68	60.68	60.69	58.96	57.55	59.52	60.47	60.27	55.59
HyenaDNA	53.77	53.99	51.30	57.10	55.75	54.30	51.32	55.36	55.34	53.22
DNABERT-2	51.82	51.92	51.73	51.83	51.92	51.73	51.83	51.92	51.73	51.83
NTv2	53.68	56.94	54.72	57.41	54.20	55.84	56.10	54.95	53.02	55.35

A.6 RELATIVE OCCUPANCY OF YEAST HISTONE MARKS




Figure 6: **Relative occupancy of histone marks on chromosome IV.** A) Histone H3K14ac vs H3 YPD ChIP chip ratios B) Histone H3K4me2 vs H3 YPD ChIP chip ratios C) Histone H3K4me3 vs H3 YPD ChIP chip ratios D) Histone H3K9ac vs H3 YPD ChIP chip ratios. Relative occupancy of histone marks on chromosome IV. Data were obtained from the Yeast Genome Database. For complete datasets and additional information, please refer to <https://www.yeastgenome.org/dataset/E-WMIT-3#resources>.