

---

# Variable Star Light Curves in Koopman Space

---

Anonymous Authors<sup>1</sup>

## Abstract

Interest for applying machine learning in astronomical object classification has been growing with forthcoming huge surveys. Here, we put forward a methodical approach to analyzing variable star light curves through the application of Koopman theory-based modern data-driven techniques for dynamical system analysis. We employ this method on light curves associated to RR Lyrae stars in the Galactic globular cluster  $\omega$  Centauri. Curves are thus summarized by a handful of complex eigenvalues, corresponding to oscillatory or fading modes. In contrast with RRab variables, we find that RRc variables are defined in terms of fewer eigenvalues, which reflects the simpler structure of their light curves. Additionally, we show how Blazhko variables may be identified using relevant eigenvalues and that a physical interpretation of the related modes may be obtained in terms of the Blazhko effect.

## 1. Introduction

In the context of time-domain astronomy and stellar variability, machine learning techniques are growing in popularity, particularly in light of the substantial amounts of data generated by current (Kaiser, 2004; Borucki et al., 2010; Drake et al., 2013; 2014; Gaia Collaboration et al., 2016; Jayasinghe et al., 2018; van Roestel et al., 2021; Coughlin et al., 2021) and planned ground- and space-based observing facilities. In the study of variable stars, these techniques have primarily been applied directly to light curves, which are represented by appropriately defined features (e.g. see Pashchenko et al. 2018; Mahabal et al. 2019; Dékány et al. 2021; Barbara et al. 2022 for optical light curves, but also e.g. Kovačević et al. 2022 for X-ray variable sources); deep neural networks have also been employed (e.g. Jamal & Bloom, 2020; Bassi et al., 2021). Although these meth-

ods produce good predictions, they are typically not interpretable and do not represent the physics behind the stellar sources that match to the observed light curves.

Here, we are interested in inherently interpretable techniques that, even though they remain fully data driven, allow us to get closer to the physics of the dynamical system producing the data. An example along these lines can be found in (Pasquato et al., 2022), where the authors use the sparse identification of nonlinear dynamics (SINDy) method to automatically identify governing equations for RRab, RRc, and  $\delta$ -Scuti variables based on observed light curves. The primary obstacle in this methodology is identifying an appropriate functional basis for the representation of the dynamical system. Indeed, this choice is frequently difficult in practice: in fact, Pasquato et al. (2022) essentially failed at modelling RRab light curves using a polynomial basis. It follows that, at this stage, featurizing light curves based on the coefficients learned by SINDy seems unfeasible.

In this paper, we go beyond this limitation employing the Dynamic Mode Decomposition (DMD) algorithm, a mathematical tool introduced by (Schmid, 2010), and to the best of our knowledge, seldom used in astronomy before (Darling & Widrow, 2019; Widrow et al., 2020; Linkmann et al., 2020; Hori et al., 2020; Albidah et al., 2021; Linkmann & Dikpati, 2021; Albidah et al., 2022). The advantage of this technique is that the extracted dynamic modes can be naturally interpreted as a generalization of global stability modes and can be potentially linked to the underlying physical mechanisms captured in the data sequence. In particular, we model the temporal evolution of the optical magnitude of a sample of RR Lyrae stars belonging to the globular cluster  $\omega$  Centauri. We automatically learn approximations for nonlinear dynamical systems encoding the time-evolution of the magnitude of each variable star and verify that the predicted evolution matches the observed data. Such approximations are expressed in terms of linear systems in suitable Koopman spaces, an eigenvector base of which is returned directly by the algorithm. The associated eigenvalues essentially summarize the light curve and can be in principle used as features for machine learning tasks. We find that the Koopman space required to describe RRc variable dynamics is smaller with respect to the one needed for encoding RRab variables dynamics. This reflects the simpler and more regular silhouettes of RRc curves and

---

<sup>1</sup>Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

Preliminary work. Under review by ICML AI4Science Workshop 2024. . Do not distribute.

provides a straightforward heuristic for RRab vs RRC classification. Finally, we identify stars affected by the Blazhko effect (long term variability; Blažko, 1907) by comparing the eigenvalues corresponding to light curves measured at different stages in the Blazhko cycle.

## 2. Photometric data

For our work, we adopted the optical (BVI-band) photometry of the RRLs in the Globular cluster  $\omega$  Centauri published by Braga et al. (2016; 2019). This data set is homogeneous in terms of photometric data reduction, even though it includes images collected during more than 20 years from seven different 1-to-8 m class telescopes, in a sky area of  $57' \times 56'$  (see Sec.2 in Braga et al. 2016 for more details). Still, to further improve the consistency of our analysis, we chose to adopt only the photometric data collected at the same telescope (Danish 1.5m, at La Silla), during five years (1995 to 1999), which constitutes the largest part (almost 60%) of the original data set. The resulting Danish subset is a truly homogeneous data set, with the exception of sporadic positional offsets in the 1998 images. As a result, the following analysis is based on light curves with 100 to over 2000 phase points in at least two of the three bands (V and B, since I-band phase points are one order of magnitude less) for 125 variables.

This subset of the data is comprised of light curves taken from 89 RRC and 70 RRab variables. Of these, 6 RRC and 2 RRab light curves were discarded because we failed to obtain an acceptable interpolation, due either to outliers or phase gaps. This leaves us with 83 RRC and 68 RRab light curves. Using the *gcv spline* Python wrapper (Le Losq & Feng, 2007) of a FORTRAN package for generalized, cross-validatory spline smoothing (Woltring, 1986), we interpolate these light curves repeated 5 times and picked equidistant (in phase) snapshots from this spline with 30 points per phase which makes a smoothed out, periodic set of 150 points from which DMD can build a model.

## 3. Methods

Dynamic Mode Decomposition (DMD) algorithm is a dimensionality reduction algorithm guaranteed in the framework of Koopman analysis for the study of nonlinear dynamical systems. The core principle of Koopman analysis is a non linear coordinate transformation that embeds a nonlinear dynamical system in an equivalent linear dynamical system of observables. The existence of such an embedding is due to an intrinsic trade-off between linearity and dimension. In this framework, it's necessary to figure out what the coordinate transformation is and what the dynamical system is turned into. For our purpose, we are interested to do this from data in one shot. This is possible thanks to

tangent linear approximation of the system and a few arguments used in iterative methods for computing solutions to linear eigenvalue problems, that DMD algorithm naturally encodes.

More precisely, given a discrete time series  $V_1^N = \{v_1, \dots, v_N\}$ , where  $v_i \in \mathbb{R}^m \ \forall i \in [m]$ , it is assumed the existence of a linear mapping  $A \in GL_n$  that associates two successive points of the time series, i.e.  $v_{i+1} = Av_i$ . This last relation defines a linear dynamical system that stays approximately the same over the sampling period, formally using the language of matrices  $V_2^N = AV_1^{N-1} + re_{N-1}^T$ , where  $r$  is the vector of residuals that encodes the behaviour not predicted by  $A$ ,  $e_{N-1}$  is the  $(N-1)$ th vector of the canonical base and  $V_1^{N-1} = \{v_1, \dots, v_{N-1}\}$  and  $V_2^N = \{v_1, \dots, v_N\}$ . Roughly speaking, DMD computes the eigenvalues and the eigenvectors of  $A$ . Computing eigenvalues and eigenvectors is an elementary task for which several subroutines can be used; commonly, DMD uses Singular Value Decomposition (SVD) algorithm. Eigenvalues and the eigenvectors of  $A$  completely define a set of modes each of which is associated with a fixed oscillation frequency and decay or growth rate. These modes are a natural generalization of the global stability modes and inherit their physical interpretation of encodings of damped or driven sinusoidal behavior in time. It's easy to deduce that the number of eigenvectors provides an important measure of the "hardness" of transforming a nonlinear system into a linear one and of the "distance" from a sinusoidal silhouette.

In this work, we deal with a low dimensional dynamical system, for this reason, we employ the extension of DMD known as *Higher Order Dynamical Mode Decomposition* (HODMD; Le Clainche & Vega, 2017). The modification introduced by HODMD is the following: instead of assuming  $v_{i+1} = Av_i$ , we consider an higher order Koopman assumption, i.e. that  $v_{i+d} = A_1v_i + A_2v_{i+1} + \dots + A_dv_{i+d-1}$  where  $d \geq 1$  is a tunable parameter. It is not difficult to notice that we are simply superimposing DMD on a sliding window over the time series. The main advantage of such an approach lies in enabling the computation of *higher order modes* enriching the space returned by classical DMD when considering dimensional systems presenting low *spatial complexity* but a large number of hidden frequencies.

In our experiments, the time series  $V_1^N = \{v_1, \dots, v_n\}$  is given by the uniformly sampled value of magnitude measured at regular intervals over the interpolated light curve as described above.

### 3.1. Implementation details

We use the library PyDMD (Demo et al., 2018). We take 30 snapshots per phase (for a total of 150 snapshots) so that enough snapshot are fed to DMD to produce a model that fits with the original data but not too much as to model

certain parts of the spline that may be due to noise.

## 4. Results

The DMD reconstruction of the light curve for an example RRab is shown in Fig. 1 and for an RRC in Fig. 2.

### 4.1. Separating RRab from RRC variables

As we vary the Koopman dimension  $d$ , which corresponds to the maximum allowable number of eigenvectors, we find that DMD models from RRC and RRab variables respond differently to increasing  $d$ . In fact, as the spatial dimension increases, the normalized mean squared error (nMSE; the definition is given in (Pasquato et al., 2022) as the ratio between the mean squared error (MSE) of the reconstructed light curve and the MSE of a constant fit) for models of RRC variables drops off steadily compared to models of RRab variables which have a more or less constant nMSE until about  $d = 25$  where we can observe a sharp drop. This is shown in Fig. 3 in the top panel and, for the number of eigenvalues, in the bottom panel. Note how nMSE starts decreasing immediately with  $d$  for the RRC and keeps decreasing in a linear fashion, while it suddenly drops at about  $d = 25$  for the RRab. From observing this phenomenon, we can choose a nMSE threshold (the nMSE is calculated 10 phases away from the last snapshot to promote stability of the DMD model) such that, between RRC and RRab variables, different numbers of spatial dimensions (and different numbers of eigenvalues) are sufficient to produce models which yield an error equal or lower than the threshold.

We find that, for the nMSE to be below 0.45, models for RRC variables require between 11 and 26 spatial dimensions ( $d$ ) and between 3 and 11 eigenvalues. For RRab variables, between 23 and 29 spatial dimensions ( $d$ ) and between 7 and 13 eigenvalues are needed to yield an nMSE below threshold. We notice that there is some overlap especially in the numbers of eigenvalues. Despite that, this difference leads to a straightforward classification algorithm. In fact, we can find that, most likely, a variable star is RRC if  $d < 24$  is sufficient to yield an nMSE below 0.45 and is RRab otherwise. The histogram of the values of  $d$  needed to achieve nMSE= 0.45 is shown in Fig. 4.

The relevant confusion matrix for the associated classifier with  $d < 24$  is shown in Tab. 1. This is calculated on the same data set on which the threshold  $d = 24$  was chosen, so the associated measure of accuracy (97%) is optimistically biased and should not be used to predict performance on unseen data. However this is still an indication that the modes identified by DMD are naturally suited to classification.

	$d < 24$	$d \geq 24$
RRab	1	67
RRC	79	4

Table 1. Confusion matrix for a classifier based on the minimum Koopman dimension  $d$  needed to achieve nMSE= 0.45.




Figure 4. Top panel: histogram of the minimum values of  $d$  needed to achieve nMSE = 0.45 (or below) for the RRab (black) and the RRC variables (red). Right panel: same, but for the number of eigenvalues.

One more point to note is that classification based on the Koopman dimension  $d$  needed to achieve a given level of mean squared error (nMSE=0.45), which depends on the shape of the light curve, overlaps quite well with the distinction between RRab and RRC variables in the Bailey diagram, as shown in Fig. 5. Discrepancies between the two could be leveraged to single out potentially interesting stars, among which perhaps mixed-mode oscillators.

A different way to look at the data presented in Fig. 5 is shown in Fig. 9, where we plot  $d$  as a function of the amplitude of the magnitude variation of each star (top panel) and as a function of the period (right panel). Additionally, in Fig. 10 we show the number of eigenvalues as a function of the same two quantities. Regarding the dependence on amplitude, both  $d$  and the number of eigenvalues increase with amplitude. Since both are measures of the complexity of the light curve shape, this is an expected result: higher amplitude oscillators become increasingly anharmonic. However,




Figure 1. DMD reconstruction (orange dashed line) of the interpolated light curve (blue points) compared with the original (red points) and time shifted (green points) raw data for RRab star V59.




Figure 2. DMD reconstruction (orange dashed line) of the interpolated light curve (blue points) compared with the original (red points) and time shifted (green points) raw data for RRc star V103.

the increase is largely driven, in both cases, by the fact that large amplitude variables are RRab and small amplitude variables are RRc. Within each class the trend is weaker but still visible for the RRab, and absent for the RRc. Regarding period, the general trend is again increasing for both quantities, but it is decreasing for the RRab. This means that even though RRab stars typically have both a longer period and a higher number of eigenvalues than RRc, within the class of the RRab an increasing period corresponds to a *lower* number of eigenvalues, i.e. a simpler light curve.

This evidence fully supports theoretical and empirical findings concerning the variation of RRab pulsation properties across the instability strip. The RRab display a steady decrease when moving from the blue to the red edge of the instability strip. The variation also applies to the shape of the light curves, indeed RRab close to the blue edge display a sawtooth shape and a more sinusoidal shape approaching the red edge. The main culprit causing this difference is convection, since it becomes more efficient when moving from hotter (bluer) to cooler (redder) effective temperatures and eventually quenches the pulsation activity (Bono & Stellingwerf, 1993).

In terms of physical interpretation, it is particularly interesting that the MSE drops for RRab when  $d$  approaches 30, the number of points sampled along a period by our interpo-

lation strategy. Since the data we fed to HODMD repeats exactly after one period, we have that  $x_{31} = x_1$  and in general  $x_{30+k} = x_k$  for every  $k$ . Thus the vector to which DMD is effectively applied at  $t + 1$  is  $\mathbf{x}_{t+1} = (x_2, x_3, \dots, x_{30}, x_1)$  and at  $t$  it is  $\mathbf{x}_t = (x_1, x_2, \dots, x_{29}, x_{30})$  and the matrix  $\mathbf{A}$  such that  $\mathbf{x}_{t+1} = \mathbf{A}\mathbf{x}_t$  becomes simply  $[\mathbf{e}_{30}, \mathbf{e}_1, \dots, \mathbf{e}_{29}]$ . The eigenvalues of this matrix are  $\lambda_k = e^{ki\pi/30}$  for  $k = 0, \dots, 29$ , so they are equally spaced on the unit circle and include  $\lambda_0 = 1$ . We can thus conclude that the DMD is predicting the future evolution of the light curve by leveraging solely the periodicity of the data and nothing else; in an intuitive sense we can claim that an important fraction of the RRab curves is ‘incompressible’.

#### 4.2. Blazhko effect

A final application of this description that goes beyond distinguishing fundamental and first overtone pulsators is related to Blazhko effect. DMD summarizes a light curve with a handful of complex numbers: do these differ over time for a Blazhko variable, and how? In Fig. 6 we show the DMD eigenvalues calculated during the first part of the observation period (top panel) and during the second part (right panel). Most eigenvalues match exactly except for a very limited number.

These eigenvalues may correspond to Koopman modes






Figure 3. Left panel: nMSE achieved by the DMD reconstruction against the original light curve as a function of the spatial dimension of the HODMD snapshot ( $d$  parameter in pyDMD). The black solid line represents the median curve for the RRab stars, the red solid line for the RRc stars. The lower and upper boundaries of the black and red shaded areas represent the first and third quartiles respectively. Right panel: Same as the left panel but nMSE is plotted as a function of the number of eigenvalues.




Figure 5. Bailey diagram with the Koopman dimension  $d$  needed to attain  $nMSE = 0.45$  color coded.




Figure 6. Top panel: DMD eigenvalues (red points) for variable V5 calculated between Julian time 2449858.7698 and 2449869.7601, corresponding to the beginning of the observation period. The green dashed line corresponds to the unit circle in the complex plane. Middle panel: DMD eigenvalues (blue points) for variable V5 calculated between Julian time 2450971.5723 and 2450985.6906, corresponding to the end of the observation period. Right panel: DMD eigenvalues that differ between the two.

specifically associated to Blazhko evolution, since V5 is a suspected Blazhko. In any case DMD has revealed a secular change in its light curve. Potentially, this may be used to automatically detect Blazhko variables without the need for direct visual inspection of the light curve. For comparison, in Fig. 8 and 7 we show the same for two more stars that are deemed affected by the Blazhko effect.




Figure 7. Top panel: DMD eigenvalues (red points) for variable V120 calculated between Julian time 2449858.7468 and 2449869.7635, corresponding to the beginning of the observation period. The green dashed line corresponds to the unit circle in the complex plane. Middle panel: DMD eigenvalues (blue points) for variable V120 calculated between Julian time 2450970.5019 and 2450985.6906, corresponding to the end of the observation period. Right panel: DMD eigenvalues that differ between the two.




Figure 8. Top panel: DMD eigenvalues (red points) for variable V115 calculated between Julian time 2449858.7468 and 2449869.7635, corresponding to the beginning of the observation period. The green dashed line corresponds to the unit circle in the complex plane. Middle panel: DMD eigenvalues (blue points) for variable V115 calculated between Julian time 2450971.5723 and 2450985.6906, corresponding to the end of the observation period. Right panel: DMD eigenvalues that differ between the two.




Figure 9. Top panel: Koopman space dimension  $d$  as a function of amplitude for RRab (black points) and RRC (red points). Right panel: Koopman space dimension  $d$  as a function of period, same color coding as the top panel.




Figure 10. Top panel: number of eigenvalues as a function of amplitude for RRab (black points) and RRC (red points). Right panel: number of eigenvalues as a function of period, same color coding as the top panel. It is known that short period, large amplitude RRab are harder to fit with fourier and in our case they also need more eigenvalues.

### 5. Conclusions

DMD and related algorithms have recently found application to fluid dynamics and other branches of physics that deal with high-dimensional snapshots either from observations or from simulations. In our application to variable stars, snapshots at a given time contain at most a few values, corresponding to magnitudes in various filters and possibly

their derivatives. We thus applied the HODMD algorithm, which essentially synthesizes a higher-dimension snapshot from delayed coordinates. This is the first application of DMD-related methods to variable star light curves to date. This was done with the goal of obtaining features from light curves in a principled way that could be amenable to physical interpretation. In DMD the observed dynamics is decomposed into modes, similar to normal modes in a coupled oscillator system; each mode corresponds to an eigenvalue, associated either to exponential decay (or increase) or to periodic oscillations.

We obtained a reconstructed light curve as a linear combination of the individual evolution of such modes, and compared it with the original data over a sample of 151 stars. We measured the discrepancy between the reconstructed light curve and the original by means of the mean squared error normalized to that of a constant fit. We are able to obtain consistently good fits (with  $nMSE$  under  $10^{-2}$ ) if we let the number of eigenvalues increase as needed. We found that simpler light curves, such as those of RRC variables, require only a limited number of eigenvalues. Conversely RRab stars, which have more complex light curve shapes, require the inclusion of more eigenvalues to be described with good accuracy. We have quantified this, finding that the Koopman dimension  $d$  needed to achieve a given normalized MSE in reconstructing the curve behaves differently for RRab and RRC variables. We used this as a basis for a classifier, choosing a threshold of 0.45 normalized MSE, to achieve which the modal RRab star requires  $d = 27$  (resulting in a mode of 11 eigenvalues) and the modal RRC star,  $d = 15$  (corresponding to a mode of 7 eigenvalues). The threshold value 0.45 corresponds to the maximum (over all the light curves) of the minimum normalized MSE reached.

We found that the number of eigenvalues that needs to be included to obtain a good representation of the curve varies with amplitude and period in accord to the fact that RRab and RRC stars occupy distinctive loci in the period-amplitude plane.

While the DMD decomposition bears some resemblance to Fourier methods, which have been extensively applied to time domain astronomy and variable star light curves in particular, the advantage of DMD is to identify modes that are potentially physically meaningful. This is the case of the long term period variability due to the Blazhko effect, where indeed a few eigenvalues found by DMD change between the beginning and the end of the observation period for the confirmed Blazhko variables in our data set. Whether this is indeed an indication of DMD finding specific modes associated to Blazhko evolution will be discussed in further detail in upcoming work.

## References

- Albidah, A. B., Brevis, W., Fedun, V., Ballai, I., Jess, D. B., Stangalini, M., Higham, J., and Verth, G. Proper orthogonal and dynamic mode decomposition of sunspot data. *Philosophical Transactions of the Royal Society of London Series A*, 379(2190):20200181, February 2021. doi: 10.1098/rsta.2020.0181.
- Albidah, A. B., Fedun, V., Aldhafeeri, A. A., Ballai, I., Brevis, W., Jess, D. B., Higham, J., Stangalini, M., Silva, S. S. A., and Verth, G. Magnetohydrodynamic Wave Mode Identification in Circular and Elliptical Sunspot Umbrae: Evidence for High-order Modes. *ApJ*, 927(2): 201, March 2022. doi: 10.3847/1538-4357/ac51d9.
- Barbara, N. H., Bedding, T. R., Fulcher, B. D., Murphy, S. J., and Van Reeth, T. Classifying Kepler light curves for 12 000 A and F stars using supervised feature-based machine learning. *MNRAS*, 514(2):2793–2804, August 2022. doi: 10.1093/mnras/stac1515.
- Bassi, S., Sharma, K., and Gomekar, A. Classification of Variable Stars Light Curves using Long Short Term Memory Network. *Frontiers in Astronomy and Space Sciences*, 8:168, November 2021. doi: 10.3389/fspas.2021.718139.
- Blažko, S. Mitteilung über veränderliche Sterne. *Astronomische Nachrichten*, 175:325, August 1907. doi: 10.1002/asna.19071752002.
- Bono, G. and Stellingwerf, R. F. Cepheid convective models. *Mem. Soc. Astron. Italiana*, 64:559, January 1993.
- Borucki, W. J., Koch, D., Basri, G., Batalha, N., Brown, T., Caldwell, D., Caldwell, J., Christensen-Dalsgaard, J., Cochran, W. D., DeVore, E., Dunham, E. W., Dupree, A. K., Gautier, T. N., Geary, J. C., Gilliland, R., Gould, A., Howell, S. B., Jenkins, J. M., Kondo, Y., Latham, D. W., Marcy, G. W., Meibom, S., Kjeldsen, H., Lissauer, J. J., Monet, D. G., Morrison, D., Sasselov, D., Tarter, J., Boss, A., Brownlee, D., Owen, T., Buzasi, D., Charbonneau, D., Doyle, L., Fortney, J., Ford, E. B., Holman, M. J., Seager, S., Steffen, J. H., Welsh, W. F., Rowe, J., Anderson, H., Buchhave, L., Ciardi, D., Walkowicz, L., Sherry, W., Horch, E., Isaacson, H., Everett, M. E., Fischer, D., Torres, G., Johnson, J. A., Endl, M., MacQueen, P., Bryson, S. T., Dotson, J., Haas, M., Kolodziejczak, J., Van Cleve, J., Chandrasekaran, H., Twicken, J. D., Quintana, E. V., Clarke, B. D., Allen, C., Li, J., Wu, H., Tenenbaum, P., Verner, E., Bruhweiler, F., Barnes, J., and Prsa, A. Kepler Planet-Detection Mission: Introduction and First Results. *Science*, 327(5968):977, February 2010. doi: 10.1126/science.1185402.
- Braga, V. F., Stetson, P. B., Bono, G., Dall’Ora, M., Ferraro, I., Fiorentino, G., Freyhammer, L. M., Iannicola, G., Marengo, M., Neeley, J., Valenti, E., Buonanno, R., Calamida, A., Castellani, M., da Silva, R., Degl’Innocenti, S., Di Cecco, A., Fabrizio, M., Freedman, W. L., Giuffrida, G., Lub, J., Madore, B. F., Marconi, M., Marinoni, S., Matsunaga, N., Monelli, M., Persson, S. E., Piersimoni, A. M., Pietrinferni, A., Prada-Moroni, P., Pulone, L., Stellingwerf, R., Tognelli, E., and Walker, A. R. On the RR Lyrae Stars in Globulars. IV.  $\omega$  Centauri Optical UBVRI Photometry. *AJ*, 152:170, December 2016. doi: 10.3847/0004-6256/152/6/170.
- Braga, V. F., Stetson, P. B., Bono, G., Dall’Ora, M., Ferraro, I., Fiorentino, G., Iannicola, G., Inno, L., Marengo, M., Neeley, J., Beaton, R. L., Buonanno, R., Calamida, A., Contreras Ramos, R., Chaboyer, B., Fabrizio, M., Freedman, W. L., Gilligan, C. K., Johnston, K. V., Lub, J., Madore, B. F., Magurno, D., Marconi, M., Marinoni, S., Marrese, P. M., Mateo, M., Matsunaga, N., Minniti, D., Monson, A. J., Monelli, M., Nonino, M., Persson, S. E., Pietrinferni, A., Sneden, C., Storm, J., Walker, A. R., Valenti, E., and Zoccali, M. New near-infrared JHK<sub>s</sub> light-curve templates for RR Lyrae variables. *A&A*, 625: A1, May 2019. doi: 10.1051/0004-6361/201834893.
- Coughlin, M. W., Burdge, K., Duev, D. A., Katz, M. L., van Roestel, J., Drake, A., Graham, M. J., Hillenbrand, L., Mahabal, A. A., Masci, F. J., Mróz, P., Prince, T. A., Yao, Y., Bellm, E. C., Burruss, R., Dekany, R., Jaodand, A., Kaplan, D. L., Kupfer, T., Laher, R. R., Riddle, R., Rigault, M., Rodriguez, H., Rusholme, B., and Zolkower, J. The ZTF Source Classification Project - II. Periodicity and variability processing metrics. *MNRAS*, 505(2): 2954–2965, August 2021. doi: 10.1093/mnras/stab1502.
- Darling, K. and Widrow, L. M. Eigenfunctions of Galactic phase space spirals from dynamic mode decomposition. *MNRAS*, 490(1):114–123, November 2019. doi: 10.1093/mnras/stz2539.
- Dékány, I., Grebel, E. K., and Pojmański, G. Metallicity Estimation of RR Lyrae Stars From Their I-Band Light Curves. *ApJ*, 920(1):33, October 2021. doi: 10.3847/1538-4357/ac106f.
- Demo, N., Tezzele, M., and Rozza, G. Pydmd: Python dynamic mode decomposition. *Journal of Open Source Software*, 3(22):530, 2018.
- Drake, A. J., Catelan, M., Djorgovski, S. G., Torrealba, G., Graham, M. J., Belokurov, V., Koposov, S. E., Mahabal, A., Prieto, J. L., Donalek, C., Williams, R., Larson, S., Christensen, E., and Beshore, E. Probing the Outer Galactic Halo with RR Lyrae from the Catalina Surveys. *ApJ*, 763(1):32, January 2013. doi: 10.1088/0004-637X/763/1/32.



- 440 Drake, A. J., Graham, M. J., Djorgovski, S. G., Catelan,  
441 M., Mahabal, A. A., Torrealba, G., García-Álvarez, D.,  
442 Donalek, C., Prieto, J. L., Williams, R., Larson, S., Chris-  
443 ten sen, E., Belokurov, V., Koposov, S. E., Beshore, E.,  
444 Boattini, A., Gibbs, A., Hill, R., Kowalski, R., John-  
445 son, J., and Shelly, F. The Catalina Surveys Periodic  
446 Variable Star Catalog. *ApJS*, 213(1):9, July 2014. doi:  
447 10.1088/0067-0049/213/1/9.
- 448
- 449 Gaia Collaboration, Brown, A. G. A., Vallenari, A., Prusti,  
450 T., de Bruijne, J. H. J., Mignard, F., Drimmel, R., Babu-  
451 siaux, C., Bailer-Jones, C. A. L., Bastian, U., Biermann,  
452 M., Evans, D. W., Eyer, L., Jansen, F., Jordi, C., Katz,  
453 D., Klioner, S. A., Lammers, U., Lindegren, L., Luri, X.,  
454 O’Mullane, W., Panem, C., Pourbaix, D., Randich, S.,  
455 Sartoretti, P., Siddiqui, H. I., Soubiran, C., Valette, V.,  
456 van Leeuwen, F., Walton, N. A., Aerts, C., Arenou, F.,  
457 Cropper, M., Høg, E., Lattanzi, M. G., Grebel, E. K., Hol-  
458 land, A. D., Huc, C., Passot, X., Perryman, M., Bramante,  
459 L., Cacciari, C., Castañeda, J., Chaoul, L., Cheek, N., De  
460 Angeli, F., Fabricius, C., Guerra, R., Hernández, J., Jean-  
461 Antoine-Piccolo, A., Masana, E., Messineo, R., Mowlavi,  
462 N., Nienartowicz, K., Ordóñez-Blanco, D., Panuzzo, P.,  
463 Portell, J., Richards, P. J., Riello, M., Seabroke, G. M.,  
464 Tanga, P., Thévenin, F., Torra, J., Els, S. G., Gracia-  
465 Abril, G., Comoretto, G., Garcia-Reinaldos, M., Lock,  
466 T., Mercier, E., Altmann, M., Andrae, R., Astraatmadja,  
467 T. L., Bellas-Velidis, I., Benson, K., Berthier, J., Blomme,  
468 R., Busso, G., Carry, B., Cellino, A., Clementini, G.,  
469 Cowell, S., Creevey, O., Cuypers, J., Davidson, M., De  
470 Ridder, J., de Torres, A., Delchambre, L., Dell’Oro, A.,  
471 Ducourant, C., Frémat, Y., García-Torres, M., Gosset, E.,  
472 Halbwachs, J. L., Hambly, N. C., Harrison, D. L., Hauser,  
473 M., Hestroffer, D., Hodgkin, S. T., Huckle, H. E., Hut-  
474 ton, A., Jasniewicz, G., Jordan, S., Kontizas, M., Korn,  
475 A. J., Lanzafame, A. C., Manteiga, M., Moitinho, A.,  
476 Muinonen, K., Osinde, J., Pancino, E., Pauwels, T., Pe-  
477 tit, J. M., Recio-Blanco, A., Robin, A. C., Sarro, L. M.,  
478 Siopis, C., Smith, M., Smith, K. W., Sozzetti, A., Thuil-  
479 lot, W., van Reeve, W., Viala, Y., Abbas, U., Abreu  
480 Aramburu, A., Accart, S., Aguado, J. J., Allan, P. M.,  
481 Allasia, W., Altavilla, G., Álvarez, M. A., Alves, J., An-  
482 derson, R. I., Andrei, A. H., Anglada Varela, E., Antiche,  
483 E., Antoja, T., Antón, S., Arcay, B., Bach, N., Baker,  
484 S. G., Balaguer-Núñez, L., Barache, C., Barata, C., Bar-  
485 bier, A., Barblan, F., Barrado y Navascués, D., Barros,  
486 M., Barstow, M. A., Becciani, U., Bellazzini, M., Bello  
487 García, A., Belokurov, V., Bendjoya, P., Berihuete, A.,  
488 Bianchi, L., Bienaymé, O., Billebaud, F., Blagorodnova,  
489 N., Blanco-Cuaresma, S., Boch, T., Bombrun, A., Bor-  
490 rachero, R., Bouquillon, S., Bourda, G., Bouy, H., Bra-  
491 gaglia, A., Breddels, M. A., Brouillet, N., Brüsemeister,  
492 T., Bucciarelli, B., Burgess, P., Burgon, R., Burlacu, A.,  
493 Busonero, D., Buzzi, R., Caffau, E., Cambras, J., Camp-  
494 bell, H., Cancelliere, R., Cantat-Gaudin, T., Carlucci,  
T., Carrasco, J. M., Castellani, M., Charlot, P., Char-  
nas, J., Chiavassa, A., Clotet, M., Cocozza, G., Collins,  
R. S., Costigan, G., Crifo, F., Cross, N. J. G., Crosta,  
M., Crowley, C., Dafonte, C., Damerджи, Y., Dapergo-  
las, A., David, P., David, M., De Cat, P., de Felice, F.,  
de Laverny, P., De Luise, F., De March, R., de Mar-  
tino, D., de Souza, R., Debosscher, J., del Pozo, E.,  
Delbo, M., Delgado, A., Delgado, H. E., Di Matteo,  
P., Diakite, S., Distefano, E., Dolding, C., Dos Anjos,  
S., Drazinos, P., Duran, J., Dzigan, Y., Edvardsson, B.,  
Enke, H., Evans, N. W., Eynard Bontemps, G., Fabre,  
C., Fabrizio, M., Faigler, S., Falcão, A. J., Farràs Casas,  
M., Federici, L., Fedorets, G., Fernández-Hernández, J.,  
Fernique, P., Fienga, A., Figueras, F., Filippi, F., Find-  
eisen, K., Fonti, A., Fouesneau, M., Fraile, E., Fraser, M.,  
Fuchs, J., Gai, M., Galleti, S., Galluccio, L., Garabato,  
D., García-Sedano, F., Garofalo, A., Garralda, N., Gavras,  
P., Gerssen, J., Geyer, R., Gilmore, G., Girona, S., Giuf-  
frida, G., Gomes, M., González-Marcos, A., González-  
Núñez, J., González-Vidal, J. J., Granvik, M., Guerrier,  
A., Guillout, P., Guiraud, J., Gúrpide, A., Gutiérrez-  
Sánchez, R., Guy, L. P., Haignon, R., Hatzidimitriou,  
D., Haywood, M., Heiter, U., Helmi, A., Hobbs, D., Hof-  
mann, W., Holl, B., Holland, G., Hunt, J. A. S., Hypki,  
A., Icardi, V., Irwin, M., Jevardat de Fombelle, G., Jofré,  
P., Jonker, P. G., Jorissen, A., Julbe, F., Karampelas,  
A., Kochoska, A., Kohley, R., Kolenberg, K., Kontizas,  
E., Koposov, S. E., Kordopatis, G., Koubisky, P., Krone-  
Martins, A., Kudryashova, M., Kull, I., Bachchan, R. K.,  
Lacoste-Seris, F., Lanza, A. F., Lavigne, J. B., Le Poncin-  
Lafitte, C., Lebreton, Y., Lebzelter, T., Leccia, S., Leclerc,  
N., Lecoœur-Taïbi, I., Lemaitre, V., Lenhardt, H., Ler-  
oux, F., Liao, S., Licata, E., Lindstrøm, H. E. P., Lister,  
T. A., Livanou, E., Lobel, A., Löffler, W., López, M.,  
Lorenz, D., MacDonald, I., Magalhães Fernandes, T.,  
Managau, S., Mann, R. G., Mantelet, G., Marchal, O.,  
Marchant, J. M., Marconi, M., Marinoni, S., Marrese,  
P. M., Marschalkó, G., Marshall, D. J., Martín-Fleitas,  
J. M., Martino, M., Mary, N., Matijevič, G., Mazeh, T.,  
McMillan, P. J., Messina, S., Michalik, D., Millar, N. R.,  
Miranda, B. M. H., Molina, D., Molinaro, R., Molinaro,  
M., Molnár, L., Moniez, M., Montegriffo, P., Mor, R.,  
Mora, A., Morbidelli, R., Morel, T., Morgenthaler, S.,  
Morris, D., Mulone, A. F., Muraveva, T., Musella, I.,  
Narbonne, J., Nelemans, G., Nicastro, L., Noval, L.,  
Ordénovic, C., Ordieres-Meré, J., Osborne, P., Pagani, C.,  
Pagano, I., Pailler, F., Palacin, H., Palaversa, L., Parsons,  
P., Pecoraro, M., Pedrosa, R., Pentikäinen, H., Pichon,  
B., Piersimoni, A. M., Pineau, F. X., Plachy, E., Plum,  
G., Poujoulet, E., Prša, A., Pulone, L., Ragaini, S., Rago,  
S., Rambaux, N., Ramos-Lerate, M., Ranalli, P., Rauw,  
G., Read, A., Regibo, S., Reylé, C., Ribeiro, R. A., Ri-  
moldini, L., Ripepi, V., Riva, A., Rixon, G., Roelens,

- 495 M., Romero-Gómez, M., Rowell, N., Royer, F., Ruiz-  
496 Dern, L., Sadowski, G., Sagristà Sellés, T., Sahlmann,  
497 J., Salgado, J., Salguero, E., Sarasso, M., Savietto, H.,  
498 Schultheis, M., Sciacca, E., Segol, M., Segovia, J. C.,  
499 Segransan, D., Shih, I. C., Smareglia, R., Smart, R. L.,  
500 Solano, E., Solitro, F., Sordo, R., Soria Nieto, S., Souchay,  
501 J., Spagna, A., Spoto, F., Stampa, U., Steele, I. A., Stei-  
502 delmüller, H., Stephenson, C. A., Stoev, H., Suess, F. F.,  
503 Stüveges, M., Surdej, J., Szabados, L., Szegedi-Elek, E.,  
504 Tapiador, D., Taris, F., Tauran, G., Taylor, M. B., Teix-  
505 eira, R., Terrett, D., Tingley, B., Trager, S. C., Turon, C.,  
506 Ulla, A., Utrilla, E., Valentini, G., van Elteren, A., Van  
507 Hemelryck, E., van Leeuwen, M., Varadi, M., Vecchiato,  
508 A., Veljanoski, J., Via, T., Vicente, D., Vogt, S., Voss,  
509 H., Votruba, V., Voutsinas, S., Walmsley, G., Weiler, M.,  
510 Weingrill, K., Wevers, T., Wyrzykowski, Ł., Yoldas, A.,  
511 Žerjal, M., Zucker, S., Zurbach, C., Zwitter, T., Alecu,  
512 A., Allen, M., Allende Prieto, C., Amorim, A., Anglada-  
513 Escudé, G., Arsenijevic, V., Azaz, S., Balm, P., Beck, M.,  
514 Bernstein, H. H., Bigot, L., Bijaoui, A., Blasco, C., Bon-  
515 figli, M., Bono, G., Boudreault, S., Bressan, A., Brown,  
516 S., Brunet, P. M., Bunclark, P., Buonanno, R., Butkevich,  
517 A. G., Carret, C., Carrion, C., Chemin, L., Chéreau, F.,  
518 Corcione, L., Darmigny, E., de Boer, K. S., de Teodoro,  
519 P., de Zeeuw, P. T., Delle Lucche, C., Domingues, C. D.,  
520 Dubath, P., Fodor, F., Frézouls, B., Fries, A., Fustes, D.,  
521 Fyfe, D., Gallardo, E., Gallegos, J., Gardiol, D., Gebran,  
522 M., Gomboc, A., Gómez, A., Grux, E., Gueguen, A.,  
523 Heyrovsky, A., Hoar, J., Iannicola, G., Isasi Parache, Y.,  
524 Janotto, A. M., Joliet, E., Jonckheere, A., Keil, R., Kim,  
525 D. W., Klagyivik, P., Klar, J., Knude, J., Kochukhov,  
526 O., Kolka, I., Kos, J., Kutka, A., Lainey, V., LeBouquin,  
527 D., Liu, C., Loreggia, D., Makarov, V. V., Marseille,  
528 M. G., Martayan, C., Martinez-Rubi, O., Massart, B.,  
529 Meynadier, F., Mignot, S., Munari, U., Nguyen, A. T.,  
530 Nordlander, T., Ocvirk, P., O’Flaherty, K. S., Olias Sanz,  
531 A., Ortiz, P., Osorio, J., Oszkiewicz, D., Ouzounis, A.,  
532 Palmer, M., Park, P., Pasquato, E., Peltzer, C., Peralta, J.,  
533 Péturaud, F., Pieniluoma, T., Pigozzi, E., Poels, J., Prat,  
534 G., Prod’homme, T., Raison, F., Rebordao, J. M., Riskey,  
535 D., Rocca-Volmerange, B., Rosen, S., Ruiz-Fuertes, M. I.,  
536 Russo, F., Sembay, S., Serraller Vizcaino, I., Short, A.,  
537 Siebert, A., Silva, H., Sinachopoulos, D., Slezak, E.,  
538 Soffel, M., Sosnowska, D., Straižys, V., ter Linden, M.,  
539 Terrell, D., Theil, S., Tiede, C., Troisi, L., Tsalmanza, P.,  
540 Tur, D., Vaccari, M., Vachier, F., Valles, P., Van Hamme,  
541 W., Veltz, L., Virtanen, J., Wallut, J. M., Wichmann, R.,  
542 Wilkinson, M. I., Ziaeeepour, H., and Zschocke, S. Gaia  
543 Data Release 1. Summary of the astrometric, photometric,  
544 and survey properties. *A&A*, 595:A2, November 2016.  
545 doi: 10.1051/0004-6361/201629512.
- 546  
547 Hori, K., Tobias, S. M., and Teed, R. J. Dynamic mode  
548 decomposition to retrieve torsional Alfvén waves. *arXiv*  
549 *e-prints*, art. arXiv:2009.13095, September 2020.
- Jamal, S. and Bloom, J. S. On Neural Architectures for  
Astronomical Time-series Classification with Application  
to Variable Stars. *ApJS*, 250(2):30, October 2020. doi:  
10.3847/1538-4365/aba8ff.
- Jayasinghe, T., Kochanek, C. S., Stanek, K. Z., Shappee,  
B. J., Holoien, T. W. S., Thompson, T. A., Prieto, J. L.,  
Dong, S., Pawlak, M., Shields, J. V., Pojmanski, G., Otero,  
S., Britt, C. A., and Will, D. The ASAS-SN catalogue of  
variable stars I: The Serendipitous Survey. *MNRAS*, 477  
(3):3145–3163, July 2018. doi: 10.1093/mnras/sty838.
- Kaiser, N. Pan-STARRS: a wide-field optical survey tele-  
scope array. In Oschmann, Jacobus M., J. (ed.), *Ground-  
based Telescopes*, volume 5489 of *Society of Photo-  
Optical Instrumentation Engineers (SPIE) Conference Se-  
ries*, pp. 11–22, October 2004. doi: 10.1117/12.552472.
- Kovačević, M., Pasquato, M., Marelli, M., De Luca, A., Sal-  
vaterra, R., and Belfiore, A. Exploring X-ray variability  
with unsupervised machine learning. I. Self-organizing  
maps applied to XMM-Newton data. *A&A*, 659:A66,  
March 2022. doi: 10.1051/0004-6361/202142444.
- Le Clainche, S. and Vega, J. M. Higher order dynamic  
mode decomposition. *SIAM Journal on Applied Dy-  
namical Systems*, 16(2):882–925, 2017. doi: 10.1137/  
15M1054924. URL [https://doi.org/10.1137/  
15M1054924](https://doi.org/10.1137/15M1054924).
- Le Losq, C. and Feng, Y. *gcv spline*. [https://github.  
com/charlesll/gcvspline](https://github.com/charlesll/gcvspline), 2007.
- Linkmann, M. and Dikpati, M. Symmetry-reduced dynamic  
mode decomposition of solar tachocline dynamics. In  
*AGU Fall Meeting Abstracts*, volume 2021, pp. SH53C–  
06, December 2021.
- Linkmann, M., Dallas, S., and Dikpati, M. Data-driven  
low-dimensional modelling of solar tachocline dynamics.  
In *AGU Fall Meeting Abstracts*, volume 2020, pp. SH006–  
02, December 2020.
- Mahabal, A., Rebbapragada, U., Walters, R., Masci, F. J.,  
Blagorodnova, N., van Roestel, J., Ye, Q.-Z., Biswas,  
R., Burdge, K., Chang, C.-K., Duev, D. A., Golkhou,  
V. Z., Miller, A. A., Nordin, J., Ward, C., Adams, S.,  
Bellm, E. C., Branton, D., Bue, B., Cannella, C., Con-  
nolly, A., Dekany, R., Feindt, U., Hung, T., Fortson,  
L., Frederick, S., Fremling, C., Gezari, S., Graham, M.,  
Groom, S., Kasliwal, M. M., Kulkarni, S., Kupfer, T., Lin,  
H. W., Lintott, C., Lunnan, R., Parejko, J., Prince, T. A.,  
Riddle, R., Rusholme, B., Saunders, N., Sedaghat, N.,  
Shupe, D. L., Singer, L. P., Soumagnac, M. T., Szkody,  
P., Tachibana, Y., Tirumala, K., van Velzen, S., and

550 Wright, D. Machine Learning for the Zwicky Transient  
551 Facility. *PASP*, 131(997):038002, March 2019. doi:  
552 10.1088/1538-3873/aaf3fa.

553 Pashchenko, I. N., Sokolovsky, K. V., and Gavras, P. Ma-  
554 chine learning search for variable stars. *MNRAS*, 475(2):  
555 2326–2343, April 2018. doi: 10.1093/mnras/stx3222.

557 Pasquato, M., Abbas, M., Trani, A. A., Nori, M., Kwiecin-  
558 ski, J. A., Trevisan, P., Braga, V. F., Bono, G., and Macciò,  
559 A. V. Sparse Identification of Variable Star Dynamics.  
560 *ApJ*, 930(2):161, May 2022. doi: 10.3847/1538-4357/  
561 ac5624.

563 Schmid, P. J. Dynamic mode decomposition of numerical  
564 and experimental data. *Journal of Fluid Mechanics*, 656:  
565 5–28, 2010. doi: 10.1017/S0022112010001217.

566 van Roestel, J., Duev, D. A., Mahabal, A. A., Coughlin,  
567 M. W., Mróz, P., Burdge, K., Drake, A., Graham, M. J.,  
568 Hillenbrand, L., Bellm, E. C., Kupfer, T., Delacroix, A.,  
569 Fremling, C., Golkhou, V. Z., Hale, D., Laher, R. R.,  
570 Masci, F. J., Riddle, R., Rosnet, P., Rusholme, B., Smith,  
571 R., Soumagnac, M. T., Walters, R., Prince, T. A., and  
572 Kulkarni, S. R. The ZTF Source Classification Project. I.  
573 Methods and Infrastructure. *AJ*, 161(6):267, June 2021.  
574 doi: 10.3847/1538-3881/abe853.

576 Widrow, L. M., Darling, K., and Li, H. Warps, Waves,  
577 and Phase Spirals in the Milky Way. In Valluri, M. and  
578 Sellwood, J. A. (eds.), *Galactic Dynamics in the Era of*  
579 *Large Surveys*, volume 353, pp. 65–70, January 2020. doi:  
580 10.1017/S1743921319009049.

581  
582 Woltring, H. J. A fortran package for generalized, cross-  
583 validatory spline smoothing and differentiation. *Advances*  
584 *in Engineering Software (1978)*, 8(2):104–113, 1986.

585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604