000 ON THE ERGODIC CONVERGENCE PROPERTIES OF THE 001 PEACEMAN-RACHFORD METHOD AND THEIR APPLI-002 003 CATIONS IN SOLVING LINEAR PROGRAMMING 004

Anonymous authors

005 006

007

008 009 010

011 012

013

014

015

016

017

018

019

021

022

023

024

025

026

027

028

029

030 031 032

033 034

035

043

046

047 048

Paper under double-blind review

ABSTRACT

In this paper, we study the ergodic convergence properties of the Peaceman-Rachford (PR) method with semi-proximal terms for solving convex optimization problems (COPs). By reformulating the PR method as a degenerate proximal point method, for the first time we establish the global convergence of the ergodic sequence generated by the PR method with broadly chosen semi-proximal terms under the assumption that there exists a Karush-Kuhn-Tucker (KKT) solution to the COPs. This result represents a significant departure from previous studies on the non-ergodic convergence of the PR method, which typically requires strong convexity (or strong monotonicity in the reformulated operator) conditions that are hardly satisfied for COPs. Moreover, we establish an ergodic iteration complexity of O(1/k) of the PR method with semi-proximal terms, measured by the objective error, the feasibility violation, and the KKT residual using the ε subdifferential. Based on these convergence properties, we introduce the solver EPR-LP, using the ergodic sequence of the PR method with semi-proximal terms for solving linear programming (LP) problems. EPR-LP incorporates an adaptive restart strategy and dynamic penalty parameter updates for efficiency and robustness. Extensive numerical experiments on LP benchmark datasets, executed on a high-performance GPU, show that our Julia-based solver outperforms the awardwinning solver PDLP at a tolerance level of 10^{-8} .

1 INTRODUCTION

In this paper, we focus on solving the following linear programming (LP) problem:

$$\min_{\substack{c \in \mathbb{R}^n}} \langle c, x \rangle$$
s.t. $A_1 x = b_1$
 $A_2 x \ge b_2$
 $x \in C,$

$$(1)$$

where $A_1 \in \mathbb{R}^{m_1 \times n}$, $A_2 \in \mathbb{R}^{m_2 \times n}$, $b_1 \in \mathbb{R}^{m_1}$, $b_2 \in \mathbb{R}^{m_2}$, and $c \in \mathbb{R}^n$. The set C is defined as $C := \{x \in \mathbb{R}^n \mid l \leq x \leq u\}$, with the vectors $l \in (\mathbb{R} \cup \{-\infty\})^n$ and $u \in (\mathbb{R} \cup \{+\infty\})^n$. Let 042 $A = [A_1; A_2] \in \mathbb{R}^{m \times n}$ with $m = m_1 + m_2$ and $b = [b_1; b_2] \in \mathbb{R}^m$. We assume that A is a non-zero 044 matrix, which is also occasionally treated as a linear operator. Then, the dual of problem (1) can be 045 expressed as:

$$\min_{\substack{y \in \mathbb{R}^n, z \in \mathbb{R}^n}} - \langle b, y \rangle + \delta_D(y) + \delta_C^*(-z)$$
s.t. $A^*y + z = c$,
(2)

049 where $\delta_D(\cdot)$ is the indicator function over $D := \{y = (y_1, y_2) \in \mathbb{R}^{m_1} \times \mathbb{R}^{m_2}_+\}$ and $\delta_C^*(\cdot)$ is the conjugate of $\delta_C(\cdot)$. LP is a fundamental optimization subject in applied mathematics, operations 051 research, and computer science, with a wide range of applications. The interior point methods and the simplex methods are the standard algorithms used in commercial LP solvers such as Gurobi 052 (Gurobi Optimization, LLC, 2024) and CPLEX (IBM, 1987), efficiently solving problems with hundreds to millions of variables and constraints. However, their performance can be inadequate for 054 problems with huge dimensions (Lu, 2024). Furthermore, these methods are difficult to parallelize, limiting their ability to leverage modern GPUs effectively. 056

Recently, the first-order methods, especially those based on the alternating direction method of 057 multipliers (ADMM) (O'donoghue et al., 2016; Stellato et al., 2020; Applegate et al., 2021; Lin et al., 2021; O'Donoghue, 2021; Applegate et al., 2023; Deng et al., 2024; Lu & Yang, 2024; Chen et al., 2024) have attracted increasing attention for solving large-scale LP problems, due to their low 060 iteration cost and ease of parallelization. Specifically, let $\sigma > 0$ be a given penalty parameter. The 061 augmented Lagrangian function associated with the dual problem (2), for any $(y, z, x) \in \mathbb{R}^m \times$ 062 $\mathbb{R}^n \times \mathbb{R}^n$, is defined as

$$L^{\mathrm{LP}}_{\sigma}(y,z;x) := -\langle b,y \rangle + \delta_D(y) + \delta^*_C(-z) + \langle x, A^*y + z - c \rangle + \frac{\sigma}{2} \|A^*y + z - c\|^2.$$

065 A preconditioned (semi-proximal) ADMM (pADMM) (Xiao et al., 2018) for solving LP is then 066 outlined in Algorithm 1. Consider the case where $S_1 = 0$. When $\rho = 1$, Algorithm 1 reduces to the 067 Douglas-Rachford method (Gabay, 1983). When $\rho = 2$, it becomes equivalent to the generalized 068 ADMM (GADMM) induced by the Peaceman-Rachford (PR) method (Eckstein & Bertsekas, 1992; 069 Lions & Mercier, 1979). For a more detailed comparison of Algorithm 1 with other algorithms, 070 refer to Xiao et al. (2018) and Sun et al. (2024). 071

Algorithm 1 A pADMM method for the LP problem (2)

- 1: Input: Set the penalty parameter $\sigma > 0$. Choose $\rho \in (0, 2]$. Let $\mathcal{S}_1 : \mathbb{R}^m \to \mathbb{R}^m$ be a selfadjoint positive semidefinite linear operator such that $S_1 + AA^*$ is positive definite. Choose an initial point $w^0 = (y^0, z^0, x^0) \in D \times \operatorname{dom}(\delta^*_C(-\cdot)) \times \mathbb{R}^n$.

2: for k = 0, 1, ..., do3: Step 1. $\bar{z}^k = \underset{z \in \mathbb{R}^n}{\operatorname{arg\,min}} \{ L^{\mathrm{LP}}_{\sigma} (y^k, z; x^k) \};$ 4: Step 2. $\bar{x}^k = x^k + \sigma(A^* y^k + \bar{z}^k - c);$ 5: Step 3. $\bar{y}^k = \underset{y \in \mathbb{R}^m}{\operatorname{arg\,min}} \{ L^{\mathrm{LP}}_{\sigma} (y, \bar{z}^k; \bar{x}^k) + \frac{\sigma}{2} \|y - y^k\|_{\mathcal{S}_1}^2 \};$ 6: Step 4. $w^{k+1} = (1 - \rho)w^k + \rho \bar{w}^k;$

082 083

063 064

072

073

074

075

076 077

079 080 081

A key difference between HPR-LP and PDLP is the choice of ρ . PDLP uses a conservative $\rho = 1$ 104 and adopts a line-search strategy to enhance performance, while HPR-LP employs a more aggressive 105

106

103

¹The authors of Applegate et al. (2021; 2023) received the Beale–Orchard-Hays Prize for Excellence in

⁰⁸⁴ Notably, Applegate et al. (Applegate et al., 2021; 2023) developed the award-winning solver PDLP¹, 085 which uses a modified version of the primal-dual hybrid gradient (PDHG) method (Zhu & Chan, 2008) as its base algorithm. This modified PDHG is a special case of Algorithm 1 with $\rho = 1$ (Esser 087 et al., 2010; Chambolle & Pock, 2011). To enhance the modified PDHG's performance, PDLP incor-880 porates several effective techniques, including using the ergodic iterate as a restart point, employing an adaptive update rule for the penalty parameter σ , and implementing a line search strategy. It is 089 worth noting that, compared to non-ergodic sequences, the ergodic sequence of the semi-proximal 090 ADMM (sPADMM) (Fazel et al., 2013), including the modified PDHG (Esser et al., 2010; Cham-091 bolle & Pock, 2011), achieves a better O(1/k) iteration complexity with respect to the objective 092 error and the feasibility violation (Cui et al., 2016). Consequently, the GPU implementation of PDLP (cuPDLP.jl (Lu & Yang, 2023) and cuPDLP-c (Lu et al., 2023)) has demonstrated advantages 094 over commercial LP solvers like Gurobi (Gurobi Optimization, LLC, 2024) and COPT (Ge et al., 2022), particularly for large-scale LP problems. 096

More recently, Chen et al. (Chen et al., 2024) introduced HPR-LP, an implementation of the Halpern Peaceman-Rachford (HPR) method (Sun et al., 2024) with semi-proximal terms for solving LP 098 problems. This method integrates the Halpern iteration (Halpern, 1967; Lieder, 2021) into the 099 PR method (Lions & Mercier, 1979; Eckstein & Bertsekas, 1992) with semi-proximal terms, cor-100 responding to Algorithm 1 using $\rho = 2$. It achieves an iteration complexity of O(1/k) for the 101 Karush-Kuhn-Tucker (KKT) residual and the objective error. The Julia implementation of HPR-LP 102 has demonstrated superior performance compared to PDLP on classical LP benchmark datasets.

Computational Mathematical Programming at the 25th International Symposium on Mathematical Program-107 ming (https://ismp2024.gerad.ca/), held from July 21-26, 2024, in Montréal, Canada.

0

-1

-2

-3

-4

-5

-6

-7 L 0

 $log_{10}(optimality)$

108 $\rho = 2$. It is well known that the PR method ($\rho = 2$) is faster than the DR method ($\rho = 1$) when the PR method converges (Lions & Mercier, 1979). This difference in ρ selection motivates our study of using the ergodic sequence of the PR method with semi-proximal terms to solve LP problems.

eraodic

nonergodic

5

imes10⁶



113 114

115

116 117

118

119 120

121 122 123

> Figure 1: The performance of PR, using ergodic and non-ergodic sequences, in solving the LP instance "datt256" from Mittelmann's LP benchmark set. The optimality is evaluated based on the relative primal-dual infeasibility and the relative duality gap.

2

iter

3

1

127

137

138

139 140

141

142 143

144

145

146

147

128 To the best of our knowledge, the ergodic convergence of the PR method with semi-proximal terms 129 (corresponding to Algorithm 1 with $\rho = 2$) is still unknown. In this paper, we address this gap by proving, for the first time, the global convergence of the ergodic sequence using the theory of the 130 degenerate proximal point method (dPPM) (Bredies et al., 2022). This result marks a significant 131 departure from previous studies on the non-ergodic convergence of the PR method, which typically 132 relies on strong convexity that is not met in LP (see Figure 1). Moreover, we also investigate the 133 ergodic iteration complexity of the pADMM in Algorithm 1 in terms of the objective error, the 134 feasibility violation, and the KKT residual based on ε -subdifferential (Rockafellar, 1970). The main 135 contributions of this paper can be summarized as follows: 136

- For a more general convex optimization problem (COP), by reformulating the pADMM with $\rho \in (0, 2]$ as a dPPM, we establish the ergodic convergence of the dPPM, thereby proving the ergodic convergence of the pADMM through this reformulation.
- For solving COP, we establish the ergodic iteration complexity of O(1/k) for the pADMM with $\rho \in (0, 2]$ in terms of the objective error, the feasibility violation, and the KKT residual based on ε -subdifferential.
- We propose the solver EPR-LP, an implementation of the ergodic sequence of **PR** method (corresponding to Algorithm 1 with $\rho = 2$) to solve **LP** problems, incorporating an adaptive restart strategy and dynamic penalty parameter updates. Extensive numerical experiments on LP benchmark datasets using a GPU demonstrate the superior performance of the Julia implementation of EPR-LP compared to PDLP (Lu & Yang, 2023).

The remainder of this paper is structured as follows: Section 2 establishes the ergodic convergence and iteration complexity of pADMM for solving convex optimization problems. The detailed implementation of EPR-LP is discussed in Section 3. Section 4 presents the results of numerical experiments on LP benchmark datasets. Finally, we conclude the paper in Section 5.

153 **Notation.** Let \mathbb{U} , \mathbb{W} , \mathbb{X} , \mathbb{Y} , and \mathbb{Z} be finite-dimensional real Euclidean spaces, each equipped 154 with an inner product $\langle \cdot, \cdot \rangle$ and its corresponding norm $\|\cdot\|$. For any convex function $f: \mathbb{X} \to \mathcal{X}$ $(-\infty, +\infty]$, we define its effective domain as dom $(f) := \{x \in \mathbb{X} : f(x) < \infty\}$, its con-155 jugate as $f^*(z) := \sup_{x \in \mathbb{X}} \{ \langle x, z \rangle - f(x) \}, z \in \mathbb{X}$, and its proximal mapping as $\operatorname{Prox}_f(x) :=$ 156 $\arg\min_{z\in\mathbb{X}} \left\{ f(z) + \frac{1}{2} \| ||z-x||^2 \right\}, x \in \mathbb{X}.$ Furthermore, consider a closed convex set $C \subseteq \mathbb{X}$. We 157 define the distance from $x \in \mathbb{X}$ to C as $\operatorname{dist}(x, C) := \inf_{z \in C} ||z - x||$, and we express the Euclidean 158 projection of x onto C as $\Pi_C(x) := \arg \min\{||x - z|| \mid z \in C\}$. Moreover, for a linear operator 159 $A: \mathbb{X} \to \mathbb{Y}$, we denote its adjoint by A^* and express its spectral norm as $||A|| := \sup_{||x|| \le 1} ||Ax||$. 160 Finally, for any self-adjoint, positive semidefinite linear operator $\mathcal{M} : \mathbb{X} \to \mathbb{X}$, we define the semi-161 norm as $||x||_{\mathcal{M}} := \sqrt{\langle x, \mathcal{M}x \rangle}$ for any $x \in \mathbb{X}$.

162 2 ERGODIC CONVERGENCE PROPERTIES OF THE PADMM

In this section, we study the ergodic convergence properties of the pADMM method, including the PR method, for solving COP. We begin by reformulating the pADMM method as a dPPM. Next, we prove the ergodic convergence of the dPPM, thereby establishing the ergodic convergence of pADMM. Finally, we analyze the ergodic iteration complexity of the pADMM method.

2.1 ERGODIC CONVERGENCE OF PADMM

171 Consider the following COP:

 $\begin{array}{ll} \min_{\substack{y \in \mathbb{Y}, z \in \mathbb{Z} \\ \text{ s.t. }}} & f_1(y) + f_2(z) \\ B_1 y + B_2 z = c, \end{array} \tag{3}$

where $f_1 : \mathbb{Y} \to (-\infty, +\infty]$ and $f_2 : \mathbb{Z} \to (-\infty, +\infty]$ are proper closed convex functions, $B_1 : \mathbb{Y} \to \mathbb{X}$ and $B_2 : \mathbb{Z} \to \mathbb{X}$ are given linear operators, and $c \in \mathbb{X}$ is a given vector. Given a penalty parameter $\sigma > 0$, the augmented Lagrangian function for problem (3) is defined, for any $(y, z, x) \in \mathbb{Y} \times \mathbb{Z} \times \mathbb{X}$, as

$$L_{\sigma}(y,z;x) := f_1(y) + f_2(z) + \langle x, B_1y + B_2z - c \rangle + \frac{\sigma}{2} ||B_1y + B_2z - c||^2.$$

The dual of problem (3) is given by

$$\max_{x \in \mathbb{X}} \left\{ -f_1^*(-B_1^*x) - f_2^*(-B_2^*x) - \langle c, x \rangle \right\}.$$
(4)

Let $w := (y, z, x) \in \mathbb{W} := \mathbb{Y} \times \mathbb{Z} \times \mathbb{X}$. The pADMM method (Xiao et al., 2018) for solving problem (3) is outlined in Algorithm 2.

Algorithm 2 A pADMM for solving COP (3)

Input: Let *T*₁ and *T*₂ be two self-adjoint, positive semidefinite operators on 𝔅 and ℤ, respectively. Select an initial point w⁰ = (y⁰, z⁰, x⁰) ∈ dom(f₁) × dom(f₂) × 𝔅. Set the parameters σ > 0 and ρ ∈ (0, 2].
 for k = 0, 1, ..., do
 Step 1. z^k = argmin {L_σ (y^k, z; x^k) + 1/2 ||z - z^k||²/_{T₂}};
 Step 2. x^k = x^k + σ(B₁y^k + B₂z^k - c);
 Step 3. y^k = argmin {L_σ (y, z^k; x^k) + 1/2 ||y - y^k||²/_{T₁}};
 Step 4. w^{k+1} = (1 - ρ)w^k + ρw^k;

As shown in Rockafellar (1970, Corollary 28.3.1), a pair $(y^*, z^*) \in \mathbb{Y} \times \mathbb{Z}$ is an optimal solution to problem (3) if and only if there exists $x^* \in \mathbb{X}$ such that (y^*, z^*, x^*) satisfies the following KKT system:

$$-B_1^* x^* \in \partial f_1(y^*), \quad -B_2^* x^* \in \partial f_2(z^*), \quad B_1 y^* + B_2 z^* - c = 0, \tag{5}$$

where ∂f_i denotes the subdifferential mapping of f_i for i = 1, 2. To discuss the ergodic convergence of pADMM as outlined in Algorithm 2, we make the following assumption:

Assumption 2.1. *The KKT system* (5) *has a nonempty solution set.*

,

Under Assumption 2.1, solving problem (3) is equivalent to finding $w \in \mathbb{W}$ such that $0 \in \mathcal{T}w$, where the maximal monotone operator \mathcal{T} is defined as

$$\mathcal{T}w = \begin{pmatrix} \partial f_1(y) + B_1^* x\\ \partial f_2(z) + B_2^* x\\ c - B_1 y - B_2 z \end{pmatrix}, \quad \forall w = (y, z, x) \in \mathbb{W}.$$
(6)

On the other hand, since f_1 and f_2 are proper closed convex functions, there exist two self-adjoint, positive semidefinite operators, Σ_{f_1} and Σ_{f_2} , such that for all $y, \hat{y} \in \text{dom}(f_1), \phi \in \partial f_1(y)$, and $\hat{\phi} \in \partial f_1(\hat{y})$, the following hold:

$$f_1(y) \ge f_1(\hat{y}) + \langle \hat{\phi}, y - \hat{y} \rangle + \frac{1}{2} \|y - \hat{y}\|_{\Sigma_{f_1}}^2, \quad \text{and} \quad \langle \phi - \hat{\phi}, y - \hat{y} \rangle \ge \|y - \hat{y}\|_{\Sigma_{f_1}}^2$$

and for all $z, \hat{z} \in \text{dom}(f_2), \varphi \in \partial f_2(z)$, and $\hat{\varphi} \in \partial f_2(\hat{z})$:

$$f_2(z) \ge f_2(\hat{z}) + \langle \hat{\varphi}, z - \hat{z} \rangle + \frac{1}{2} \| z - \hat{z} \|_{\Sigma_{f_2}}^2, \text{ and } \langle \varphi - \hat{\varphi}, z - \hat{z} \rangle \ge \| z - \hat{z} \|_{\Sigma_{f_2}}^2.$$

To ensure that each step of the pADMM is well-defined, we also make the following assumption: **Assumption 2.2.** Both $\Sigma_{f_1} + B_1^*B_1 + \mathcal{T}_1$ and $\Sigma_{f_2} + B_2^*B_2 + \mathcal{T}_2$ are positive definite.

Define the self-adjoint linear operator $\mathcal{M} : \mathbb{W} \to \mathbb{W}$ as follows:

$$\mathcal{M} = \begin{bmatrix} \sigma B_1^* B_1 + \mathcal{T}_1 & 0 & B_1^* \\ 0 & \mathcal{T}_2 & 0 \\ B_1 & 0 & \sigma^{-1} \mathcal{I} \end{bmatrix}.$$
 (7)

According to Sun et al. (2024, Proposition 3.2), we have the following equivalence between Algorithm 2 and the dPPM (Bredies et al., 2022).

Proposition 2.1. Suppose that Assumption 2.2 holds. Consider the operators \mathcal{T} defined in (6) and \mathcal{M} defined in (7). Then the sequence $\{w^k\}$ generated by the pADMM in Algorithm 2 coincides with the sequence $\{w^k\}$ generated by the dPPM as follows:

$$\bar{w}^k = \widehat{\mathcal{T}} w^k = (\mathcal{M} + \mathcal{T})^{-1} \mathcal{M} w^k, \quad w^{k+1} = (1-\rho) w^k + \rho \bar{w}^k \tag{8}$$

with the same initial point $w^0 \in \mathbb{W}$. Additionally, \mathcal{M} is an admissible preconditioner² such that $(\mathcal{M} + \mathcal{T})^{-1}$ is Lipschitz continuous.

We now analyze the ergodic convergence of the dPPM for a general maximal monotone operator \mathcal{T} and an admissible preconditioner \mathcal{M} , encompassing Algorithm 2 for solving the COP problem (3). To this end, we define the following two ergodic sequences:

$$w_a^k := \frac{1}{k+1} \sum_{t=0}^k w^t, \quad \bar{w}_a^k := \frac{1}{k+1} \sum_{t=0}^k \bar{w}^t, \quad \forall k \ge 0,$$
(9)

where the sequences $\{w^t\}$ and $\{\bar{w}^t\}$ are generated by the dPPM in (8). Note that, for a maximal monotone operator $\mathcal{T}: \mathbb{W} \to 2^{\mathbb{W}}$ and $\varepsilon \ge 0$, the ε -enlargement of \mathcal{T} at w (Burachik et al., 1997) is defined as

$$\mathcal{T}^{\varepsilon}(w) = \{ v \in \mathbb{W} : \langle w - w', v - v' \rangle \ge -\varepsilon, \, \forall (w', v') \in \mathsf{gph}(\mathcal{T}) \},\$$

where $gph(\mathcal{T}) = \{(w, v) \in \mathbb{W} \times \mathbb{W} \mid v \in \mathcal{T}w\}$. Using the ε -enlargement of \mathcal{T} , we can derive the following proposition regarding the ergodic convergence properties of the dPPM.

Proposition 2.2. Let $\mathcal{T} : \mathbb{W} \to 2^{\mathbb{W}}$ be a maximal monotone operator with $\mathcal{T}^{-1}(0) \neq \emptyset$, and let *M* be an admissible preconditioner. Then the ergodic sequences $\{\bar{w}_a^k\}$ and $\{w_a^k\}$, generated by the *dPPM* in (8) with $\rho \in (0, 2]$, satisfy the following properties for any $k \ge 0$:

(a)
$$\|\bar{w}_a^k - w_a^k\|_{\mathcal{M}} \le \frac{2}{\rho(k+1)} \|w^0 - w^*\|_{\mathcal{M}}, \quad \forall w^* \in \mathcal{T}^{-1}(0);$$

(b)
$$\mathcal{M}(w_a^k - \bar{w}_a^k) \in \mathcal{T}^{\bar{\varepsilon}_a^k}(\bar{w}_a^k)$$
, where $\bar{\varepsilon}_a^k := \frac{1}{k+1} \sum_{t=0}^k \langle \bar{w}^t - \bar{w}_a^k, w^t - \bar{w}^t \rangle_{\mathcal{M}}$ and

257 258 259

260 261

262

263

264

265

266

267

268

269

255 256

216

217 218

219 220

221

222 223

229

234 235

236

237

242 243 244

248

$$0 \le \bar{\varepsilon}_a^k \le \frac{1}{2\rho(k+1)} \|w^0 - w^*\|_{\mathcal{M}}^2, \quad \forall w^* \in \mathcal{T}^{-1}(0).$$

Remark 2.1. Monteiro & Sim (2018) used a non-Euclidean hybrid proximal extragradient framework to obtain a similar result to Proposition 2.2, which could be adapted to analyze the ergodic iteration complexity of Algorithm 2 with $T_1 = 0$, $T_2 = 0$, and $\rho = 2$. In contrast, Proposition 2.2, developed using the dPPM framework, is more general as it only requires \mathcal{M} to be positive semidefinite. This broader setting allows for the analysis of the ergodic iteration complexity of Algorithm 2 for positive semidefinite linear operators T_1 and T_2 .

²In Bredies et al. (2022), an admissible preconditioner for the operator $\mathcal{T} : \mathbb{W} \to 2^{\mathbb{W}}$ is a linear, bounded, self-adjoint, and positive semidefinite operator $\mathcal{M} : \mathbb{W} \to \mathbb{W}$ such that $\widehat{\mathcal{T}} = (\mathcal{M} + \mathcal{T})^{-1}\mathcal{M}$ is single-valued and has full domain.

Theorem 2.1. Let $\mathcal{T} : \mathbb{W} \to 2^{\mathbb{W}}$ be a maximal monotone operator with $\mathcal{T}^{-1}(0) \neq \emptyset$, and let \mathcal{M} be an admissible preconditioner such that $(\mathcal{M} + \mathcal{T})^{-1}$ is L-Lipschitz continuous. Then, the ergodic sequence $\{\bar{w}_a^k\}$ generated by the dPPM in (8) with $\rho \in (0, 2]$ converges to a point in $\mathcal{T}^{-1}(0)$.

Remark 2.2. Note that we only assume *M* to be positive semidefinite in Theorem 2.1. If *M* is positive definite, one can directly apply Baillon's nonlinear ergodic theorem (Baillon, 1975) to obtain the ergodic convergence.

Consider the ergodic sequence of the pADMM:

$$(y_a^k, z_a^k, x_a^k) := \frac{1}{k+1} \sum_{t=0}^k (y^t, z^t, x^t), \quad (\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k) := \frac{1}{k+1} \sum_{t=0}^k (\bar{y}^t, \bar{z}^t, \bar{x}^t), \quad k \ge 0,$$

where the sequences $\{w^t\} = \{(y^t, z^t, x^t)\}$ and $\{\bar{w}^t\} = \{(\bar{y}^t, \bar{z}^t, \bar{x}^t)\}$ are generated by the pADMM in Algorithm 2. The equivalence established in Proposition 2.1 demonstrates that the ergodic convergence of the dPPM in Theorem 2.1 can be leveraged to derive the ergodic convergence of the pADMM, as stated in the following corollary.

Corollary 2.1. Suppose that Assumptions 2.1 and 2.2 hold. Then the ergodic sequence $\{\bar{w}_a^k\} = \{(\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k)\}$, generated by the pADMM in Algorithm 2, converges to the point $w^* = (y^*, z^*, x^*)$, where (y^*, z^*) is a solution to problem (3), and x^* is a solution to problem (4).

2.2 ERGODIC ITERATION COMPLEXITY OF PADMM

We introduce the concept of the ε -subgradient of a convex function f (Rockafellar, 1970):

Definition 2.1. Let $f : \mathbb{X} \to (-\infty, +\infty]$ be a proper convex function, and let $\bar{x} \in \text{dom}(f)$. Given $\varepsilon \ge 0$, the ε -subgradient of f at \bar{x} is defined as

$$\partial_{\varepsilon} f(\bar{x}) := \{ x^* \in \mathbb{X}^* \mid \langle x^*, x - \bar{x} \rangle \le f(x) - f(\bar{x}) + \varepsilon, \, \forall x \in \mathbb{X} \}$$

Based on the optimality conditions of each subproblem in the pADMM, we derive the following lemma regarding the ergodic sequence $\{\bar{w}_a^k\}$ using the ε -subgradient.

Lemma 2.1. Suppose that Assumptions 2.1 and 2.2 hold. Let $\{(\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k)\}$ be the sequence generated by Algorithm 2 and Let $w^* = (y^*, z^*, x^*)$ be a solution to the KKT system (5). The following things hold: for any $k \ge 0$,

$$\begin{cases} -B_{2}^{*}\bar{x}_{a}^{k} - \mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k}) \in \partial_{\bar{\varepsilon}_{z}^{k}} f_{2}(\bar{z}_{a}^{k}), \\ -B_{1}^{*}(\bar{x}_{a}^{k} + \sigma(B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c)) - \mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k}) \in \partial_{\bar{\varepsilon}_{a}^{k}} f_{1}(\bar{y}_{a}^{k}), \end{cases}$$
(10)

where

$$\begin{cases} \bar{\varepsilon}_{z}^{k} = \frac{1}{k+1} \sum_{t=0}^{k} \langle -B_{2}^{*} \bar{x}^{t} - \mathcal{T}_{2}(\bar{z}^{t} - z^{t}), \bar{z}^{t} - \bar{z}_{a}^{k} \rangle \geq 0, \\ \bar{\varepsilon}_{y}^{k} = \frac{1}{k+1} \sum_{t=0}^{k} \langle -B_{1}^{*}(\bar{x}^{t} + \sigma(B_{1}\bar{y}^{t} + B_{2}\bar{z}^{t} - c)) - \mathcal{T}_{1}(\bar{y}^{t} - y^{t}), \bar{y}^{t} - \bar{y}_{a}^{k} \rangle \geq 0, \end{cases}$$
(11)

and

$$\bar{\varepsilon}_{z}^{k} + \bar{\varepsilon}_{y}^{k} \le \frac{1}{2\rho(k+1)} \|w^{0} - w^{*}\|_{\mathcal{M}}^{2}.$$
(12)

Furthermore, to estimate the objective error, we define

$$h(\bar{y}_a^k, \bar{z}_a^k) := f_1(\bar{y}_a^k) + f_2(\bar{z}_a^k) - f_1(y^*) - f_2(z^*), \quad \forall k \ge 0,$$

where (y^*, z^*) is the limit point of the sequence $\{(\bar{y}_a^k, \bar{z}_a^k)\}$. Then the ergodic iteration complexity of pADMM is established in Theorem 2.2.

Theorem 2.2. Suppose that Assumptions 2.1 and 2.2 hold. Let $\{(\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k)\}$ be the ergodic sequence generated by Algorithm 2 with $\rho \in (0, 2]$. Let $w^* = (y^*, z^*, x^*)$ be a solution to the KKT system (5), and $R_0 = ||w^0 - w^*||_{\mathcal{M}}$. For all $k \ge 0$, the following bound holds:

$$dist \left(0, \partial_{\bar{\varepsilon}_{y}^{k}} f_{1}(\bar{y}_{a}^{k}) + B_{1}^{*} \bar{x}_{a}^{k} \right) + dist \left(0, \partial_{\bar{\varepsilon}_{z}^{k}} f_{2}(\bar{z}_{a}^{k}) + B_{2}^{*} \bar{x}_{a}^{k} \right) + \|B_{1} \bar{y}_{a}^{k} + B_{2} \bar{z}_{a}^{k} - c \|$$

$$\leq \left(\frac{\sigma \|B_{1}^{*}\| + 1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_{2}}\| + \|\sqrt{\mathcal{T}_{1}}\| \right) \frac{2R_{0}}{\rho(k+1)},$$

$$(13)$$

where $ar{arepsilon}_z^k + ar{arepsilon}_y^k \leq rac{1}{2
ho(k+1)} \|w^0 - w^*\|_{\mathcal{M}}^2$. Moreover,

$$\begin{array}{l} 321 \\ 322 \\ 323 \end{array} \qquad \left(\frac{-1}{\sqrt{\sigma}}\|x^*\|\right) \frac{2R_0}{\rho(k+1)} \le h(\bar{y}_a^k, \bar{z}_a^k) \le (R_0 + 4\sqrt{\sigma}\|B_1y^*\|) \frac{R_0}{2\rho(k+1)} + \frac{\|x^0 + \sigma B_1y^0\|^2}{2\rho(k+1)}. \end{array}$$
(14)

324

325

326

327

328

330

331 332 333

334

335 336

337

338

367

372

373

Remark 2.3. For the ergodic iteration complexity about LP, please refer to Appendix B. The pADMM with $\rho \in (0,2)$ achieves only a non-ergodic iteration complexity of $o(1/\sqrt{k})$ with respect to the objective error, feasibility violation, and KKT residual, as established in Appendix C. A detailed comparison of the iteration complexities of related algorithms can be found in Appendix D.

Based on the ergodic complexity results in Theorem 2.2, the optimal choice for ρ is 2, resulting in an ergodic PR (EPR) method with semi-proximal terms. In the next section, we apply this EPR method to solve large-scale LP problems.

3 A PEACEMAN-RACHFORD METHOD USING ERGODIC SEQUENCE FOR SOLVING LP

In this section, we introduce the solver EPR-LP for large-scale LP problems (Algorithm 3), which incorporates a restart strategy and adaptive updates of the penalty parameter σ into the EPR method with semi-proximal terms.

Alg	gorithm 3 EPR-LP: A Peaceman-Rachford method using ergodic sequence for the problem (2)
1:	Input: Let $S_1 : \mathbb{R}^m \to \mathbb{R}^m$ be a self-adjoint, positive semidefinite linear operator such that $S_1 + AA^*$ is positive definite. Choose an initial point $w^{0,0} = (y^{0,0}, z^{0,0}, x^{0,0}) \in D \times \text{dom}(\delta^*_C(-\cdot)) \times \mathbb{R}^n$.
2:	Initialization : Set the outer loop counter $r = 0$, the total loop counter $k = 0$, and the initial penalty parameter $\sigma_0 > 0$.
3:	repeat
4:	initialize the inner loop: set inner loop counter $t = 0$;
5:	repeat
6:	$ar{z}^{r,t} = rgmin\left\{L_{\sigma_r}^{ ext{LP}}\left(y^{r,t},z;x^{r,t} ight) ight\};$
	$z \in \mathbb{R}^n$
7:	$ar{x}^{r,\iota}=x^{r,\iota}+\sigma_r(A^*y^{r,\iota}+ar{z}^{r,\iota}-c);$
8:	$\bar{y}^{r,t} = \operatorname*{argmin}_{y \in \mathbb{R}^m} \left\{ L^{\mathrm{LP}}_{\sigma_r} \left(y, \bar{z}^{r,t}; \bar{x}^{r,t} \right) + \frac{\sigma_r}{2} \ y - y^{r,t} \ _{\mathcal{S}_1}^2 \right\};$
9:	$w^{r,t+1} = 2\bar{w}^{r,t} - w^{r,t};$
10.	$_{aar{v}}r,t = \sum_{t=1}^{t} \frac{1}{aar{v}}r,i.$
10.	$w_a = \sum_{i=0}^{\infty} \frac{1}{i+1} w^{-i}$
11:	t = t + 1, k = k + 1:
12:	until one of the restart criteria holds or termination criteria hold
13:	restart the inner loop: $ au_r = t, w^{r+1,0} = \bar{w}_a^{r, au_r}$,
14:	$\sigma_{r+1} = $ SigmaUpdate $(\bar{w}_a^{r,\tau_r}, w^{r,0}, \mathcal{S}_1, A), r = r+1;$
15:	until termination criteria hold
16.	Output: $\{\bar{w}^{r,t}\}$.

Remark 3.1. If line 9 in Algorithm 3 is modified to $w^{r,t+1} = \bar{w}^{r,t}$, the resulting method is referred to as EDR-LP, which uses the ergodic sequence of the DR method with semi-proximal terms for solving LP. In the numerical experiments, we compare the performance of EPR-LP and EDR-LP to evaluate the impact of the parameter ρ .

3.1 RESTART STRATEGY

The EPR method with semi-proximal terms achieves an ergodic iteration complexity of O(1/k) for the objective error, the feasibility violation, and the KKT condition, as shown in Theorem 2.2. This result, derived from Proposition 2.2, guides the restart criteria using the merit function:

 $R_{r,t} := \|w^{r,t} - w^*\|_{\mathcal{M}}, \quad \forall r \ge 0, \ t \ge 0,$

where w^* is a solution to the KKT system (5). $R_{r,0}$ gives the upper bound from Proposition 2.2. A natural restart occurs when $R_{r,t} \leq \alpha_1 R_{r,0}$, with $\alpha_1 \in (0,1)$. In practice, if α_1 is too small, the algorithm may fail to reduce sufficiently, so we also consider the inner loop length and oscillation in $R_{r,t}$. These ideas are implemented in PDLP with other merit functions (Applegate et al., 2021; Lu & Yang, 2023). In addition, since w^* is unknown, we replace $R_{r,t}$ with the following weighted primal and dual infeasibility:

$$\widetilde{R}_{r,t} = \sqrt{\sigma_r^{-1} \| \Pi_D (b - A \bar{x}_a^{r,t}) \|^2} + \sigma_r \| c - A^* \bar{y}_a^{r,t} - \bar{z}_a^{r,t} \|^2.$$

Consequently, the restart criteria in EPR-LP are defined as follows:

1. Sufficient decay of $R_{r,t}$:

$$\widetilde{R}_{r,t} \le \alpha_1 \widetilde{R}_{r,0}; \tag{15}$$

2. Necessary decay + no local progress of $R_{r,t}$:

$$\widetilde{R}_{r,t} \le \alpha_2 \widetilde{R}_{r,0}$$
 and $\widetilde{R}_{r,t+1} > \widetilde{R}_{r,t};$ (16)

3. Long inner loop:

$$t \ge \alpha_3 k,\tag{17}$$

where $\alpha_1 \in (0, \alpha_2), \, \alpha_2 \in (0, 1)$, and $\alpha_3 \in (0, 1)$. In EPR-LP, we set $\alpha_1 = 0.2, \, \alpha_2 = 0.6$, and $\alpha_3 = 0.2$. Once any of the three restart criteria is met, we restart the inner loop for the (r+1)-th iteration, set $w^{r+1,0} = \bar{w}_a^{r,\tau_r}$, and update σ_{r+1} .

3.2 Update rule for σ

Motivated by the update rule for σ based on the iteration complexity of pADMM proposed in Chen et al. (2024), we update σ_{r+1} at the (r+1)-th restart for any $r \ge 0$ by solving the following optimization problem:

$$\sigma_{r+1} := \arg\min_{\sigma} \left\| w^{r+1,0} - w^* \right\|_{\mathcal{M}}^2,$$
(18)

where $||w^{r+1,0} - w^*||_{\mathcal{M}}$ represents the upper bound of the complexity results in Proposition 2.2 at the (r+1)-th outer loop. A smaller upper bound is expected to lead to a smaller residual $\|\bar{w}_a^{r+1,t} - w^{r+1,t}\|$ $w_a^{r+1,t} \parallel_{\mathcal{M}}$ for any $t \ge 0$. Specializing \mathcal{M} in (7) to Algorithm 1, we derive the following:

$$\sigma_{r+1} = \arg\min_{\sigma} \left\| w^{r+1,0} - w^* \right\|_{\mathcal{M}}^2$$

$$= \sqrt{\frac{\|x^{r+1,0} - x^*\|^2}{\|y^{r+1,0} - y^*\|_{\mathcal{S}_1}^2 + \|A^*(y^{r+1,0} - y^*)\|^2}}.$$
(19)

Since computing $||x^{r+1,0} - x^*||$ and $||y^{r+1,0} - y^*||_{\mathcal{S}_1}^2 + ||A^*(y^{r+1,0} - y^*)||^2$ is not implementable, we approximate these terms in EPR-LP using:

$$\Delta_x := \|\bar{x}_a^{r,\tau_r} - x^{r,0}\| \quad \text{and} \quad \Delta_y := \sqrt{\|\bar{y}_a^{r,\tau_r} - y^{r,0}\|_{\mathcal{S}_1}^2 + \|A^*(\bar{y}_a^{r,\tau_r} - y^{r,0})\|^2}, \tag{20}$$

respectively. Consequently, we update σ_{r+1} as follows:

$$\sigma_{r+1} = \frac{\Delta_x}{\Delta_y}.$$
(21)

Because the approximations Δ_x and Δ_y may deviate significantly from their true values, we im-plement some safeguards for the σ update rule. For more details, see Appendix E. Furthermore, to ensure that the EPR-LP method has an explicit update formula for $\bar{y}^{r,t}$ for solving general LP problems, we select

$$S_1 = \lambda I_m - AA^*$$

with $\lambda \geq \lambda_1(AA^*)$, as proposed in Esser et al. (2010); Chambolle & Pock (2011); Xu & Wu (2011). In this case, the updates of $\bar{y}^{r,t}$ are given by:

426
427
428

$$\bar{y}_{1}^{r,t} = y_{1}^{r,t} + \frac{1}{\lambda} \left(\frac{b_{1}}{\sigma} - A_{1}R_{y} \right),$$

 $\bar{y}_{2}^{r,t} = \Pi_{\mathbb{R}^{m_{2}}_{+}} \left(y_{2}^{r,t} + \frac{1}{\lambda} \left(\frac{b_{2}}{\sigma} - A_{2}R_{y} \right) \right)$

where
$$R_y := \bar{x}^{r,t}/\sigma + (A^*y^{r,t} + \bar{z}^{r,t} - c)$$
. Thus, the penalty parameter σ_{r+1} is updated as:

431
$$\sigma_{r+1} = \frac{1}{\sqrt{\lambda}} \frac{\|\bar{x}_a^{r,r} - x^{r,0}\|}{\|\bar{y}_a^{r,\tau_r} - y^{r,0}\|}.$$

432 4 NUMERICAL EXPERIMENT 433

434 In this section, we evaluate the performance of EPR-LP, EDR-LP, and cuPDLP (Lu & Yang, 2023) 435 on Mittelmann's LP benchmark set³ and LP instances relaxed from MIP problems in the MIPLIB 436 2017 collection (Gleixner et al., 2021). All algorithms are implemented in Julia, with experiments conducted on an NVIDIA A100-SXM4-80GB GPU running CUDA 12.3. Each solver is terminated when the relative primal and dual infeasibility errors, as well as the relative duality gap, reach a 438 tolerance of 10^{-8} , or when the time limit is exceeded. For more details on the experimental setup, 439 please refer to Appendix F. 440

441 442

443

444

445

446

447

448

449

450 451

452

453 454

455

456

472

473 474

475

481

482 483

484

485

437

4.1 MITTELMANN'S LP BENCHMARK SET

Mittelmann's LP benchmark set is a standard benchmark for evaluating the numerical performance of LP solvers. In this experiment, we compare the performance of EPR-LP, EDR-LP, and cuPDLP on 49 publicly available instances from Mittelmann's LP benchmark. The performance profiles (Dolan & Moré, 2002) for solving time on the presolved and unpresolved datasets are shown in Figures 2a and 2b, respectively. The key observations are summarized as follows:

- Compared to EDR-LP and cuPDLP, EPR-LP is the fastest solver on approximately 80% of the problems in the presolved dataset and 65% in the unpresolved dataset.
- Among all the algorithms, EPR-LP demonstrates the best success rate, solving the highest percentage of problems across both the presolved and unpresolved datasets. In particular, EPR-LP solves about 8% more problems than cuPDLP on the presolved dataset.
- To solve 90% of the problems in the presolved dataset, EPR-LP requires twice the time of the best solver, while EDR-LP takes four times as long. This difference likely stems from the ratio of ρ , as suggested by the iteration complexity results in (13) and (14).



Figure 2: Performance profiles of solving time for 49 instances of Mittelmann's LP benchmark set with Gurobi's presolve (a) and without presolve (b).

4.2 MIP RELAXATIONS

476 In this experiment, we evaluate the performance of EPR-LP, EDR-LP, and cuPDLP on 380 LP 477 instances relaxed from MIPLIB 2017 collection (Gleixner et al., 2021), both with and without pre-478 solve. Figures 3a and 3b show the performance profiles for solving time on the presolved and 479 unpresolved datasets, respectively. The key observations are listed below: 480

- Compared to EDR-LP and cuPDLP, EPR-LP is the fastest solver on approximately 85% of the problems in the presolved dataset and 80% in the unpresolved dataset.
- EPR-LP demonstrated a slightly higher success rate than cuPDLP on the presolved dataset and a nearly comparable success rate on the unpresolved dataset.

³https://plato.asu.edu/ftp/lpfeas.html.



Figure 3: Performance profiles of solving time for LP instances relaxed from MIP, with Gurobi's presolve (a) and without presolve (b).

4.3 SUMMARY OF EXPERIMENTS

The previous numerical experiments show that EPR-LP outperforms both EDR-LP and cuPDLP. Specifically, EPR-LP achieves the highest success rates across nearly all datasets. Moreover, EPR-LP emerges as the fastest solver for approximately 65% to 85% of the problems. Since the periteration times of EDR-LP and EPR-LP are nearly the same, EPR-LP's advantage over EDR-LP stems from using $\rho = 2$, leading to fewer iterations. To further underscore EPR-LP's advantages over cuPDLP, we present the per-iteration time ratio between these two solvers in Table 1. The median per-iteration time ratio, ranging from 2.7 to 4.0 across datasets, highlights EPR-LP's lower iteration cost compared to cuPDLP, likely attributed to cuPDLP's dependence on a more time-intensive heuristic line search.

Table 1: Per-iteration time ratio (cuPDLP/EPR-LP) for different datasets with and without presolve.

Dataset	Median	Mean	Standard deviation
Mittelmann's LP benchmark set without presolve	2.7	3.5	3.0
Mittelmann's LP benchmark set with presolve	3.2	3.9	4.0
MIP relaxations without presolve	3.9	8.2	14.5
MIP relaxations with presolve	4.0	8.8	15.5

5 CONCLUSION

In this paper, we proved the ergodic convergence of the PR method with semi-proximal terms for solving convex optimization problems. We established the ergodic iteration complexity of O(1/k)with respect to the objective error, the feasibility violation, and the KKT residual based on ε subdifferential. Building on these results, we developed the solver EPR-LP for solving large-scale LP problems, which incorporates adaptive restart and penalty parameter updates. Extensive numerical experiments on LP benchmark datasets highlighted the advantages of EPR-LP compared to PDLP. In the future, it would be interesting to explore combining the ergodic sequence with acceleration techniques, such as Halpern's iteration (Halpern, 1967; Lieder, 2021; Sun et al., 2024), to design a more efficient algorithm with an iteration complexity better than O(1/k).

540 REFERENCES 541

34 I	
542 543	Vando A Adona, Max LN Gonçalves, and Jefferson G Melo. Iteration-complexity analysis of a generalized alternating direction method of multipliers. <i>J. Global Optim.</i> , 73:331–348, 2019.
544 545 546	David Applegate, Mateo Díaz, Oliver Hinder, Haihao Lu, Miles Lubin, Brendan O'Donoghue, and Warren Schudy. Practical large-scale linear programming using primal-dual hybrid gradient. In <i>Advances in Neural Information Processing System</i> , volume 34, pp. 20243–20257, 2021.
547 548 549	David Applegate, Oliver Hinder, Haihao Lu, and Miles Lubin. Faster first-order primal-dual methods for linear programming using restarts and sharpness. <i>Math. Program.</i> , 201(1):133–184, 2023.
550 551	Jean-Bernard Baillon. Un théorème de type ergodique pour les contractions non linéaires dans un espace de hilbert. <i>CR Acad. Sci. Paris Sér. AB</i> , 280:1511–1514, 1975.
552 553 554	Heinz H Bauschke and Patrick L Combettes. <i>Convex Analysis and Monotone Operator Theory in Hilbert Spaces</i> , 2nd edn. Springer, New York, 2017.
555 556	Jeff Bezanson, Alan Edelman, Stefan Karpinski, and Viral B Shah. Julia: A fresh approach to numerical computing. <i>SIAM REV</i> , 59(1):65–98, 2017.
557 558 559	Kristian Bredies, Enis Chenchene, Dirk A Lorenz, and Emanuele Naldi. Degenerate preconditioned proximal point algorithms. <i>SIAM J. Optim.</i> , 32(3):2376–2401, 2022.
560 561	Regina S Burachik, Alfredo N Iusem, and Benar Fux Svaiter. Enlargement of monotone operators with applications to variational inequalities. <i>Set-Valued Analysis</i> , 5:159–180, 1997.
562 563	Antonin Chambolle and Thomas Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. <i>J. Math. Imaging Vision</i> , 40:120–145, 2011.
565 566 567	Kaihuang Chen, Defeng Sun, Yancheng Yuan, Guojun Zhang, and Xinyuan Zhao. HPR- LP: An implementation of an HPR method for solving linear programming. <i>arXiv preprint</i> <i>arXiv:2408.12179</i> , 2024.
568 569 570	Ying Cui, Xudong Li, Defeng Sun, and Kim-Chuan Toh. On the convergence properties of a majorized alternating direction method of multipliers for linearly constrained convex optimization problems with coupled objective functions. <i>J. Optim. Theory Appl.</i> , 169(3):1013–1041, 2016.
571 572 573	Damek Davis and Wotao Yin. Convergence rate analysis of several splitting schemes. In <i>Splitting Methods in Communication, Imaging, Science, and Engineering</i> , pp. 115–163. Springer, 2016.
574 575 576	Qi Deng, Qing Feng, Wenzhi Gao, Dongdong Ge, Bo Jiang, Yuntian Jiang, Jingsong Liu, Tianhao Liu, Chenyu Xue, Yinyu Ye, et al. An enhanced alternating direction method of multipliers-based interior point method for linear and conic optimization. <i>INFORMS J. Comput</i> , 2024.
577 578 579	Elizabeth D Dolan and Jorge J Moré. Benchmarking optimization software with performance pro- files. <i>Math. Program</i> , 91:201–213, 2002.
580 581 582	Jonathan Eckstein and Dimitri P Bertsekas. On the Douglas—Rachford splitting method and the proximal point algorithm for maximal monotone operators. <i>Math. Program.</i> , 55(1):293–318, 1992.
583 584 585 586	Ernie Esser, Xiaoqun Zhang, and Tony F Chan. A general framework for a class of first order primal-dual algorithms for convex optimization in imaging science. <i>SIAM J. Imaging Sci</i> , 3(4): 1015–1046, 2010.
587 588 589	Maryam Fazel, Ting Kei Pong, Defeng Sun, and Paul Tseng. Hankel matrix rank minimization with applications to system identification and realization. <i>SIAM J. Matrix Anal. Appl.</i> , 34(3):946–977, 2013.
590 591	Daniel Gabay. Applications of the method of multipliers to variational inequalities. In <i>Studies in Mathematics and Its Applications</i> , volume 15, pp. 299–331. Elsevier, 1983.
593	Dongdong Ge, Qi Huangfu, Zizhuo Wang, Jian Wu, and Yinyu Ye. Cardinal Optimizer (COPT) User Guide. <i>arXiv preprint arXiv:2208.14314</i> , 2022.

594 595 596 597	Ambros Gleixner, Gregor Hendel, Gerald Gamrath, Tobias Achterberg, Michael Bastubbe, Timo Berthold, Philipp Christophel, Kati Jarck, Thorsten Koch, Jeff Linderoth, et al. MIPLIB 2017: data-driven compilation of the 6th mixed-integer programming library. <i>Math. Program. Comput.</i> , 13(3):443–490, 2021.
598 599	Gene H Golub and Charles F Van Loan. Matrix Computations. JHU press, 2013.
600 601	Gurobi Optimization, LLC. Gurobi Optimizer Reference Manual, 2024. URL https://www.gurobi.com.
602 603	Benjamin Halpern. Fixed points of nonexpanding maps. Bull. Am. Math. Soc., 73(6):957–961, 1967.
604 605 606	Deren Han, Defeng Sun, and Liwei Zhang. Linear rate convergence of the alternating direction method of multipliers for convex composite programming. <i>Math. Oper. Res.</i> , 43(2):622–637, 2018.
607 608 609	Jean-Baptiste Hiriart-Urruty and Claude Lemaréchal. <i>Convex Analysis and Minimization Algorithms</i> <i>II: Advanced Theory and Bundle Methods</i> , volume 306. Springer science & business media, 1993.
610	IBM. IBM ILOG CPLEX Optimization Studio CPLEX User's Manual, 1987.
611 612 613	Leonid Kantorovich. On the transfer of masses (in Russian). <i>Doklady Akademii Nauk</i> , 37(2): 227–229, 1942.
614	Felix Lieder. On the convergence rate of the Halpern-iteration. Optim. Lett., 15(2):405-418, 2021.
615 616 617	Tianyi Lin, Shiqian Ma, Yinyu Ye, and Shuzhong Zhang. An ADMM-based interior-point method for large-scale linear programming. <i>Optim. Methods Softw.</i> , 36(2-3):389–424, 2021.
618 619	Pierre-Louis Lions and Bertrand Mercier. Splitting algorithms for the sum of two nonlinear opera- tors. <i>SIAM J. Numer. Anal.</i> , 16(6):964–979, 1979.
620 621	Haihao Lu. First-order methods for linear programming. arXiv preprint arXiv:2403.14535, 2024.
622 623	Haihao Lu and Jinwen Yang. cuPDLP.jl: A GPU implementation of restarted primal-dual hybrid gradient for linear programming in julia. <i>arXiv preprint arXiv:2311.12180</i> , 2023.
624 625	Haihao Lu and Jinwen Yang. Restarted Halpern PDHG for linear programming. <i>arXiv preprint arXiv:2407.16144</i> , 2024.
627 628 629	Haihao Lu, Jinwen Yang, Haodong Hu, Qi Huangfu, Jinsong Liu, Tianhao Liu, Yinyu Ye, Chuwen Zhang, and Dongdong Ge. cuPDLP-C: A strengthened implementation of cuPDLP for linear programming by C language. <i>arXiv preprint arXiv:2312.14832</i> , 2023.
630 631	Gaspard Monge. Mémoire sur la théorie des déblais et des remblais. <i>Histoire de l'Académie Royale des Sciences</i> , pp. 666–704, 1781.
632 633 634 635	Renato DC Monteiro and Chee-Khian Sim. Complexity of the relaxed Peaceman–Rachford splitting method for the sum of two maximal strongly monotone operators. <i>Computational Optimization and Applications</i> , 70:763–790, 2018.
636 637	Renato DC Monteiro and Benar F Svaiter. Iteration-complexity of block-decomposition algorithms and the alternating direction method of multipliers. <i>SIAM J. Optim.</i> , 23(1):475–507, 2013.
638 639 640	Brendan O'Donoghue. Operator splitting for a homogeneous embedding of the linear complemen- tarity problem. <i>SIAM J. Optim.</i> , 31(3):1999–2023, 2021.
641 642	Zdzisław Opial. Weak convergence of the sequence of successive approximations for nonexpansive mappings. 1967.
643 644 645	Brendan O'donoghue, Eric Chu, Neal Parikh, and Stephen Boyd. Conic optimization via operator splitting and homogeneous self-dual embedding. <i>J. Optim. Theory Appl.</i> , 169:1042–1068, 2016.
646 647	Thomas Pock and Antonin Chambolle. Diagonal preconditioning for first order primal-dual al- gorithms in convex optimization. In 2011 International Conference on Computer Vision, pp. 1762–1769. IEEE, 2011.

648 640	R Tyrrell Rockafellar. Convex Analysis, volume 18. Princeton University Press, 1970.
650	
000	Daniel Ruiz. A scaling algorithm to equilibrate both rows and columns norms in matrices. Technical
100	report, Rutherford Appleton Laboratory, 2001.
052	
003	Jörn Schrieber, Dominic Schuhmacher, and Carsten Gottschlich, Dotmark-a benchmark for discrete
054	optimal transport. <i>IEEE Access</i> , 5:271–282, 2016.
655	· · · · · · · · · · · · · · · · · · ·
000	Li Shan and Shaahua Dan. Waightad iteration complexity of the sDADMM on the KKT residuals
057	for convex composite optimization arXiv preprint arXiv:1611.03167, 2016
658	for convex composite optimization. <i>urxiv preprint urxiv.1011.05107</i> , 2010.
659	
660	Bartolomeo Stellato, Goran Banjac, Paul Goulart, Alberto Bemporad, and Stephen Boyd. OSQP:
661	An operator splitting solver for quadratic programs. <i>Math. Program. Comput.</i> , 12(4):637–672,
662	2020.
663	
664	Defeng Sun, Yancheng Yuan, Guojun Zhang, and Xinyuan Zhao. Accelerating preconditioned
665	ADMM via degenerate proximal point mappings. <i>arXiv preprint arXiv:2403.18618</i> , 2024.
666	
667	Yunhai Xiao, Liang Chen, and Donghui Li. A generalized alternating direction method of multipliers
668	with semi-proximal terms for convex composite conic programming. Math. Program. Comput.,
669	10:533–555, 2018.
670	
671	MH Xu and Ting Wu. A class of linearized proximal alternating direction methods. J. Optim. Theory
672	Appl., 151:321–337, 2011.
673	
674	Minggiang Zhu and Tony Chan. An efficient primal-dual hybrid gradient algorithm for total variation
675	image restoration. UCLA Cam Report. 34(2), 2008.
676	
677	
678	
679	A PROOFS FOR SECTION 2
680	
681	A.1 PROOF OF PROPOSITION 2.2
682	
683	<i>Proof.</i> Since 7 is <i>M</i> -firmly nonexpansive, as stated in Sun et al. (2024, Proposition 2.3), we obtain
684	
685	
686	$\ w^{k+1} - w^*\ _{\mathcal{M}}^2 \le \ w^k - w^*\ _{\mathcal{M}}^2 - \rho(2-\rho) \ w^k - \bar{w}^k\ _{\mathcal{M}}^2, \forall k \ge 0, w^* \in \mathcal{T}^{-1}(0), (22)$
687	
688	which together with the iteration scheme (8) implies that
689	when, together with the neration scheme (0), implies that
690	1 k $ $
691	$\ \bar{w}_{a}^{k} - w_{a}^{k}\ _{\mathcal{M}} = \left\ \frac{1}{1-1}\sum_{k=1}^{\infty} (\bar{w}^{t} - w^{t})\right\ $
692	$\ k+1 \underset{t=0}{\swarrow} \ $
693	

$$= \left\| \frac{1}{k+1} \sum_{t=0}^{k} \frac{(w^{t+1} - w^{t})}{\rho} \right\|_{\mathcal{M}}$$

$$= \frac{1}{\rho(k+1)} \left\| (w^{k+1} - w^{0}) \right\|_{\mathcal{M}}$$

$$\leq \frac{1}{\rho(k+1)} \left\| (w^{k+1} - w^{*}) - (w^{0} - w^{*}) \right\|_{\mathcal{M}}$$

$$\leq \frac{2}{\rho(k+1)} \left\| (w^{0} - w^{*}) \right\|_{\mathcal{M}}, \quad \forall k \ge 0, w^{*} \in \mathcal{T}^{-1}(0).$$

This concludes the proof of statement (a). Moreover, for any $(w', v') \in \operatorname{gph}(\mathcal{T})$, and by the defini-tions of w_a^k and \bar{w}_a^k in (9), we have, for any $k \ge 0$,

 $\begin{array}{l} \langle w_{a}^{n} - w', \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - v' \rangle \\ = & \frac{1}{k+1} \sum_{t=0}^{k} \langle \bar{w}^{t} - w', \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - v' \rangle \\ = & \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \bar{w}^{t} - w', \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - \mathcal{M}(w^{t} - \bar{w}^{t}) \rangle + \langle \bar{w}^{t} - w', \mathcal{M}(w^{t} - \bar{w}^{t}) - v' \rangle \right) \\ \geq & \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \bar{w}^{t} - w', \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - \mathcal{M}(w^{t} - \bar{w}^{t}) \rangle \right) \\ = & \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \bar{w}^{t} - \bar{w}_{a}^{k}, \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - \mathcal{M}(w^{t} - \bar{w}^{t}) \rangle + \langle \bar{w}_{a}^{k} - w', \mathcal{M}(w_{a}^{k} - \bar{w}_{a}^{k}) - \mathcal{M}(w^{t} - \bar{w}^{t}) \rangle \right) \\ = & \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \bar{w}^{t} - \bar{w}_{a}^{k}, (w_{a}^{k} - \bar{w}_{a}^{k}) - (w^{t} - \bar{w}^{t}) \rangle_{\mathcal{M}} \right) \\ = & -\frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \bar{w}^{t} - \bar{w}_{a}^{k}, (w^{t} - \bar{w}^{t}) \rangle_{\mathcal{M}} \right) \\ = & -\bar{\varepsilon}_{a}^{k}. \end{array}$

Next, we prove that $\bar{\varepsilon}_a^k \ge 0$ for all $k \ge 0$ by contradiction. Suppose $\bar{\varepsilon}_a^k < 0$ for some $k \ge 0$. Then, for any $(w', v') \in gph(\mathcal{T})$, we have

$$\langle \bar{w}_a^k - w', \mathcal{M}(w_a^k - \bar{w}_a^k) - v' \rangle > 0,$$

which, combined with the maximality of \mathcal{T} , implies that $(\bar{w}_a^k, \mathcal{M}(w_a^k - \bar{w}_a^k)) \in \operatorname{gph}(\mathcal{T})$. Taking $(w', v') = (\bar{w}_a^k, \mathcal{M}(w_a^k - \bar{w}_a^k))$, we obtain $0 \ge -\bar{\varepsilon}_a^k$, which contradicts our assumption. Thus, $\bar{\varepsilon}_a^k \ge 0$ for all $k \ge 0$. Next, we establish an upper bound for $\bar{\varepsilon}_a^k$ for any $k \ge 0$. Indeed, we have

$$\begin{aligned} \vec{r}_{21} & \vec{\varepsilon}_{a}^{k} &= \frac{1}{k+1} \sum_{t=0}^{k} \langle w^{t} - \bar{w}^{t}, \bar{w}^{t} - \bar{w}_{a}^{k} \rangle_{\mathcal{M}} \\ &= \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \frac{1}{\rho} (w^{t} - w^{t+1}), w^{t} - \frac{1}{\rho} (w^{t} - w^{t+1}) - \bar{w}_{a}^{k} \rangle_{\mathcal{M}} \right) \\ \vec{r}_{23} & = \frac{1}{k+1} \sum_{t=0}^{k} \left(-\frac{1}{\rho^{2}} \| w^{t} - w^{t+1} \|_{\mathcal{M}}^{2} + \frac{1}{2\rho} (\| w^{t} - w^{t+1} \|_{\mathcal{M}}^{2} + \| w^{t} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2} - \| w^{t+1} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2}) \right) \\ \vec{r}_{25} & \leq \frac{1}{k+1} \sum_{t=0}^{k} \left(\frac{1}{2\rho} (\| w^{t} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2} - \| w^{t+1} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2}) \right) \\ \vec{r}_{26} & = \frac{1}{2\rho(k+1)} \left(\| w^{0} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2} - \| w^{k+1} - \bar{w}_{a}^{k} \|_{\mathcal{M}}^{2}) \right) \\ \vec{r}_{27} & = \frac{1}{2\rho(k+1)} \left(-\| w^{0} - w^{k+1} \|_{\mathcal{M}}^{2} - 2 \langle w^{k+1} - w^{0}, w^{0} - \bar{w}_{a}^{k} \rangle_{\mathcal{M}} \right) \\ \vec{r}_{28} & = \frac{1}{2\rho(k+1)} \left(-\| w^{0} - w^{k+1} \|_{\mathcal{M}}^{2} + 2\| w^{k+1} - w^{0} \|_{\mathcal{M}} \| w^{0} - \bar{w}_{a}^{k} \|_{\mathcal{M}} \right). \end{aligned}$$

$$(23)$$

Using the convexity of $\|\cdot\|_{\mathcal{M}}$, we can obtain

$$\|w^{0} - w_{a}^{k}\|_{\mathcal{M}} = \|\frac{1}{k+1}\sum_{t=0}^{k}(w^{0} - w^{t})\|_{\mathcal{M}} \le \frac{1}{k+1}\sum_{t=0}^{k}\|w^{0} - w^{t}\|_{\mathcal{M}} \le 2\|w^{0} - w^{*}\|_{\mathcal{M}}.$$

Combining this with (23), we derive that for any $k \ge 0$,

$$\bar{\bar{z}}_{a}^{k} \leq \frac{1}{2\rho(k+1)} \left(-\|w^{0} - w^{k+1}\|_{\mathcal{M}}^{2} + 2\|w^{k+1} - w^{0}\|_{\mathcal{M}}\|w^{0} - w^{*}\|_{\mathcal{M}}^{2} \right)$$

$$\leq \frac{1}{2\rho(k+1)}\|w^{0} - w^{*}\|_{\mathcal{M}}^{2}.$$

This completes the proof.

A.2 PROOF OF THEOREM 2.1

Proof. Suppose that $\mathcal{M} = \mathcal{CC}^*$ is a decomposition of \mathcal{M} according to Bredies et al. (2022, Proposition 2.3), where $\mathcal{C}: \mathbb{U} \to \mathbb{W}$. Since $(\mathcal{M} + \mathcal{T})^{-1}$ is L-Lipschitz continuous and $\|\mathcal{C}^* w\| = \|w\|_{\mathcal{M}}$ for every $w \in \mathcal{H}$, we have, for all $w' \in \mathcal{H}$ and $w^* \in \mathcal{T}^{-1}(\hat{0})$,

$$\|\widehat{\mathcal{T}}w' - \widehat{\mathcal{T}}w^*\| = \|(\mathcal{M} + \mathcal{T})^{-1}\mathcal{C}\mathcal{C}^*w' - (\mathcal{M} + \mathcal{T})^{-1}\mathcal{C}\mathcal{C}^*w^*\| \le L\|\mathcal{C}\|\|w' - w^*\|_{\mathcal{M}},$$

which, together with (22), implies that

$$\|\bar{w}^{k} - w^{*}\| = \|\widehat{\mathcal{T}}w^{k} - w^{*}\| \le L\|\mathcal{C}\|\|w^{k} - w^{*}\|_{\mathcal{M}} \le L\|\mathcal{C}\|\|w^{0} - w^{*}\|_{\mathcal{M}}.$$

Thus, both sequences $\{\bar{w}^k\}$ and $\{\bar{w}^k_a\}$ are bounded. According to Proposition 2.2 and the maximal-ity of \mathcal{T} , any cluster point of $\{\bar{w}_a^k\}$ belongs to $\mathcal{T}^{-1}(0)$.

To establish the uniqueness of cluster points, we define two shadow sequences as follows:

$$u^{k} := \mathcal{C}^{*} w^{k}$$
 and $u^{k}_{a} := \frac{1}{k+1} \sum_{t=0}^{k} u^{t}, \quad \forall k \ge 0.$ (24)

Through straightforward calculations, we obtain

$$u^{k+1} = \widehat{\mathcal{F}}_{\rho} u^k, \quad \forall k \ge 0,$$

where $\widehat{\mathcal{F}}_{\rho} := (1-\rho)\mathcal{I} + \rho(\mathcal{C}^*(\mathcal{M}+\mathcal{T})^{-1}\mathcal{C})$ with $\rho \in (0,2]$ is a nonexpansive operator, according to Sun et al. (2024, Proposition 2.5). By Baillon's nonlinear ergodic theorem (Baillon, 1975) and (Bauschke & Combettes, 2017, Example 5.38), the sequence $\{u_a^k\}$ converges to a point in Fix $(\widehat{\mathcal{F}}_{\rho})$, where Fix $(\widehat{\mathcal{F}}_{\rho})$ represents the set of fixed points of the nonexpansive operator $\widehat{\mathcal{F}}_{\rho}$. Given the equivalence between Fix $(\widehat{\mathcal{F}}_{\rho})$ and $\mathcal{C}^*\mathcal{T}^{-1}(0)$ as stated in Sun et al. (2024, Proposition 2.5), we conclude that there exists $w_a^* \in \mathcal{T}^{-1}(0)$ such that

$$\|u_a^k - \mathcal{C}^* w_a^*\| \to 0.$$

Therefore, by the definition of $\{u_a^k\}$ in (24), we have

$$\|w_a^k - w_a^*\|_{\mathcal{M}} = \left\|\frac{1}{k+1}\sum_{t=0}^k \mathcal{C}^* w^t - \mathcal{C}^* w_a^*\right\| = \|u_a^k - \mathcal{C}^* w_a^*\| \to 0,$$

which, together with part (a) of Proposition 2.2, implies that

$$\|\bar{w}_{a}^{k} - w_{a}^{*}\|_{\mathcal{M}}^{2} = \|\bar{w}_{a}^{k} - w_{a}^{k}\|_{\mathcal{M}}^{2} + \|w_{a}^{k} - w_{a}^{*}\|_{\mathcal{M}}^{2} + 2\langle \bar{w}_{a}^{k} - w_{a}^{k}, w^{k} - w_{a}^{*}\rangle_{\mathcal{M}} \to 0.$$
(25)

Since the sequence $\{\bar{w}_a^k\}$ is bounded, it must have at least one cluster point. Assume a subsequence $\{\bar{w}_a^{k_i}\}$ converges to w^* . Suppose $||w^* - w_a^*||_{\mathcal{M}} > 0$. By an Opial-type argument (Opial, 1967, Lemma 1), we have

$$\liminf_{i \to \infty} \|\bar{w}_a^{k_i} - w^*\|_{\mathcal{M}} < \liminf_{i \to \infty} \|\bar{w}_a^{k_i} - w_a^*\|_{\mathcal{M}}$$

which, combined with (25), implies $\liminf_{i \to \infty} \|\bar{w}_a^{k_i} - w^*\|_{\mathcal{M}} < 0$, a contradiction to the semi-positive definiteness of \mathcal{M} . Hence, $\|w^* - w_a^*\|_{\mathcal{M}} = 0$. It follows that

 $w^* = (\mathcal{M} + \mathcal{T})^{-1} \mathcal{M} w^* = (\mathcal{M} + \mathcal{T})^{-1} \mathcal{M} w^*_a = w^*_a.$

Taking any other cluster point w^{**} , we can similarly show that $w^{**} = w_a^*$. Therefore, the cluster point is unique, and the sequence $\{\bar{w}_a^k\}$ converges to w_a^* .

A.3 PROOF OF LEMMA 2.1

Proof. From the optimality conditions of the subproblems in Algorithm 2, we have, for any $t \ge 0$,

$$\begin{cases} f_2(z) \ge f_2(\bar{z}^t) + \langle -B_2^* \bar{x}^t - \mathcal{T}_2(\bar{z}^t - z^t), z - \bar{z}^t \rangle, & \forall z \in \mathbb{Z}, \\ f_1(y) \ge f_1(\bar{y}^t) + \langle -B_1^* (\bar{x}^t + \sigma(B_1 \bar{y}^t + B_2 \bar{z}^t - c)) - \mathcal{T}_1(\bar{y}^t - y^t), y - \bar{y}^t \rangle, & \forall y \in \mathbb{Y} \end{cases}$$

Summing these from t = 0 to k, and dividing by k + 1, we obtain

$$\begin{cases} f_2(z) \geq \frac{1}{k+1} \sum_{t=0}^k f_2(\bar{z}^t) + \frac{1}{k+1} \sum_{t=0}^k \langle -B_2^* \bar{x}^t - \mathcal{T}_2(\bar{z}^t - z^t), z - \bar{z}^t \rangle, & \forall z \in \mathbb{Z}, \\ f_1(y) \geq \frac{1}{k+1} \sum_{t=0}^k f_1(\bar{y}^t) + \frac{1}{k+1} \sum_{t=0}^k \langle -B_1^*(\bar{x}^t + \sigma(B_1\bar{y}^t + B_2\bar{z}^t - c)) \\ -\mathcal{T}_1(\bar{y}^t - y^t), y - \bar{y}^t \rangle, & \forall y \in \mathbb{Y}. \end{cases}$$

By the convexity of f_2 , we have

$$\begin{aligned} f_2(z) &\geq f_2(\bar{z}_a^k) + \frac{1}{k+1} \sum_{t=0}^k \langle -B_2^* \bar{x}^t - \mathcal{T}_2(\bar{z}^t - z^t), z - \bar{z}^t \rangle, \, \forall z \in \mathbb{Z} \\ &= f_2(\bar{z}_a^k) + \frac{1}{k+1} \sum_{t=0}^k \langle -B_2^* \bar{x}^t - \mathcal{T}_2(\bar{z}^t - z^t), z - \bar{z}_a^k \rangle - \bar{\varepsilon}_z^k, \, \forall z \in \mathbb{Z}, \end{aligned}$$

where

$$\bar{\varepsilon}_z^k := \frac{1}{k+1} \sum_{t=0}^k \langle -B_2^* \bar{x}^t - \mathcal{T}_2(\bar{z}^t - z^t), \bar{z}^t - \bar{z}_a^k \rangle$$

is non-negative by substituting $z = \overline{z}_a^k$ in the first inequality. Hence, we have

$$-(B_2^*\bar{x}_a^k + \mathcal{T}_2(\bar{z}_a^k - z_a^k)) = \frac{1}{k+1}\sum_{t=0}^k -(B_2^*\bar{x}^t + \mathcal{T}_2(\bar{z}^t - z^t)) \in \partial_{\bar{z}_z^k} f_2(\bar{z}_a^k), \ \forall k \ge 0.$$

810 Similarly, we obtain 811

$$-B_1^*(\bar{x}_a^k+\sigma(B_1\bar{y}_a^k+B_2\bar{z}_a^k-c))-\mathcal{T}_1(\bar{y}_a^k-y_a^k)\in\partial_{\bar{\varepsilon}_y^k}f_1(\bar{y}_a^k),\,\forall k\geq 0,$$

where

$$\bar{\varepsilon}_y^k := \frac{1}{k+1} \sum_{t=0}^k \langle -B_1^*(\bar{x}^t + \sigma(B_1\bar{y}^t + B_2\bar{z}^t - c)) - \mathcal{T}_1(\bar{y}^t - y^t), \bar{y}^t - \bar{y}_a^k \rangle \ge 0.$$

Now, we show the upper bound of $\bar{\varepsilon}_z^k + \bar{\varepsilon}_y^k$. According to definitions of $\bar{\varepsilon}_z^k$ and $\bar{\varepsilon}_y^k$ in (11), we have

$$\begin{split} & \bar{\varepsilon}_{z}^{k} + \bar{\varepsilon}_{y}^{k} \\ &= \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle -B_{2}^{*} \bar{x}^{t} - \mathcal{T}_{2}(\bar{z}^{t} - z^{t}), \bar{z}^{t} - \bar{z}_{a}^{k} \rangle + \langle -B_{1}^{*}(\bar{x}^{t} + \sigma(B_{1}\bar{y}^{t} + B_{2}\bar{z}^{t} - c)) - \mathcal{T}_{1}(\bar{y}^{t} - y^{t}), \bar{y}^{t} - \bar{y}_{a}^{k} \rangle \right) \\ &= \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \mathcal{M}(w^{t} - \bar{w}^{t}), \bar{w}^{t} - \bar{w}_{a}^{k} \rangle - \langle B_{1}^{*}\bar{x}^{t}, y^{t} - \bar{y}_{a}^{k} \rangle - \langle B_{2}^{*}\bar{x}^{t}, z^{t} - \bar{z}_{a}^{k} \rangle - \langle c - B_{1}\bar{y}^{t} - B_{2}\bar{z}^{t}, \bar{x}^{t} - \bar{x}_{a}^{k} \rangle \right) \\ &= \frac{1}{k+1} \sum_{t=0}^{k} \left(\langle \mathcal{M}(w^{t} - \bar{w}^{t}), \bar{w}^{t} - \bar{w}_{a}^{k} \rangle \right). \end{split}$$

Thus, by the definition of $\bar{\varepsilon}_a^k$ in Proposition 2.2, and using the equivalence between the pADMM and the dPPM in Proposition 2.1, we can derive

$$\bar{\varepsilon}_z^k + \bar{\varepsilon}_y^k = \bar{\varepsilon}_a^k \le \frac{1}{2\rho(k+1)} \|w^0 - w^*\|_{\mathcal{M}}^2$$

This completes the proof.

A.4 PROOF OF THEOREM 2.2

Proof. According to Propositions 2.1 and 2.2, we have

$$\|\bar{w}_a^k - w_a^k\|_{\mathcal{M}}^2 \le \frac{4R_0^2}{\rho^2(k+1)^2}, \quad \forall k \ge 0.$$

By the definition of \mathcal{M} in (7), this can be rewritten as

$$\|\bar{y}_{a}^{k} - y_{a}^{k}\|_{\mathcal{T}_{1}}^{2} + \frac{1}{\sigma} \|\sigma B_{1}(\bar{y}_{a}^{k} - y_{a}^{k}) + (\bar{x}_{a}^{k} - x_{a}^{k})\|^{2} + \|\bar{z}_{a}^{k} - z_{a}^{k}\|_{\mathcal{T}_{2}}^{2} \le \frac{4R_{0}^{2}}{\rho^{2}(k+1)^{2}}, \ \forall k \ge 0.$$
 (26)

From Step 2 of Algorithm 2, we can deduce that for any $k \ge 0$,

$$\begin{aligned} \|\sigma B_1(\bar{y}_a^k - y_a^k) + (\bar{x}_a^k - x_a^k)\| &= \|\sigma B_1(\bar{y}_a^k - y_a^k) + \sigma (B_1 y_a^k + B_2 \bar{z}_a^k - c)\| \\ &= \sigma \|B_1 \bar{y}_a^k + B_2 \bar{z}_a^k - c\|, \end{aligned}$$

which, together with (26), yields that

$$\|B_1 \bar{y}_a^k + B_2 \bar{z}_a^k - c\| \le \frac{2R_0}{\sqrt{\sigma}\rho(k+1)}, \quad \forall k \ge 0.$$
⁽²⁷⁾

Furthermore, according to the Lemma 2.1, we have for $k \ge 0$,

$$\begin{cases} -B_2^* \bar{x}_a^k - \mathcal{T}_2(\bar{z}_a^k - z_a^k) \in \partial_{\bar{z}_z^k} f_2(\bar{z}_a^k), \\ -B_1^*(\bar{x}_a^k + \sigma(B_1 \bar{y}_a^k + B_2 \bar{z}_a^k - c)) - \mathcal{T}_1(\bar{y}_a^k - y_a^k) \in \partial_{\bar{z}_y^k} f_1(\bar{y}_a^k), \end{cases}$$

which, together with (26) and (27), implies

$$\operatorname{dist}\left(0, \partial_{\bar{z}_{a}^{k}} f_{2}(\bar{z}_{a}^{k}) + B_{2}^{*} \bar{x}_{a}^{k}\right) \leq \|\mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k})\| \leq \|\sqrt{\mathcal{T}_{2}}\| \|(\bar{z}_{a}^{k} - z_{a}^{k})\|_{\mathcal{T}_{2}} \leq \|\sqrt{\mathcal{T}_{2}}\| \frac{2R_{0}}{\rho(k+1)}$$
(28)

and

dist
$$\left(0, \partial_{\bar{\varepsilon}_{y}^{k}} f_{1}(\bar{y}_{a}^{k}) + B_{1}^{*} \bar{x}_{a}^{k}\right) \leq \sigma \|B_{1}^{*}\| \|(B_{1} \bar{y}_{a}^{k} + B_{2} \bar{z}_{a}^{k} - c)\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \leq \left(\frac{\sigma \|B_{1}^{*}\|}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_{1}}\|\right) \frac{2R_{0}}{\rho(k+1)}.$$
 (29)

Thus, combining (27), (28), (29) and Lemma 2.1, we derive the iteration complexity bound in (13).

We now estimate the ergodic iteration complexity results for the objective error. From the KKT conditions in (5), we have, for any $k \ge 0$,

$$f_1(\bar{y}_a^k) - f_1(y^*) \ge \langle -B_1^* x^*, \bar{y}_a^k - y^* \rangle, \quad f_2(\bar{z}_a^k) - f_2(z^*) \ge \langle -B_2^* x^*, \bar{z}_a^k - z^* \rangle.$$

Thus, it follows from (27) that for all $k \ge 0$,

$$h(\bar{y}_{a}^{k}, \bar{z}_{a}^{k}) \geq \langle B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c, -x^{*} \rangle$$
$$\geq -\|x^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\|$$

$$\geq -\frac{2R_0\|x^*\|}{\sqrt{\sigma}\rho(k+1)}.$$

> For the upper bound of the objective error, from Sun et al. (2024, Lemma 3.6), we first have the following upper bounds:

$$\begin{aligned} h(\bar{y}^k, \bar{z}^k) &\leq \left\langle \sigma B_1(y^* - \bar{y}^k) - \bar{x}^k, (B_1 \bar{y}^k + B_2 \bar{z}^k - c) \right\rangle \\ &+ \left\langle y^* - \bar{y}^k, \mathcal{T}_1(\bar{y}^k - y^k) \right\rangle + \left\langle z^* - \bar{z}^k, \mathcal{T}_2(\bar{z}^k - z^k) \right\rangle. \end{aligned} \tag{30}$$

Note that from Step 4 of Algorithm 2, we have for any $k \ge 0$,

$$\begin{cases} y^{*} - \bar{y}^{k}, \mathcal{T}_{1}(\bar{y}^{k} - y^{k}) \rangle \\ = \left\langle y^{*} - (y^{k} + \frac{y^{k+1} - y^{k}}{\rho}), \mathcal{T}_{1}(\frac{y^{k+1} - y^{k}}{\rho}) \right\rangle \\ = \frac{1}{2\rho} (\|y^{k} - y^{*}\|_{\mathcal{T}_{1}}^{2} - \|y^{k+1} - y^{*}\|_{\mathcal{T}_{1}}^{2}) + \frac{\rho - 2}{2\rho^{2}} \|y^{k+1} - y^{k}\|_{\mathcal{T}_{1}}^{2} \\ \leq \frac{1}{2\rho} (\|y^{k} - y^{*}\|_{\mathcal{T}_{1}}^{2} - \|y^{k+1} - y^{*}\|_{\mathcal{T}_{1}}^{2}). \end{cases}$$

$$(31)$$

Similarly, we also have

$$\left\langle z^* - \bar{z}^k, \mathcal{T}_2(\bar{z}^k - z^k) \right\rangle \le \frac{1}{2\rho} (\|z^k - z^*\|_{\mathcal{T}_1}^2 - \|z^{k+1} - z^*\|_{\mathcal{T}_1}^2), \quad \forall k \ge 0.$$
 (32)

Additionally, define

$$\Delta_k := x^k + \sigma B_1 y^k, \quad \forall k \ge 0$$

From Step 4 of Algorithm 2, we can derive that

=

$$\begin{cases}
\langle \sigma B_{1}(y^{*} - \bar{y}^{k}) - \bar{x}^{k}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle \\
= \langle \sigma B_{1}y^{*}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle - \langle \bar{x}^{k} + \sigma B_{1}\bar{y}^{k}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle \\
= \langle \sigma B_{1}y^{*}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle - \langle \Delta_{k} + \frac{\Delta_{k+1} - \Delta_{k}}{\rho}, \frac{\Delta_{k+1} - \Delta_{k}}{\rho} \rangle \\
= \langle \sigma B_{1}y^{*}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle - \frac{1}{2\rho} (\|\Delta_{k+1}\|^{2} - \|\Delta_{k}\|^{2}) + \frac{\rho - 2}{2\rho^{2}} \|\Delta_{k+1}\|^{2} \\
\leq \langle \sigma B_{1}y^{*}, (B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) \rangle - \frac{1}{2\rho} (\|\Delta_{k+1}\|^{2} - \|\Delta_{k}\|^{2}), \quad \forall k \ge 0.
\end{cases}$$
(33)

Thus, combing with (30), (31), (32), and (33), we conclude that for all $k \ge 0$,

$$h(\bar{y}^k, \bar{z}^k) \leq \frac{1}{2\rho} (\|y^k - y^*\|_{\mathcal{T}_1}^2 - \|y^{k+1} - y^*\|_{\mathcal{T}_1}^2) + \frac{1}{2\rho} (\|z^k - z^*\|_{\mathcal{T}_1}^2 - \|z^{k+1} - z^*\|_{\mathcal{T}_1}^2) \\ + \langle \sigma B_1 y^*, (B_1 \bar{y}^k + B_2 \bar{z}^k - c) \rangle - \frac{1}{2\rho} (\|\Delta_{k+1}\|^2 - \|\Delta_k\|^2).$$

It follows from the convexity of h and (27) that for any $k \ge 0$,

$$\begin{split} h(\bar{y}_{a}^{k},\bar{z}_{a}^{k}) &\leq \frac{1}{k+1}\sum_{t=0}^{k}h(\bar{y}^{t},\bar{z}^{t}) \\ &\leq \frac{1}{2\rho(k+1)}\left(\|y^{0}-y^{*}\|_{\mathcal{T}_{1}}^{2}+\|z^{0}-z^{*}\|_{\mathcal{T}_{2}}^{2}\right) + \left\langle\sigma B_{1}y^{*},\left(B_{1}\bar{y}_{a}^{k}+B_{2}\bar{z}_{a}^{k}-c\right)\right\rangle + \frac{1}{2\rho(k+1)}\|\Delta_{0}\|^{2} \\ &\leq \frac{R_{0}^{2}}{2\rho(k+1)} + \sqrt{\sigma}\|B_{1}y^{*}\|\frac{2R_{0}}{\rho(k+1)} + \frac{1}{2\rho(k+1)}\|x^{0}+\sigma B_{1}y^{0}\|^{2}. \end{split}$$

This completes the proof.

В THE ERGODIC ITERATION COMPLEXITY OF PADMM FOR SOLVING LP

In this section, without loss of generality, we consider the LP problem without inequality constraints (i.e., $m_2 = 0$). Inequality constraints can be easily converted into equality constraints by introducing slack variables. The KKT system of the LP problem (2) is

$$0 \in Ax^* - b, \quad 0 \in z^* + \partial \delta_C(x^*), \quad A^*y^* + z^* - c = 0.$$
(34)

919 Let $f_1(y) = \langle b, y \rangle$, $f_2(z) = \delta_C^*(-z)$, $B_1 = A^*, B_2 = I_n$, $\mathcal{T}_1 = \sigma(\lambda_1(AA^*)I_m - AA^*)$, $\mathcal{T}_2 = 0$. 920 Then, for LP we define

$$\mathcal{M}^{LP} = \begin{bmatrix} \sigma \lambda_1 (AA^*) I_m & 0 & A^* \\ 0 & 0 & 0 \\ A & 0 & \sigma^{-1} I_n \end{bmatrix},$$

and

$$h^{LP}(\bar{y}_a^k, \bar{z}_a^k) := \langle b, y_a^k \rangle + \delta_C^*(-z_a^k) - (\langle b, y^* \rangle + \delta_C^*(-z^*)), \quad \forall k \ge 0,$$

where (y^*, z^*) is a solution to the problem (2). Thus, the ergodic complexity of pADMM for solving the LP problem (2) is presented in the following corollary.

Corollary B.1. Suppose the solution set to the KKT system (34) is nonempty. Let $\{(\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k)\}$ be the ergodic sequence generated by Algorithm 2 with $\mathcal{T}_1 = \sigma(\lambda_1(AA^*)I_m - AA^*), \mathcal{T}_2 = 0, \text{ and } \rho \in (0, 2]$. Let $w^* = (y^*, z^*, x^*)$ be a solution to the KKT system (34), and define $R_0 = ||w^0 - w^*||_{\mathcal{M}^{LP}}$. For all $k \ge 0$, the following ergodic iteration complexity bound holds:

$$\begin{cases} \|A\bar{x}_a^k - b\| + \|A^*\bar{y}_a^k + \bar{z}_a^k - c\| \le \left(\frac{2\sigma\sqrt{\lambda_1(AA^*)} + 1}{\sqrt{\sigma}}\right)\frac{2R_0}{\rho(k+1)},\\ -\bar{z}_a^k \in \partial_{\bar{\varepsilon}_a^k}\delta_C(\bar{x}_a^k), \end{cases}$$

where $\partial_{\bar{\varepsilon}_z^k} \delta_C(\bar{x}_a^k) = \{ z \in \mathbb{R}^n \mid \langle z, x' - \bar{x}_a^k \rangle \leq \bar{\varepsilon}_z^k, \ \forall x' \in C \}$ and $\bar{\varepsilon}_z^k \leq \frac{1}{2\rho(k+1)} R_0^2$. Moreover,

$$\left(-\frac{1}{\sqrt{\sigma}}\|x^*\|\right)\frac{2R_0}{\rho(k+1)} \le h(\bar{y}_a^k, \bar{z}_a^k) \le \left(R_0 + 4\sqrt{\sigma}\|A^*y^*\|\right)\frac{R_0}{2\rho(k+1)} + \frac{\|x^0 + \sigma A^*y^0\|^2}{2\rho(k+1)}.$$

Proof. Furthermore, since $f_1(y) = \langle b, y \rangle$ for any $y \in \mathbb{R}^m$, it follows from (Hiriart-Urruty & Lemaréchal, 1993, Proposition 1.3.1) that, for any $\varepsilon \ge 0$,

$$\partial_{\varepsilon} f_1(y) = b. \tag{35}$$

On the other hand, from (10) and $T_2 = 0$, we have

$$\bar{x}_a^k \in \partial_{\bar{\varepsilon}_a^k} \delta_C^*(-\bar{z}_a^k),$$

which, by (Hiriart-Urruty & Lemaréchal, 1993, Proposition 1.2.1), is equivalent to

$$-\bar{z}_a^k \in \partial_{\bar{z}_a^k} \delta_C(\bar{x}_a^k). \tag{36}$$

Combing (35),(36), and Theorem 2.2, we complete the proof.

C THE NON-ERGODIC ITERATION COMPLEXITY OF THE PADMM

To establish the non-ergodic iteration complexity of the pADMM, we first establish the non-ergodic iteration complexity of the dPPM.

Proposition C.1. Assume that $\mathcal{T}^{-1}(0)$ is nonempty and $\rho \in (0, 2)$, then the sequences $\{w^k\}$ and $\{\bar{w}^k\}$ generated by the dPPM scheme (8) satisfy

$$\left\|w^{k} - \bar{w}^{k}\right\|_{\mathcal{M}}^{2} \leq \frac{1}{\rho(2-\rho)(k+1)} \left\|w^{0} - w^{*}\right\|_{\mathcal{M}}^{2}, \quad \forall k \geq 0, w^{*} \in \mathcal{T}^{-1}(0)$$

Furthermore, $\|w^t - \bar{w}^t\|_{\mathcal{M}}^2 = o(1/(k+1))$ as $k \to \infty$.

Proof. Since $\hat{\mathcal{T}}$ is \mathcal{M} -firmly nonexpansive in Sun et al. (2024, Proposition 2.3), we can obtain

$$\|w^{k+1} - w^*\|_{\mathcal{M}}^2 \le \|w^k - w^*\|_{\mathcal{M}}^2 - \rho(2-\rho) \|w^k - \bar{w}^k\|_{\mathcal{M}}^2, \quad \forall k \ge 0, w^* \in \mathcal{T}^{-1}(0).$$
(37)

972 Summing this inequality from 0 to k, we have

$$\sum_{t=0}^{k} \rho(2-\rho) \|w^{t} - \bar{w}^{t}\|_{\mathcal{M}}^{2} \le \|w^{0} - w^{*}\|_{\mathcal{M}}^{2} - \|w^{k+1} - w^{*}\|_{\mathcal{M}}^{2} \le \|w^{0} - w^{*}\|_{\mathcal{M}}^{2}, \quad (38)$$

which implies that $||w^t - \bar{w}^t||_{\mathcal{M}}^2 = o(1/(k+1))$ as $k \to \infty$. Now, we claim that the sequence $\{||w^t - \bar{w}^t||_{\mathcal{M}}\}$ is monotonically nonincreasing. Indeed,

$$= \frac{\|w^{t+1} - \bar{w}^{t+1}\|_{\mathcal{M}}^{2}}{\|(w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t})\|_{\mathcal{M}}^{2} + \|(w^{t} - \bar{w}^{t})\|_{\mathcal{M}}^{2}}{+2\langle (w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t}), (w^{t} - \bar{w}^{t})\rangle_{\mathcal{M}}}$$
(39)
$$= \|(w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t})\|_{\mathcal{M}}^{2} + \|(w^{t} - \bar{w}^{t})\|_{\mathcal{M}}^{2}}{-\frac{2}{\rho}\langle (w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t}), (w^{t+1} - w^{t})\rangle_{\mathcal{M}}}.$$

 Since

$$\begin{cases} \langle (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t), (w^{t+1} - w^t) \rangle_{\mathcal{M}} \\ = & \langle (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t), (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t) \rangle_{\mathcal{M}} \\ & + \langle (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t), \bar{w}^{t+1} - \bar{w}^t \rangle_{\mathcal{M}} \\ = & \| (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t) \|_{\mathcal{M}}^2 + \langle (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t), \bar{w}^{t+1} - \bar{w}^t \rangle_{\mathcal{M}}, \end{cases}$$

we can obtain from (39) that

 $= \frac{\|w^{t+1} - \bar{w}^{t+1}\|_{\mathcal{M}}^{2}}{\|w^{t} - \bar{w}^{t}\|_{\mathcal{M}}^{2} - \frac{2-\rho}{\rho}\|(w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t})\|_{\mathcal{M}}^{2}}{-\frac{2}{\rho}\langle (w^{t+1} - \bar{w}^{t+1}) - (w^{t} - \bar{w}^{t}), \bar{w}^{t+1} - \bar{w}^{t}\rangle_{\mathcal{M}}}.$ (40)

By using the monotonicity of \mathcal{T} , we have

$$\langle (w^{t+1} - \bar{w}^{t+1}) - (w^t - \bar{w}^t), \bar{w}^{t+1} - \bar{w}^t \rangle_{\mathcal{M}} \ge 0.$$

Hence,

$$\|w^{t+1} - \bar{w}^{t+1}\|_{\mathcal{M}}^2 \le \|w^t - \bar{w}^t\|_{\mathcal{M}}^2$$

which together with (38) implies

$$||w^k - \bar{w}^k||_{\mathcal{M}}^2 \le \frac{1}{\rho(2-\rho)(k+1)} ||w^0 - w^*||_{\mathcal{M}}^2, \ \forall k \ge 0.$$

1003 This completes the proof.

To analyze the iteration complexity of Algorithm 2, we begin by considering the residual mapping associated with the KKT system (5), as introduced in Han et al. (2018):

$$\mathcal{R}(w) = \begin{pmatrix} y - \operatorname{Prox}_{f_1}(y - B_1^*x) \\ z - \operatorname{Prox}_{f_2}(z - B_2^*x) \\ c - B_1y - B_2z \end{pmatrix}, \quad \forall w = (y, z, x) \in \mathbb{W}.$$
(41)

1011 It is clear that $w^* = (y^*, z^*, x^*)$ satisfies the KKT system (5) if and only if $\mathcal{R}(w^*) = 0$. Now, we are ready to present the non-ergodic iteration complexity for Algorithm 2.

Theorem C.1. Suppose that Assumptions 2.1 and 2.2 hold. Let $\{(\bar{y}^k, \bar{z}^k, \bar{x}^k)\}$ be the sequence generated by Algorithm 2 with $\rho \in (0, 2)$, and let $w^* = (y^*, z^*, x^*)$ be the limit point of the sequence $\{(\bar{y}^k, \bar{z}^k, \bar{x}^k)\}$ and $R_0 = ||w^0 - w^*||_{\mathcal{M}}$. For all $k \ge 0$, we have the following iteration complexity bounds:

$$\|\mathcal{R}(\bar{w}^k)\| \le \left(\frac{\sigma \|B_1^*\| + 1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_2}\| + \|\sqrt{\mathcal{T}_1}\|\right) \frac{R_0}{\sqrt{\rho(2-\rho)(k+1)}}$$
(42)

¹⁰¹⁹ and

$$\left(\frac{-1}{\sqrt{\sigma}} \|x^*\|\right) \frac{R_0}{\sqrt{\rho(2-\rho)(k+1)}} \le h(\bar{y}^k, \bar{z}^k) \le \left(3R_0 + \frac{1}{\sqrt{\sigma}} \|x^*\|\right) \frac{R_0}{\sqrt{\rho(2-\rho)(k+1)}}$$

1022 *Furthermore, as* $k \to \infty$ *,*

1023
1024
$$\|\mathcal{R}(\bar{w}^k)\| \le \left(\frac{\sigma \|B_1^*\| + 1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_2}\| + \|\sqrt{\mathcal{T}_1}\|\right) o(\frac{1}{\sqrt{k+1}})$$

and $|h(\bar{y}^k, \bar{z}^k)| = o(\frac{1}{\sqrt{k+1}}).$

Proof. We first estimate the convergence rate of $\mathcal{R}(\bar{w}^k)$ for any $k \geq 0$. According to Proposition C.1, we have

$$\|w^k - \bar{w}^k\|_{\mathcal{M}}^2 \le \frac{R_0^2}{\rho(2-\rho)(k+1)}, \quad \forall k \ge 0.$$

By the definition of \mathcal{M} in (7), this can be rewritten as

$$\|\bar{y}^{k} - y^{k}\|_{\mathcal{T}_{1}}^{2} + \frac{1}{\sigma} \|\sigma B_{1}(\bar{y}^{k} - y^{k}) + (\bar{x}^{k} - x^{k})\|^{2} + \|\bar{z}^{k} - z^{k}\|_{\mathcal{T}_{2}}^{2} \le \frac{R_{0}^{2}}{\rho(2 - \rho)(k + 1)}, \forall k \ge 0.$$
(43)

Due to Step 2 in Algorithm 2, we can deduce that for any $k \ge 0$,

$$\begin{aligned} \|\sigma B_1(\bar{y}^k - y^k) + (\bar{x}^k - x^k)\| &= \|\sigma B_1(\bar{y}^k - y^k) + \sigma(B_1y^k + B_2\bar{z}^k - c)\| \\ &= \sigma \|B_1\bar{y}^k + B_2\bar{z}^k - c\|, \end{aligned}$$

which together with (43) yields that

$$\|B_1 \bar{y}^k + B_2 \bar{z}^k - c\| \le \frac{R_0}{\sqrt{\sigma \rho (2 - \rho)(k + 1)}}, \quad \forall k \ge 0.$$
(44)

Moreover, from the optimality conditions of the subproblems in Algorithm 2, we have for any $k \ge 0$,

> $\begin{cases} \bar{z}^k = \operatorname{Prox}_{f_2}(\bar{z}^k - B_2^* \bar{x}^k - \mathcal{T}_2(\bar{z}^k - z^k)), \\ \bar{y}^k = \operatorname{Prox}_{f_1}(\bar{y}^k - B_1^* (\bar{x}^k + \sigma (B_1 \bar{y}^k + B_2 \bar{z}^k - c)) - \mathcal{T}_1(\bar{y}^k - y^k)), \end{cases}$ (45)

which together with (43) yields that for any $k \ge 0$,

$$\begin{aligned} \|\bar{z}^{k} - \operatorname{Prox}_{f_{2}}(\bar{z}^{k} - B_{2}^{*}\bar{x}^{k})\| \\ \|\operatorname{Prox}_{f_{2}}(\bar{z}^{k} - B_{2}^{*}\bar{x}^{k} - \mathcal{T}_{2}(\bar{z}^{k} - z^{k})) - \operatorname{Prox}_{f_{2}}(\bar{z}^{k} - B_{2}^{*}\bar{x}^{k})\| \\ & \leq \|\mathcal{T}_{2}(\bar{z}^{k} - z^{k})\| \\ \|\bar{z}^{k} - z^{k}\|_{\mathcal{T}_{2}} \\ \|\sqrt{\mathcal{T}_{2}}\| \|\bar{z}^{k} - z^{k}\|_{\mathcal{T}_{2}} \\ & \leq \|\sqrt{\mathcal{T}_{2}}\| \frac{R_{0}}{\sqrt{\rho(2-\rho)(k+1)}}. \end{aligned}$$

$$(46)$$

Similarly, from (43), (44), and (45), we also have for any k > 0,

 $\begin{aligned} &\|\bar{y}^{k} - \operatorname{Prox}_{f_{1}}(\bar{y}^{k} - B_{1}^{*}\bar{x}^{k})\| \\ &\leq &\|B_{1}^{*}\sigma(B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c) + \mathcal{T}_{1}(\bar{y}^{k} - y^{k})\| \\ &\leq &\sigma\|B_{1}^{*}\|\|B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}^{k} - y^{k})\| \\ &\leq &(\sqrt{\sigma}\|B_{1}^{*}\| + \|\sqrt{\mathcal{T}_{1}}\|)\frac{R_{0}}{\sqrt{\rho(2-\rho)(k+1)}}. \end{aligned}$ (47)

Therefore, by (44), (46), and (47), we can obtain that for any $k \ge 0$,

$$\begin{aligned} \|\mathcal{R}(\bar{w}^{k})\| &\leq \sqrt{\left(\frac{1}{\sigma} + \|\sqrt{\mathcal{T}_{2}}\|^{2} + (\sqrt{\sigma}\|B_{1}^{*}\| + \|\sqrt{\mathcal{T}_{1}}\|)^{2}\right)} \frac{R_{0}}{\sqrt{\rho(2-\rho)(k+1)}} \\ &\leq \left(\frac{\sigma\|B_{1}^{*}\|+1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_{2}}\| + \|\sqrt{\mathcal{T}_{1}}\|\right) \frac{R_{0}}{\sqrt{\rho(2-\rho)(k+1)}}. \end{aligned}$$
(48)

Now, we estimate the complexity result concerning the objective error. For the lower bound of the objective error, from Sun et al. (2024, Lemma 3.6) and (44), we have for all $k \ge 0$,

$$h(\bar{y}^{k}, \bar{z}^{k}) \ge \langle B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c, -x^{*} \rangle$$

$$\ge -\|x^{*}\|\|B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c\|$$

1071
$$\geq -\|x^*\|\|B_1\bar{y}^k + B_2\bar{z}^k\|$$

1072
$$\geq -\frac{R_0 \|x^*\|}{\sqrt{-x^2/2-x^2/(k+1)}}$$

1073
$$\geq -\frac{1}{\sqrt{\sigma\rho(2-\rho)(k+1)}}$$

On the other hand, from the (22), we can obtain

$$||w^{k+1} - w^*||_{\mathcal{M}} \le ||w^k - w^*||_{\mathcal{M}} \le \dots \le R_0, \quad \forall k \ge 0.$$

It follows from the \mathcal{M} -nonexpansiveness of $\widehat{\mathcal{T}}$ by Sun et al. (2024, Proposition 2.3) that

$$\|\bar{w}^k - w^*\|_{\mathcal{M}} = \|\widehat{\mathcal{T}}w^k - w^*\|_{\mathcal{M}} \le \|w^k - w^*\|_{\mathcal{M}} \le R_0, \quad \forall k \ge 0,$$

which implies

$$\|\bar{y}^k - y^*\|_{\mathcal{T}_1}^2 + \frac{1}{\sigma} \|\sigma B_1(\bar{y}^k - y^*) + (\bar{x}^k - x^*)\|^2 + \|\bar{z}^k - z^*\|_{\mathcal{T}_2}^2 \le R_0^2, \quad \forall k \ge 0$$

This inequality together with (30), (43), and (44) yields that for all $k \ge 0$,

$$\begin{aligned} h(\bar{y}^{k}, \bar{z}^{k}) &\leq \left(\|\sigma B_{1}(\bar{y}^{k} - y^{*}) + (\bar{x}^{k} - x^{*})\| + \|x^{*}\| \right) \|B_{1}\bar{y}^{k} + B_{2}\bar{z}^{k} - c\| \\ &+ \|y^{*} - \bar{y}^{k}\|_{\mathcal{T}_{1}} \|\bar{y}^{k} - y^{k}\|_{\mathcal{T}_{1}} + \|z^{*} - \bar{z}^{k}\|_{\mathcal{T}_{2}} \|\bar{z}^{k} - z^{k}\|_{\mathcal{T}_{2}} \\ &\leq \left(\sqrt{\sigma}R_{0} + \|x^{*}\|\right) \frac{R_{0}}{\sqrt{\sigma\rho(2-\rho)(k+1)}} + \frac{2R_{0}^{2}}{\sqrt{\rho(2-\rho)(k+1)}} \\ &= \left(3R_{0} + \frac{1}{\sqrt{\sigma}}\|x^{*}\|\right) \frac{R_{0}}{\sqrt{\rho(2-\rho)(k+1)}}. \end{aligned}$$

We now establish the complexity results as $k \to \infty$. According to Proposition C.1, we have

$$\left\|w^{k}-\bar{w}^{k}\right\|_{\mathcal{M}}^{2}=o\left(\frac{1}{k+1}\right).$$

Following a similar approach as in the previous proof, we obtain

$$\|\mathcal{R}(\bar{w}^k)\| \le \left(\frac{\sigma \|B_1^*\| + 1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_2}\| + \|\sqrt{\mathcal{T}_1}\|\right) o\left(\frac{1}{\sqrt{k+1}}\right),$$

and

$$|h(\bar{y}^k, \bar{z}^k)| = o\left(\frac{1}{\sqrt{k+1}}\right)$$

This completes the proof.

D COMPARISON OF THE ITERATION COMPLEXITY

In addition to the pADMM in Algorithm 2, another widely used variant of the semi-proximal ADMM for solving COPs, as introduced by Fazel et al. (2013), is presented in Algorithm 4.

Algorithm 4 A semi-proximal ADMM (sPADMM) for solving COP (3) 1: Input: Let \mathcal{T}_1 and \mathcal{T}_2 be two self-adjoint, positive semidefinite operators on \mathbb{Y} and \mathbb{Z} , respec-tively. Select an initial point $(y^0, z^0, x^0) \in \text{dom}(f_1) \times \text{dom}(f_2) \times \mathbb{X}$. Set the parameters $\sigma > 0$ and $\tau \in (0, \frac{1+\sqrt{5}}{2})$. 2: for k = 0, 1, ..., doStep 1. $y^{k+1} = \arg\min_{\tau} \left\{ L_{\sigma} \left(y, z^k; x^k \right) + \frac{1}{2} \| y - y^k \|_{\mathcal{T}_1}^2 \right\};$ 3: Step 2. $z^{k+1} = \operatorname*{argmin}_{z \in \mathcal{T}} \left\{ L_{\sigma} \left(y^{k+1}, z; x^k \right) + \frac{1}{2} \| z - z^k \|_{\mathcal{T}_2}^2 \right\};$ 4: Step 3. $x^{k+1} = x^k + \tau \sigma (B_1 y^{k+1} + B_2 z^{k+1} - c);$ 5:

Unlike Algorithm 4, the pADMM in Algorithm 2 can be reformulated as a dPPM, facilitating the analysis of its ergodic convergence properties. We summarize some iteration complexity results of these two algorithms in Table 2.

Paper	Algorithm	Proximal operators	ρ	au	Feasibility violation	Objective error	KKT residual	Ту
Davis & Yin (2016)	GADMM	$\mathcal{T}_1 = 0, \mathcal{T}_2 = 0$	(0, 2)	1	$o(1/\sqrt{k})$	$o(1/\sqrt{k})$	-	non-e
Cui et al. (2016)	(majorized) sPADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0^{a}$	1	$(0, \frac{1+\sqrt{5}}{2})$	$o(1/\sqrt{k})$	-	$o(1/\sqrt{k})$	non-e
Ours	pADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0$	(0, 2)	1	$o(1/\sqrt{k})$	$o(1/\sqrt{k})$	$o(1/\sqrt{k})$	non-
Monteiro & Svaiter (2013)	ADMM	$\mathcal{T}_1 = 0, \mathcal{T}_2 = 0$	1	1	O(1/k)	-	$O_{\varepsilon}(1/k)^{\rm b}$	erg
Davis & Yin (2016)	GADMM	$\mathcal{T}_1 = 0, \mathcal{T}_2 = 0$	(0, 2]	1	O(1/k)	O(1/k)	-	erg
Cui et al. (2016)	(majorized) sPADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0$	1	$(0, \tfrac{1+\sqrt{5}}{2})$	O(1/k)	O(1/k)	-	erg
Adona et al. (2019)	GADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0$	(0, 2]	1	O(1/k)	-	$O_{\varepsilon}(1/k)$	er
Shen & Pan (2016)	sPADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0$	1	$(0, \frac{1+\sqrt{5}}{2})$	O(1/k)	-	$O_{\varepsilon}(1/k)$	er
Ours	pADMM	$\mathcal{T}_1 \succeq 0, \mathcal{T}_2 \succeq 0$	(0, 2]	1	O(1/k)	O(1/k)	$O_{\varepsilon}(1/k)$	er

Table 2: The iteration complexity result of pADMM/sPADMM

^a $\mathcal{T}_1 \succeq 0$ denotes that \mathcal{T}_1 is positive semi-definite.

^b $O_{\varepsilon}(1/k)$ of the KKT residual: an O(1/k) iteration complexity of the KKT residual based on ε subdifferntial in (13).

1152 **Remark D.1.** Note that pADMM with $T_1 = 0$, $T_2 = 0$, and $\rho = 2$ is equivalent to GADMM. 1153 In comparison to the results of Davis & Yin (2016), our work not only establishes both ergodic and non-ergodic iteration complexities for the KKT residual—an aspect not addressed in Davis & 1154 Yin (2016)—but also extends the analysis to incorporate general \mathcal{T}_1 and \mathcal{T}_2 . This generalization is 1155 particularly important, as suitable choices of T_1 and T_2 can simplify the solution of subproblems in 1156 solving key convex optimization problems, such as general LP. Furthermore, the GADMM algorithm 1157 with semi-proximal terms studied in (Adona et al., 2019) differs from pADMM due to the way the 1158 proximal terms are incorporated. 1159

1160 To further discuss the relationship between the KKT residual based on ε -subdifferential in (13) and 1161 the KKT residual defined in (41), we first present the following lemma to highlight the difference 1162 between $\partial_{\varepsilon} f(\cdot)$ and $\operatorname{Prox}_{f}(\cdot)$ for a proper closed convex function f.

1163 **Lemma D.1.** Let $f: \mathbb{X} \to (-\infty, +\infty]$ be a proper closed convex function, and let $\bar{x} \in \text{dom}(f)$. 1164 Given $\varepsilon \geq 0$, if $v \in \partial_{\varepsilon} f(\bar{x})$, then 1165

$$\|\bar{x} - \operatorname{Prox}_f(\bar{x} + v)\|^2 \le \varepsilon.$$
(49)

1168 *Proof.* For notational convenience, denote $\operatorname{Proof}_{f}(\bar{x}+v)$ by \tilde{x} . According to the definition of 1169 $\operatorname{Prox}_{f}(\bar{x}+v)$, we have 1170

$$0 \in \partial f(\tilde{x}) + (\tilde{x} - (\bar{x} + v)),$$

 $f(x) > f(\widetilde{x}) + \langle \overline{x} + v - \widetilde{x}, x - \widetilde{x} \rangle, \forall x \in \mathbb{X}.$

1171 which implies 1172

1173 1174 It follows that

1175 1176

1179

1182

1184

1187

1166

1167

 $\begin{aligned} f(\bar{x}) &\geq f(\widetilde{x}) + \langle \bar{x} + v - \widetilde{x}, \bar{x} - \widetilde{x} \rangle, \\ &= f(\widetilde{x}) + \| \bar{x} - \widetilde{x} \|^2 + \langle v, \bar{x} - \widetilde{x} \rangle. \end{aligned}$

1177 In addition, according to assumption that $v \in \partial_{\varepsilon} f(\bar{x})$, we have 1178

$$f(\tilde{x}) \ge f(\bar{x}) + \langle v, \tilde{x} - \bar{x} \rangle - \varepsilon.$$
(51)

1180 Summing (50) and (51), we have 1181

$$\|\widetilde{x} - \bar{x}\|^2 \le \varepsilon$$

This completes the proof. 1183

Remark D.2. The following strongly convex quadratic function demonstrates that the inequality in 1185 Lemma D.1 is tight up to a constant factor of 2: 1186

$$f(x) := \frac{1}{2} \langle x, x \rangle + \langle b, x \rangle,$$

(50)

1134

where $b \in \mathbb{R}^n$. As shown in Hiriart-Urruty & Lemaréchal (1993, Example 1.2.2), it follows that

$$\partial_{\varepsilon}f(x) = \left\{x + b + u : \frac{1}{2}\langle u, u \rangle \le \varepsilon \right\}.$$

1192 1193 Consider $\bar{x} \in \text{dom}(f)$. For any $\varepsilon \ge 0$, we take the ε -subgradient v of $f(\cdot)$ at \bar{x} as follows:

$$v = \bar{x} + b + u$$

1195 1196 where $\frac{1}{2} ||u||^2 = \varepsilon$. Through direct calculation, we obtain

$$\|\bar{x} - \operatorname{Prox}_f(\bar{x} + v)\|^2 = \|\bar{x} - \bar{x} - \frac{u}{2}\|^2 = \frac{1}{4}\|u\|^2 = \frac{1}{2}\varepsilon.$$

1199 1200 This demonstrates that the inequality in (49) is tight, up to the constant factor of 2.

Now, we are ready to give the iteration complexity of the ergodic sequence $\{\bar{w}_a^k\}$ generated by Algorithm 2 in the following theorem.

Theorem D.1. Suppose that Assumptions 2.1 and 2.2 hold. Let $\{(\bar{y}_a^k, \bar{z}_a^k, \bar{x}_a^k)\}$ be the sequence generated by Algorithm 2 with $\rho \in (0, 2]$. Let $w^* = (y^*, z^*, x^*)$ be a solution to the KKT system (5), and $R_0 = ||w^0 - w^*||_{\mathcal{M}}$. For all $k \ge 0$, we have the following iteration complexity bound:

 $\left\{\begin{array}{l} -B_2^*\bar{x}_a^k-\mathcal{T}_2(\bar{z}_a^k-z_a^k)\in\partial_{\bar{\varepsilon}_z^k}f_2(\bar{z}_a^k),\\ -B_1^*(\bar{x}_a^k+\sigma(B_1\bar{y}_a^k+B_2\bar{z}_a^k-c))-\mathcal{T}_1(\bar{y}_a^k-y_a^k)\in\partial_{\bar{\varepsilon}_y^k}f_1(\bar{y}_a^k), \end{array}\right.$

$$\|\mathcal{R}(\bar{w}_{a}^{k})\| \leq \frac{2R_{0}}{\sqrt{2\rho(k+1)}} + \left(\frac{\sigma\|B_{1}^{*}\| + 1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_{2}}\| + \|\sqrt{\mathcal{T}_{1}}\|\right)\frac{2R_{0}}{\rho(k+1)}.$$
(52)

¹²¹⁰ *Proof.* According to the Lemma 2.1, we have for $k \ge 0$,

1207 1208 1209

1190 1191

1194

1197 1198

1214

1216 1217 1218

1226 1227 1228

1229

1215 which together with Lemma D.1 implies that

$$\begin{cases} \|\bar{z}_{a}^{k} - \operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k} - \mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k}))\|^{2} \leq \bar{\varepsilon}_{z}^{k}, \\ \|\bar{y}_{a}^{k} - \operatorname{Prox}_{f_{1}}(\bar{y}_{a}^{k} - B_{1}^{*}(\bar{x}_{a}^{k} + \sigma(B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c)) - \mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\|^{2} \leq \bar{\varepsilon}_{y}^{k}. \end{cases}$$
(53)

1219 It follows that for any $k \ge 0$,

$$\begin{aligned} &\|\bar{z}_{a}^{k} - \operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k})\| \\ &\leq &\|\bar{z}_{a}^{k} - \operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k} - \mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k}))\| \\ &\quad + \|\operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k} - \mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k})) - \operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k})\| \\ &\leq &\sqrt{\bar{\varepsilon}_{z}^{k}} + \|\mathcal{T}_{2}(\bar{z}_{a}^{k} - z_{a}^{k})\|. \end{aligned}$$
(54)

Hence, from (12) and (26), we can obtain

$$\|\bar{z}_{a}^{k} - \operatorname{Prox}_{f_{2}}(\bar{z}_{a}^{k} - B_{2}^{*}\bar{x}_{a}^{k})\| \leq \frac{R_{0}}{\sqrt{2\rho(k+1)}} + \|\sqrt{\mathcal{T}_{2}}\|\frac{2R_{0}}{\rho(k+1)}.$$
(55)

Similarly, from (12), (26), (27), and (53), we also have for any $k \ge 0$,

$$\begin{aligned} \|\bar{y}_{a}^{k} - \operatorname{Prox}_{f_{1}}(\bar{y}_{a}^{k} - B_{1}^{*}\bar{x}_{a}^{k})\| \\ & \|\bar{y}_{a}^{k} - \operatorname{Prox}_{f_{1}}(\bar{y}_{a}^{k} - B_{1}^{*}\bar{x}_{a}^{k})\| \\ & \|\bar{y}_{a}^{k} - \operatorname{Prox}_{f_{1}}(\bar{y}_{a}^{k} - B_{1}^{*}\bar{x}_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \|\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}(\bar{y}_{a}^{k} - y_{a}^{k})\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{y}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + B_{2}\bar{z}_{a}^{k} - c\| + \|\mathcal{T}_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\| \\ & \|\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}^{*}\|\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{1}\bar{z}_{a}^{k} + \sigma\|B_{$$

Therefore, by (27), (55), and (56), we can obtain that for any $k \ge 0$, $k \ge 0$,

1238
1239
1240
1241

$$\|\mathcal{R}(\bar{w}_{a}^{k})\| \leq \frac{2R_{0}}{\sqrt{2\rho(k+1)}} + \left(\frac{1}{\sqrt{\sigma}} + \sqrt{\sigma}\|B_{1}^{*}\| + \|\sqrt{\mathcal{T}_{1}}\| + \|\sqrt{\mathcal{T}_{2}}\|\right)\frac{2R_{0}}{\rho(k+1)}$$

$$\leq \frac{2R_{0}}{\sqrt{2\rho(k+1)}} + \left(\frac{\sigma\|B_{1}^{*}\|+1}{\sqrt{\sigma}} + \|\sqrt{\mathcal{T}_{2}}\| + \|\sqrt{\mathcal{T}_{1}}\|\right)\frac{2R_{0}}{\rho(k+1)}.$$
(57)

This completes the proof.

¹²⁴² E UPDATE RULE FOR σ WITH SAFEGUARDS

Since the approximations Δ_x and Δ_y may significantly deviate from their true values, we update σ using formula (21) only when the following conditions are met; otherwise, we reset σ to 1:

1. Δ_x and Δ_y are within the range:

$$\Delta_x, \Delta_y \in (10^{-16}, 10^{12}); \tag{58}$$

2. The ratio of relative primal and dual infeasibility errors is within acceptable bounds:

$$\frac{\text{error}_d}{\text{error}} \in (10^{-8}, 10^8),$$
 (59)

where

e

$$\operatorname{error}_{p} := \frac{\|\Pi_{D}(b - A\bar{x}_{a}^{r,\tau_{r}})\|}{1 + \|b\|} \quad \text{ and } \quad \operatorname{error}_{d} := \frac{\|c - A^{*}\bar{y}_{a}^{r,\tau_{r}} - \bar{z}_{a}^{r,\tau_{r}}\|}{1 + \|c\|}.$$

1256 In summary, the update rule for σ is presented in Algorithm 5.

Algorithm 5 SigmaUpdate

1: Input: $(\bar{w}^{r,\tau_r}, w^{r,0}, \mathcal{S}_1, A)$. 2: Calculate Δ_x and Δ_y defined in (20); 3: if conditions (58) and (59) are satisfied then 4: $\sigma_{r+1} = \frac{\Delta_x}{\Delta_y}$; 5: else 6: $\sigma_{r+1} = 1$; 7. Output:

1266 7: **Output:** σ_{r+1} .

1268 1269

1270

1247 1248 1249

1250

1251 1252

1253

1255

1257

1259

1260

1261

1262

1263 1264

1265

F EXPERIMENTAL SETUP

Benchmark datasets. Our benchmark datasets include Mittelmann's LP benchmark set and LP
relaxations of instances from the MIPLIB 2017 collection. We test the algorithms on 49 publicly
available instances from Mittelmann's LP benchmark. From the MIPLIB 2017 collection, we select
383 instances following the criteria in Lu & Yang (2023). Of these, two are reported as unbounded,
and one is solved by Gurobi's presolve (Gurobi Optimization, LLC, 2024), leaving 380 instances
for testing.

Software and computing environment. EPR-LP and EDR-LP are implemented in Julia (Bezanson et al., 2017). For a fair comparison, the infeasibility detection of cuPDLP (Lu & Yang, 2023) is disabled. All algorithms are tested on an NVIDIA A100-SXM4-80GB GPU with CUDA 12.3.

Presolve and preconditioning. We compare all algorithms across all datasets, both with and without presolve (using Gurobi 11.0.3, academic license). Before running EPR-LP and EDR-LP, all problems are preconditioned for numerical stability, involving 10 steps of Ruiz scaling (Ruiz, 2001), followed by bidiagonal preconditioning as described in Pock & Chambolle (2011) with $\alpha = 1$. Finally, the vectors *b* and *c* are normalized by ||b|| + 1 and ||c|| + 1, respectively. cuPDLP (Lu & Yang, 2023) uses its default settings.

1286 1287 1288 Initialization and parameter setting. The initial points of EPR-LP and EDR-LP are the origin. We set the penalty parameter $\sigma_0 = 1$. After preconditioning, we estimate $\lambda_1(AA^*)$ by the power method (Golub & Van Loan, 2013).

Termination criteria. We check the stopping criteria for the sequence $\{\bar{w}_a^{r,t}\}$ for EPR-LP and EDR-LP. The feasibilities $\bar{x}_a^{r,t} \in C$ and $\bar{y}_a^{r,t} \in D$ are satisfied for any $r \ge 0$, and $t \ge 1$. We terminate the algorithms when the following stopping criteria (used in Applegate et al. (2021); Lu & Yang (2023); Lu et al. (2023)) are satisfied for the tolerance $\varepsilon \in (0, \infty)$:

1293 1294 1295 $|\langle b, y \rangle - \delta_C^*(-z) - \langle c, x \rangle| \le \varepsilon \left(1 + |\langle b, y \rangle - \delta_C^*(-z)| + |\langle c, x \rangle|\right),$ 1295 $\|\Pi_D(b - Ax)\| \le \varepsilon \left(1 + \|b\|\right),$

1295
$$||\Pi_D(o - \Pi_L)|| \le \varepsilon (1 + ||o||),$$

 $||c - A^*y - z|| \le \varepsilon (1 + ||c||).$

We test all algorithms with $\varepsilon = 10^{-8}$ for all the datasets.

Time limit. We set a time limit of 15,000 seconds for Mittelmann's benchmark dataset. For the LP relaxations of MIP problems, the time limit is 3,600 seconds if the number of nonzero elements in A is less than 10 million; otherwise, the limit is 18,000 seconds.

SUPPLEMENTARY EXPERIMENTS G

SPARSITY OF THE SOLUTION G.1

In this subsection, we use the optimal transport (OT) problem (Monge, 1781; Kantorovich, 1942) as an example to demonstrate that EPR-LP, combined with a restart strategy, effectively preserves the sparsity of the solution. For the experiment, we calculate the optimal transport mapping be-tween two 64x64 pixel images selected from the shape category in the DOTmark dataset (Schrieber et al., 2016). The sparsity of the variable x is shown in Figure 4. It can be observed that, with the restart strategy, EPR-LP quickly maintains the sparsity of x similar to the nonergodic version. Consequently, the sparsity has little effect on the computational efficiency of EPR-LP.



Algorithm	Iterations	Time (s)
Nonergodic pADMM	8,529,300	19,514.3
EPR-LP	220,050	514.9

(b) Iteration number and time

Figure 4: (a) The sparsity of x. (b) Comparison of iteration count and runtime between nonergodic pADMM ($\rho = 1.6$) and EPR-LP with a tolerance of 10^{-8} .