DIRECTIONAL CONVERGENCE, BENIGN OVERFITTING OF GRADIENT DESCENT IN LEAKY RELU TWO-LAYER NEURAL NETWORKS

Anonymous authors

Paper under double-blind review

ABSTRACT

In this paper, we provide sufficient conditions of benign overfitting of fixed width leaky ReLU two-layer neural network classifiers trained on mixture data via gradient descent. Our results are derived by establishing directional convergence of the network parameters and classification error bound of the convergent direction. Our classification error bound also lead to the discovery of a newly identified phase transition. Previously, directional convergence in (leaky) ReLU neural networks was established only for gradient flow. Due to the lack of directional convergence, previous results on benign overfitting were limited to those trained on nearly orthogonal data. All of our results hold on mixture data, which is a broader data setting than the nearly orthogonal data setting in prior work. We demonstrate our findings by showing that benign overfitting occurs with high probability in a much wider range of scenarios than previously known. Our results also allow us to characterize cases when benign overfitting provably fails even if directional convergence occurs. Our work thus provides a more complete picture of benign overfitting in leaky ReLU two-layer neural networks.

1 Introduction

The practical success of deep learning has revealed surprising theoretical phenomena. One such phenomenon, which has inspired intense theoretical research, is benign overfitting, where overparametrized neural network models can achieve arbitrarily small test errors while perfectly interpolating training data—even in the presence of label noise (Zhang et al. (2017); Belkin et al. (2019)). Benign overfitting has attracted broad attention since it seems to conflict with the classical statistical understanding that there should be a trade-off between fitting the training data and generalization on the test data. This led to intense theoretical research which has revealed that benign overfitting also occurs in several classical statistical models, such as linear regression (Bartlett et al. (2020), Muthukumar et al. (2020), Mei and Montanari (2022), Hastie et al. (2022)), ridge regression (Tsigler and Bartlett (2024)), binary linear classification (Chatterji and Long (2021); Wang and Thrampoulidis (2022); Cao et al. (2021); Hashimoto et al. (2025)), and kernel-based estimators (Liang and Rakhlin (2020), Liang et al. (2020)), to name a few. However, our understanding of the phenomenon for neural networks remains limited.

It is widely believed that the implicit bias, which results from implicit regularization of gradient based optimization, is the key to understand generalization performance in the over-parametrized regime. In fact, for binary linear classification, it is known that linear classifiers trained under gradient descent with a loss function with a tight exponential tail, e.g. the exponential loss and logistic loss, converge in direction to the maximum margin classifier (Soudry et al. (2018)). Benign overfitting in binary linear classification has been studied thoroughly by analyzing the convergent direction (Wang and Thrampoulidis (2022); Cao et al. (2021); Hashimoto et al. (2025)).

In this work, we investigate benign overfitting in two-layer leaky ReLU neural networks with fixed width for binary classification whose network parameters are trained on mixture data by gradient descent with exponential loss. We show that benign overfitting occurs in a much wider scenarios than previously known.

Our results are derived by establishing directional convergence of the network parameters with precise characterization of the convergent direction and by obtaining a classification error bound of that direction. Due to the difficulty in establishing directional convergence of gradient descent for ReLU type networks, prior work for leaky ReLU networks were limited to either gradient flow (Frei et al. (2023a)), smoothed approximation of leaky ReLU (Frei et al. (2022)), or nearly orthogonal data regime (Xu and Gu (2023)), i.e. the predictors $\{x_i\}_{i=1,\ldots,n} \subset \mathbb{R}^p$ satisfy $\|x_i\|^2 \gg \max_{i\neq k} |\langle x_i, x_k \rangle|$.

The precise characterization of convergent direction not only allows us to prove benign overfitting beyond nearly orthogonal data regime but also leads to a novel lower bound of the classification error in the case of Gaussian mixture. The lower bound shows our bound is tight and allows us to characterize cases when benign overfitting provably fails even if directional convergence occurs.

Moreover, sufficient conditions of directional convergence and classification error bound are obtained in a deterministic manner, which allows us to show that benign overfitting occurs with high probability on polynomially tailed mixture models while the prior results are limited to sub-Gaussian mixture.

2 RELATED WORK

Benign overfitting in neural networks for binary classification. As noted earlier, benign overfitting in neural networks remains poorly understood. In fact, very few theoretical results are established beyond the so called neural tangent kernel (NTK) regime. Cao et al. (2022) studied benign overfitting for two-layer convolutional neural networks with polynomial ReLU activation (ReLU q , q > 2). Their results were strengthened by Kou et al. (2023a) by relaxing the polynomial ReLU assumption. Both Cao et al. (2022) and Kou et al. (2023a) are not applicable to our setting since our work considers fully connected neural networks.

More closely related work is Frei et al. (2022), which studied benign overfitting in smoothed leaky ReLU two-layer neural networks on mixture data. Their work was extended by Xu and Gu (2023) to non-smooth leaky ReLU like activation. While Xu and Gu (2023)'s work is not limited to (leaky) ReLU, their work is limited to the nearly orthogonal data regime and only holds for isotropic mixture with bounded log-Sobolev constant, which is a stronger assumption than sub-Gaussian mixture. Our work establishes benign overfitting beyond the nearly orthogonal data regime and can be applied to a wider class of mixtures than the prior work such as polynomially tailed mixture. Moreover, we also establish a novel lower bound of the classification error in the case of Gaussian mixture, which shows tightness of our result and helps characterize cases when benign overfitting provably fails. Even with gradient flow, benign overfitting in leaky ReLU neural networks is limited to nearly orthogonal data setting and sub-Gaussian mixture model (Frei et al. (2023a)).

Lastly, our result is shown for fixed width two-layer neural networks, which relaxes Xu and Gu (2023) who required m to grow as n, and thus is also beyond the NTK regime or the lazy training regime (Jacot et al. (2018); Arora et al. (2019)).

Implicit bias in neural networks for binary classification. Here, we focus on works on leaky ReLU homogeneous neural networks which are closely related to ours. We refer readers to Vardi (2022) for a comprehensive survey on implicit bias in neural networks.

Most prior work on the implicit bias of (leaky) ReLU neural networks for binary classification has considered gradient flow. For gradient flow, Lyu and Li (2020) showed that any limit point of the network parameter of homogeneous neural networks trained with exponential type loss is a KKT point of a constrained optimization problem and that margin maximization occurs. Ji and Telgarsky (2020) further established directional convergence of the parameter of homogeneous neural networks trained with the exponential type loss. Taking advantage of the directional convergence of gradient flow, Lyu et al. (2021); Bui Thi Mai and Lampert (2021); Sarussi et al. (2021); Frei et al. (2023b); Min et al. (2024) obtained detailed characterizations of the convergent direction of the parameters of (leaky) ReLU two-layer neural networks under further assumptions on the training data.

In contrast to gradient flow, directional convergence of network parameters for ReLU type neural networks was not proven for gradient descent with exponential type loss, even under strong assumptions. As a result, only weaker forms of implicit bias have been obtained for gradient descent. Lyu and Li (2020) showed that margin maximization also occurs for the network parameters of a homogeneous network trained by gradient descent, but their assumptions include smoothness of the network, and

hence rule out (leaky) ReLU networks. Frei et al. (2023b) showed that the stable rank of network parameters stays bounded by a constant for two-layer neural networks, but their result is limited to the smoothed approximation of leaky ReLU. Kou et al. (2023b) proved similar results for (leaky) ReLU two-layer neural networks assuming that the number of positive and negative neurons are equal. Moreover, Frei et al. (2023b); Kou et al. (2023b) are limited to the rather specialized scenario of nearly orthogonal data regime. Cai et al. (2025) showed directional convergence of gradient descent, but their result is not applicable to ReLU type neural networks since their result requires networks to be twice differentiable with respect to the parameters. Schechtman and Schreuder (2025) extended Lyu and Li (2020) by removing twice differentiability but their result does not guarantee directional convergence and assumes that training is already in the late-stage.

To the best of our knowledge, our work is the first to show directional convergence of gradient descent in ReLU type neural networks with precise characterization of the convergent direction.

A concise summary of related prior work is given in Table 1.

Table 1: Summary of Existing Studies of leaky ReLU two-layer neural networks

	Directional Convergence		Benign Overfitting	
	Nearly Orthogonal Data	Other regime	Nearly Orthogonal Data	Other regime
Gradient Flow	Ji and Telgarsky (2020)		Frei et al. (2023a)	NA
Gradient Descent	NA*		Frei et al. (2022)**,	NA
			Xu and Gu (2023)	

^{*} Cai et al. (2025) showed directional convergence assuming twice differentiability

3 Preliminaries

We consider two-layer neural networks

$$f(\boldsymbol{x}; W) = \sum_{j=1}^{m} a_j \phi(\langle \boldsymbol{x}, \boldsymbol{w}_j \rangle), \tag{1}$$

where $\boldsymbol{x} \in \mathbb{R}^p$, $\boldsymbol{w}_j \in \mathbb{R}^p$, $W = (\boldsymbol{w}_1, \boldsymbol{w}_2, \dots, \boldsymbol{w}_m) \in \mathbb{R}^{p \times m}$, $|a_j| = \pm \frac{1}{\sqrt{m}}$ and ϕ is γ -leaky ReLU activation, i.e. $\phi(x) = \max\{x, \gamma x\}$ for $\gamma \in (0, 1)$. Letting $\zeta(x) = 1$ if $x \geq 0$ and $\zeta(x) = \gamma$ if x < 0, we may conveniently write $\phi(x) = \zeta(x)x$.

Letting $J = \{1, 2, \dots, m\}$, $J_+ = \{j \in J : a_j = \frac{1}{\sqrt{m}}\}$, and $J_- = J \setminus J_+$, we can rewrite the network (1) as

$$f(\boldsymbol{x};W) = \sum_{j \in J_+} \frac{1}{\sqrt{m}} \phi(\langle \boldsymbol{x}, \boldsymbol{w}_j \rangle) - \sum_{j \in J_-} \frac{1}{\sqrt{m}} \phi(\langle \boldsymbol{x}, \boldsymbol{w}_j \rangle).$$

We consider the exponential loss $\ell(u) = \exp(-u)$ and define the empirical loss of the dataset $\{(\boldsymbol{x}_i, y_i)\}_{i \in I}$, where $I := \{1, 2, \dots, n\}$, by

$$\mathcal{L}(W) = \sum_{i=1}^{n} \ell(y_i f(\boldsymbol{x}_i; W)) = \sum_{i=1}^{n} \exp(-y_i f(\boldsymbol{x}_i; W)).$$

We train the first layer weights w_j by gradient descent initialized at $W^{(0)}$, i.e.

$$W^{(t+1)} = W^{(t)} - \alpha \nabla_W \mathcal{L}(W^{(t)}), \quad t = 0, 1, 2, \dots,$$
(2)

where $\alpha>0$ is a fixed constant and we assume $\phi'(0)$ is any number in $[\gamma,1]$. The second layer weights a_j are fixed throughout the training. Fixing the second layer weights $a_j\in\{\pm 1/\sqrt{m}\}$ and only optimizing the first layer weights \boldsymbol{w}_j is standard in the existing literature, see e.g. Arora et al. (2019); Cao et al. (2022); Frei et al. (2022); Xu and Gu (2023); Kou et al. (2023b). We do not consider random initialization of $W^{(0)}$ since only the magnitude of $W^{(0)}$ plays a role in our analysis.

Assume that we observe i.i.d. copies $\{(x_i, y_i)\}_{i=1,...,n}$ from a mixture distribution on $\mathbb{R}^p \times \{\pm 1\}$ which is defined as follows

^{**} Frei et al. (2022) studied smoothed approximation of leaky ReLU instead.

(M) The observations consist of n i.i.d copies $(x_i, y_i), i = 1, ..., n$ of the pair (x, y). Here for a random variable $y \in \{-1, 1\}$ satisfying P(y = 1) = P(y = -1) = 1/2, a random vector $z \in \mathbb{R}^p$ independent of y, and a deterministic $\mu \in \mathbb{R}^p$, we have

$$x = y\mu + z. (3)$$

The sub-Gaussian norm of a random variable X, say $||X||_{\psi_2}$, is defined as

$$||X||_{\psi_2} := \inf\{t > 0 : \mathbb{E}\exp(X^2/t^2) \le 2\}.$$

The sub-Gaussian norm of a random vector $m{X} \in \mathbb{R}^p$ is denoted by $\|m{X}\|_{\psi_2}$ and defined as

$$\|oldsymbol{X}\|_{\psi_2} := \sup_{oldsymbol{v} \in S^{p-1}} \|\langle oldsymbol{v}, oldsymbol{X}
angle \|_{\psi_2}.$$

As special case of the model (M), we consider sub-Gaussian mixture model (sG) and polynomially tailed mixture model (PM):

- (sG) Suppose in model (M) $z = \Sigma^{\frac{1}{2}} \xi$, where $\xi \in \mathbb{R}^p$ has independent entries ξ_j that have mean zero, unit variance, and satisfy $\|\xi_j\|_{\psi_2} \leq L$ for all $j = 1, \ldots, p$ for $L < \infty$,
- (PM) Suppose in model (M) $z = \Sigma^{\frac{1}{2}} \xi$, where $\xi \in \mathbb{R}^p$ has independent entries ξ_j that have mean zero, unit variance, and satisfy $\mathbb{E}|\xi_i|^r \leq K$ for all $j = 1, \ldots, p$ for $r \in (2, 4]$ and $K < \infty$.

The mixture model (sG) have been studied for linear classifiers by Chatterji and Long (2021); Wang and Thrampoulidis (2022); Hashimoto et al. (2025) and for two-layer neural networks with $\Sigma = I_p$ by Frei et al. (2022); Xu and Gu (2023). The model (PM) has been studied for linear classifiers by Hashimoto et al. (2025).

We denote by ||A|| and $||A||_F$ the spectral norm and the Frobenius norm of a matrix A, respectively.

4 Directional Convergence of Gradient Descent

We start from our main result on directional convergence of gradient descent in leaky ReLU twolayer neural networks trained on mixture data. Our result provides precise characterization of the convergent direction, which is a key for our analysis on benign overfitting.

In the following, we let $R_{max}(z) := \max_i ||z_i||$ and $R_{min}(z) := \min_i ||z_i||$, and define an event that capture deterministic conditions the training data need to satisfy for the directional convergence:

$$E(\theta_1, \theta_2) = \left\{ \max_{i \neq k} \left| \left\langle \frac{\boldsymbol{z}_i}{\|\boldsymbol{z}_i\|}, \frac{\boldsymbol{z}_k}{\|\boldsymbol{z}_k\|} \right\rangle \right| \leq \theta_1 \quad \text{and} \quad \max_i \left| \left\langle \frac{\boldsymbol{z}_i}{\|\boldsymbol{z}_i\|}, \frac{\boldsymbol{\mu}}{\|\boldsymbol{\mu}\|} \right\rangle \right| \leq \theta_2 \right\}, \tag{4}$$

for some $\theta_1, \theta_2 \in [0,1]$ (take $\theta_1 = 0$ if $\boldsymbol{\mu} = 0$ or $R_{max}(\boldsymbol{z}) = 0$ and take $\theta_2 = 0$ if $\boldsymbol{\mu} = 0$ or $R_{max}(\boldsymbol{z}) = 0$). Event E allows us to control $\|\boldsymbol{x}_i\|$ and $\langle \boldsymbol{x}_i, \boldsymbol{x}_k \rangle, i \neq k$ in terms of $\|\boldsymbol{\mu}\|, R_{min}(\boldsymbol{z}), R_{max}(\boldsymbol{z}), \theta_1$, and θ_2 .

In addition, we denote by $\sigma := \max_j \| \boldsymbol{w}_j \|$ the size of initialization. It is also convenient to let $\rho := \sigma \sqrt{m(1+\theta_2)\{\|\boldsymbol{\mu}\|^2 + R_{max}^2(\boldsymbol{z})\}}$ to keep the assumptions of our main result concise.

We present sufficient conditions of directional convergence for the following two cases:

Case 1: This is the case when $\langle y_i x_i, y_k x_k \rangle \ge 0$ is guaranteed for any $i \ne k$. In this case, we make the following assumptions:

Assumption 4.1 (Positive Correlation).

$$\|\boldsymbol{\mu}\|^2 \ge 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z}).$$

Assumption 4.2 (Small Initialization).

$$\alpha > \frac{\rho \exp(\rho)}{\gamma (1 - \theta_2) \{ \|\boldsymbol{\mu}\|^2 + R_{min}^2(\boldsymbol{z}) \}}$$

Assumption 4.3 (Small Step Size).

$$\alpha(n\|\boldsymbol{\mu}\|^2 + R_{max}(\boldsymbol{z})^2) < 1.$$

The intuition behind these assumptions is as follows: It can be shown that Assumption 4.1 ensures $\langle y_i x_i, y_k x_k \rangle \geq 0$ for any $i \neq k$. Assumption 4.2 ensures the step size sufficiently larger than initialization so that neuron activation occurs immediately after one step of (2). Assumption 4.3 ensures the step size is small enough for directional convergence to hold.

Case 2: This is when $\langle y_k x_k, y_i x_i \rangle$, $k \neq i$ could be negative. The assumptions we make are:

Assumption 4.4 (Near Orthogonality).

$$2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z}) \leq \frac{\varepsilon_1 \gamma (1 - \theta_2) R_{min}^2(\boldsymbol{z})}{n \exp(2\rho)} \quad \textit{for some } \varepsilon_1 \in [0, 1].$$

Assumption 4.5 (Small Initialization).

$$\alpha \geq \frac{\rho \exp(\rho)}{\varepsilon_2 \gamma (1 - \theta_2) R_{min}^2(z)} \quad \textit{for some } \varepsilon_2 \in (0, 1].$$

Assumption 4.6 (Weak Signal).

$$n\|\boldsymbol{\mu}\|^2 \le \varepsilon_3 R_{min}^2(\boldsymbol{z})$$
 for some $\varepsilon_3 \in [0,1)$.

Assumption 4.7 (Small Step Size).

$$6\alpha R_{max}^2(\boldsymbol{z})\exp(\rho) < 1.$$

Assumption 4.4 and 4.6 allow us to carefully control $\langle y_k x_k, y_i x_i \rangle$. Small $\varepsilon_1, \varepsilon_3$ ensure that the training data are nearly orthogonal. By taking $\mu = 0$, this case includes the nearly orthogonal data regime studied in the prior work (Frei et al. (2023b); Kou et al. (2023b)). The other assumptions are analogous to Case 1.

We are now ready to state our main result on directional convergence, which not only establishes directional convergence but also provides detailed characterization of the convergent direction:

Theorem 4.8 (Directional Convergence on Mixture Data). Suppose event E holds under one of the following conditions:

- (i) Assumptions 4.1, 4.2 and 4.3 with $\theta_2 < 1$,
- (ii) Assumption 4.4, 4.5, 4.6, and 4.7 with $\varepsilon_1 + \varepsilon_2 + \varepsilon_3 < 1$, $C_w \varepsilon_i \le \frac{1}{2}, i = 1, 3$, and $\theta_2 \le \frac{1}{2}$, where $C_w := 24 \exp(1) \gamma^{-2} R_{max}^2(\boldsymbol{z}) R_{min}^{-2}(\boldsymbol{z})$.

Then, the gradient descent iterate (2) keeps all the neurons activated for $t \ge 1$, satisfies $\mathcal{L}(W^{(t)}) = O(t^{-1})$, and converges in direction.

Furthermore, the convergent direction $\{\hat{w}_j\}_{j\in J}$ can also be given by $\hat{w}_j = w_+$ for $j \in J_+$ and $\hat{w}_j = w_-$ for $j \in J_-$, which are the solution to the following optimization problem:

Minimize
$$|J_{+}| ||\boldsymbol{w}_{+}||^{2} + |J_{-}| ||\boldsymbol{w}_{-}||^{2}$$

s.t. $\frac{|J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}_{i}, \boldsymbol{w}_{+} \rangle - \frac{\gamma |J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}_{i}, \boldsymbol{w}_{-} \rangle \geq 1, \forall i \in I_{+},$ (5)
 $\frac{|J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}_{i}, \boldsymbol{w}_{-} \rangle - \frac{\gamma |J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}_{i}, \boldsymbol{w}_{+} \rangle \geq 1, \forall i \in I_{-}.$

Lastly, the resulting network $f(\cdot; \hat{W})$ has the linear decision boundary defined by $\bar{\boldsymbol{w}} := \frac{|J_+|}{\sqrt{m}} \boldsymbol{w}_+ - \frac{|J_-|}{\sqrt{m}} \boldsymbol{w}_-$, i.e. $\operatorname{sign}(f(\boldsymbol{x}:\hat{W})) = \operatorname{sign}(\langle \boldsymbol{x}, \bar{\boldsymbol{w}} \rangle)$ for any $\boldsymbol{x} \in \mathbb{R}^p$.

We note that condition (i) goes beyond nearly orthogonal data regime studied in the existing literature since there is no upper bound to the magnitude of the signal μ , while condition (ii) contains the nearly orthogonal data regime studied in Frei et al. (2023b); Kou et al. (2023b). As discussed in Introduction, directional convergence of gradient descent in ReLU type neural networks was not established in the existing literature even under nearly orthogonal data regime. In addition, the characterizations of the convergent direction in Theorem 4.8 were obtained only for gradient flow trained on nearly orthogonal data by Frei et al. (2023b).

We show in Section 6 that these deterministic conditions on training data can be achieved with high probability under sufficient over-parametrization not only by (sG) but also by (PM).

The proof of Theorem 4.8 is given in Appendix B.

5 CLASSIFICATION ERROR OF THE CONVERGENT DIRECTION

In this section, we discuss the classification performance of the convergent direction \hat{W} .

Near orthogonality of z_i 's to each other and to μ is again the key to obtain the closed form expression and to establish an error bound. Letting $Z=(z_1,\ldots,z_n)^{\top}\in\mathbb{R}^{n\times p}, n_+=|\{i\in I:y_i=1\}|$, and $n_-=n-n_+$, the near orthogonality condition is characterized by the parameters from the following events which are defined for $R\geq 0, \tilde{\varepsilon}_1\geq 0, \tilde{\varepsilon}_2\geq 0$, and $\tilde{\varepsilon}_3\in [0,1]$:

$$\tilde{E}_1(\tilde{\varepsilon}_1, R) := \left\{ \|ZZ^\top - RI_n\| \le \frac{\tilde{\varepsilon}_1}{3}R \right\},\tag{6}$$

$$\tilde{E}_2(\tilde{\varepsilon}_2, R) := \left\{ \| Z \boldsymbol{\mu} \| \le \frac{\tilde{\varepsilon}_2}{3} R \right\},\tag{7}$$

$$\tilde{E}_3(\tilde{\varepsilon}_3) := \left\{ |n_+ - \frac{n}{2}| \le \tilde{\varepsilon}_3 \frac{n}{2} \right\}. \tag{8}$$

Event \tilde{E}_1 captures near orthogonality of z_i 's to each other and uniformity of their norms, and event \tilde{E}_2 captures near orthogonality of z_i 's to μ . Although we could use the same parameters as in the analysis of directional convergence, these events \tilde{E}_1, \tilde{E}_2 makes the analysis of \hat{W} simpler.

With $\tilde{\varepsilon} := \max\{\tilde{\varepsilon}_1, \sqrt{n}\tilde{\varepsilon}_2\}$, $q_+ = |J_+|/m$, $q_- = |J_-|/m$, and $q_\gamma = \min\{q_+ + \gamma^2 q_-, q_- + \gamma^2 q_+\}$, we can now present our main result on the classification performance of \hat{W} :

Theorem 5.1 (Classification Error Bounds). Suppose event $\bigcap_{i=1,2,3} \tilde{E}_i$ holds with $\tilde{\varepsilon} \leq \frac{q_{\gamma}}{2}$, $\tilde{\varepsilon}_3 \leq \frac{1}{2}$, and further assume one of the following condition holds for a constant $C = C(\gamma, q_+)$:

(i)
$$Cn\|\boldsymbol{\mu}\|^2 \le R$$
, $C\tilde{\varepsilon} \le n^{-\frac{1}{2}}$, and $C\tilde{\varepsilon}_2 \le \frac{\sqrt{n}\|\boldsymbol{\mu}\|^2}{R}$,

(ii)
$$q_{\gamma} > \gamma$$
, $n \|\boldsymbol{\mu}\|^2 \ge CR$, $C\tilde{\varepsilon} \le \frac{R}{n^{\frac{3}{2}} \|\boldsymbol{\mu}\|^2}$.

Then, there exists constants c, N which depend on γ, q_+ such that $n \geq N$ implies

$$\mathbb{P}_{(\boldsymbol{x},y)}(yf(\boldsymbol{x};\hat{W})<0) \leq \begin{cases} \exp\left(-c\frac{n\|\boldsymbol{\mu}\|^4}{\|\boldsymbol{z}\|_{\psi_2}^2\{n\|\boldsymbol{\mu}\|^2+R\}}\right) & \textit{for model (sG)}, \\ c\|\boldsymbol{\Sigma}\|\left(\frac{1}{\|\boldsymbol{\mu}\|^2}+\frac{R}{n\|\boldsymbol{\mu}\|^4}\right) & \textit{for model (PM)}. \end{cases}$$

If $z \sim \mathcal{N}(0, \Sigma)$, then we have instead

$$\kappa \left\{ c^{-1} \beta_{\min}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R} \right)^{\frac{1}{2}} \right\} \leq \mathbb{P}_{(\boldsymbol{x}, y)}(y f(\boldsymbol{x}; \hat{W}) < 0) \leq \kappa \left\{ c \beta_{\max}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R} \right)^{\frac{1}{2}} \right\},$$

where $\kappa(t) = \mathbb{P}(|\xi_1| \geq t)$ and β_{min} is the smallest eigenvalue of Σ and $\beta_{max} = ||\Sigma||$.

This is a concise version for the purpose of presentation. A detailed version of the theorem and the proof are given in Theorem C.13, Appendix C.

Theorem 5.1 shows that there is a phase transition between the weak signal regime, i.e. $n\|\mu\|^2 \lesssim R$ and the strong signal regime, i.e. $n\|\mu\|^2 \gtrsim R$. The weak signal regime corresponds to the nearly

orthogonal data regime studied in prior work under specific distributional settings (Frei et al. (2022); Xu and Gu (2023)). The lower bound in the case of Gaussian mixture implies that the phase transition is indeed a feature of the model. Previously, this type of phase transition was not shown even with gradient flow. Moreover, it was only known to occur for binary linear classification (Cao et al. (2021); Wang and Thrampoulidis (2022); Hashimoto et al. (2025)).

Moreover, the assumptions of Theorem 5.1 are given in a deterministic manner. We confirm in the next section that these assumptions are satisfied with high probability under model (sG) and (PM), and conjecture that they can also be verified in other settings.

Lastly, we note that the condition $q_{\gamma} > \gamma$ in (iii) is necessary to ensure the equivalence of \hat{w} and the minimum norm least square estimator. We can even prove that the equivalence fails if $q_{\gamma} < \gamma$, which can be viewed as another form of the phase transition in the behavior of the convergent direction. Additional details are provided in Appendix C.

6 BENIGN OVERFITTING ON MIXTURE MODELS

We here demonstrate that the results of Theorem 4.8 and 5.1 hold with high probability in model (sG) and (PM) under sufficient over-parametrization. For (sG) we make the following assumptions:

Assumption 6.1. One of the following conditions holds for a constant C

$$\begin{split} (a) \;\; n \geq C, \, \alpha \leq \frac{1}{8 \mathrm{tr}(\Sigma)}, \, \sigma \leq 0.1 \gamma \alpha \sqrt{\frac{\mathrm{tr}(\Sigma)}{m}}, \, \| \pmb{\mu} \|^2 \geq C \| \Sigma^{]} f rac 12 \pmb{\mu} \|, \\ \mathrm{tr}(\Sigma) \geq C \max \left\{ n \| \pmb{\mu} \|^2, n \| \Sigma \|_F, n^{\frac{3}{2}} \| \Sigma \|, n^{\frac{3}{2}} \| \Sigma^{\frac{1}{2}} \pmb{\mu} \|, n \frac{\| \Sigma^{\frac{1}{2}} \pmb{\mu} \|^2}{\| \pmb{\mu} \|^2} \right\}, \end{split}$$

and further suppose one of the following:

(i)
$$\|\boldsymbol{\mu}\|^2 \ge C \max \left\{ \sqrt{n} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\|, \sqrt{n} \|\Sigma\|_F, n \|\Sigma\| \right\}$$

(ii)
$$\operatorname{tr}(\Sigma) \ge C \max \left\{ n^{\frac{3}{2}} \| \Sigma^{\frac{1}{2}} \boldsymbol{\mu} \|, n^{\frac{3}{2}} \| \Sigma \|_F, n^2 \| \Sigma \| \right\}$$

$$\begin{split} (b) \ \ q_{\gamma} > \gamma, \, n \geq C, \, \alpha \leq \frac{1}{2.4n\|\boldsymbol{\mu}\|^2}, \, \sigma \leq 0.2\gamma\alpha\sqrt{\frac{\|\boldsymbol{\mu}\|^2 + \text{tr}(\Sigma)}{m}}, \\ \text{tr}(\Sigma) \geq C \max\Big\{n\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|_F^{\frac{1}{2}}, n^{5/4}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{5}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{2}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}, n^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}\|\|\boldsymbol{\mu}\|\|^{\frac{1}{4}}\|\|\boldsymbol{\mu}\|\|\boldsymbol{\Sigma}\|\|^{\frac{1}{4}}\|\|\boldsymbol{$$

We note that condition (a) above corresponds to the weak signal regime and (b) to the strong signal regime in Theorem 5.1. We are now ready to present our main classification error bound on (sG):

Theorem 6.2. Consider model (sG). Let $\delta \in (0, \frac{1}{4})$. Suppose Assumption 6.1 holds for a sufficiently large constant C which depends only on δ, γ, q_+ . Then, we have with probably at least $1 - 3\delta$: the gradient descent iterate (2) keeps all the neurons activated for $t \geq 1$, satisfies $\mathcal{L}(W^{(t)}) = O(t^{-1})$, converges in direction, and the convergent direction \hat{W} attains

$$\mathbb{P}_{(\boldsymbol{x},y)}(yf(x;\hat{W})<0) \le \exp\left(-\frac{c}{L^2} \frac{n\|\boldsymbol{\mu}\|^4}{\|\boldsymbol{\Sigma}\|\{n\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})\}}\right),$$

where c is a constant which depends only on γ and q_+ .

If $z \sim \mathcal{N}(0, \Sigma)$, then we have instead

$$\kappa \left\{ c^{-1} \beta_{min}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})} \right)^{\frac{1}{2}} \right\} \leq \mathbb{P}_{(\boldsymbol{x},y)}(y f(\boldsymbol{x}; \hat{W}) < 0) \leq \kappa \left\{ c \beta_{max}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})} \right)^{\frac{1}{2}} \right\}.$$

Theorem 6.2 implies that benign overfitting occurs if $n\|\boldsymbol{\mu}\|^4 \gg \|\boldsymbol{\Sigma}\| \operatorname{tr}(\boldsymbol{\Sigma})$ under (a) or $\|\boldsymbol{\mu}\|^2 \gg \|\boldsymbol{\Sigma}\|$ under (b) is satisfied. For Gaussian mixture model, we see that benign overfitting provably fails even though directional convergence occurs if $n\|\boldsymbol{\mu}\|^4 \lesssim \beta_{min}\operatorname{tr}(\boldsymbol{\Sigma})$ under (a) or $\|\boldsymbol{\mu}\|^2 \lesssim \beta_{min}$ under (b).

We note that Theorem 6.2 under Assumption 6.1 (a) (i) and (b) are novel regimes which were not identified in the existing literature (Frei et al. (2022); Xu and Gu (2023)). Only the result under Assumption 6.1 (a) (ii) with $\Sigma = I_p$ was previously studied. A more detailed comparison with the prior work is given in Appendix A.

Moreover, Theorem 4.8 and 5.1 allow us to extend the above result to a heavier tailed model (PM). This is a significant distributional relaxation from the existing results which assumed z is sub-Gaussian and strongly log-concave (Frei et al. (2022)) or z has a bounded log-Sobolev constant (Xu and Gu (2023)). For brevity, we only present a result for $\Sigma = I_p$:

Theorem 6.3. Consider model (PM) with $\Sigma = I_p$. Then, for sufficiently large n, benign overfitting occurs with the convergent direction \hat{W} with high probability if either one of the following holds:

(i)
$$\|\mu\| \gtrsim \sqrt{n} \text{ and } \max\{n\|\mu\|^2, n^{\frac{r+4}{2r-4}}, n^{\frac{8}{r}+1}\} \lesssim p \lesssim \min\left\{\frac{\|\mu\|^4}{n^{\frac{8}{r}}}, \frac{\|\mu\|^r}{n}\right\}$$

$$\begin{aligned} & (ii) \;\; \| \boldsymbol{\mu} \| \gtrsim \sqrt{n} \; \text{and} \\ & \max \{ n^{\frac{5}{4}} \| \boldsymbol{\mu} \|^{\frac{3}{2}}, n^{\frac{8}{3r}+1} \| \boldsymbol{\mu} \|^{\frac{4}{3}}, n^{\frac{3r+4}{3r-4}} \| \boldsymbol{\mu} \|^{\frac{4r}{3r-4}} \} \lesssim p \lesssim \min \Big\{ n \| \boldsymbol{\mu} \|^2, \frac{\| \boldsymbol{\mu} \|^4}{n^{\frac{8}{3r}}}, \frac{\| \boldsymbol{\mu} \|^r}{n} \Big\}, \end{aligned}$$

(iii)
$$\max\{n\|{\pmb{\mu}}\|^2, n^{\frac{r+4}{2r-4}}, n^{\frac{8}{r}+2}\} \lesssim p \ll n\|{\pmb{\mu}}\|^4.$$

Detailed versions of the theorems and the proofs are given in Appendix D.

7 Proof Sketch

In this section, we provide the proof sketch of our main theorems. The key to establish the classification error bound in Theorem 5.1 is the precise characterization of the convergent direction in Theorem 4.8. The precise characterization allows us to obtain both upper and lower bounds of the key quantity $\frac{\|\bar{w}\|}{\langle \bar{w}, \mu \rangle}$ in classification error bounds (98), which results in the novel lower bound in the case of Gaussian mixture. We also note that the conditions in Theorem 4.8 and 5.1 are given in a deterministic manner. The separation of deterministic and distributional arguments allows us to substantially relax distributional assumptions made in prior work. In fact, Theorem 6.2 and 6.3 are proved by showing that all the conditions in Theorem 4.8 and 5.1 are achieved with high probability under (sG) and (PM) with sufficient overparametrization.

Proof sketch of Theorem 4.8 The key to show directional convergence of the gradient descent is activation of neurons. We say that j-th neuron $\phi(\langle \cdot, \boldsymbol{w}_j \rangle)$ is activated if $a_j y_i \phi(\langle \boldsymbol{x}_i, \boldsymbol{w}_j \rangle) > 0$ holds for any $i \in I$, which can be equivalently written as $a_j y_i \langle \boldsymbol{x}_i, \boldsymbol{w}_j \rangle > 0$ for any $i \in I$.

The next lemma shows that if all the neurons stay activated after some step of gradient descent (2), directional convergence occurs towards the maximum margin vector $\hat{\boldsymbol{w}} := \text{vec}(\hat{W})$ in a transformed sample space, i.e. \hat{W} is the solution to the following optimization problem:

Minimize
$$\|\boldsymbol{w}\|^2$$
 s.t. $\langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w} \rangle \geq 1$ for all $i \in I$, where $\boldsymbol{w} = \text{vec}(W)$, (9)

where $\tilde{\boldsymbol{x}}_{i}^{\top} = (\tilde{\boldsymbol{x}}_{i,1}^{\top}, \tilde{\boldsymbol{x}}_{i,2}^{\top}, \dots, \tilde{\boldsymbol{x}}_{i,m}^{\top}) \in \mathbb{R}^{1 \times mp}$ with $\tilde{\boldsymbol{x}}_{i,j} = a_{j}\zeta(a_{j}y_{i})\boldsymbol{x}_{i}$. Letting $\sigma_{max}(\tilde{X})$ be the maximum singular value of $\tilde{X} \in \mathbb{R}^{n \times mp}$ whose rows consist of $\tilde{\boldsymbol{x}}_{i}^{\top}$'s, we have

Lemma 7.1. Suppose all the neurons are activated for $t \ge T$ and $\alpha \sigma_{max}^2(\tilde{X}) < 2$. Then, the gradient descent iterate (2) satisfies $\mathcal{L}(W^{(t)}) = O(t^{-1})$ and $W^{(t)}$ converges in the direction characterized by Theorem 4.8.

The proof is done by reducing the argument to that of binary linear classification, i.e. classifiers of the form $\hat{y} = \text{sgn}(\langle \boldsymbol{x}, \boldsymbol{w} \rangle)$. In fact, if all the neurons are activated for all (\boldsymbol{x}_i, y_i) , $i = 1, \ldots, n$ at step t, then we have

$$y_i f(\boldsymbol{x}_i; W^{(t)}) = \langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w}^{(t)} \rangle,$$

where $\boldsymbol{w}^{(t)} = \text{vec}(W^{(t)})$. Therefore, if all the neurons stay activated after $t \geq T$, gradient descent iterate (2) becomes that of linear classifier in a transformed space, and the directional convergence follows from Theorem 3, Soudry et al. (2018).

To establish sufficient conditions for the neuron activation, observe that, for any i, j, we have

$$a_{j}\langle y_{i}\boldsymbol{x}_{i}, \boldsymbol{w}_{j}^{(t+1)} - \boldsymbol{w}_{j}^{(t)} \rangle$$

$$= \frac{\alpha}{m} \Big[\zeta(\langle \boldsymbol{x}_{i}, \boldsymbol{w}_{j}^{(t)} \rangle) \|\boldsymbol{x}_{i}\|^{2} \exp(-y_{i}f(\boldsymbol{x}_{i}; W^{(t)})) +$$

$$+ \sum_{k \neq i} \zeta(\langle \boldsymbol{x}_{k}, \boldsymbol{w}_{j}^{(t)} \rangle) \langle y_{k}\boldsymbol{x}_{k}, y_{i}\boldsymbol{x}_{i} \rangle \exp(-y_{k}f(\boldsymbol{x}_{k}; W^{(t)})) \Big].$$

$$(10)$$

The only source of negativity in (10) are $\langle y_i x_i, y_k x_k \rangle$, $i \neq k$. We note that event E implies

$$\left| \langle y_i \boldsymbol{x}_i, y_k \boldsymbol{x}_k \rangle - \|\boldsymbol{\mu}\|^2 \right| \le 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z}), \forall i \ne k.$$
 (11)

So, by keeping $2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$ small (Assumption 4.1 and 4.4), the update (10) should stay positive. Then, it should suffice to make all the neurons activated at step t=1, which can be done by taking initialization σ sufficiently smaller than the step size α (Assumption 4.2 and 4.5).

Proof sketch of Theorem 5.1. The detailed characterizations of the convergent direction given by Theorem 4.8 allows us to establish the desired classification error bound. This is done by obtaining a closed form expression of $\hat{\boldsymbol{w}} = \text{vec}(\hat{W})$, which is $\hat{\boldsymbol{w}} = \tilde{X}^{\top}(\tilde{X}\tilde{X}^{\top})^{-1}\boldsymbol{y}$, where $\boldsymbol{y} = (y_1,\ldots,y_n)^{\top} \in \mathbb{R}^n$. To establish $\hat{\boldsymbol{w}} = \tilde{X}^{\top}(\tilde{X}\tilde{X}^{\top})^{-1}\boldsymbol{y}$, it is necessary to understand the behavior of $\boldsymbol{y}^{\top}(\tilde{X}\tilde{X}^{\top})^{-1}$. Letting $X = (\boldsymbol{x}_1^{\top}, \boldsymbol{x}_2^{\top}, \ldots, \boldsymbol{x}_n^{\top})^{\top} \in R^{n \times p}$ and recalling $\tilde{\boldsymbol{x}}_{ij} = a_j \zeta(y_i a_j) \boldsymbol{x}_i$, we can write

$$\tilde{X} = (a_1 B_1 X \quad a_2 B_2 X \quad \cdots \quad a_m B_m X), \tag{12}$$

where $B_j \in \mathbb{R}^{n \times n}$ is the diagonal matrix whose diagonal entry $(B_j)_{kk} = \zeta(a_j y_k)$. Then we have

$$\tilde{X}\tilde{X}^{\top} = \sum_{j \in J} a_j^2 B_j X X^{\top} B_j. \tag{13}$$

Notice that B_j 's are the same among $j \in J_+$ and $j \in J_-$, respectively. So, we can write $B_+ = B_j, j \in J_+$ and $B_- = B_j, j \in J_-$. With $a_j^2 = \frac{1}{m}$, we have

$$\tilde{X}\tilde{X}^{\top} = \frac{|J_{+}|}{m} B_{+} X X^{\top} B_{+} + \frac{|J_{-}|}{m} B_{-} X X^{\top} B_{-}. \tag{14}$$

The analysis of $y(\tilde{X}\tilde{X}^{\top})^{-1}$ will be done through $XX^{\top} = \|\mu\|^2 yy^{\top} + y(Z\mu)^{\top} + Z\mu y^{\top} + ZZ^{\top} \approx \|\mu\|^2 yy^{\top} + RI_n$ under sufficient near orthogonality of z_i 's to each other and to μ (events \tilde{E}_1, \tilde{E}_2).

The rest of the argument is split into the strong signal regime and the weak signal regime. The latter is easier since we have $XX^{\top} \approx RI_n$, which further implies $\tilde{X}\tilde{X}^{\top} \approx R(q_+B_+^2 + q_-B_-^2)$, which is a diagonal matrix. The strong signal case is more difficult to analyze, and relies on a careful characterization of the eigenvectors of $\|\boldsymbol{\mu}\|^2(q_+B_+\boldsymbol{y}\boldsymbol{y}^{\top}B_+ + q_-B_-\boldsymbol{y}\boldsymbol{y}^{\top}B_-) + R(q_+B_+^2 + q_-B_-^2)$.

8 DISCUSSION

In this paper, we have established sufficient conditions of benign overfitting of gradient descent in leaky ReLU two-layer neural networks. Here we discuss directions for future research other than the obvious direction of extending to deep neural networks:

Not only showing directional convergence in more generality but also obtaining detailed characterization of the convergent direction remains an important open question. Even for two-layer neural networks we studied, our setting is rather restricted considering a more general result obtained by Ji and Telgarsky (2020) for gradient flow.

In this study, we did not introduce noise to the label y as was done in some prior work on binary classification (Chatterji and Long (2021); Wang and Thrampoulidis (2022); Frei et al. (2022); Xu and Gu (2023); Hashimoto et al. (2025)). We conjecture that introducing noise will not change the result for the weak signal regime since in this regime the signal μ is negligible. However, the situation is likely to change substantially in the strong signal regime with noisy data as was observed in Hashimoto et al. (2025) for binary linear classification.

REFERENCES

- Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 322–332. PMLR, 09–15 Jun 2019. URL https://proceedings.mlr.press/v97/arora19a.html.
- Peter L. Bartlett, Philip M. Long, Gábor Lugosi, and Alexander Tsigler. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020. doi: 10.1073/pnas.1907378117. URL https://www.pnas.org/doi/abs/10.1073/pnas.1907378117.
- Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias-variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019. doi: 10.1073/pnas.1903070116. URL https://www.pnas.org/doi/abs/10.1073/pnas.1903070116.
- Phuong Bui Thi Mai and Christoph Lampert. The inductive bias of ReLU networks on orthogonally separable data. In 9th International Conference on Learning Representations, 2021.
- Yuhang Cai, Kangjie Zhou, Jingfeng Wu, Song Mei, Michael Lindsey, and Peter L. Bartlett. Implicit bias of gradient descent for non-homogeneous deep networks, 2025. URL https://arxiv.org/abs/2502.16075.
- Yuan Cao, Quanquan Gu, and Mikhail Belkin. Risk bounds for over-parameterized maximum margin classification on sub-gaussian mixtures. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 8407–8418. Curran Associates, Inc., 2021. URL https://proceedings.neurips.cc/paper_files/paper/2021/file/46e0eae7d5217c79c3ef6b4c212b8c6f-Paper.pdf.
- Yuan Cao, Zixiang Chen, Misha Belkin, and Quanquan Gu. Benign overfitting in two-layer convolutional neural networks. In S. Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, *Advances in Neural Information Processing Systems*, volume 35, pages 25237–25250. Curran Associates, Inc., 2022. URL https://proceedings.neurips.cc/paper_files/paper/2022/file/a12c999be280372b157294e72a4bbc8b-Paper-Conference.pdf.
- Niladri S. Chatterji and Philip M. Long. Finite-sample analysis of interpolating linear classifiers in the overparameterized regime. *Journal of Machine Learning Research*, 22(129):1–30, 2021. URL http://jmlr.org/papers/v22/20-974.html.
- Spencer Frei, Niladri S Chatterji, and Peter Bartlett. Benign overfitting without linearity: Neural network classifiers trained by gradient descent for noisy linear data. In Po-Ling Loh and Maxim Raginsky, editors, *Proceedings of Thirty Fifth Conference on Learning Theory*, volume 178 of *Proceedings of Machine Learning Research*, pages 2668–2703. PMLR, 02–05 Jul 2022. URL https://proceedings.mlr.press/v178/frei22a.html.
- Spencer Frei, Gal Vardi, Peter Bartlett, and Nathan Srebro. Benign overfitting in linear classifiers and leaky relu networks from kkt conditions for margin maximization. In Gergely Neu and Lorenzo Rosasco, editors, *Proceedings of Thirty Sixth Conference on Learning Theory*, volume 195 of *Proceedings of Machine Learning Research*, pages 3173–3228. PMLR, 12–15 Jul 2023a. URL https://proceedings.mlr.press/v195/frei23a.html.
- Spencer Frei, Gal Vardi, Peter L. Bartlett, Nathan Srebro, and Wei Hu. Implicit bias in leaky relu networks trained on high-dimensional data. 2023b. 11th International Conference on Learning Representations, ICLR 2023; Conference date: 01-05-2023 Through 05-05-2023.
- Ichiro Hashimoto, Stanislav Volgushev, and Piotr Zwiernik. Universality of benign overfitting in binary linear classification, 2025. URL https://arxiv.org/abs/2501.10538.

Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *The Annals of Statistics*, 50(2):949–986, 2022. doi: 10.1214/21-aos2133. URL https://doi.org/10.1214/21-aos2133.

- Daniel Hsu, Vidya Muthukumar, and Ji Xu. On the proliferation of support vectors in high dimensions. In Arindam Banerjee and Kenji Fukumizu, editors, *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 91–99. PMLR, 13–15 Apr 2021. URL https://proceedings.mlr.press/v130/hsu21a.html.
- Arthur Jacot, Franck Gabriel, and Clement Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper_files/paper/2018/file/5a4be1fa34e62bb8a6ec6b91d2462f5a-Paper.pdf.
- Ziwei Ji and Matus Telgarsky. Directional convergence and alignment in deep learning. In H. Larochelle, M. Ranzato, R. Hadsell, M.F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 17176–17186. Curran Associates, Inc., 2020. URL https://proceedings.neurips.cc/paper_files/paper/2020/file/c76e4b2fa54f8506719a5c0dc14c2eb9-Paper.pdf.
- Yiwen Kou, Zixiang Chen, Yuanzhou Chen, and Quanquan Gu. Benign overfitting in two-layer ReLU convolutional neural networks. In Andreas Krause, Emma Brunskill, Kyunghyun Cho, Barbara Engelhardt, Sivan Sabato, and Jonathan Scarlett, editors, *Proceedings of the 40th International Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pages 17615–17659. PMLR, 23–29 Jul 2023a. URL https://proceedings.mlr.press/v202/kou23a.html.
- Yiwen Kou, Zixiang Chen, and Quanquan Gu. Implicit bias of gradient descent for two-layer relu and leaky relu networks on nearly-orthogonal data. In A. Oh, T. Naumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 30167–30221. Curran Associates, Inc., 2023b. URL https://proceedings.neurips.cc/paper_files/paper/2023/file/602f5c1b803c53b2aaf0b3864bf3383a-Paper-Conference.pdf.
- Tengyuan Liang and Alexander Rakhlin. Just interpolate: Kernel "ridgeless" regression can generalize. *The Annals of Statistics*, 48(3):1329–1347, June 2020. doi: 10.1214/19-aos1849. URL https://doi.org/10.1214/19-AOS1849.
- Tengyuan Liang, Alexander Rakhlin, and Xiyu Zhai. On the multiple descent of minimum-norm interpolants and restricted lower isometry of kernels. In Jacob Abernethy and Shivani Agarwal, editors, *Proceedings of Thirty Third Conference on Learning Theory*, volume 125 of *Proceedings of Machine Learning Research*, pages 2683–2711. PMLR, 09–12 Jul 2020. URL https://proceedings.mlr.press/v125/liang20a.html.
- Kaifeng Lyu and Jian Li. Gradient descent maximizes the margin of homogeneous neural networks. In *International Conference on Learning Representations*, 2020. URL https://openreview.net/forum?id=SJeLIgBKPS.
- Kaifeng Lyu, Zhiyuan Li, Runzhe Wang, and Sanjeev Arora. Gradient descent on two-layer nets: Margin maximization and simplicity bias. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 12978–12991. Curran Associates, Inc., 2021. URL https://proceedings.neurips.cc/paper_files/paper/2021/file/6c351da15b5e8a743a21ee96a86e25df-Paper.pdf.
- Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and the double descent curve. *Communications on Pure and Applied Mathematics*, 75(4):667–766, 2022. doi: 10.1002/cpa.22008. URL https://doi.org/10.1002/cpa.22008.

Hancheng Min, Enrique Mallada, and Rene Vidal. Early neuron alignment in two-layer reLU networks with small initialization. In *The Twelfth International Conference on Learning Representations*, 2024. URL https://openreview.net/forum?id=QibPzdVrRu.

- Vidya Muthukumar, Kailas Vodrahalli, Vignesh Subramanian, and Anant Sahai. Harmless interpolation of noisy data in regression. *IEEE Journal on Selected Areas in Information Theory*, 1 (1):67–83, 2020. doi: 10.1109/JSAIT.2020.2984716. URL https://doi.org/10.1109/JSAIT.2020.2984716.
- Roei Sarussi, Alon Brutzkus, and Amir Globerson. Towards understanding learning in neural networks with linear teachers. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 9313–9322. PMLR, 18–24 Jul 2021. URL https://proceedings.mlr.press/v139/sarussi21a.html.
- Sholom Schechtman and Nicolas Schreuder. The late-stage training dynamics of (stochastic) subgradient descent on homogeneous neural networks, 2025. URL https://arxiv.org/abs/2502.05668.
- Daniel Soudry, Elad Hoffer, Mor Shpigel Nacson, Suriya Gunasekar, and Nathan Srebro. The implicit bias of gradient descent on separable data. *Journal of Machine Learning Research*, 19(70):1–57, 2018.
- Alexander Tsigler and Peter L. Bartlett. Benign overfitting in ridge regression. *Journal of Machine Learning Research*, 24(1):123, 2024. ISSN 1532-4435.
- Gal Vardi. On the implicit bias in deep-learning algorithms, 2022. URL https://arxiv.org/abs/2208.12591.
- Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2018. doi: 10.1017/9781108231596.
- Ke Wang and Christos Thrampoulidis. Binary classification of gaussian mixtures: Abundance of support vectors, benign overfitting, and regularization. *SIAM Journal on Mathematics of Data Science*, 4(1):260–284, 2022. doi: 10.1137/21M1415121. URL https://doi.org/10.1137/21M1415121.
- Xingyu Xu and Yuantao Gu. Benign overfitting of non-smooth neural networks beyond lazy training. In Francisco Ruiz, Jennifer Dy, and Jan-Willem van de Meent, editors, *Proceedings of The 26th International Conference on Artificial Intelligence and Statistics*, volume 206 of *Proceedings of Machine Learning Research*, pages 11094–11117. PMLR, 25–27 Apr 2023. URL https://proceedings.mlr.press/v206/xu23k.html.
- Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *International Conference on Learning Representations*, 2017. URL https://openreview.net/forum?id=Sy8qdB9xx.

A COMPARISON WITH THE EXISTING RESULTS

 In this section, we provide more detailed comparison of our main results with the existing results (Frei et al. (2022); Xu and Gu (2023)). Since Frei et al. (2022) only considers smoothed approximation of leaky ReLU, our comparison is mostly with Xu and Gu (2023).

Since only isotropic sub-Gaussian mixture was studied in the existing literature, we present here sufficient conditions of benign overfitting derived from Theorem 6.2 in the case of $\Sigma = I_P$ for the purpose of direct comparison:

Corollary A.1. Consider model (sG) with $\Sigma = I_p$. Then, for sufficiently large n, benign overfitting occurs with the convergent direction \hat{W} with high probability if either one of the following holds:

(i)
$$\|\boldsymbol{\mu}\| \gtrsim \sqrt{n}$$
 and $n\|\boldsymbol{\mu}\|^2 \lesssim p \lesssim \frac{\|\boldsymbol{\mu}\|^4}{n}$,

(ii)
$$\|\mu\| \gtrsim \sqrt{n}$$
 and $n^{\frac{5}{4}} \|\mu\|^{\frac{3}{2}} \lesssim p \lesssim \min\{n\|\mu\|^2, \frac{\|\mu\|^4}{n}\},$

(iii)
$$\max\{n\|\mu\|^2, n^3\} \lesssim p \ll n\|\mu\|^4$$
.

We first note that conditions (i) and (ii) are newly identified regimes on which benign overfitting occurs, and only condition (iii) is directly comparable to the results studied in the prior work (Frei et al. (2022); Xu and Gu (2023)). As shown in Theorem 6.3, we have further extended the result in the case of (PM).

Data Regime: Our work considers both weak and strong signal regimes and identified the surprising phase transition in classification error occurring at $n\|\mu\|^2 \approx \operatorname{tr}(\Sigma)$. Frei et al. (2022); Xu and Gu (2023) only studied nearly orthogonal data regime, which corresponds to the weak signal regime in this paper, and thus did not identify the phase transition. Even in the weak signal regime, condition (i) in Corollary A.1 was not studied.

Distributional setting: Our results significantly relax distributional conditions in prior work by extending to anisotropic polynomially tailed mixture (PM). Both Frei et al. (2022); Xu and Gu (2023) considered mixtures even stronger than (sG). Specifically, Frei et al. (2022) assumed z is not only sub-Gaussian but also strongly log-concave. Xu and Gu (2023) assumed z has a bounded log-Sobolev constant, which is also stronger than just assuming z is sub-Gaussian. Moreover, both work only considered isotropic mixture.

Comparing condition (iii) with that of Xu and Gu (2023), we have a better upper bound on p (Xu and Gu (2023) requires $p\log(mp) \lesssim n\|\boldsymbol{\mu}\|^4$). We allow the width m to be arbitrary while Xu and Gu (2023) requires $m \gtrsim \log n$. We note that Xu and Gu (2023)'s result holds for non-smooth leaky ReLU like activation, which is not limited to (leaky) ReLU. We also note that our lower bound on p is slightly stronger than theirs which requires $p \gtrsim \max\{n\|\boldsymbol{\mu}\|^2, n^2\log n\}$.

B Proof of Directional Convergence

B.1 Proof of Lemma 7.1

Lemma B.1. Consider (M). If all the neurons are activated at step t, i.e. $a_j \langle \boldsymbol{w}_j^{(t)}, y_i \boldsymbol{x}_i \rangle > 0$ holds for all $i \in I$ and $j \in J$, then

$$\boldsymbol{w}_{j}^{(t+1)} - \boldsymbol{w}_{j}^{(t)} = \alpha \sum_{i \in I} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(t)})) a_{j} \zeta(a_{j} y_{i}) y_{i} \boldsymbol{x}_{i}.$$

Proof. By (2), we have

$$\begin{aligned} \boldsymbol{w}_{j}^{(t+1)} - \boldsymbol{w}_{j}^{(t)} &= -\alpha \sum_{i} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(t)}))(-y_{i}) \nabla_{\boldsymbol{w}_{j}} \left\{ \sum_{k} a_{k} \phi(\langle \boldsymbol{w}_{k}, \boldsymbol{x}_{i} \rangle) \right\}_{W = W^{(t)}} \\ &= \alpha \sum_{i} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(t)})) y_{i} a_{j} \nabla_{\boldsymbol{w}_{j}} \phi(\langle \boldsymbol{w}_{j}, \boldsymbol{x}_{i} \rangle)_{\boldsymbol{w}_{j} = \boldsymbol{w}_{j}^{(t)}} \\ &= \alpha \sum_{i} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(t)})) a_{j} \zeta(a_{j} y_{i}) y_{i} \boldsymbol{x}_{i}, \end{aligned}$$

where the last equality is due to the assumption $a_j y_i \langle \boldsymbol{w}_j^{(t)}, \boldsymbol{x}_i \rangle > 0$ which implies $\mathrm{sgn}(\langle \boldsymbol{w}_j^{(t)}, \boldsymbol{x}_i \rangle) = \mathrm{sgn}(a_j y_i)$.

Remark. Lemma B.1 implies that the gradient descent update is same within $j \in J_+$ and $j \in J_-$, respectively.

Lemma B.2. Suppose $W=(\boldsymbol{w}_1,\ldots,\boldsymbol{w}_m)$ has $\boldsymbol{w}_j=\boldsymbol{w}_+$ if $j\in J_+$ and $\boldsymbol{w}_j=\boldsymbol{w}_-$ if $j\in J_-$. Then, $f(\boldsymbol{x};W)$ has the linear decision boundary defined by $\bar{\boldsymbol{w}}:=\frac{|J_+|}{\sqrt{m}}\boldsymbol{w}_+-\frac{|J_-|}{\sqrt{m}}\boldsymbol{w}_-$, i.e. $\mathrm{sign}(f(\boldsymbol{x};W))=\mathrm{sign}(\langle \boldsymbol{x},\bar{\boldsymbol{w}}\rangle)$.

Proof. We consider several cases separately:

Case $\langle \boldsymbol{x}, \boldsymbol{w}_+ \rangle \geq 0$ and $\langle \boldsymbol{x}, \boldsymbol{w}_- \rangle \geq 0$: we have

$$f(\boldsymbol{x}; \hat{W}) = \frac{|J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle - \frac{|J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle = \langle \boldsymbol{x}, \overline{\boldsymbol{w}} \rangle.$$

Case $\langle \boldsymbol{x}, \boldsymbol{w}_+ \rangle \geq 0$ and $\langle \boldsymbol{x}, \boldsymbol{w}_- \rangle < 0$: we have

$$f(\boldsymbol{x}; \hat{W}) = \frac{|J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle - \frac{\gamma |J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle \geq \frac{|J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle - \frac{|J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle = \langle \boldsymbol{x}, \bar{\boldsymbol{w}} \rangle \geq 0.$$

Case $\langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle < 0$ and $\langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle \geq 0$: we have

$$0 > \langle \boldsymbol{x}, \overline{\boldsymbol{w}} \rangle = \frac{|J_+|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_+ \rangle - \frac{|J_-|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_- \rangle \ge \frac{\gamma |J_+|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_+ \rangle - \frac{|J_-|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_- \rangle = f(\boldsymbol{x}; \hat{W}).$$

Case $\langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle < 0$ and $\langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle < 0$: we have

$$f(\boldsymbol{x}; \hat{W}) = \gamma \left(\frac{|J_{+}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{+} \rangle - \frac{|J_{-}|}{\sqrt{m}} \langle \boldsymbol{x}, \boldsymbol{w}_{-} \rangle \right) = \gamma \langle \boldsymbol{x}, \bar{\boldsymbol{w}} \rangle.$$

Therefore, we have confirmed $\operatorname{sign}(f(\boldsymbol{x}:\hat{W})) = \operatorname{sign}(\langle \boldsymbol{x}, \bar{\boldsymbol{w}} \rangle)$ holds for all the cases above. \Box

Proof of Lemma 7.1. By the assumption that all the neurons stay activated after $t \ge T$, the directional convergence follows from Corollary 8 of Soudry et al. (2018).

For completeness and to show the remaining claims, we provide the complete proof of the directional convergence here.

First note that $a_j\phi(\langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)})\rangle = a_j\zeta(a_jy_i)\langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)}\rangle$ holds for all $i\in I, j\in J, t\geq T$ by the assumption that all the neurons are activated after $t\geq T$. Then, we have for $t\geq T$,

$$y_i f(\boldsymbol{x}_i; W^{(t)}) = \sum_j a_j y_i \phi(\langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle)$$

$$= \sum_j a_j y_i \zeta(a_j y_i) \langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle$$

$$= \sum_j \langle a_j \zeta(a_j y_i) y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle$$

$$= \sum_j \langle y_i \tilde{\boldsymbol{x}}_{ij}, \boldsymbol{w}_j^{(t)} \rangle$$

$$= \langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w}^{(t)} \rangle,$$

where the fourth equality follows from the definition of \tilde{x}_i and $w^{(t)} = \text{vec}(W^{(t)})$.

With this, the gradient descent updates in Lemma B.1 can be rewritten as

$$\boldsymbol{w}_{j}^{(t+1)} - \boldsymbol{w}_{j}^{(t)} = \alpha \sum_{i} \exp\left(-\langle y_{i}\tilde{\boldsymbol{x}}_{i}, \boldsymbol{w}^{(t)}\rangle\right) a_{j} \zeta(a_{j}y_{i}) y_{i} \boldsymbol{x}_{i}.$$

Therefore, we have

$$\boldsymbol{w}^{(t+1)} - \boldsymbol{w}^{(t)} = \alpha \sum_{i \in I} \exp(-\langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w}^{(t)} \rangle) y_i \tilde{\boldsymbol{x}}_i,$$
(15)

which can be viewed as the gradient descent iterates of binary linear classifier of the dataset $\{(\tilde{\boldsymbol{x}}_i,y_i)\}_{i\in I}\subset\mathbb{R}^{mp}\times\{\pm 1\}.$

Therefore, gradient descent iterates $\boldsymbol{w}^{(t)}$ converges in direction if the dataset $\{(\tilde{\boldsymbol{x}}_i, y_i)\}_{i \in I}$ is linearly separable and α is sufficiently small due to Theorem 3 of Soudry et al. (2018). We now provide additional details verifying that this result is applicable in our setting.

It is easy to see linear separability of $\{(\tilde{x}_i, y_i)\}_{i \in I}$ by expanding the following inner product

$$\langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w}^{(t)} \rangle = \sum_{j \in J} \langle y_i a_j \zeta(a_j y_i) \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle = \sum_{j \in J} \zeta(a_j y_i) a_j \langle y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle.$$
 (16)

Now it is clear that the inner product is positive by the assumption that all the neurons are activated at time $t \geq T$.

In addition, $u \mapsto e^{-u}$ is 1-Lipschitz on $(0, \infty)$ and (16) implies that $\langle y_i \tilde{x}_i, w^{(t)} \rangle$ stays in this region. Therefore, Theorem 3 of Soudry et al. (2018) can be used with $\beta = 1$.

Since $\{(\tilde{\boldsymbol{x}}_i,y_i)\}_{i\in I}$ is linearly separable and satisfies $\alpha\sigma_{max}^2(\tilde{X}) < 2$, $\lim_{t\to\infty} \|\boldsymbol{w}^{(t)}\| = \infty$ follows from Lemma 1 of Soudry et al. (2018), $\mathcal{L}(W^{(t)}) = O(t^{-1})$ follows from Theorem 5 of Soudry et al. (2018), and the directional convergence follows from Theorem 3 of Soudry et al. (2018), respectively.

Now we turn to the characterization of the convergent direction. First, we show that not only $\|\boldsymbol{w}^{(t)}\|$ but also $\|\boldsymbol{w}_j^{(t)}\|$ diverges to infinity for all $j \in J$. To see this, note that by Lemma B.1, we have for t > T

$$\boldsymbol{w}_{j}^{(t)} - \boldsymbol{w}_{j}^{(T)} = \alpha \sum_{i} \left(\sum_{s=T}^{t-1} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(s)})) \right) \zeta(a_{j} y_{i}) a_{j} y_{i} \boldsymbol{x}_{i}.$$
(17)

Since $\|\boldsymbol{w}^{(t)}\|$ is diverging to infinity, we must have $\|\boldsymbol{w}_j^{(t)}\|$ diverging to infinity for some j, and hence $\sum_{s=T}^{t-1} \exp(-y_i f(\boldsymbol{x}_i; W^{(s)})) \to \infty$ as $t \to \infty$ for some i. Denote such i as i_* .

Again by Lemma B.1, we have for any j,

$$\langle \boldsymbol{w}_{j}^{(t)} - \boldsymbol{w}_{j}^{(T)}, \boldsymbol{w}_{j}^{(T)} \rangle = \alpha \sum_{i} \left(\sum_{s=T}^{t-1} \exp(-y_{i} f(\boldsymbol{x}_{i}; W^{(s)})) \right) \zeta(a_{j} y_{i}) a_{j} \langle y_{i} \boldsymbol{x}_{i}, \boldsymbol{w}_{j}^{(T)} \rangle$$

$$\geq \alpha \sum_{s=T}^{t-1} \exp(-y_{i_{*}} f(\boldsymbol{x}_{i_{*}}; W^{(s)})) \zeta(a_{j} y_{i_{*}}) a_{j} \langle y_{i_{*}} \boldsymbol{x}_{i_{*}}, \boldsymbol{w}_{j}^{(T)} \rangle \to \infty,$$
(18)

where the inequality is due to the assumption that all the neurons are activated at $s \geq T$ and the divergence is due to $\sum_{s=T}^{t-1} \exp(-y_{i_*} f(\boldsymbol{x}_{i_*}; W^{(s)})) \to \infty$ and the fact that all neurons are activated at time T, implying $\langle y_{i_*} \boldsymbol{x}_{i_*}, \boldsymbol{w}_j^{(T)} \rangle \neq 0$. Since this holds for any j, we must have $\|\boldsymbol{w}_j\|$ diverging to infinity for all $j \in J$.

By Theorem 3 of Soudry et al. (2018), the convergent direction is equal to that of the solution to the following optimization problem:

Minimize
$$\|\boldsymbol{w}\|^2$$
 s.t. $\langle y_i \tilde{\boldsymbol{x}}_i, \boldsymbol{w} \rangle \ge 1, \forall i \in I,$ (19)

which is equivalently written as

Minimize
$$\sum_{j \in J} \|\boldsymbol{w}_j\|^2$$
 s.t. $\sum_{j \in J} \langle a_j y_i \zeta(a_j y_i) \boldsymbol{x}_i, \boldsymbol{w}_j \rangle \ge 1, \forall i \in I.$ (20)

Recalling that the gradient descent updates are same within $j \in J_+$ and $j \in J_-$, respectively, and that $\|\boldsymbol{w}_j^{(t)}\|$ diverges to infinity for all $j \in J$, the direction that $\boldsymbol{w}_j^{(t)}$ converges to must be the same for all $j \in J_+$ and the same for all $j \in J_-$. Thus the solution to the optimization problem (20) must have $\boldsymbol{w}_j = \boldsymbol{w}_+$ for all $j \in J_+$ and $\boldsymbol{w}_j = \boldsymbol{w}_-$ for all $j \in J_-$. With this, (20) can be further rewritten as the desired optimization problem (5).

Finally, the claim about the linear decision boundary follows from Lemma B.2

Before proceeding to the subsequent subsections where we prove Theorem 4.8, we introduce some ancillary results:

We first note that event E implies (11) and the following inequality:

$$(1 - \theta_2)\{\|\boldsymbol{\mu}\|^2 + R_{min}^2(\boldsymbol{z})\} \le \|y_i \boldsymbol{x}_i\|^2 \le (1 + \theta_2)\{\|\boldsymbol{\mu}\|^2 + R_{max}^2(\boldsymbol{z})\}, \forall i.$$
 (21)

We also note that the definition of ρ implies

$$\max|f(\boldsymbol{x}_i; W^{(0)})| \le \rho. \tag{22}$$

By (11) and (21), we obtain the following lemma, which is going to be useful to establish sufficient conditions for $\alpha \sigma_{max}^2(\tilde{X}) < 2$:

Lemma B.3.

$$\sigma_{max}^2(\tilde{X}) \le (1 + \theta_2)\{\|\boldsymbol{\mu}\|^2 + R_{max}^2(\boldsymbol{z})\} + (n - 1)\left[\|\boldsymbol{\mu}\|^2 + \{2\theta_2\|\boldsymbol{\mu}\|R_{max}(\boldsymbol{z}) + \theta_1R_{max}^2(\boldsymbol{z})\}\right]$$

Proof. Note that

$$|\langle \tilde{\boldsymbol{x}}_{i}, \tilde{\boldsymbol{x}}_{k} \rangle| = \left| \sum_{j} a_{j}^{2} \zeta(a_{j}y_{i}) \zeta(a_{j}y_{k}) \langle \boldsymbol{x}_{i}, \boldsymbol{x}_{k} \rangle \right|$$

$$\leq \sum_{j} \frac{\zeta(a_{j}y_{i}) \zeta(a_{j}y_{k})}{m} |\langle \boldsymbol{x}_{i}, \boldsymbol{x}_{k} \rangle|$$

$$\leq |\langle \boldsymbol{x}_{i}, \boldsymbol{x}_{k} \rangle|,$$
(23)

where the last inequality is due to $\gamma^2 \le \zeta(a_i y_i) \zeta(a_i y_k) \le 1$.

With this, (11), and (21), for any $v \in S^{n-1}$, we have

$$\begin{split} \|\tilde{X}^{\top}\boldsymbol{v}\|^{2} &= \sum_{i} v_{i}^{2} \|\tilde{\boldsymbol{x}}_{i}\|^{2} + \sum_{i,k:i\neq k} v_{i}v_{k}\langle \tilde{\boldsymbol{x}}_{i}, \tilde{\boldsymbol{x}}_{k}\rangle \\ &\leq (1+\theta_{2})\{\|\boldsymbol{\mu}\|^{2} + R_{max}^{2}(\boldsymbol{z})\} \sum_{i} v_{i}^{2} \\ &+ \left[\|\boldsymbol{\mu}\|^{2} + \{2\theta_{2}\|\boldsymbol{\mu}\|R_{max}(\boldsymbol{z}) + \theta_{1}R_{max}^{2}(\boldsymbol{z})\}\right] \sum_{i,k:i\neq k} |v_{i}v_{k}| \\ &\leq (1+\theta_{2})\{\|\boldsymbol{\mu}\|^{2} + R_{max}^{2}(\boldsymbol{z})\} \\ &+ (n-1)\left[\|\boldsymbol{\mu}\|^{2} + \{2\theta_{2}\|\boldsymbol{\mu}\|R_{max}(\boldsymbol{z}) + \theta_{1}R_{max}^{2}(\boldsymbol{z})\}\right], \end{split}$$

where in the last inequality we used $\sum_{i,k:i\neq k} |v_iv_k| \le n-1$, which follows from the Cauchy-Schwartz inequality.

B.2 PROOF OF THEOREM 4.8 UNDER CONDITION (I)

 Lemma B.4. Suppose event E holds with $\|\boldsymbol{\mu}\|^2 \ge 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$ and Assumption 4.2. Then, all the neurons stay activated for $t \ge 1$.

Proof of Lemma B.4. Under $\|\boldsymbol{\mu}\|^2 \ge 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$, the gradient descent updates (10) stays positive. Therefore, it is enough to confirm all the neurons are activated at t=1.

Again by (10), (21), (22), $\|\boldsymbol{\mu}\|^2 \ge 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$,

$$a_{j}\langle y_{i}\boldsymbol{x}_{i},\boldsymbol{w}_{j}^{(1)}\rangle$$

$$\geq \frac{1}{m} \left[\alpha\gamma(1-\theta_{2})\{\|\boldsymbol{\mu}\|^{2}+R_{min}^{2}(\boldsymbol{z})\}\right]$$

$$\times \exp\{-\sqrt{m}\sigma\max_{i}\|\boldsymbol{x}_{i}\|\}-\sqrt{m}\sigma\max_{i}\|\boldsymbol{x}_{i}\|\right]$$

$$\geq \frac{1}{m} \left[\alpha\gamma(1-\theta_{2})\{\|\boldsymbol{\mu}\|^{2}+R_{min}^{2}(\boldsymbol{z})\}\exp(-\rho)-\rho\right]>0.$$
(24)

Proof of Theorem 4.8 under condition (i). By Lemma B.4, all the neurons stay activated after $t \ge 1$.

By Assumption 4.3, it suffices to show $\sigma_{max}^2(\tilde{X}) \leq 2(n\|\boldsymbol{\mu}\|^2 + R_{max}^2(\boldsymbol{z}))$ to ensure $\alpha \sigma_{max}^2(\tilde{X}) < 2$.

By Lemma B.3 and $\|\boldsymbol{\mu}\|^2 \geq 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$ we have

$$\sigma_{max}^{2}(\tilde{X}) \leq (1 + \theta_{2})(\|\boldsymbol{\mu}\|^{2} + R_{max}^{2}(\boldsymbol{z})) + 2(n - 1)\|\boldsymbol{\mu}\|^{2}$$

$$\leq 2(\|\boldsymbol{\mu}\|^{2} + R_{max}^{2}(\boldsymbol{z})) + 2(n - 1)\|\boldsymbol{\mu}\|^{2}$$

$$\leq 2(n\|\boldsymbol{\mu}\|^{2} + R_{max}^{2}(\boldsymbol{z})).$$
(25)

Therefore, the directional convergence follows by Lemma 7.1.

B.3 Proof of Theorem 4.8 under condition (II)

First, it is useful to recognize the following inequality which follows from (10), (11), and (21):

$$\frac{\alpha}{m} \left[\gamma (1 - \theta_2) \{ \| \boldsymbol{\mu} \|^2 + R_{min}^2(\boldsymbol{z}) \} \exp(-y_i f(\boldsymbol{x}_i; W^{(t)})) \right. \\
+ \sum_{k:k \neq i} \zeta(\langle \boldsymbol{x}_k, \boldsymbol{w}_j^{(t)} \rangle) \{ \| \boldsymbol{\mu} \|^2 - 2\theta_2 \| \boldsymbol{\mu} \| R_{max}(\boldsymbol{z}) - \theta_1 R_{max}^2(\boldsymbol{z}) \} \exp(-y_k f(\boldsymbol{x}_k; W^{(t)})) \right] \\
\leq a_j \langle y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(t+1)} - \boldsymbol{w}_j^{(t)} \rangle \\
\leq \frac{\alpha}{m} \left[(1 + \theta_2) \{ \| \boldsymbol{\mu} \|^2 + R_{max}^2(\boldsymbol{z}) \} \exp(-y_i f(\boldsymbol{x}_i; W^{(t)})) \right. \\
+ \sum_{k:k \neq i} \{ \| \boldsymbol{\mu} \|^2 + 2\theta_2 \| \boldsymbol{\mu} \| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z}) \} \exp(-y_k f(\boldsymbol{x}_k; W^{(t)})) \right].$$
(26)

We start from establishing neuron activation at t = 1.

Lemma B.5. Suppose event E, Assumption 4.4 and 4.5 with $\varepsilon_1 + \varepsilon_2 < 1$. Then, we have for any i, j,

$$a_j \langle y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(1)} \rangle > 0,$$

and hence all the neurons are activated at t = 1.

Proof. By (22), (26), and the assumptions, we have

$$a_{j}\langle y_{i}\boldsymbol{x}_{i},\boldsymbol{w}_{j}^{(1)}\rangle \geq \frac{1}{m} \left[\alpha \left\{\gamma(1-\theta_{2})R_{min}^{2}(\boldsymbol{z})\exp(-\rho)\right.\right.$$
$$\left.\left.\left.\left.\left.\left(2\theta_{2}\|\boldsymbol{\mu}\|R_{max}(\boldsymbol{z})+\theta_{1}R_{max}^{2}(\boldsymbol{z})\right)\exp(\rho)\right\}-\rho\right]\right.$$
$$\geq \frac{1}{m} \left[(1-\varepsilon_{1})\alpha\gamma(1-\theta_{2})R_{min}^{2}(\boldsymbol{z})\exp(-\rho)-\rho\right]$$
$$\geq \frac{1}{m} (1-\varepsilon_{1}-\varepsilon_{2})\alpha\gamma(1-\theta_{2})R_{min}^{2}(\boldsymbol{z})\exp(-\rho)>0.$$

Thus, all the neurons are activated at t = 1 if $\varepsilon_1 + \varepsilon_2 < 1$.

Lemma B.6. Suppose event E, Assumption 4.4, 4.5, 4.6, and 4.7 with $\varepsilon_1 + \varepsilon_2 + \varepsilon_3 < 1$. Then we have

$$\max_{k \neq i} \frac{\exp(-y_i f(\boldsymbol{x}_i; W^{(1)}))}{\exp(-y_k f(\boldsymbol{x}_k; W^{(1)}))} \leq \exp(1) \quad \textit{and} \quad \alpha \sigma_{max}^2(\tilde{X}) < 2.$$

Proof. By (22), (26), and the assumptions, we have

$$a_{j}\langle y_{i}\boldsymbol{x}_{i},\boldsymbol{w}_{j}^{(1)}\rangle \leq \frac{1}{m} \left[\alpha\left\{(1+\theta_{2})\{n\|\boldsymbol{\mu}\|^{2}+R_{max}^{2}(\boldsymbol{z})\}\exp(\rho)\right.\right.$$
$$\left.+n\left\{2\theta_{2}\|\boldsymbol{\mu}\|R_{max}(\boldsymbol{z})+\theta_{1}R_{max}^{2}(\boldsymbol{z})\right\}\exp(\rho)\right\}+\rho\left]\right]$$
$$\leq \frac{1}{m} \left[\alpha\left\{(1+\theta_{2})(1+\varepsilon_{3})R_{max}^{2}(\boldsymbol{z})+\varepsilon_{1}R_{max}^{2}(\boldsymbol{z})\right\}\exp(\rho)+\rho\right]$$
$$\leq \frac{\alpha}{m}(1+\theta_{2})(1+\varepsilon_{1}+\varepsilon_{2}+\varepsilon_{3})R_{max}^{2}(\boldsymbol{z})\exp(\rho)$$
$$\leq \frac{4\alpha R_{max}^{2}(\boldsymbol{z})\exp(\rho)}{m}$$
$$\leq \frac{1}{m},$$

where the second inequality is due to Assumption 4.4 and 4.6, the third is due to Assumption 4.5, the fourth is due to $\varepsilon_1 + \varepsilon_2 < 1$, and the last inequality is due to Assumption 4.7.

Therefore, with Lemma B.5, we have $0 < y_i f(x_i; W^{(1)}) \le 1$, and hence

$$\max_{k \neq i} \frac{\exp(-y_i f(\boldsymbol{x}_i; W^{(1)}))}{\exp(-y_k f(\boldsymbol{x}_k; W^{(1)}))} \le \max_k \exp(y_k f(\boldsymbol{x}_k; W^{(1)})) \le \exp(1).$$

We now show $\alpha \sigma_{max}^2(\tilde{X}) < 2$. By Lemma B.3, we have

$$\alpha \sigma_{max}^{2}(\tilde{X}) \leq \alpha \left\{ (1 + \theta_{2}) \{n \| \boldsymbol{\mu} \|^{2} + R_{max}^{2}(\boldsymbol{z}) \} + \varepsilon_{1} (1 + \theta_{2}) R_{max}^{2}(\boldsymbol{z}) \right\}$$

$$\leq \alpha (1 + \theta_{2}) (1 + \varepsilon_{1} + \varepsilon_{3}) R_{max}^{2}(\boldsymbol{z})$$

$$\leq 4\alpha R_{max}^{2}(\boldsymbol{z}) < 2,$$

where the first inequality is due to Assumption 4.4, the second inequality is due to Assumption 4.6, and the last line is due to $\varepsilon_1 + \varepsilon_3 < 1$ and Assumption 4.7.

Finally, we prove that all the neurons stay activated after $t \geq 1$.

Lemma B.7. Suppose event E, Assumption 4.4, 4.5, 4.6, and 4.7 with $\varepsilon_1 + \varepsilon_2 + \varepsilon_3 < 1$, $C_w \varepsilon_i < \frac{1}{2}$, i = 1, 3, and $\theta_2 \leq \frac{1}{2}$. Then we have for any i, j

$$a_j \langle y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle > 0$$

and

$$\max_{k \neq i} \frac{\exp(-y_i f(\boldsymbol{x}_i; W^{(t)}))}{\exp(-y_k f(\boldsymbol{x}_k; W^{(t)}))} \leq C_w.$$
(27)

Proof. We prove both claims simultaneously by induction. By Lemma B.5 and B.6, both claims hold at step t = 1. Suppose both claims hold for $t \ge 1$.

We first show the first claim also holds for t + 1. By (26), Assumption 4.4, and the induction hypothesis, we have

$$a_{j}\langle y_{i}\boldsymbol{x}_{i},\boldsymbol{w}_{j}^{(t+1)}-\boldsymbol{w}_{j}^{(t)}\rangle \geq \frac{\alpha}{m}\exp(-y_{i}f(\boldsymbol{x}_{i};W^{(t)}))$$

$$\times \left\{\gamma(1-\theta_{2})R_{min}^{2}(\boldsymbol{z})-C_{w}\varepsilon_{1}\gamma(1-\theta_{2})R_{min}^{2}(\boldsymbol{z})\right\}$$

$$=\frac{1}{m}\gamma(1-C_{w}\varepsilon_{1})(1-\theta_{2})\alpha R_{min}^{2}(\boldsymbol{z})\exp(-y_{i}f(\boldsymbol{x}_{i};W^{(t)}))$$

$$\geq \frac{1}{4m}\gamma\alpha R_{min}^{2}(\boldsymbol{z})\exp(-y_{i}f(\boldsymbol{x}_{i};W^{(t)}))>0.$$
(28)

where the last line is due to $C_w \varepsilon_1 \leq \frac{1}{2}$ and $\theta_2 \leq \frac{1}{2}$. Therefore, all the neurons are active at step t+1.

Next, we prove (27). By (26), the assumptions, and the induction hypothesis, we have

$$a_{j}\langle y_{i}\boldsymbol{x}_{i}, \boldsymbol{w}_{j}^{(t+1)} - \boldsymbol{w}_{j}^{(t)}\rangle$$

$$\leq \frac{\alpha}{m} \exp(-y_{i}f(\boldsymbol{x}_{i}; W^{(t)}))$$

$$\times \left[(1+\theta_{2})\left(1+\frac{\varepsilon_{3}}{n}\right)R_{max}^{2}(\boldsymbol{z}) + C_{w}\varepsilon_{3}R_{max}^{2}(\boldsymbol{z}) + C_{w}\varepsilon_{1}(1+\theta_{2})R_{max}^{2}(\boldsymbol{z}) \right]$$

$$\leq \frac{1}{m} \left\{ 1+\frac{\varepsilon_{3}}{n} + C_{w}\varepsilon_{3} + C_{w}\varepsilon_{1} \right\} (1+\theta_{2})\alpha R_{max}^{2}(\boldsymbol{z}) \exp(-y_{i}f(\boldsymbol{x}_{i}; W^{(t)}))$$

$$\leq \frac{1}{m} 6\alpha R_{max}^{2}(\boldsymbol{z}) \exp(-y_{i}f(\boldsymbol{x}_{i}; W^{(t)})),$$
(29)

where the first inequality is due to Assumption 4.4 and 4.6, the last is due to the assumptions on ε_i .

Since all the neurons are activated at both t and t + 1, we have

$$a_j y_i \phi(\langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t+1)} \rangle) - a_j y_i \phi(\langle \boldsymbol{x}_i, \boldsymbol{w}_j^{(t)} \rangle) = \zeta(a_j y_i) a_j \langle y_i \boldsymbol{x}_i, \boldsymbol{w}_j^{(t+1)} - \boldsymbol{w}_j^{(t)} \rangle.$$

With this, (28) and (29), we have

$$\frac{\gamma^2}{4} \alpha R_{min}^2(\boldsymbol{z}) \exp(-y_i f(\boldsymbol{x}_i; W^{(t)}))$$

$$\leq y_i f(\boldsymbol{x}_i; W^{(t+1)}) - y_i f(\boldsymbol{x}_i; W^{(t)})$$

$$\leq 6\alpha R_{max}^2(\boldsymbol{z}) \exp(-y_i f(\boldsymbol{x}_i; W^{(t)})).$$
(30)

Now we fix $i \neq k$ and let $A_t = \frac{\exp(-y_i f(\boldsymbol{x}_i; W^{(t)}))}{\exp(-y_k f(\boldsymbol{x}_k; W^{(t)}))}$. By induction hypothesis, we have $A_t \leq C_w$. By (30), we have

$$A_{t+1} \leq A_t \exp\left[-\frac{\gamma^2}{4}\alpha R_{min}^2(\boldsymbol{z}) \exp(-y_i f(\boldsymbol{x}_i; W^{(t)})) + 6\alpha R_{max}^2(\boldsymbol{z}) \exp(-y_k f(\boldsymbol{x}_k; W^{(t)}))\right]$$

$$= A_t \exp\left[-\frac{\gamma^2}{4}\alpha R_{min}^2(\boldsymbol{z}) \exp(-y_k f(\boldsymbol{x}_k; W^{(t)})) \left\{A_t - \frac{24R_{max}^2(\boldsymbol{z})}{\gamma^2 R^2 + (\boldsymbol{z})}\right\}\right].$$
(32)

If $A_t \geq \frac{24R_{max}^2(z)}{\gamma^2 R_{min}^2(z)}$, then $A_{t+1} \leq A_t$ immediately follows from (32).

If $A_t < \frac{24R_{max}^2(z)}{\gamma^2 R_{min}^2(z)}$, then (31) and the fact that all neurons are activated at time t, yielding $y_i f(x_i; W^{(t)}) > 0$ for all i, implies

$$A_{t+1} \le \frac{24R_{max}^2(z)}{\gamma^2 R_{min}^2(z)} \exp\left[6\alpha R_{max}^2(z)\right]$$
$$\le \frac{24R_{max}^2(z)}{\gamma^2 R_{min}^2(z)} \exp(1) \le C_w,$$

where the last line holds from Assumption 4.7. This concludes the induction argument proving the desired claims. \Box

Proof of Theorem 4.8 under condition (ii). The conclusion now follows immediately from Lemma B.6, B.7, and Lemma 7.1. □

C Proof of Theorem 5.1

 The proof of Theorem 5.1 is done by first establishing the equivalence of the maximum margin vector and the minimum norm least square estimator, i.e. $\hat{\boldsymbol{w}} = \tilde{X}^{\top} (\tilde{X} \tilde{X}^{\top})^{-1} \boldsymbol{y}$, and carefully analyzing the behavior.

To prove $\hat{\boldsymbol{w}} = \tilde{X}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} \boldsymbol{y}$, we rely on the "proliferation of support vector" phenomenon due to Lemma 1, Hsu et al. (2021).

Lemma C.1. If $\mathbf{y}^{\top}(\tilde{X}\tilde{X}^{\top})^{-1}y_i\mathbf{e}_i > 0$ for all i, then

$$\hat{\boldsymbol{w}} = \tilde{X}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} \boldsymbol{y}.$$

Proof. By Lemma 1, Hsu et al. (2021), the assumption implies all (\tilde{x}_i, y_i) , i = 1, 2, ..., n are support vectors. Under this condition, the optimization problem defining \hat{w} is equivalent to

$$\hat{\boldsymbol{w}} = \arg\min \|\boldsymbol{w}\|^2$$
, subject to $\langle \boldsymbol{w}, y_i \tilde{\boldsymbol{x}}_i \rangle = 1$,

and is also equivalent to the optimization problem defining the least square estimator. \Box

As noted in Section 7, the analysis is done by establishing $\tilde{X}\tilde{X}^{\top} \approx \|\boldsymbol{\mu}\|^2 (q_+ B_+ \boldsymbol{y} \boldsymbol{y}^{\top} B_+ + q_- B_- \boldsymbol{y} \boldsymbol{y}^{\top} B_-) + R(q_+ B_+^2 + q_- B_-^2)$.

Lemma C.2. If event $\tilde{E}_1 \cap \tilde{E}_2$ holds, then the following inequality holds

$$||XX^{\top} - (||\boldsymbol{\mu}||^2 \boldsymbol{y} \boldsymbol{y}^{\top} + RI_n)|| \leq \tilde{\varepsilon} R,$$

where $\tilde{\varepsilon} := \max\{\tilde{\varepsilon}_1, \sqrt{n}\tilde{\varepsilon}_2\}.$

Proof. Noting event \tilde{E}_2 implies $\|\boldsymbol{y}(Z\boldsymbol{\mu})^\top\| \leq \frac{\tilde{\varepsilon}_2}{3}R$ and $XX^\top = \|\boldsymbol{\mu}\|^2\boldsymbol{y}\boldsymbol{y}^\top + \boldsymbol{y}(Z\boldsymbol{\mu})^\top + (Z\boldsymbol{\mu})\boldsymbol{y}^\top + ZZ^\top$, we have

$$||XX^{\top} - (||\boldsymbol{\mu}||^2 \boldsymbol{y} \boldsymbol{y}^{\top} + RI_n)|| \le ||\boldsymbol{y}(Z\boldsymbol{\mu})^{\top}|| + ||(Z\boldsymbol{\mu})\boldsymbol{y}^{\top}|| + ||ZZ^{\top} - RI_n||$$

$$\le \frac{\tilde{\varepsilon}}{3}R + \frac{\tilde{\varepsilon}}{3}R + \frac{\tilde{\varepsilon}}{3}R = \tilde{\varepsilon}R.$$

With (14), this establishes $\tilde{X}\tilde{X}^{\top} \approx \|\boldsymbol{\mu}\|^2 (q_+B_+\boldsymbol{y}\boldsymbol{y}^{\top}B_+ + q_-B_-\boldsymbol{y}\boldsymbol{y}^{\top}B_-) + R(q_+B_+^2 + q_-B_-^2)$. It is convenient to introduce $B^2 = q_+B_+^2 + q_-B_-^2$ where $B \in \mathbb{R}^{n \times n}$ is a diagonal matrix with

$$B_{ii} = \begin{cases} \sqrt{q_{+} + \gamma^{2} q_{-}} & \text{if } i \in I_{+} \\ \sqrt{q_{-} + \gamma^{2} q_{+}} & \text{if } i \in I_{-}. \end{cases}$$

Lemma C.3. If event $\tilde{E}_1 \cap \tilde{E}_2$ holds, then we have

$$\left\| B^{-1} \tilde{X} \tilde{X}^{\top} B^{-1} - A \right\| \le \tilde{\varepsilon} q_{\gamma}^{-1} R$$

where

$$A := \|\boldsymbol{\mu}\|^2 B^{-1} (q_+ B_+ \boldsymbol{y} \boldsymbol{y}^\top B_+ + q_- B_- \boldsymbol{y} \boldsymbol{y}^\top B_-) B^{-1} + R I_n.$$

To obtain approximation of $(\tilde{X}\tilde{X}^{\top})^{-1}$ from Lemma C.3, we need the following ancillary lemma:

Lemma C.4. If $||U-V|| \le sL$ and $V \succeq LI_n$ for some L > 0 and $s \in [0, \frac{1}{2}]$, then

$$||U^{-1} - V^{-1}|| \le 2sL^{-1}.$$

Proof. Since $V \succeq LI_n$ for L > 0, V is invertible and $||V^{-1}|| \leq L^{-1}$.

Then we have

$$||I_n - V^{-1}U|| \le ||V^{-1}|| ||U - V|| \le s.$$

Letting $S = I_n - V^{-1}U$, we have $||S|| \le s \le \frac{1}{2}$. So, U is invertible and

$$||(V^{-1}U)^{-1} - I_n|| \le ||S|| + ||S||^2 + \dots \le \frac{s}{1-s} \le 2s.$$

Combining with $||V^{-1}|| \le L^{-1}$, we conclude

$$||U^{-1} - V^{-1}|| \le ||V^{-1}|| ||(V^{-1}U)^{-1} - I_n|| \le 2sL^{-1}.$$

Now we are ready to obtain the following approximation of $(\tilde{X}\tilde{X}^{\top})^{-1}$, which our remaining arguments heavily relies on:

Lemma C.5. If event $\tilde{E}_1 \cap \tilde{E}_2$ holds with $\tilde{\varepsilon} \leq \frac{q_{\gamma}}{2}$, then we have

$$||B(\tilde{X}\tilde{X}^{\top})^{-1}B - A^{-1}|| \le \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R}.$$

Proof. Note that the additional assumption implies $\tilde{\epsilon}q_{\gamma}^{-1} \leq \frac{1}{2}$. Then, the desired conclusion follows from Lemma C.3 and C.4 by taking $U = B^{-1}\tilde{X}\tilde{X}^{\top}B^{-1}, V = A, s = \tilde{\epsilon}q_{\gamma}^{-1}$, and L = R.

By Lemma C.1 and C.5, the rest of our arguments is reduced to the analysis of $(B^{-1}y)^{\top}A^{-1}$. Here, we present technical results regarding this quantity which are used frequently later.

Lemma C.6. Let $y = \mathbf{1}_+ - \mathbf{1}_-$, where $(\mathbf{1}_+)_i = 1$ if $i \in I_+$ and $(\mathbf{1}_+)_i = 0$ if $i \in I_-$ and $(\mathbf{1}_-)_i = 0$ if $i \in I_+$. We have the following equalities:

1. if
$$q_+ = 1$$
, $q_- = 0$,

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1} = R^{-1}\left\{ \left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n} \right) \mathbf{1}_{+} - \left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n} \right) \mathbf{1}_{-} \right\}^{\top},$$
(33)

2. if
$$q_+ = 0$$
, $q_- = 1$,

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1} = R^{-1}\left\{ \left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1}n_{+} + n_{-}}{n} \right) \mathbf{1}_{+} - \left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1}n_{+} + n_{-}}{n} \right) \mathbf{1}_{-} \right\}^{\top},$$
(34)

3. if
$$q_+q_- \neq 0$$
,

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}$$

$$= \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}$$

$$\times \left[\sqrt{q_{+} + \gamma^{2}q_{-}}\{(q_{-} + \gamma^{2}q_{+} - \gamma)n_{-}\|\boldsymbol{\mu}\|^{2} + (q_{-} + \gamma^{2}q_{+})R\}\mathbf{1}_{+}\right]^{\top}$$

$$-\sqrt{q_{-} + \gamma^{2}q_{+}}\{(q_{+} + \gamma^{2}q_{-} - \gamma)n_{+}\|\boldsymbol{\mu}\|^{2} + (q_{+} + \gamma^{2}q_{-})R\}\mathbf{1}_{-}\right]^{\top},$$
(35)

where

$$d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})$$

$$= (1 - \gamma^{2})^{2}q_{+}q_{-}n_{+}n_{-}\|\boldsymbol{\mu}\|^{4} + (q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\{n\|\boldsymbol{\mu}\|^{2} + R\}R.$$
(36)

Proof. Let $A = \sum_{i=1}^{n} s_i \boldsymbol{u}_i \boldsymbol{u}_i^{\top}$ be the singular value decomposition of A with $s_1 \geq s_2 \geq \ldots \geq s_n$. We split in cases.

We first consider the case either q_+ or $q_-=0$, i.e. either $|J_+|$ or $|J_-|=0$. Then we have $B=B_+$ if $q_-=0$ and $B=B_-$ if $q_+=0$, and hence $A=\|\boldsymbol{\mu}\|^2\boldsymbol{y}\boldsymbol{y}^\top+RI_n$ either way. Then, we have $s_1=n\|\boldsymbol{\mu}\|^2+R,$ $s_i=R,$ $i\geq 2$, and can take $\boldsymbol{u}_1=\frac{\boldsymbol{y}}{\sqrt{n}}$. Thus,

$$A^{-1} = [n\|\boldsymbol{\mu}\|^{2} + R]^{-1} \frac{\boldsymbol{y}}{\sqrt{n}} \frac{\boldsymbol{y}^{\top}}{\sqrt{n}} + R^{-1} \sum_{i \geq 2} \boldsymbol{u}_{i} \boldsymbol{u}_{i}^{\top}$$

$$= R^{-1} \sum_{i=1}^{n} \boldsymbol{u}_{i} \boldsymbol{u}_{i}^{\top} - \frac{n\|\boldsymbol{\mu}\|^{2}}{[n\|\boldsymbol{\mu}\|^{2} + R]R} \frac{\boldsymbol{y}}{\sqrt{n}} \frac{\boldsymbol{y}^{\top}}{\sqrt{n}}$$

$$= R^{-1} \left(I_{n} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\boldsymbol{y}}{\sqrt{n}} \frac{\boldsymbol{y}^{\top}}{\sqrt{n}} \right).$$
(37)

Case $q_+=1, q_-=0$: In this case, we have $B=B_+$ as noted earlier. Since ${\boldsymbol y}={\bf 1}_+-{\bf 1}_-$ and $B_+^{-1}{\boldsymbol y}={\bf 1}_+-\gamma^{-1}{\bf 1}_-$, we have that

$$\begin{split} &(B_{+}^{-1}\boldsymbol{y})^{\top}A^{-1} \\ = &R^{-1}\left\{B_{+}^{-1}\boldsymbol{y} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\langle B_{+}^{-1}\boldsymbol{y}, \boldsymbol{y} \rangle}{n} \boldsymbol{y}\right\}^{\top} \\ = &R^{-1}\left\{\left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n}\right) \mathbf{1}_{+} \right. \\ &\left. - \left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n}\right) \mathbf{1}_{-}\right\}^{\top}. \end{split}$$

Case $q_+=0, q_-=1$: In this case, we have $B=B_-$ as noted earlier. Since ${\boldsymbol y}={\bf 1}_+-{\bf 1}_-$ and $B_-^{-1}{\boldsymbol y}=\gamma^{-1}{\bf 1}_+-{\bf 1}_-$, we have that

$$(B_{-}^{-1}\boldsymbol{y})^{\top}A^{-1}$$

$$=R^{-1}\left\{B_{-}^{-1}\boldsymbol{y} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\langle B_{-}^{-1}\boldsymbol{y}, \boldsymbol{y} \rangle}{n} \boldsymbol{y}\right\}^{\top}$$

$$=R^{-1}\left\{\left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1}n_{+} + n_{-}}{n}\right) \mathbf{1}_{+} - \left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1}n_{+} + n_{-}}{n}\right) \mathbf{1}_{-}\right\}^{\top}.$$

Case $q_+q_- \neq 0$: In this case, we have the span of $\{B^{-1}B_+\boldsymbol{y}, B^{-1}B_-\boldsymbol{y}\}$ equal to the span of $\{\boldsymbol{u}_1, \boldsymbol{u}_2\}$, and $s_i = R$ for $i \geq 3$.

To determine $\sum_{i=1,2} s_i^{-1} \boldsymbol{u}_i \boldsymbol{u}_i^{\top}$, it is convenient to use isomorphism between the span of $\{B^{-1}B_+\boldsymbol{y}, B^{-1}B_-\boldsymbol{y}\}$ and \mathbb{R}^2 by $\mathbf{1}_+ \leftrightarrow \sqrt{n_+}\boldsymbol{e}_1$ and $\mathbf{1}_- \leftrightarrow \sqrt{n_-}\boldsymbol{e}_2$. Under this isomorphism, we have

$$B^{-1}B_{+}y = \frac{1}{\sqrt{q_{+} + \gamma^{2}q_{-}}} \mathbf{1}_{+} - \frac{\gamma}{\sqrt{q_{-} + \gamma^{2}q_{+}}} \mathbf{1}_{-} \leftrightarrow \frac{\sqrt{n_{+}}}{\sqrt{q_{+} + \gamma^{2}q_{-}}} e_{1} - \frac{\gamma\sqrt{n_{-}}}{\sqrt{q_{-} + \gamma^{2}q_{+}}} e_{2},$$

$$B^{-1}B_{-}y = \frac{\gamma}{\sqrt{q_{+} + \gamma^{2}q_{-}}} \mathbf{1}_{+} - \frac{1}{\sqrt{q_{-} + \gamma^{2}q_{+}}} \mathbf{1}_{-} \leftrightarrow \frac{\gamma\sqrt{n_{+}}}{\sqrt{q_{+} + \gamma^{2}q_{-}}} e_{1} - \frac{\sqrt{n_{-}}}{\sqrt{q_{-} + \gamma^{2}q_{+}}} e_{2}.$$
(38)

Thus, $\sum_{i=1,2} s_i u_i u_i^{\top}$ is regarded as a 2×2 matrix

$$\begin{pmatrix}
n_{+} \|\boldsymbol{\mu}\|^{2} + R & -\frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} \\
-\frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} & n_{-} \|\boldsymbol{\mu}\|^{2} + R
\end{pmatrix}.$$
(39)

Thus, $\sum_{i=1,2} s_i^{-1} \boldsymbol{u}_i \boldsymbol{u}_i^{\top}$ is regarded as

$$\begin{pmatrix}
n_{+} \|\boldsymbol{\mu}\|^{2} + R & -\frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} \\
-\frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} & n_{-} \|\boldsymbol{\mu}\|^{2} + R
\end{pmatrix}^{-1}$$

$$= d^{-1} \begin{pmatrix}
n_{-} \|\boldsymbol{\mu}\|^{2} + R & \frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} \\
\frac{\gamma\sqrt{n_{+}n_{-}}\|\boldsymbol{\mu}\|^{2}}{\sqrt{(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})}} & n_{+} \|\boldsymbol{\mu}\|^{2} + R
\end{pmatrix}, (40)$$

where

$$d = \{n_{+} \|\boldsymbol{\mu}\|^{2} + R\} \{n_{-} \|\boldsymbol{\mu}\|^{2} + R\} - \frac{\gamma^{2} n_{+} n_{-} \|\boldsymbol{\mu}\|^{4}}{(q_{+} + \gamma^{2} q_{-})(q_{-} + \gamma^{2} q_{+})}.$$

Since $B^{-1}\boldsymbol{y} \leftrightarrow \frac{\sqrt{n_+}}{\sqrt{q_+ + \gamma^2 q_-}}\boldsymbol{e}_1 - \frac{\sqrt{n_-}}{\sqrt{q_- + \gamma^2 q_+}}\boldsymbol{e}_2$ and $B^{-1}\boldsymbol{y} \perp \boldsymbol{u}_i$ for all $i \geq 3$, we have

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}$$

$$=d^{-1}\left[\left\{\frac{n_{-}\|\boldsymbol{\mu}\|^{2}+R}{\sqrt{q_{+}+\gamma^{2}q_{-}}}-\frac{\gamma n_{-}\|\boldsymbol{\mu}\|^{2}}{\sqrt{q_{+}+\gamma^{2}q_{-}}(q_{-}+\gamma^{2}q_{+})}\right\}\mathbf{1}_{+}\right.$$

$$\left.-\left\{\frac{n_{+}\|\boldsymbol{\mu}\|^{2}+R}{\sqrt{q_{-}+\gamma^{2}q_{+}}}-\frac{\gamma n_{+}\|\boldsymbol{\mu}\|^{2}}{\sqrt{q_{-}+\gamma^{2}q_{+}}(q_{+}+\gamma^{2}q_{-})}\right\}\mathbf{1}_{-}\right]^{\top}$$

$$=\left\{d(q_{+}+\gamma^{2}q_{-})(q_{-}+\gamma^{2}q_{+})\right\}^{-1}$$

$$\times\left[\sqrt{q_{+}+\gamma^{2}q_{-}}\left\{(q_{-}+\gamma^{2}q_{+}-\gamma)n_{-}\|\boldsymbol{\mu}\|^{2}+(q_{-}+\gamma^{2}q_{+})R\right\}\mathbf{1}_{+}\right.$$

$$\left.-\sqrt{q_{-}+\gamma^{2}q_{+}}\left\{(q_{+}+\gamma^{2}q_{-}-\gamma)n_{+}\|\boldsymbol{\mu}\|^{2}+(q_{+}+\gamma^{2}q_{-})R\right\}\mathbf{1}_{-}\right]^{\top}.$$

$$(41)$$

 Before proceeding to the next lemma, we note that the following inequality holds under E_3 :

$$(1 - \hat{\varepsilon}_3^2) \frac{n^2}{4} \le n_+ n_- \le \frac{n^2}{4}. \tag{42}$$

The next lemma establishes useful bounds on d.

Lemma C.7. We have

$$d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \leq \left\{\gamma^{2} + \frac{5(1 - \gamma^{2})^{2}}{4}q_{+}q_{-}\right\} \left(n\|\boldsymbol{\mu}\|^{2} + R\right)^{2}$$

$$\leq \frac{5q_{\gamma}}{4} \left(n\|\boldsymbol{\mu}\|^{2} + R\right)^{2}.$$
(43)

If additionally $\tilde{E}_3(\tilde{\epsilon}_3)$ holds for some $\tilde{\epsilon}_3 \in [0,1]$, then we also have

$$d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \ge \frac{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}}{8} \left(n\|\boldsymbol{\mu}\|^{2} + R\right)^{2} + \gamma^{2}R^{2}.$$
(44)

Proof. We first note that

$$q_{\gamma} \ge (q_{+} + \gamma^{2} q_{-})(q_{-} + \gamma^{2} q_{+}) = (1 + \gamma^{4})q_{+}q_{-} + \gamma^{2}(q_{+}^{2} + q_{-}^{2}) = \gamma^{2} + (1 - \gamma^{2})^{2}q_{+}q_{-}.$$
 (45)

With (45) and also noting that $n_+ n_- \le \frac{n^2}{4}$, it is easy to see (43) follows from (36).

By (36), (45), and (42), we have

$$\begin{split} &d(q_{+}+\gamma^{2}q_{-})(q_{-}+\gamma^{2}q_{+})\\ \geq &\frac{(1-\tilde{\varepsilon}_{3}^{2})(1-\gamma^{2})^{2}}{4}q_{+}q_{-}\left(n\|\boldsymbol{\mu}\|^{2}\right)^{2}+\left\{\gamma^{2}+(1-\gamma^{2})^{2}q_{+}q_{-}\right\}R^{2}\\ \geq &\frac{(1-\tilde{\varepsilon}_{3}^{2})(1-\gamma^{2})^{2}}{4}q_{+}q_{-}\left\{\left(n\|\boldsymbol{\mu}\|^{2}\right)^{2}+R^{2}\right\}+\gamma^{2}R^{2}\\ \geq &\frac{(1-\tilde{\varepsilon}_{3}^{2})(1-\gamma^{2})^{2}}{8}q_{+}q_{-}\left(n\|\boldsymbol{\mu}\|^{2}+R\right)^{2}+\gamma^{2}R^{2}, \end{split}$$

where the last inequality is due to $a^2 + b^2 \ge \frac{(a+b)^2}{2}$.

Now we are ready to establish sufficient conditions of $\hat{\boldsymbol{w}} = \tilde{X}(\tilde{X}\tilde{X}^{\top})^{-1}\boldsymbol{y}$.

Lemma C.8. Suppose event $\bigcap_{i=1,2,3} \tilde{E}_i$ holds with $\tilde{\varepsilon} \leq \frac{q_{\gamma}}{2}$ and further assume one of the following conditions holds:

(i)
$$q_+q_-=0$$
, $n\|\mu\|^2<\frac{\gamma}{1+(1+\gamma)\tilde{\varepsilon}_3}R$, and $\sqrt{n}\tilde{\varepsilon}\leq\frac{\gamma^3}{4}$,

(ii)
$$q_+q_- \neq 0$$
, $n\|\boldsymbol{\mu}\|^2 \leq \lambda R$, and $\sqrt{n}\tilde{\varepsilon} \leq \frac{q_\gamma^2}{5(1+\lambda)^2}$ for some $\lambda \leq \frac{q_\gamma}{2\gamma(1-\gamma)}$,

(iii)
$$q_{\gamma} > \gamma$$
, $n\|\boldsymbol{\mu}\|^2 \geq \lambda R$, and $\sqrt{n}\tilde{\varepsilon} \leq \tilde{C}(\gamma, q_+, \tilde{\varepsilon}_3, \lambda) \frac{R}{n\|\boldsymbol{\mu}\|^2}$ for some $\lambda > 0$, where

$$\tilde{C}(\gamma, q_+, \tilde{\varepsilon}_3, \lambda) := \frac{(1 - \tilde{\varepsilon}_3)q_{\gamma}^2(q_{\gamma} - \gamma)}{5(1 + \lambda^{-1})^2}.$$

Then, we have

$$\hat{\boldsymbol{w}} = \tilde{X}(\tilde{X}\tilde{X}^{\top})^{-1}y.$$

Proof. By Lemma C.1, it suffices to show $\mathbf{y}^{\top}(\tilde{X}\tilde{X}^{\top})^{-1}y_i\mathbf{e}_i > 0$ for all i.

By Lemma C.5, we have

$$\mathbf{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_{i} \mathbf{e}_{i}
= (B^{-1}\mathbf{y})^{\top} B (\tilde{X}\tilde{X}^{\top})^{-1} B B^{-1} y_{i} \mathbf{e}_{i}
\geq (B^{-1}\mathbf{y})^{\top} A^{-1} B^{-1} y_{i} \mathbf{e}_{i} - \|B^{-1}\mathbf{y}\| B_{ii}^{-1} 2\tilde{\epsilon} q_{\gamma}^{-1} R^{-1}
\geq B_{ii}^{-1} (B^{-1}\mathbf{y})^{\top} A^{-1} y_{i} \mathbf{e}_{i} - 2B_{ii}^{-1} q_{\gamma}^{-1} \tilde{\epsilon} R^{-1} \sqrt{\frac{n_{+}}{q_{+} + \gamma^{2} q_{-}}} + \frac{n_{-}}{q_{-} + \gamma^{2} q_{+}}
\geq B_{ii}^{-1} \left\{ (B^{-1}\mathbf{y})^{\top} A^{-1} y_{i} \mathbf{e}_{i} - 2q_{\gamma}^{-3/2} \tilde{\epsilon} R^{-1} \sqrt{n} \right\},$$
(46)

where the last inequality follows from the definition of q_{γ} and $n_{+} + n_{-} = n$. The rest of the argument is split in cases.

Case $q_{+} = 1, q_{-} = 0$: By (33), we have

$$(B_{+}^{-1}\boldsymbol{y})^{\top}A^{-1}B_{+}^{-1}y_{i}\boldsymbol{e}_{i}$$

$$=\begin{cases}R^{-1}\left(1-\frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2}+R}\frac{n_{+}+\gamma^{-1}n_{-}}{n}\right), & \text{if } i \in I_{+},\\ \gamma^{-1}R^{-1}\left(\gamma^{-1}-\frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2}+R}\frac{n_{+}+\gamma^{-1}n_{-}}{n}\right), & \text{if } i \in I_{-}.\end{cases}$$

$$(47)$$

Therefore, by the definition of \tilde{E}_3 , (46) and (47), we have

$$\mathbf{y}^{\top} (\tilde{X} \tilde{X}^{\top})^{-1} y_{i} \mathbf{e}_{i} \\
\geq \begin{cases}
R^{-1} \left(1 - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \gamma^{-1})(1 + \tilde{\epsilon}_{3})}{2} - 2\gamma^{-3} \sqrt{n} \tilde{\epsilon} \right), & \text{if } i \in I_{+}, \\
\gamma^{-1} R^{-1} \left(\gamma^{-1} - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \gamma^{-1})(1 + \tilde{\epsilon}_{3})}{2} - 2\gamma^{-3} \sqrt{n} \tilde{\epsilon} \right), & \text{if } i \in I_{-}.
\end{cases} (48)$$

Case $q_+ = 0, q_- = 1$: By (34), we have

$$(B_{-}^{-1}\boldsymbol{y})^{\top} A^{-1} B_{-}^{-1} y_{i} \boldsymbol{e}_{i}$$

$$= \begin{cases} \gamma^{-1} R^{-1} \left(\gamma^{-1} - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1} n_{+} + n_{-}}{n} \right), & \text{if } i \in I_{+} \\ R^{-1} \left(1 - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{\gamma^{-1} n_{+} + n_{-}}{n} \right), & \text{if } i \in I_{-}. \end{cases}$$

$$(49)$$

Therefore, by the definition of \tilde{E}_3 , (46) and (49), we have

$$\mathbf{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_{i} \mathbf{e}_{i}$$

$$\geq \begin{cases} \gamma^{-1} R^{-1} \left(\gamma^{-1} - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \gamma^{-1})(1 + \tilde{\varepsilon}_{3})}{2} - 2\gamma^{-3} \sqrt{n} \tilde{\varepsilon} \right), & \text{if } i \in I_{+}, \\ R^{-1} \left(1 - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \gamma^{-1})(1 + \tilde{\varepsilon}_{3})}{2} - 2\gamma^{-3} \sqrt{n} \tilde{\varepsilon} \right), & \text{if } i \in I_{-}. \end{cases}$$
(50)

Therefore, for both the cases above, it suffices to have

$$\frac{n\|\boldsymbol{\mu}\|^2}{n\|\boldsymbol{\mu}\|^2+R}\frac{(1+\gamma^{-1})(1+\tilde{\varepsilon}_3)}{2}<\frac{1}{2},\quad \text{and}\quad 2\gamma^{-3}\sqrt{n}\tilde{\varepsilon}\leq\frac{1}{2},$$

which follows from the condition (i).

Case $q_{+}q_{-} \neq 0$: By (41) and (46), we have for $i \in I_{+}$

$$\mathbf{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_{i} \mathbf{e}_{i}$$

$$\geq B_{ii}^{-1} \Big[\{ d(q_{+} + \gamma^{2} q_{-})(q_{-} + \gamma^{2} q_{+}) \}^{-1} \\
\times \Big\{ \sqrt{q_{+} + \gamma^{2} q_{-}} \{ (q_{-} + \gamma^{2} q_{+} - \gamma) n_{-} \|\boldsymbol{\mu}\|^{2} + (q_{-} + \gamma^{2} q_{+}) R \Big\} \\
- 2q_{\gamma}^{-3/2} R^{-1} \sqrt{n} \tilde{\varepsilon} \Big],$$
(51)

and for $i \in I_-$

$$\mathbf{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_{i} \mathbf{e}_{i}$$

$$\geq B_{ii}^{-1} \Big[\{ d(q_{+} + \gamma^{2} q_{-}) (q_{-} + \gamma^{2} q_{+}) \}^{-1}$$

$$\times \Big\{ \sqrt{q_{-} + \gamma^{2} q_{+}} \Big\{ (q_{+} + \gamma^{2} q_{-} - \gamma) n_{+} \|\boldsymbol{\mu}\|^{2} + (q_{+} + \gamma^{2} q_{-}) R \Big\}$$

$$- 2q_{\gamma}^{-3/2} R^{-1} \sqrt{n} \tilde{\varepsilon} \Big].$$
(52)

Under regime (ii), (43) implies

$$d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \le \frac{5q_{\gamma}}{4}(1+\lambda)^{2}R^{2}.$$
 (53)

By $q_{\gamma} > \gamma^2$, we have

$$(q_{-} + \gamma^{2} q_{+} - \gamma) n_{-} \|\boldsymbol{\mu}\|^{2} + (q_{-} + \gamma^{2} q_{+}) R > q_{\gamma} R - \gamma (1 - \gamma) n_{-} \|\boldsymbol{\mu}\|^{2}$$

$$\geq q_{\gamma} R - \gamma (1 - \gamma) n \|\boldsymbol{\mu}\|^{2}$$

$$\geq \frac{q_{\gamma}}{2} R,$$
(54)

where the last inequality is due to the assumption on λ .

1348 Similarly, we have

$$(q_{+} + \gamma^{2}q_{-} - \gamma)n_{+}\|\boldsymbol{\mu}\|^{2} + (q_{+} + \gamma^{2}q_{-})R > \frac{q_{\gamma}}{2}R.$$
 (55)

Therefore, for $i \in I_+$, (51), (53), and (54) imply

$$\mathbf{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_i \mathbf{e}_i > B_{ii}^{-1} R^{-1} \left[\frac{2q_{\gamma}^{\frac{1}{2}}}{5(1+\lambda)^2} - 2q_{\gamma}^{-\frac{3}{2}} \sqrt{n}\tilde{\varepsilon} \right] \ge 0, \tag{56}$$

where the last inequality is due to the assumption on $\sqrt{n}\tilde{\varepsilon}$.

The same conclusion analogously holds for $i \in I_{-}$ as well by (52), (53), and (54).

Under regime (iii), by (43), we have that

$$d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \le \frac{5q_{\gamma}}{4}(1 + \lambda^{-1})^{2} \left(n\|\boldsymbol{\mu}\|^{2}\right)^{2}.$$
 (57)

With (51) and (57), we have for $i \in I_+$

$$\boldsymbol{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y_i \boldsymbol{e}_i$$

$$>B_{ii}^{-1} \left[\frac{q_{\gamma}^{\frac{1}{2}} \{ (q_{\gamma} - \gamma)(1 - \tilde{\varepsilon}_{3}) \frac{n}{2} \|\boldsymbol{\mu}\|^{2} \}}{\frac{5q_{\gamma}}{4} (1 + \lambda^{-1})^{2} (n \|\boldsymbol{\mu}\|^{2})^{2}} - 2q_{\gamma}^{-\frac{3}{2}} \sqrt{n} \tilde{\varepsilon} R^{-1} \right]$$

$$= (B_{ii}n \|\boldsymbol{\mu}\|^{2})^{-1} q_{\gamma}^{-\frac{3}{2}} \left[\frac{q_{\gamma}^{2} (q_{\gamma} - \gamma)(1 - \tilde{\varepsilon}_{3})}{5(1 + \lambda^{-1})^{2}} - 2\sqrt{n} \tilde{\varepsilon} \frac{n \|\boldsymbol{\mu}\|^{2}}{R} \right] \geq 0.$$
(58)

Similarly, the same lower bound holds for $i \in I_{-}$ from (52).

Remark. By similar argument we can show under (iii), an upper bound of the similar form as (58) for $i \in I_+$ whose dominant term is $(q_+ + \gamma^2 q_+ - \gamma) n \|\boldsymbol{\mu}\|^2$. Similarly, for $i \in I_-$, we have an upper bound whose dominant term is $(q_+ + \gamma^2 q_- - \gamma) n \|\boldsymbol{\mu}\|^2$. If we have $q_\gamma < \gamma$, exactly one of $q_- + \gamma^2 q_+ - \gamma$ and $q_+ + \gamma^2 q_- - \gamma$ becomes negative and the other stays positive. Suppose $q_- + \gamma^2 q_+ - \gamma < 0$, then we have $\boldsymbol{y}^\top (\tilde{X} \tilde{X}^\top)^{-1} y_i \boldsymbol{e}_i < 0$ for all $i \in I_+$ when $n \|\boldsymbol{\mu}\|^2$ is sufficiently larger than R, which means the equivalence of $\hat{\boldsymbol{w}} = \tilde{X}^\top (\tilde{X} \tilde{X}^\top)^{-1} \boldsymbol{y}$ provably fails in such case. The same applies with the case $q_+ + \gamma^2 q_- - \gamma < 0$.

With $\hat{\boldsymbol{w}} = \tilde{X}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} y$, by defining

$$\mathbf{w}_{+} = \frac{1}{\sqrt{m}} X^{\top} B_{+} (\tilde{X} \tilde{X}^{\top})^{-1} \mathbf{y}$$

$$= \frac{1}{\sqrt{m}} X^{\top} B_{+} (q_{+} B_{+} X X^{\top} B_{+} + q_{-} B_{-} X X^{\top} B_{-})^{-1} \mathbf{y},$$

$$\mathbf{w}_{-} = -\frac{1}{\sqrt{m}} X^{\top} B_{-} (\tilde{X} \tilde{X}^{\top})^{-1} \mathbf{y}$$

$$= -\frac{1}{\sqrt{m}} X^{\top} B_{-} (q_{+} B_{+} X X^{\top} B_{+} + q_{-} B_{-} X X^{\top} B_{-})^{-1} \mathbf{y},$$

$$(59)$$

we have $\hat{\boldsymbol{w}}_j = \boldsymbol{w}_+$ if $j \in J_+$ and $\hat{\boldsymbol{w}}_j = \boldsymbol{w}_-$ if $j \in J_-$.

By Lemma B.2, the network $f(x; \hat{W})$ has the linear decision boundary defined by

$$\bar{w} = \frac{|J_{+}|}{\sqrt{m}} w_{+} - \frac{|J_{-}|}{\sqrt{m}} w_{-}. \tag{60}$$

By (59) and (60), we have

$$\bar{\boldsymbol{w}} = \boldsymbol{X}^{\top} \left(q_{+} \boldsymbol{B}_{+} + q_{-} \boldsymbol{B}_{-} \right) (\tilde{\boldsymbol{X}} \tilde{\boldsymbol{X}}^{\top})^{-1} \boldsymbol{y}.$$

This implies

$$\|\bar{\boldsymbol{w}}\|^{2} = \boldsymbol{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} (q_{+}B_{+} + q_{-}B_{-}) XX^{\top} (q_{+}B_{+} + q_{-}B_{-}) (\tilde{X}\tilde{X}^{\top})^{-1} \boldsymbol{y},$$
(61)

$$\langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle = \boldsymbol{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} (q_{+}B_{+} + q_{-}B_{-}) X\boldsymbol{\mu}$$

$$= \boldsymbol{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} (q_{+}B_{+} + q_{-}B_{-}) (\|\boldsymbol{\mu}\|^{2} \boldsymbol{y} + Z\boldsymbol{\mu}).$$
(62)

We need to obtain bounds of $\|\bar{w}\|^2$ and $\langle \bar{w}, \mu \rangle$, respectively, in order to later establish a classification error bound of \bar{w} .

To obtain bounds of $\|\bar{\boldsymbol{w}}\|^2$, the following inequality is useful.

Lemma C.9. Suppose that the conclusions of Lemma C.2, C.5, and C.8 hold. Then, we have

$$\left| \|\bar{\boldsymbol{w}}\|^2 - (\|\boldsymbol{\mu}\|^2 K_1 + RK_2) \right| \le \tilde{\varepsilon} RK_2,$$

and

$$\left| \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle - \|\boldsymbol{\mu}\|^2 \sqrt{K_1} \right| \leq \frac{q_{\gamma}^{-\frac{1}{2}}}{3} \tilde{\varepsilon}_2 \| (B^{-1}y)^{\top} A^{-1} \| R + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_2,$$

where

$$K_{1} := \left\{ \boldsymbol{y}^{\top} (\tilde{X}\tilde{X}^{\top})^{-1} (q_{+}B_{+} + q_{-}B_{-}) \boldsymbol{y} \right\}^{2}$$

$$= \left\{ (B^{-1}\boldsymbol{y})^{\top} B (\tilde{X}\tilde{X}^{\top})^{-1} B \cdot B^{-1} (q_{+}B_{+} + q_{-}B_{-}) \boldsymbol{y} \right\}^{2} \text{ and}$$

$$K_{2} := \left\| (q_{+}B_{+} + q_{-}B_{-}) (\tilde{X}\tilde{X}^{\top})^{-1} \boldsymbol{y} \right\|^{2}$$
(63)

$$\mathbf{x}_{2} := \| (q_{+}B_{+} + q_{-}B_{-})(XX^{-}) \cdot \mathbf{y} \|$$

$$= \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot B(\tilde{X}\tilde{X}^{\top})^{-1}B \cdot B^{-1}\mathbf{y} \|^{2}.$$

$$(64)$$

Proof. Letting $\boldsymbol{v} = (q_+ B_+ + q_- B_-) (\tilde{X} \tilde{X}^\top)^{-1} \boldsymbol{y}$, we have $\|\bar{\boldsymbol{w}}\|^2 = \boldsymbol{v}^\top X X^\top \boldsymbol{v}$.

Then, by (61) and the conclusion of Lemma C.2, we have

$$\left| \|\bar{\boldsymbol{w}}\|^2 - \left(\|\boldsymbol{\mu}\|^2 \langle \boldsymbol{v}, \boldsymbol{y} \rangle^2 + R \|\boldsymbol{v}\|^2 \right) \right| \leq \tilde{\varepsilon} R \|\boldsymbol{v}\|^2.$$

The first desired inequality now follows by noting $\langle \boldsymbol{v}, \boldsymbol{y} \rangle^2 = K_1$ and $\|\boldsymbol{v}\|^2 = K_2$.

To show the second desired inequality, we note that (62) implies

$$\langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle = \|\boldsymbol{\mu}\|^2 \sqrt{K_1} + (B^{-1}\boldsymbol{y})^{\top} A^{-1} B^{-1} (q_+ B_+ + q_- B_-) Z \boldsymbol{\mu} + (B^{-1}\boldsymbol{y})^{\top} \left[B(\tilde{X}\tilde{X}^{\top})^{-1} - A^{-1} \right] B^{-1} (q_+ B_+ + q_- B_-) Z \boldsymbol{\mu}.$$
(65)

By Lemma C.5 and (65), we have

$$\begin{split} \left| \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle - \| \boldsymbol{\mu} \|^2 \sqrt{K_1} \right| &\leq \| (B^{-1} \boldsymbol{y})^\top A^{-1} \| \| B^{-1} \| \| q_+ B_+ + q_- B_- \| \| Z \boldsymbol{\mu} \| \\ &+ \| B^{-1} \| \| \boldsymbol{y} \| \frac{2q_\gamma^{-1} \tilde{\varepsilon}}{R} \| B^{-1} \| \| q_+ B_+ + q_- B_- \| \| \| Z \boldsymbol{\mu} \| \\ &\leq \frac{q_\gamma^{-\frac{1}{2}}}{3} \tilde{\varepsilon}_2 \| (B^{-1} \boldsymbol{y})^\top A^{-1} \| R + \frac{2q_\gamma^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_2, \end{split}$$

where the second inequality follows from the definition of $\tilde{\varepsilon}_2$, $||B^{-1}|| \leq q_{\gamma}^{-\frac{1}{2}}$, and $||q_+B_+| + q_-B_-|| \leq 1$.

Lemma C.10. Suppose that all the assumptions of Lemma C.8 hold with $\tilde{\varepsilon}_3 \leq \frac{1}{2}$ in (ii) and (iii). Then there exists constant \tilde{c}_1 which depends only on γ and q_+ such that

$$\|\bar{\boldsymbol{w}}\|^2 \le \tilde{c}_1 \frac{n}{n\|\boldsymbol{\mu}\|^2 + R}.$$

Furthermore, there exist constants \hat{c}_1 and N which only depends on γ , q_+ (and λ under (ii) and (iii)) such that $n \geq N$ implies

$$\|\bar{\boldsymbol{w}}\|^2 \ge \hat{c}_1 \frac{n}{n\|\boldsymbol{\mu}\|^2 + R}.$$

Proof. By Lemma C.9, the proof is reduced to bounding K_i , i = 1, 2.

By Lemma C.5, we have

$$\begin{vmatrix}
(B^{-1}\boldsymbol{y})^{\top}B(\tilde{X}\tilde{X}^{\top})^{-1}B \cdot B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y} - (B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}
\end{vmatrix}$$

$$\begin{vmatrix}
(B^{-1}\boldsymbol{y})^{\top}B(\tilde{X}\tilde{X}^{\top})^{-1}B \cdot B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y} - (B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}
\end{vmatrix}$$

$$\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \|B^{-1}\boldsymbol{y}\|\|B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}\|$$

$$\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \|\frac{1}{\sqrt{q_{+} + \gamma^{2}q_{-}}}\mathbf{1}_{+} - \frac{1}{\sqrt{q_{-} + \gamma^{2}q_{+}}}\mathbf{1}_{-} \| \cdot \|\frac{q_{+} + \gamma q_{-}}{\sqrt{q_{+} + \gamma^{2}q_{-}}}\mathbf{1}_{+} - \frac{q_{-} + \gamma q_{+}}{\sqrt{q_{-} + \gamma^{2}q_{+}}}\mathbf{1}_{-} \|$$

$$\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \sqrt{\frac{1}{q_{+} + \gamma^{2}q_{-}}}n_{+} + \frac{1}{q_{-} + \gamma^{2}q_{+}}n_{-} \cdot \sqrt{\frac{(q_{+} + \gamma q_{-})^{2}}{q_{+} + \gamma^{2}q_{-}}}n_{+} + \frac{(q_{-} + \gamma q_{+})^{2}}{q_{-} + \gamma^{2}q_{+}}n_{-} \\
\end{cases}$$

$$\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \sqrt{\frac{n}{q_{\gamma}}} \cdot \sqrt{\frac{n}{q_{\gamma}}}$$

$$\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \sqrt{\frac{n}{q_{\gamma}}} \cdot \sqrt{\frac{n}{q_{\gamma}}}$$

$$\leq \frac{2q_{\gamma}^{-2}n\tilde{\varepsilon}}{R}$$

Similarly, we have

$$\left| \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot B(\tilde{X}\tilde{X}^{\top})^{-1}B \cdot B^{-1}\boldsymbol{y} \| - \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1}\boldsymbol{y} \| \right| \\
\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \| \cdot \| B^{-1}\boldsymbol{y} \| \\
\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \max \left\{ \frac{q_{+} + \gamma q_{-}}{\sqrt{q_{+} + \gamma^{2}q_{-}}}, \frac{q_{-} + \gamma q_{+}}{\sqrt{q_{-} + \gamma^{2}q_{+}}} \right\} \cdot \sqrt{\frac{n}{q_{\gamma}}} \\
\leq \frac{2q_{\gamma}^{-1}\tilde{\varepsilon}}{R} \frac{1}{\sqrt{q_{\gamma}}} \cdot \sqrt{\frac{n}{q_{\gamma}}} \\
\leq \frac{2q_{\gamma}^{-2}\sqrt{n}\tilde{\varepsilon}}{R}.$$
(67)

We split in cases.

Regime (i): We first note that $n\|\mu\|^2 < \frac{\gamma}{1+(1+\gamma)\tilde{\varepsilon}_3}R \leq \gamma R$ implies

$$R^{-1} \le (1+\gamma)(n\|\boldsymbol{\mu}\|^2 + R)^{-1}. (68)$$

We first consider the case when $q_+ = 1, q_- = 0$. By (33), we have

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}$$

$$=(B_{+}^{-1}\boldsymbol{y})^{\top}A^{-1}(B_{+}^{-1}\boldsymbol{y})$$

$$=R^{-1}\left\{\left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n}\right)n_{+}\right.$$

$$\left. + \gamma^{-1}\left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1}n_{-}}{n}\right)n_{-}\right\}$$

$$\leq \frac{n_{+} + \gamma^{-2}n_{-}}{R}$$

$$\leq (1 + \gamma)\gamma^{-2}\frac{n}{n\|\boldsymbol{\mu}\|^{2} + R},$$
(69)

where the last line is due to (68).

We also have

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}$$

$$\geq R^{-1}\left\{\left(1 - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \tilde{\varepsilon}_{3})(1 + \gamma^{-1})}{2}\right)n_{+} + \gamma^{-1}\left(\gamma^{-1} - \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \frac{(1 + \tilde{\varepsilon}_{3})(1 + \gamma^{-1})}{2}\right)n_{-}\right\}$$

$$\geq R^{-1}\left\{\left(1 - \frac{1}{2}\right)n_{+} + \gamma^{-1}\left(\gamma^{-1} - \frac{1}{2}\right)n_{-}\right\}$$

$$\geq \{2(1 + \gamma)\}^{-1}\frac{n}{n\|\boldsymbol{\mu}\|^{2} + R},$$

$$(70)$$

where the first inequality is due to \tilde{E}_3 , the second inequality is due to the assumption $n\|\mu\|^2 < \frac{\gamma}{1+(1+\gamma)\tilde{\varepsilon}_3}R$, and the last inequality is due to $\gamma \in (0,1)$.

Analogously, we have the same results for $q_+ = 0, q_- = 1$.

With these, (66) with (68), and noting that $q_{\gamma} = \gamma^2$ when $q_{+}q_{-} = 0$, we have under the regime (i)

$$\left\{ \frac{3}{8(1+\gamma)} \right\}^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} \leq \left(\frac{1}{2(1+\gamma)} - 2(1+\gamma)\gamma^{-4}\tilde{\varepsilon} \right)^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} \\
\leq K_{1} \leq (1+\gamma)^{2} \left(\gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon} \right)^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} \tag{71}$$

for sufficiently large n.

Similarly, (33) implies

$$\begin{aligned} & \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1} \boldsymbol{y} \| | \\ &= \| A^{-1} \cdot B_{+}^{-1} \boldsymbol{y} \| | \\ &= R^{-1} \left\{ \left(1 - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1} n_{-}}{n} \right)^{2} n_{+} \right. \\ & \left. + \gamma^{-2} \left(\gamma^{-1} - \frac{n \|\boldsymbol{\mu}\|^{2}}{n \|\boldsymbol{\mu}\|^{2} + R} \frac{n_{+} + \gamma^{-1} n_{-}}{n} \right)^{2} n_{-} \right\}^{\frac{1}{2}}, \end{aligned}$$
(72)

and so under the regime (i), we have

$$\frac{1}{2(1+\gamma)} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^2 + R} \le \|(q_+ B_+ + q_- B_-)B^{-1} \cdot A^{-1} \cdot B^{-1} \boldsymbol{y}\| \le (1+\gamma)\gamma^{-2} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^2 + R}.$$
(73)

With (67), we have

$$\left\{ \frac{3}{8(1+\gamma)} \right\}^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} \leq \left(\frac{1}{2(1+\gamma)} - 2(1+\gamma)\gamma^{-4}\tilde{\varepsilon} \right)^{2} \frac{n}{R^{2}} \\
\leq K_{2} \leq (1+\gamma)^{2} \left(\gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon} \right)^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}}$$
(74)

for sufficiently large n.

Thus, by Lemma C.9, (71), and (74), we have

$$\left[\left\{ \frac{3}{8(1+\gamma)} \right\}^2 - (1+\gamma)^2 \left(\gamma^{-2} + 2\gamma^{-4} \tilde{\varepsilon} \right)^2 \tilde{\varepsilon} \right] \frac{n}{n \|\boldsymbol{\mu}\|^2 + R} \\
\leq \|\boldsymbol{\bar{w}}\|^2 \leq (1+\tilde{\varepsilon})(1+\gamma)^2 \left(\gamma^{-2} + 2\gamma^{-4} \tilde{\varepsilon} \right)^2 \frac{n}{n \|\boldsymbol{\mu}\|^2 + R},$$

with the left hand side being positive for sufficiently large n.

1569 **Case** $q_+q_- \neq 0$: By (35), we have

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}$$

$$= \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}$$

$$\times \left[(q_{+} + \gamma q_{-})\{(q_{-} + \gamma^{2}q_{+} - \gamma)n_{-}\|\boldsymbol{\mu}\|^{2} + (q_{-} + \gamma^{2}q_{+})R\}n_{+}\right]$$

$$+ (q_{-} + \gamma q_{+})\{(q_{+} + \gamma^{2}q_{-} - \gamma)n_{+}\|\boldsymbol{\mu}\|^{2} + (q_{+} + \gamma^{2}q_{-})R\}n_{-}\right].$$

$$(75)$$

Thus, we have with Lemma C.7

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y} \leq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}n(n\|\boldsymbol{\mu}\|^{2} + R)$$

$$\leq \frac{8}{(1 - \tilde{\varepsilon}_{2}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}} \frac{n}{n\|\boldsymbol{\mu}\|^{2} + R}.$$
(76)

Similarly, (35) implies

$$\begin{aligned} & \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1} \boldsymbol{y} \| | \\ &= \{ d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \}^{-1} \\ &\times \left[(q_{+} + \gamma q_{-})^{2} \left\{ (q_{-} + \gamma^{2}q_{+} - \gamma)n_{-} \| \boldsymbol{\mu} \|^{2} + (q_{-} + \gamma^{2}q_{+})R \right\}^{2} n_{+} \right. \end{aligned}$$

$$\left. + (q_{-} + \gamma q_{+})^{2} \left\{ (q_{+} + \gamma^{2}q_{-} - \gamma)n_{+} \| \boldsymbol{\mu} \|^{2} + (q_{+} + \gamma^{2}q_{-})R \right\}^{2} n_{-} \right]^{\frac{1}{2}}$$

$$(77)$$

Thus, we have with Lemma C.7

$$\begin{aligned} \left\| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1} \boldsymbol{y} \right\| & \leq \left\{ d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \right\}^{-1} \sqrt{n} (n\|\boldsymbol{\mu}\|^{2} + R^{2}) \\ & \leq \frac{8}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^{2} + R}. \end{aligned}$$

$$(78)$$

We now consider regime (ii) and (iii) separately.

Regime (ii): By (66) and (76), we have

$$K_1 \le \left\{ \frac{8}{(1 - \tilde{\varepsilon}_3^2)(1 - \gamma^2)^2 q_+ q_-} + 2(1 + \lambda) q_{\gamma}^{-2} \tilde{\varepsilon} \right\}^2 \frac{n^2}{(n||\boldsymbol{\mu}||^2 + R)^2}.$$
 (79)

Similarly, (67) and (78) imply

$$K_2 \le \left\{ \frac{8}{(1 - \tilde{\varepsilon}_3^2)(1 - \gamma^2)^2 q_+ q_-} + 2(1 + \lambda) q_{\gamma}^{-2} \tilde{\varepsilon} \right\}^2 \frac{n}{(n \|\boldsymbol{\mu}\|^2 + R)^2}.$$
 (80)

(54) and (75) imply

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}\frac{q_{\gamma}}{2}Rn$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}\frac{q_{\gamma}^{2}}{2(1 + \lambda)}n(n\|\boldsymbol{\mu}\|^{2} + R).$$
(81)

By Lemma C.7, (66), (81), and the assumption on $\tilde{\varepsilon}$, we have for sufficiently large n

$$K_1 \ge \left\{ \frac{2q_{\gamma}}{5(1+\lambda)} - 2(1+\lambda)q_{\gamma}^{-2}\tilde{\varepsilon} \right\}^2 \frac{n^2}{(n\|\boldsymbol{\mu}\|^2 + R)^2} \ge \left\{ \frac{q_{\gamma}}{5(1+\lambda)} \right\}^2 \frac{n^2}{(n\|\boldsymbol{\mu}\|^2 + R)^2}. \tag{82}$$

Similarly, (54) and (77) imply

$$\|(q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1}\boldsymbol{y}\|\|$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1} \left[q_{\gamma}^{2} \left\{\frac{q_{\gamma}}{2}R\right\}^{2} n\right]^{\frac{1}{2}}$$

$$= \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1} \frac{q_{\gamma}^{2}}{2} \sqrt{n}R$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1} \frac{q_{\gamma}^{2}}{2(1 + \lambda)} \sqrt{n}(n\|\boldsymbol{\mu}\|^{2} + R).$$
(83)

By Lemma C.7, (67), (83), and the assumption on $\tilde{\varepsilon}$, we have for sufficiently large n

$$K_2 \ge \left\{ \frac{2q_{\gamma}}{5(1+\lambda)} - 2(1+\lambda)q_{\gamma}^{-2}\tilde{\varepsilon} \right\}^2 \frac{n}{(n\|\boldsymbol{\mu}\|^2 + R)^2} \ge \left\{ \frac{q_{\gamma}}{5(1+\lambda)} \right\}^2 \frac{n^2}{(n\|\boldsymbol{\mu}\|^2 + R)^2}.$$
 (84)

By Lemma C.9, (79), (80), (82), and (84), we have

$$\left[\left\{ \frac{q_{\gamma}}{5(1+\lambda)} \right\}^{2} - \left\{ \frac{8}{(1-\tilde{\varepsilon}_{3}^{2})(1-\gamma^{2})^{2}q_{+}q_{-}} + 2q_{\gamma}^{-2}\tilde{\varepsilon} \right\}^{2} \tilde{\varepsilon} \right] \frac{n}{n\|\boldsymbol{\mu}\|^{2} + R} \\
\leq \|\bar{\boldsymbol{w}}\|^{2} \leq (1+\tilde{\varepsilon}) \left\{ \frac{8}{(1-\tilde{\varepsilon}_{3}^{2})(1-\gamma^{2})^{2}q_{+}q_{-}} + 2q_{\gamma}^{-2}\tilde{\varepsilon} \right\}^{2} \frac{n}{n\|\boldsymbol{\mu}\|^{2} + R}, \tag{85}$$

where the left hand side is positive for sufficiently large n due to the assumption on $\tilde{\varepsilon}$.

Regime (iii): By $n\|\mu\|^2 \ge \lambda R$, $\sqrt{n}\tilde{\varepsilon} \le \tilde{C} \frac{R}{n\|\mu\|^2} \le \frac{\tilde{C}}{\lambda}$, (66), and (76), we have

$$K_{1} \leq \left\{ \frac{8}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}} + 2q_{\gamma}^{-2} \left(1 + \frac{n\|\boldsymbol{\mu}\|^{2}}{R} \right) \tilde{\varepsilon} \right\}^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2} + R)^{2}}$$

$$\leq \left\{ \frac{8}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}} + 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1} + 1}{\sqrt{n}} \right\}^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2} + R)^{2}}.$$

$$(86)$$

Similarly, $n\|\boldsymbol{\mu}\|^2 \geq \lambda R$, $\sqrt{n}\tilde{\varepsilon} \leq \tilde{C}\frac{R}{n\|\boldsymbol{\mu}\|^2} \leq \frac{\tilde{C}}{\lambda}$, (67), and (78) imply

$$K_{2} \leq \left\{ \frac{8}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}} + \left(1 + \frac{n\|\boldsymbol{\mu}\|^{2}}{R}\right)\tilde{\varepsilon} \right\}^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2} + R)^{2}}$$

$$\leq \left\{ \frac{8}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}} + 2q_{\gamma}^{-2}\tilde{C}\frac{\lambda^{-1} + 1}{\sqrt{n}} \right\}^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2} + R)^{2}}.$$

$$(87)$$

By (42) and (75), we have

$$(B^{-1}\boldsymbol{y})^{\top}A^{-1}B^{-1}(q_{+}B_{+} + q_{-}B_{-})\boldsymbol{y}$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}q_{\gamma}(q_{\gamma} - \gamma)\frac{1 - \tilde{\varepsilon}_{3}}{2}n(n_{+} + n_{-})\|\boldsymbol{\mu}\|^{2}$$

$$\geq \{d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+})\}^{-1}\frac{(1 - \tilde{\varepsilon}_{3})q_{\gamma}(q_{\gamma} - \gamma)}{2}n^{2}\|\boldsymbol{\mu}\|^{2}.$$
(88)

By Lemma C.7, (66), and (88), we have for sufficiently large n

$$K_{1} \geq \left\{ \frac{2(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5} \frac{n^{2} \|\boldsymbol{\mu}\|^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} - 2q_{\gamma}^{-2} \left(1 + \frac{n\|\boldsymbol{\mu}\|^{2}}{R}\right) \tilde{\varepsilon} \frac{n}{n\|\boldsymbol{\mu}\|^{2}+R} \right\}^{2}$$

$$\geq \left\{ \frac{2(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5(1+\lambda^{-1})} - 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1}+1}{\sqrt{n}} \right\}^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}}$$

$$\geq \left\{ \frac{(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5(1+\lambda^{-1})} \right\}^{2} \frac{n^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}}.$$
(89)

Similarly, (42) and (77) imply

$$\begin{aligned} & \| (q_{+}B_{+} + q_{-}B_{-})B^{-1} \cdot A^{-1} \cdot B^{-1}\boldsymbol{y} \| | \\ & \geq \{ d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \}^{-1} \left[q_{\gamma}^{2}(q_{\gamma} - \gamma)^{2}n_{+}n_{-}(n_{+} + n_{-}) \|\boldsymbol{\mu}\|^{4} \right]^{\frac{1}{2}} \\ & \geq \{ d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \}^{-1} q_{\gamma}(q_{\gamma} - \gamma) \sqrt{\frac{(1 - \tilde{\varepsilon}_{3}^{2})}{4}n^{3}} \|\boldsymbol{\mu}\|^{2} \\ & \geq \{ d(q_{+} + \gamma^{2}q_{-})(q_{-} + \gamma^{2}q_{+}) \}^{-1} q_{\gamma}(q_{\gamma} - \gamma) \frac{(1 - \tilde{\varepsilon}_{3})}{2} n^{\frac{3}{2}} \|\boldsymbol{\mu}\|^{2} \end{aligned}$$

$$(90)$$

By Lemma C.7, (66), and (90), we have for sufficiently large n

$$K_{2} \geq \left\{ \frac{2(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5} \frac{n^{\frac{3}{2}} \|\boldsymbol{\mu}\|^{2}}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}} - 2q_{\gamma}^{-2} \left(1 + \frac{n\|\boldsymbol{\mu}\|^{2}}{R}\right) \tilde{\varepsilon} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^{2}+R} \right\}^{2}$$

$$\geq \left\{ \frac{2(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5(1+\lambda^{-1})} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^{2}+R} - 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1}+1}{\sqrt{n}} \frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^{2}+R} \right\}^{2}$$

$$= \left\{ \frac{2(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5(1+\lambda^{-1})} - 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1}+1}{\sqrt{n}} \right\}^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}}$$

$$\geq \left\{ \frac{(1-\tilde{\varepsilon}_{3})(q_{\gamma}-\gamma)}{5(1+\lambda^{-1})} \right\}^{2} \frac{n}{(n\|\boldsymbol{\mu}\|^{2}+R)^{2}}.$$
(91)

By Lemma C.9, (79), (80), (89), and (91), we have

$$\left[\left\{ \frac{(1 - \tilde{\varepsilon}_3)(q_{\gamma} - \gamma)}{5(1 + \lambda^{-1})} \right\}^2 - \left\{ \frac{8}{(1 - \tilde{\varepsilon}_3^2)(1 - \gamma^2)^2 q_+ q_-} + 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1} + 1}{\sqrt{n}} \right\}^2 \tilde{\varepsilon} \right] \frac{n}{n \|\boldsymbol{\mu}\|^2 + R} \\
\leq \|\bar{\boldsymbol{w}}\|^2 \leq (1 + \tilde{\varepsilon}) \left\{ \frac{8}{(1 - \tilde{\varepsilon}_3^2)(1 - \gamma^2)^2 q_+ q_-} + 2q_{\gamma}^{-2} \tilde{C} \frac{\lambda^{-1} + 1}{\sqrt{n}} \right\}^2 \frac{n}{n \|\boldsymbol{\mu}\|^2 + R}, \tag{92}$$

where the left hand side is positive for sufficiently large n.

We now turn to obtain bounds of $\langle \bar{w}, \mu \rangle$.

Lemma C.11. Suppose that all the assumptions of Lemma C.8 hold with $\tilde{\varepsilon}_3 \leq \frac{1}{2}$ in regime (ii) and (iii). Further suppose $\tilde{\varepsilon}_2 \leq \tilde{D} \frac{\sqrt{n}\|\boldsymbol{\mu}\|^2}{R}$ in (i) and (ii), where \tilde{D} is a constant which depends only on γ , q_+ (and λ under regime (ii)). Then, there exist constants \hat{c}_2 , \tilde{c}_2 , N which depend only on γ , q_+ (and λ under regime (ii) and (iii)) such that $n \geq N$ imply

$$\hat{c}_2 \frac{n \|\boldsymbol{\mu}\|^2}{n \|\boldsymbol{\mu}\|^2 + R} \le \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle \le \tilde{c}_2 \frac{n \|\boldsymbol{\mu}\|^2}{n \|\boldsymbol{\mu}\|^2 + R}.$$

Proof. By Lemma C.6 and C.7, we have

$$\|(B^{-1}\boldsymbol{y})^{\top}A^{-1}\| \leq \begin{cases} \gamma^{-1}\frac{\sqrt{n}}{R}, & \text{if } q_{+}q_{-} = 0, \\ \frac{8q_{\gamma}^{\frac{1}{2}}}{(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2}q_{+}q_{-}}\frac{\sqrt{n}}{n\|\boldsymbol{\mu}\|^{2} + R}, & \text{if } q_{+}q_{-} \neq 0. \end{cases}$$
(93)

The remaining argument is split in cases.

Regime (i):

We first note (71) and $n\|\mu\|^2 \le \gamma R$ imply

$$\frac{9}{64(1+\gamma)^3} \frac{n\|\boldsymbol{\mu}\|^2}{R} \le \|\boldsymbol{\mu}\|^2 \sqrt{K_1} \le (1+\gamma)^2 \left(\gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon}\right) \frac{n\|\boldsymbol{\mu}\|^2}{R}.$$
 (94)

Furthermore, Lemma C.9 and (93) imply

$$\left| \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle - \|\boldsymbol{\mu}\|^{2} \sqrt{K_{1}} \right| \leq \frac{\gamma^{-1} q_{\gamma}^{-\frac{1}{2}}}{3} \sqrt{n} \tilde{\varepsilon}_{2} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_{2}$$

$$= \frac{\gamma^{-1} q_{\gamma}^{-\frac{1}{2}} + 2q_{\gamma}^{-2} \tilde{\varepsilon}}{3} \sqrt{n} \tilde{\varepsilon}_{2}$$

$$= \frac{\gamma^{-2} + 2\gamma^{-4} \tilde{\varepsilon}}{3} \sqrt{n} \tilde{\varepsilon}_{2}$$

$$\leq \frac{\gamma^{-2} + 2\gamma^{-4} \tilde{\varepsilon}}{3} \tilde{D} \frac{n \|\boldsymbol{\mu}\|^{2}}{R},$$

$$(95)$$

where the last line is due to the assumption on $\tilde{\varepsilon}_2$.

By (94) and (95), it suffices to choose \tilde{D} so that

$$\frac{\gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon}}{3}\tilde{D} \le \frac{1}{8(1+\gamma)^3}.$$

Then, we have

$$\begin{split} &\frac{1}{64(1+\gamma)^3}\frac{n\|\pmb{\mu}\|^2}{n\|\pmb{\mu}\|^2+R} \leq \frac{1}{64(1+\gamma)^3}\frac{n\|\pmb{\mu}\|^2}{R} \leq \langle \bar{\pmb{w}}, \pmb{\mu} \rangle \\ &\leq \left\{ \gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon} + \frac{1}{8(1+\gamma)^3} \right\} \frac{n\|\pmb{\mu}\|^2}{R} \leq (1+\gamma) \left\{ \gamma^{-2} + 2\gamma^{-4}\tilde{\varepsilon} + \frac{1}{8(1+\gamma)^3} \right\} \frac{n\|\pmb{\mu}\|^2}{n\|\pmb{\mu}\|^2+R}. \end{split}$$

Regime (ii): We note that under regime (ii), we have $\|\mu\|^2 \sqrt{K_1} \approx \frac{n\|\mu\|^2}{n\|\mu\|^2 + R}$ by (79) and (82). On the other hand, Lemma C.9 and (93) imply

$$\left| \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle - \|\boldsymbol{\mu}\|^{2} \sqrt{K_{1}} \right| \leq \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \tilde{\varepsilon}_{2} \frac{\sqrt{n}R}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_{2}
= \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \frac{\tilde{\varepsilon}_{2}R}{\sqrt{n}\|\boldsymbol{\mu}\|^{2}} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_{2}
\leq \frac{8\tilde{D}}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \tilde{D} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R}
= \left\{ \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \right\} \tilde{D} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R}.$$
(96)

Therefore, choosing an appropriate \hat{D} suffices to establish the desired bounds.

Specifically, under regime (ii), we have by (82)

$$\|\boldsymbol{\mu}\|^2 \sqrt{K_1} \ge \frac{q_{\gamma}}{5(1+\lambda)} \frac{n\|\boldsymbol{\mu}\|^2}{n\|\boldsymbol{\mu}\|^2 + R}$$

and hence it suffices to choose D so that

$$\left\{ \frac{8}{3(1-\tilde{\varepsilon}_3^2)(1-\gamma^2)^2 q_+ q_-} + \frac{2q_{\gamma}^{-2}}{3}\tilde{\varepsilon} \right\} \tilde{D} \le \frac{q_{\gamma}}{10(1+\lambda)}.$$

Regime (iii): Similarly, we note that under regime (iii), we have $\|\mu\|^2 \sqrt{K_1} \approx \frac{n\|\mu\|^2}{n\|\mu\|^2 + R} \approx 1$ by (86) and (89). On the other hand, Lemma C.9 and (93) imply

$$\left| \langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle - \|\boldsymbol{\mu}\|^{2} \sqrt{K_{1}} \right| \\
\leq \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \tilde{\varepsilon}_{2} \frac{\sqrt{n}R}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_{2} \\
= \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \frac{\tilde{\varepsilon}_{2}R}{\sqrt{n}\|\boldsymbol{\mu}\|^{2}} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} \cdot \sqrt{n} \tilde{\varepsilon}_{2} \\
\leq \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \frac{\tilde{\varepsilon}R}{n\|\boldsymbol{\mu}\|^{2}} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon}^{2} \\
\leq \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \lambda^{-1} \tilde{\varepsilon} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon}^{2} (1 + \lambda^{-1})^{-1} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R} \\
= \left\{ \frac{8}{3(1 - \tilde{\varepsilon}_{3}^{2})(1 - \gamma^{2})^{2} q_{+} q_{-}} \lambda^{-1} + \frac{2q_{\gamma}^{-2}}{3} \tilde{\varepsilon} (1 + \lambda^{-1})^{-1} \right\} \tilde{\varepsilon} \frac{n\|\boldsymbol{\mu}\|^{2}}{n\|\boldsymbol{\mu}\|^{2} + R}. \tag{97}$$

By (86), (89), and (97), we have the desired bound for sufficiently large n.

Lemma C.12. Consider model (M). If $w \in \mathbb{R}^p$ is such that $\langle w, \mu \rangle > 0$, Then, we have

$$\mathbb{P}(\langle \boldsymbol{w}, y\boldsymbol{x}\rangle \leq 0) \leq \exp\left\{-c\left(\frac{\langle \boldsymbol{w}, \boldsymbol{\mu}\rangle}{\|\boldsymbol{w}\| \|\boldsymbol{z}\|_{\psi_2}}\right)^2\right\},\,$$

where c is a universal constant and $\|z\|_{\psi_2}$ is the sub-Gaussian norm of z.

If $\mathbf{z} = \Sigma^{\frac{1}{2}} \boldsymbol{\xi}$ with $\boldsymbol{\xi} \sim \mathcal{N}(0, I_p)$, then we further have

$$\kappa\left(\beta_{min}^{-\frac{1}{2}}\frac{\langle \boldsymbol{w},\boldsymbol{\mu}\rangle}{\|\boldsymbol{w}\|}\right) \leq \mathbb{P}(\langle \boldsymbol{w},y\boldsymbol{x}\rangle \leq 0) = \kappa\left(\frac{\langle \boldsymbol{w},\boldsymbol{\mu}\rangle}{\|\Sigma^{\frac{1}{2}}\boldsymbol{w}\|}\right) \leq \kappa\left(\beta_{max}^{-\frac{1}{2}}\frac{\langle \boldsymbol{w},\boldsymbol{\mu}\rangle}{\|\boldsymbol{w}\|}\right), \tag{98}$$

where $\kappa(t) = \mathbb{P}(|\xi_1| \geq t)$ and β_{min} is the smallest eigenvalue of Σ and $\beta_{max} = ||\Sigma||$.

Proof. Since $yx = \mu + yz$, we have

$$\begin{split} \mathbb{P}(\langle \boldsymbol{w}, y \boldsymbol{x} \rangle \leq 0) &= \mathbb{P}(y \langle \boldsymbol{w}, \boldsymbol{z} \rangle \leq -\langle \boldsymbol{w}, \boldsymbol{\mu} \rangle) \\ &= \frac{1}{2} \mathbb{P}(\langle \boldsymbol{w}, \boldsymbol{z} \rangle \leq -\langle \boldsymbol{w}, \boldsymbol{\mu} \rangle) + \frac{1}{2} \mathbb{P}(-\langle \boldsymbol{w}, \boldsymbol{z} \rangle \leq -\langle \boldsymbol{w}, \boldsymbol{\mu} \rangle) \\ &= \mathbb{P}(|\langle \boldsymbol{w}, \boldsymbol{z} \rangle| > \langle \boldsymbol{w}, \boldsymbol{\mu} \rangle). \end{split}$$

Then the desired conclusion follows from the basic property of sub-Gaussian distributions (see Proposition 2.5.2, Vershynin (2018)).

The second result follows since
$$\langle \boldsymbol{w}, \boldsymbol{z} \rangle = \langle \boldsymbol{w}, \Sigma^{\frac{1}{2}} \boldsymbol{\xi} \rangle = \langle \Sigma^{\frac{1}{2}} \boldsymbol{w}, \boldsymbol{\xi} \rangle \sim \mathcal{N}(0, \boldsymbol{w}^{\top} \Sigma \boldsymbol{w}) \text{ if } \boldsymbol{\xi} \sim \mathcal{N}(0, \Sigma).$$

Theorem C.13 (Detailed version of Theorem 5.1). Suppose event $\bigcap_{i=1,2,3} \tilde{E}_i$ holds with one of the following conditions:

(i)
$$q_+q_- = 0$$
, $n\|\mu\|^2 < \frac{\gamma}{1 + (1 + \gamma)\tilde{\epsilon}_3}R$, and $\sqrt{n}\tilde{\epsilon} \leq \frac{\gamma^3}{4}$,

(ii)
$$q_+q_- \neq 0$$
, $n\|\boldsymbol{\mu}\|^2 \leq \frac{q_{\gamma}}{2\gamma(1-\gamma)}R$, $\sqrt{n}\tilde{\varepsilon} \leq \frac{q_{\gamma}^2}{20}$, and $\tilde{\varepsilon}_3 \leq \frac{1}{2}$,

$$\textit{(iii)} \ \ q_{\gamma} > \gamma, \ n\|\boldsymbol{\mu}\|^2 \geq \frac{q_{\gamma}}{2\gamma(1-\gamma)}R, \ \tilde{\varepsilon}_3 \leq \frac{1}{2}, \\ \textit{and} \ \sqrt{n}\tilde{\varepsilon} \leq \frac{q_{\gamma}^2(q_{\gamma}-\gamma)}{40}\frac{R}{n\|\boldsymbol{\mu}\|^2}$$

Further suppose $\tilde{\varepsilon}_2 \leq \tilde{D} \frac{\sqrt{n} \|\mu\|^2}{R}$ in (i) and (ii), where \tilde{D} is a constant which depends only on γ , q_+ .

Then, there exist constants c and N, which depend on γ and q_+ , such that $n \geq N$ imply

$$\mathbb{P}_{(\boldsymbol{x},y)}(yf(\boldsymbol{x};\hat{W})<0) \leq \exp\left(-c\frac{n\|\boldsymbol{\mu}\|^4}{\|\boldsymbol{z}\|_{\psi_2}^2\{n\|\boldsymbol{\mu}\|^2+R\}}\right).$$

If $z = \sum_{1}^{1} \xi$ with $\xi \sim \mathcal{N}(0, I_p)$, then we have instead

$$\kappa \left\{ c^{-1} \beta_{min}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R} \right)^{\frac{1}{2}} \right\} \leq \mathbb{P}_{(\boldsymbol{x}, y)}(y f(\boldsymbol{x}; \hat{W}) < 0) \leq \kappa \left\{ c \beta_{max}^{-\frac{1}{2}} \left(\frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R} \right)^{\frac{1}{2}} \right\},$$

where $\kappa(t) = \mathbb{P}(|\xi_1| \geq t)$ and β_{min} is the smallest eigenvalue of Σ and $\beta_{max} = ||\Sigma||$.

Proof. Since we assume $\sqrt{n}\tilde{\varepsilon}\lesssim 1$ in all the regimes, we have $\tilde{\varepsilon}\leq \frac{q_{\gamma}}{2}$ for sufficiently large n.

By Lemma C.10 and C.11 with $\lambda = \frac{q_{\gamma}}{2\gamma(1-\gamma)}$, we have

$$\frac{\hat{c}_2^2}{\tilde{c}_1} \frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R} \le \left(\frac{\langle \bar{\boldsymbol{w}}, \boldsymbol{\mu} \rangle}{\|\bar{\boldsymbol{w}}\|}\right)^2 \le \frac{\hat{c}_2^2}{\hat{c}_1} \frac{n \|\boldsymbol{\mu}\|^4}{n \|\boldsymbol{\mu}\|^2 + R}.$$

Then, the desired conclusion holds from Lemma C.12.

D PROOF OF THEOREM 6.2 AND THEOREM 6.3

First we establish connection between events $E(\theta_1, \theta_2)$ and $\tilde{E}_1(\tilde{\epsilon}_1, R), \tilde{E}_2(\epsilon_2, R)$.

Lemma D.1. Suppose $\tilde{E}_1(\tilde{\varepsilon}_1, R) \cap \tilde{E}_2(\varepsilon_2, R)$ hold with $\tilde{\varepsilon}_1 \leq \frac{1}{2}$, then event $E(\theta_1, \theta_2)$ holds with

$$\theta_1 = rac{ ilde{arepsilon}_1}{2} \quad and \quad heta_2 = rac{ ilde{arepsilon}_2 \sqrt{R}}{2 \|oldsymbol{\mu}\|}.$$

Proof. By $\tilde{E}_1(\tilde{\varepsilon}_1,R)$, we have $|\|\boldsymbol{z}_i\|^2 - R| \leq \frac{\tilde{\varepsilon}_1}{3}R$ and $|\langle \boldsymbol{z}_i, \boldsymbol{z}_k \rangle| \leq \frac{\tilde{\varepsilon}_1}{3}R$ for any $i \neq k$. Then, we have

$$\left|\left\langle \frac{oldsymbol{z}_i}{\|oldsymbol{z}_i\|}, \frac{oldsymbol{z}_k}{\|oldsymbol{z}_k\|} \right
ight
angle
ight| \leq rac{ ilde{arepsilon}_1 R}{(1 - ilde{rac{ ilde{arepsilon}_1}{3}}) R} \ \leq rac{ ilde{arepsilon}_1}{2},$$

where the last line follows from the assumption $\tilde{\varepsilon}_1 \leq \frac{1}{2}$.

Similarly, by $\tilde{E}_1(ilde{arepsilon}_1,R)\cap \tilde{E}_2(arepsilon_2,R)$, we have

$$\left|\left\langle \frac{\boldsymbol{z}_i}{\|\boldsymbol{z}_i\|}, \frac{\boldsymbol{\mu}}{\|\boldsymbol{\mu}\|} \right\rangle \right| \leq \frac{\frac{\tilde{\varepsilon}_2}{3}R}{\sqrt{(1-\frac{\tilde{\varepsilon}_1}{3})R}\|\boldsymbol{\mu}\|} \leq \frac{\tilde{\varepsilon}_2\sqrt{R}}{2\|\boldsymbol{\mu}\|},$$

where the last inequality is due to $\tilde{\varepsilon}_1 \leq \frac{1}{2}$.

D.1 Proof or Theorem 6.2

For model (sG), we have

Lemma D.2 (Lemma S4.1, S4.2, Hashimoto et al. (2025)). Consider model (sG). There exist universal constants C_i , i=1,2 such that for any $\delta \in (0,\frac{1}{2})$, event $\tilde{E}_1 \cap \tilde{E}_2$ holds with probability at least $1-2\delta$, where

$$R = \operatorname{tr}(\Sigma),$$

$$\tilde{\varepsilon}_1 := C_1 \max \left\{ \sqrt{n \log\left(\frac{1}{\delta}\right)} \|\Sigma\|_F, n \log\left(\frac{1}{\delta}\right) \|\Sigma\| \right\} / \operatorname{tr}(\Sigma)$$

$$\tilde{\varepsilon}_2 := C_2 \sqrt{n \log\left(\frac{1}{\delta}\right)} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\| / \operatorname{tr}(\Sigma).$$

Lemma D.3. There exists a universal constant C_3 such that we have with probability at least $1-\delta$

$$|n_+ - n_-| \le C_3 \sqrt{n \log(\frac{1}{\delta})}.$$

Proof. By independence of y_i and $y_i \in \{\pm 1\}$, Hoeffding's inequality implies that

$$\left| \sum_{i} y_i \right| \le C_3 \sqrt{n \log(\frac{1}{\delta})}$$

holds with probability at least $1 - \delta$, where C_3 is a universal constant.

The desired result follows by noting that $\sum_i y_i = n_+ - n_-$.

Corollary D.4. Event \tilde{E}_3 holds with $\tilde{\varepsilon}_3 := C_3 \sqrt{\frac{\log(\frac{1}{\delta})}{n}}$ with probability at least $1 - \delta$.

With these, we can rewrite assumptions of our main theorems, namely Theorem 4.8 and C.13, using parameters defining the model (sG).

From Theorem 4.8, we have the following result on (sG).

Theorem D.5. Consider model (sG). Suppose

$$\operatorname{tr}(\Sigma) \ge \max \left\{ 2C_1 \sqrt{n \log(\frac{1}{\delta})} \|\Sigma\|_F, 2C_1 n \log(\frac{1}{\delta}) \|\Sigma\|, C_2^2 n \log(\frac{1}{\delta}) \frac{\|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\|^2}{\|\boldsymbol{\mu}\|^2} \right\}, \tag{99}$$

and further suppose one of the following conditions:

$$\begin{aligned} \text{(i)} & & \|\boldsymbol{\mu}\|^2 \geq 2.18 \max \big\{ C_1 \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}\|_F, C_1 n \log(\frac{1}{\delta}) \|\boldsymbol{\Sigma}\|, C_2 \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu}\| \big\}, \\ & & \alpha \leq \left[1.2 \{ n \|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma}) \} \right]^{-1}, \text{ and } \sigma \leq 0.2 \gamma \alpha \sqrt{\frac{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})}{m}}, \end{aligned}$$

$$\begin{split} (ii) & \operatorname{tr}(\Sigma) \geq \frac{67.2e}{\gamma^2} n \|\boldsymbol{\mu}\|^2, \\ & \operatorname{tr}(\Sigma) \geq \frac{453e}{\gamma^3} n \max \big\{ C_1 \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}\|_F, C_1 n \log(\frac{1}{\delta}) \|\boldsymbol{\Sigma}\|, C_2 \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu}\| \big\}, \\ & \alpha \leq \{8 \operatorname{tr}(\Sigma)\}^{-1}, \text{ and } \sigma \leq 0.1 \gamma \alpha \sqrt{\frac{\operatorname{tr}(\Sigma)}{m}}. \end{split}$$

Then, we have with probability at least $1-2\delta$, the gradient descent iterate (2) keeps all the neurons activated for $t \geq 1$, satisfies $\mathcal{L}(W^{(t)}) = O(t^{-1})$, and converges in the direction given in Theorem 4.8.

Proof. By Lemma D.1, D.2, and the assumption (99), all the conclusions of Lemma D.1 and D.2 hold with probability at least $1-2\delta$ with

$$\begin{split} R &= \operatorname{tr}(\Sigma), \\ \theta_1 &= \frac{\tilde{\varepsilon}_1}{2} = \frac{C_1 \max\left\{\sqrt{n \log(\frac{1}{\delta})} \|\Sigma\|_F, n \log(\frac{1}{\delta}) \|\Sigma\|\right\}}{2 \operatorname{tr}(\Sigma)} \leq \frac{1}{4}, \\ \theta_2 &= \frac{\tilde{\varepsilon}_2 \sqrt{R}}{2 \|\boldsymbol{\mu}\|} = \frac{C_2 \sqrt{n \log(\frac{1}{\delta})} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\|}{2 \sqrt{\operatorname{tr}(\Sigma)} \|\boldsymbol{\mu}\|} \leq \frac{1}{2}. \end{split}$$

Then, we have $0.91\sqrt{\mathrm{tr}(\Sigma)} \leq R_{min}(\boldsymbol{z}) \leq R_{max}(\boldsymbol{z}) \leq 1.09\sqrt{\mathrm{tr}(\Sigma)}$.

From these, we have

$$2\theta_{2} \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_{1} R_{max}^{2}(\boldsymbol{z})$$

$$\leq 1.09 C_{2} \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu}\|$$

$$+ 0.6 C_{1} \max \left\{ \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}\|_{F}, n \log(\frac{1}{\delta}) \|\boldsymbol{\Sigma}\| \right\}$$

$$\leq 2.18 \max \{ C_{1} \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}\|_{F}, C_{1} n \log(\frac{1}{\delta}) \|\boldsymbol{\Sigma}\|, C_{2} \sqrt{n \log(\frac{1}{\delta})} \|\boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu}\| \right\}.$$

$$(100)$$

We first show the desired conclusions under condition (i).

First, it is immediate from (100) to see $\|\boldsymbol{\mu}\|^2 \ge 2\theta_2 \|\boldsymbol{\mu}\| R_{max}(\boldsymbol{z}) + \theta_1 R_{max}^2(\boldsymbol{z})$ holds under condition (i).

Furthermore, condition (i) also implies that Assumption 4.3 is satisfied since

$$\alpha \{n \|\boldsymbol{\mu}\|^2 + R_{max}^2(\boldsymbol{z})\} < 1.2\alpha \{n \|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\Sigma)\} \le 1.$$

We note that $\exp(\rho) < 1.26$ holds under condition (i) since

$$\rho \le \sigma \sqrt{m \cdot \frac{3}{2} \cdot 1.2\{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\Sigma)\}}$$

$$\le 1.35\sigma \sqrt{m\{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\Sigma)\}}$$

$$\le 0.27\gamma \alpha\{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\Sigma)\}$$

$$\le \frac{0.27}{1.2}\gamma \le 0.225.$$

where the first inequality follows from the definition of ρ and $\theta_2 \leq \frac{1}{2}$ and $R_{max}(z) \leq 1.2\sqrt{\text{tr}(\Sigma)}$, the remaining inequalities follow from condition (i).

Noting $\exp(\rho) < 1.26$ and $\rho \le 0.27\gamma\alpha\{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\Sigma)\}$, we also see that Assumption 4.2 is satisfied since

$$\begin{split} \rho \exp(\rho) &< 1.26 \rho \\ &\leq 1.26 \cdot 0.27 \gamma \alpha \{ \| \boldsymbol{\mu} \|^2 + \mathrm{tr}(\Sigma) \} \\ &\leq \frac{1.26 \cdot 0.27}{0.8} \gamma \alpha \{ \| \boldsymbol{\mu} \|^2 + R_{min}^2(\boldsymbol{z}) \} \\ &\leq \gamma (1 - \theta_2) \alpha \{ \| \boldsymbol{\mu} \|^2 + R_{min}^2(\boldsymbol{z}) \}. \end{split}$$

Since all the assumptions of (i) of Theorem 4.8 are satisfied, the desired conclusions hold with probability at least $1 - 2\delta$ under condition (i).

Next we show the desired conclusions under condition (ii). To see this, first note that $\tilde{\varepsilon}_1 \leq \frac{1}{2}$ implies that $\frac{R_{max}^2(\mathbf{z})}{R_{min}^2(\mathbf{z})} \leq \frac{1+\frac{\tilde{\varepsilon}_1}{3}}{1-\frac{\tilde{\varepsilon}_1}{3}} \leq \frac{7}{5}$. So, we have C_w in Theorem 4.8 with $C_w \leq \frac{168e}{5\gamma^2}$. So, by setting $\varepsilon_i = \frac{5\gamma^2}{368e}, i = 1, 3, C_w \varepsilon_i \leq \frac{1}{2}$ is ensured. Moreover, by setting $\varepsilon_2 = \frac{1}{2}, \varepsilon_1 + \varepsilon_2 + \varepsilon_3 < 1$ is ensured.

We note that $\exp(\rho) < \frac{10}{9}$ holds under condition (ii) since

$$\begin{split} \rho &\leq 1.35\sigma\sqrt{m\{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})\}} \\ &\leq 1.35\sigma\sqrt{m(1.0055)\operatorname{tr}(\boldsymbol{\Sigma})} \\ &\leq 1.36\sigma\sqrt{m\operatorname{tr}(\boldsymbol{\Sigma})} \\ &\leq 1.36\cdot 0.1\gamma\alpha\operatorname{tr}(\boldsymbol{\Sigma}) \\ &\leq \frac{1.36\cdot 0.1}{8}\gamma \leq 0.017 < \log(\frac{10}{9}). \end{split}$$

By noting $\rho \leq 0.136 \gamma \alpha \mathrm{tr}(\Sigma)$ and $\exp(\rho) < \frac{10}{9}$, condition (ii) implies that Assumption 4.5 is satisfied since

$$\rho \exp(\rho) \leq \frac{10}{9} \cdot 0.136 \gamma \alpha \operatorname{tr}(\Sigma)
\leq \frac{10}{9} \cdot 0.136 \cdot 0.8^{-1} \gamma \alpha R_{min}^{2}(\boldsymbol{z})
\leq 0.19 \gamma \alpha R_{min}^{2}(\boldsymbol{z})
\leq \varepsilon_{2} \gamma (1 - \theta_{2}) \alpha R_{min}^{2}(\boldsymbol{z}).$$

Similarly, by noting $\exp(\rho) < \frac{10}{9}$, condition (ii) also implies that Assumption 4.7 is satisfied since

$$6\alpha R_{min}^2(\boldsymbol{z}) \exp(\rho) < 7.2 \cdot \frac{10}{9} \alpha R_{min}^2(\boldsymbol{z})$$
$$= 8\alpha \operatorname{tr}(\Sigma) \le 1.$$

By (100), condition (ii) implies that Assumption 4.4 is satisfied since

$$2\theta_{2} \| \boldsymbol{\mu} \| R_{max}(\boldsymbol{z}) + \theta_{1} R_{max}^{2}(\boldsymbol{z})$$

$$\leq 2.18 \max\{C_{1} \sqrt{n \log(\frac{1}{\delta})} \| \boldsymbol{\Sigma} \|_{F}, C_{1} n \log(\frac{1}{\delta}) \| \boldsymbol{\Sigma} \|, C_{2} \sqrt{n \log(\frac{1}{\delta})} \| \boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu} \| \}$$

$$\leq \frac{\gamma^{3} \operatorname{tr}(\boldsymbol{\Sigma})}{453en}$$

$$\leq \frac{\frac{5\gamma^{2}}{336e} \cdot \frac{1}{2} \cdot 0.8 \gamma \operatorname{tr}(\boldsymbol{\Sigma})}{n(10/9)^{2}}$$

$$\leq \frac{\varepsilon_{1} \gamma(1 - \theta_{2}) R_{min}^{2}(\boldsymbol{z})}{n \exp(2\rho)} .$$

Since all the assumptions of (ii) of Theorem 4.8 are satisfied, the desired conclusions hold with with probability at least $1-2\delta$ under condition (ii).

Application of Theorem C.13 on the model (sG) is given below:

Theorem D.6. Consider model (sG). Suppose $n \ge \max \left\{ 4C_3^2 \log(\frac{1}{\delta}), N(\gamma, q_+) \right\}$ and one of the following holds:

$$\begin{split} (i) \ \ q_{+}q_{-} &= 0, \, n \| \boldsymbol{\mu} \|^{2} < \frac{\gamma}{1 + (1 + \gamma)/2} \mathrm{tr}(\boldsymbol{\Sigma}), \\ \\ \mathrm{tr}(\boldsymbol{\Sigma}) &\geq 4 \gamma^{-3} n \sqrt{\log(\frac{1}{\delta})} \max \left\{ C_{1} \| \boldsymbol{\Sigma} \|_{F}, C_{1} \| \boldsymbol{\Sigma} \| \sqrt{n \log(\frac{1}{\delta})}, C_{2} \sqrt{n} \| \boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu} \| \right\}, \text{and} \\ \\ \| \boldsymbol{\mu} \|^{2} &\geq \tilde{D}^{-1} C_{2} \| \boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu} \| \sqrt{\log(\frac{1}{\delta})}, \end{split}$$

(ii)
$$q_+q_- \neq 0$$
, $n\|\boldsymbol{\mu}\|^2 \leq \frac{q_{\gamma}}{2\gamma(1-\gamma)} \mathrm{tr}(\Sigma)$,
$$\mathrm{tr}(\Sigma) \geq 20q_{\gamma}^{-2} n \sqrt{\log(\frac{1}{\delta})} \max\left\{ C_1 \|\Sigma\|_F, C_1 \|\Sigma\| \sqrt{n\log(\frac{1}{\delta})}, C_2 \sqrt{n} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\| \right\}, and$$
$$\|\boldsymbol{\mu}\|^2 \geq \tilde{D}^{-1} C_2 \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\| \sqrt{\log(\frac{1}{\delta})},$$

(iii)
$$q_{\gamma} > \gamma$$
, $n \|\boldsymbol{\mu}\|^2 \ge \frac{q_{\gamma}}{2\gamma(1-\gamma)} \operatorname{tr}(\Sigma)$,

$$\operatorname{tr}(\Sigma) \ge \frac{2\sqrt{10}}{q_{\gamma}\sqrt{q_{\gamma}-\gamma}} n \|\boldsymbol{\mu}\| \{\log(\frac{1}{\delta})\}^{1/4}$$

$$\times \sqrt{\max\left\{C_1 \|\Sigma\|_F, C_1 \|\Sigma\|\sqrt{n\log(\frac{1}{\delta})}, C_2\sqrt{n}\|\Sigma^{\frac{1}{2}}\boldsymbol{\mu}\|\right\}}.$$

Then, we have with probability at least $1-3\delta$

$$\mathbb{P}_{(\boldsymbol{x},y)}(yf(\boldsymbol{x};\hat{W})<0) \leq \exp\left(-c\frac{n\|\boldsymbol{\mu}\|^4}{L^2\|\boldsymbol{\Sigma}\|\{n\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})\}}\right),$$

where c is a constant which depends only on γ and q_+ .

Proof. By Lemma D.2 and Corollary D.4, event $\bigcap_{i=1,2,3} \tilde{E}_i$ holds with probability at least $1-3\delta$ with

$$R = \operatorname{tr}(\Sigma),$$

$$\tilde{\varepsilon}_1 = C_1 \max \left\{ \sqrt{n \log\left(\frac{1}{\delta}\right)} \|\Sigma\|_F, n \log\left(\frac{1}{\delta}\right) \|\Sigma\| \right\} / \operatorname{tr}(\Sigma)$$

$$\tilde{\varepsilon}_2 = C_2 \sqrt{n \log\left(\frac{1}{\delta}\right)} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\| / \operatorname{tr}(\Sigma),$$

$$\tilde{\varepsilon}_3 = C_3 \sqrt{\frac{\log(\frac{1}{\delta})}{n}}.$$

Note that the assumption $n \geq 4C_3^2 \log(\frac{1}{\delta})$ implies $\tilde{\varepsilon}_3 \leq \frac{1}{2}$.

Then, the desired conclusion follows by simply rewriting assumptions of Theorem C.13.

We can now see that Theorem 6.2 follow from Theorem D.5 and D.6 by ensuring both assumptions are satisfied. Specifically, Theorem D.5 under condition (i) and Theorem D.6 under condition (ii) and (ii) imply Theorem 6.2 under (a)-(i). Similarly, Theorem D.5 under condition (ii) and Theorem D.6 under condition (i) and (ii) imply Theorem 6.2 under condition (a)-(ii). Lastly, Theorem D.5 under condition (i) and Theorem D.6 under condition (iii) imply Theorem 6.2 under condition (b).

D.2 PROOF OF THEOREM 6.3

For model (PM), we have instead of Lemma D.2

Lemma D.7 (Lemma S3.4, S3.5, Hashimoto et al. (2025)). Consider model (PM). There exist a constant C(r, K) such that for any $\delta \in (0, \frac{1}{2})$, event $\tilde{E}_1 \cap \tilde{E}_2$ holds with probability at least $1 - 2\delta$, where

$$\begin{split} R &= \operatorname{tr}(\Sigma), \\ \tilde{\varepsilon}_1 &:= C(r, K) \delta^{-\frac{2}{r}} \max \left\{ n^{\frac{2}{r}} p^{\frac{2}{r} - \frac{1}{2}}, n^{\frac{4}{r}} \right\} \frac{\|\Sigma\|_F}{\operatorname{tr}(\Sigma)}, \\ \tilde{\varepsilon}_2 &:= \sqrt{\frac{n}{\delta}} \frac{\|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\|}{\operatorname{tr}(\Sigma)}. \end{split}$$

Similar arguments using D.7 instead of D.2 allows us to establish results analogous to Theorem D.5 and D.6.

Theorem D.8. Consider model (PM). Let $\delta \in (0, \frac{1}{2})$. Suppose, for sufficiently large constant C which depends on $\delta, \gamma, q_+, r, K$,

$$\operatorname{tr}(\Sigma) \ge C \max \left\{ n^{\frac{2}{r}} p^{\frac{2}{r} - \frac{1}{2}} \|\Sigma\|_F, n^{\frac{4}{r}} \|\Sigma\|_F, n^{\frac{1}{2}} \frac{\mu}{\|\mu\|^2} \right\}$$
(101)

and further suppose one of the following conditions:

$$\begin{split} (i) & \ \|\boldsymbol{\mu}\|^2 \geq C \max \big\{ n^{\frac{2}{r}} p^{\frac{2}{r} - \frac{1}{2}} \|\boldsymbol{\Sigma}\|_F, n^{\frac{4}{r}} \|\boldsymbol{\Sigma}\|_F, \sqrt{n} \|\boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\mu}\| \big\}, \\ & \alpha \leq \left[1.2 \{ n \|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma}) \} \right]^{-1}, \text{ and } \sigma \leq 0.2 \gamma \alpha \sqrt{\frac{\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})}{m}}, \end{split}$$

$$\begin{split} \text{(ii)} \ \ & \text{tr}(\Sigma) \geq C \max \big\{ n \| \pmb{\mu} \|^2, n^{\frac{2}{r}+1} p^{\frac{2}{r}-\frac{1}{2}} \| \Sigma \|_F, n^{\frac{4}{r}+1} \| \Sigma \|_F, n^{\frac{3}{2}} \| \Sigma^{\frac{1}{2}} \pmb{\mu} \| \big\}, \\ & \alpha \leq \{8 \text{tr}(\Sigma)\}^{-1}, \text{ and } \sigma \leq 0.1 \gamma \alpha \sqrt{\frac{\text{tr}(\Sigma)}{m}}. \end{split}$$

Then, we have with probability at least $1-2\delta$, the gradient descent iterate (2) keeps all the neurons activated for $t \geq 1$, satisfies $\mathcal{L}(W^{(t)}) = O(t^{-1})$, and converges in the direction given in Theorem 4.8.

Proof. The theorem follows from Theorem 4.8, Lemma D.1, and Lemma D.7 by repeating the same argument from the proof of Theorem D.5 by replacing Lemma D.2 with Lemma D.7. □

Similarly, we obtain a result analogous to Theorem D.6.

Theorem D.9. Consider model (PM). Let $\delta \in (0, \frac{1}{3})$. Suppose $n \ge \max\left\{4C_3^2\log(\frac{1}{\delta}), N(\gamma, q_+)\right\}$ and one of the following holds for sufficiently large constant C which depends on $\delta, \gamma, q_+, r, K$:

(i)
$$\operatorname{tr}(\Sigma) \ge C \max\{n\|\boldsymbol{\mu}\|^2, n^{\frac{2}{r} + \frac{1}{2}} p^{\frac{2}{r} - \frac{1}{2}} \|\Sigma\|_F, n^{\frac{4}{r} + \frac{1}{2}} \|\Sigma\|_F, n^{\frac{3}{2}} \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\| \}$$
 and
$$\|\boldsymbol{\mu}\|^2 \ge C \|\Sigma^{\frac{1}{2}} \boldsymbol{\mu}\|,$$

$$\begin{aligned} \text{(ii)} \ \ q_{\gamma} > \gamma, \, n \| \pmb{\mu} \|^2 &\geq \frac{q_{\gamma}}{2\gamma(1-\gamma)} \mathrm{tr}(\Sigma), \, and \\ \\ \mathrm{tr}(\Sigma) &\geq C \| \pmb{\mu} \| \max \left\{ n^{\frac{1}{r} + \frac{3}{4}} p^{\frac{1}{r} - \frac{1}{4}} \| \Sigma \|_F^{\frac{1}{2}}, n^{\frac{2}{r} + \frac{3}{4}} \| \Sigma \|_F^{\frac{1}{2}}, n^{\frac{5}{4}} \| \Sigma^{\frac{1}{2}} \pmb{\mu} \|^{\frac{1}{2}} \right\}. \end{aligned}$$

Then, we have with probability at least $1-3\delta$

$$\mathbb{P}_{(\boldsymbol{x},y)}(yf(\boldsymbol{x};\hat{W})<0) \leq \exp\left(-c\frac{n\|\boldsymbol{\mu}\|^4}{L^2\|\boldsymbol{\Sigma}\|\{n\|\boldsymbol{\mu}\|^2 + \operatorname{tr}(\boldsymbol{\Sigma})\}}\right),$$

where c is a constant which depends only on γ and q_+ .

Proof. The theorem follows from Theorem C.13, Lemma D.1, Lemma D.7, and Lemma D.4 by repeating the same argument from the proof of Theorem D.6 by replacing Lemma D.2 with Lemma D.7.

Putting these together with $\Sigma = I_p$, we obtain Theorem 6.3. Specifically, Theorem 6.3 under condition (i) follows from Theorem D.8 under (i) and Theorem D.9 under (i), Theorem 6.3 under condition (ii) follows from Theorem D.8 under (i) and Theorem D.9 under (ii), and Theorem 6.3 under condition (iii) follows from Theorem D.8 under (ii) and Theorem D.9 under (i).