

Tweedie Moment Projected Diffusions for Inverse Problems

Anonymous authors

Paper under double-blind review

Abstract

Diffusion generative models unlock new possibilities for inverse problems as they allow for the incorporation of strong empirical priors into the process of scientific inference. Recently, diffusion models are repurposed for solving inverse problems using Gaussian approximations to conditional densities of the reverse process via Tweedie’s formula to parameterise the mean, complemented with various heuristics. To address various challenges arising from these approximations, we leverage higher order information using Tweedie’s formula and obtain a statistically principled approximation. We further provide a theoretical guarantee specifically for posterior sampling which can lead to better theoretical understanding of diffusion-based conditional sampling. Finally, we illustrate the empirical effectiveness of our approach for general linear inverse problems on toy synthetic examples as well as image restoration. We show that our method (i) removes any time-dependent step-size hyperparameters required by earlier methods, (ii) brings stability and better sample quality across multiple noise levels, (iii) is the only method that works in a stable way with variance exploding (VE) forward processes as opposed to earlier works.

1 Introduction

Solving inverse problems is one of the most crucial goals in scientific inference (Tarantola, 2005; Stuart, 2010). Diffusion models (Song et al., 2020; Ho et al., 2020) recently emerged as a strong alternative to classical Bayesian approaches to solve inverse problems to build samplers for complicated and implicit conditional distributions. With their flexibility, these models replace handcrafted priors on the latent signal with pretrained and strong empirical priors. For example, given a latent signal \mathbf{x}_0 (say a face image), we can *train* a diffusion model to sample from the prior $p(\mathbf{x}_0)$. The idea for solving inverse problems is to leverage extraordinary modelling power of diffusion models to learn samplers for priors and couple this with a given likelihood $p(\mathbf{y}|\mathbf{x}_0)$ for a given data \mathbf{y} to sample from posterior densities for inverse problems. However, designing a diffusion model for the posterior comes with its own challenges due to intractability. Despite its challenges, this approach has been recently taking off with lots of activity in the field, e.g. for compressed sensing (Bora et al., 2017; Kadkhodaie & Simoncelli, 2021), projecting score-based stochastic differential equations (SDEs) (Song et al., 2021b), gradient-based approaches (Daras et al., 2022; Chung et al., 2022b), magnetic resonance imaging (MRI) by approximating annealed Langevin dynamics with approximate scores (Jalal et al., 2021), image restoration (Kawar et al., 2022), score-based models as priors but with a normalising flow approach (Feng et al., 2023), variational approaches (Mardani et al., 2023; Feng & Bouman, 2023). Most relevant ideas to us, which we will review in detail in Section 5, use Tweedie’s formula (Efron, 2011; Laumont et al., 2022) to approximate the smoothed likelihood, e.g. Diffusion posterior sampling (DPS) (Chung et al., 2022a; 2023) and pseudo-guided diffusion models (PIGDM) (Song et al., 2023). Similar approaches are also exploited using singular-value decomposition (SVD) based approaches (Kawar et al., 2021). In this work, we develop an approach which builds a tighter approximation to optimal formulae for approximating the scores of the posterior diffusion model.

This paper is devoted developing novel methods to solving inverse problems, given a latent (target) signal $\mathbf{x}_0 \in \mathbb{R}^{d_x}$, noisy observed data $\mathbf{y} \in \mathbb{R}^{d_y}$, a known linear observation map \mathbf{H} , and a *pretrained* diffusion prior. The main tool we use is Tweedie’s formula

to obtain both the mean and the covariance for approximating diffused likelihoods, to be used for building the final posterior score approximation. This is as opposed to previous works which only utilised first moment approximations using Tweedie’s formula (Chung et al., 2022a; Song et al., 2023). We show that utilising covariance approximation, with moment projections, provides a principled scheme with improved performance, which we term *Tweedie Moment Projected Diffusions* (TMPD).

To demonstrate our method briefly, Figure 1 demonstrates a sampling scenario of a *Gaussian random field* (GRF) whose mean and variance entries are plotted under “Analytic” column¹. We demonstrate the approximations under this setting provided by our method (TMPD) and its diagonal (cheaper) version (DTMPD), compared with Π GDM (Song et al., 2023), and DPS (Chung et al., 2022a). The figure demonstrates the optimality of our method: Our first and second moment approximations become exact in this case. This results in a drastic performance improvement stemming from the statistical optimality of our method for near-Gaussian settings and also unlocks a possible line for theoretical research for understanding similar diffusion models for inverse problems.

In what follows, we will first introduce the technical background in Section 2 and then describe TMPD in detail in Section 3. We will then provide some theoretical results about our method in Section 4 and provide a discussion to closely related work in literature in Section 5. Finally, Section 6 will present experiments on Gaussian mixtures, image inpainting and super-resolution, demonstrating quantitative and qualitative improvements provided by TMPD.

2 Technical Background

There are two main diffusion modelling paradigms, namely, score-based models (SGMs) (Song et al., 2020) and denoising diffusion probabilistic models (DDPM) (Ho et al., 2020). In both, the goal is to sample from a target distribution $p_0 := p_{\text{data}}$. To that end, an artificial path p_t is introduced, with the property that p_t will approach $\mathcal{N}(0, I)$ for large t , i.e., $p_t \rightarrow \mathcal{N}(0, I_d)$ as $t \rightarrow \infty$. Then, one learns to reverse this process, so that one can sample from a standard normal distribution and transform these samples to samples from p_{data} . In the SDE paradigm, the interpolation parameter t will take continuous values in $t \in [0, T]$, whereas in the DDPM setting, t is discrete. However, the DDPM setting can be seen as discretization of the SDE (Song et al., 2020).

In SGMs, a stochastic differential equation (SDE) is used to noise the data. A very common choice is to use the time-rescaled Ornstein-Uhlenbeck process:

$$\begin{aligned} d\mathbf{x}_t &= -\frac{1}{2}\beta(t)\mathbf{x}_t dt + \sqrt{\beta(t)}d\mathbf{w}_t, \\ \mathbf{x}_0 &\sim p_0 = p_{\text{data}}. \end{aligned} \tag{1}$$

¹We only plot the variances for visualisation while the GRF has a full covariance.




Figure 1: Error to target posterior for a *Gaussian random field*. (Top row) visualisation of the empirical mean and variance of the 1500 samples that were used to compute this error against the analytical moments. (Bottom) Wasserstein distances of different methods w.r.t. sample size. For details, see Appendix E.2.

The corresponding reverse SDE is then given by

$$\begin{aligned} d\mathbf{z}_t &= \frac{1}{2}\beta(T-t)\mathbf{z}_t dt + \beta(T-t)\nabla_{\mathbf{z}_t} \log p_{T-t}(\mathbf{z}_t)dt + \sqrt{\beta(T-t)}d\bar{\mathbf{w}}_t, \\ \mathbf{z}_0 &\sim p_T. \end{aligned} \quad (2)$$

The transition kernels for the time-rescaled SDE can be derived from the transition kernels of a normal OU process $p_t(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_t; \sqrt{\alpha_t}\mathbf{x}_0, v_t\mathbf{I}_{d_x})$ where $\alpha_t := \exp\left(-\int_0^t \beta(s)ds\right)$ and $v_t := 1 - \alpha_t$. A parameterisation that performs well in practice is $\beta(t) = \beta_{\min} + t(\beta_{\max} - \beta_{\min})$. In the diffusion modelling literature, the time-rescaled OU process is also sometimes called a Variance Preserving SDE. However, this is not the only SDE that is suitable for the forward process. See Appendix C for details on a time-rescaled Brownian motion (Variance Exploding SDE (Song et al., 2020)).

There are two usual approximations to solve the SDE in equation 2. First, we do not know p_T , since it is a noised version of the distribution p_{data} . However for T large enough, we can approximate $p_T \approx p_{\text{ref}} = \mathcal{N}(0, I_d)$. We also do not have $\nabla \log p_{T-t}$ which we need for the drift in equation 2. This can be circumvented by approximating the drift using score-matching techniques (Hyvärinen, 2005; Ho et al., 2020). These methods construct an estimate of the score function by solving the score matching problem in the form of $\mathbf{s}_\theta(\mathbf{x}_t, t) \approx \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$. This score can also be used in the setting of DDPM (Ho et al., 2020).

2.1 Conditional sampling for the linear inverse problem

In the preceding section we introduced diffusion models as a method to sample from a target distribution p_{data} . We now suppose that we have access to measurements, or observations $\mathbf{y} \in \mathbb{R}^{d_y}$ of $\mathbf{x}_0 \in \mathbb{R}^{d_x}$:

$$\mathbf{y} = \mathbf{H}\mathbf{x}_0 + \mathbf{u}, \quad \mathbf{u} \sim \mathcal{N}(0, \sigma_y^2 \mathbf{I}_{d_y}). \quad (3)$$

We would then be interested in sampling from the conditional distribution of \mathbf{x}_0 given \mathbf{y} , i.e., $p_{\text{data}}(\cdot | \mathbf{y})$. To that end, we have to modify the reverse SDE. Optimally, we would like to sample from the reverse SDE to the forward SDE started in $p_{\text{data}}(\cdot | \mathbf{y})$, instead of the one started in p_{data} .

Optimally, we would want to replace the score $\nabla_{\mathbf{z}_t} \log p_{T-t}(\mathbf{z}_t)$ in equation 2 with the posterior score $\nabla_{\mathbf{z}_t} \log p_{T-t|\mathbf{y}}(\mathbf{z}_t|\mathbf{y})$. Written in terms of the forward process, this coincides with

$$\nabla_{\mathbf{x}_t} \log p_{t|\mathbf{y}}(\mathbf{x}_t|\mathbf{y}) = \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) + \nabla_{\mathbf{x}_t} \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t). \quad (4)$$

The term $p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t)$ is given by the integral

$$p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) = \int p_{\mathbf{y}|0}(\mathbf{y}|\mathbf{x}_0)p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)d\mathbf{x}_0, \quad (5)$$

which involves a marginalization over \mathbf{x}_0 . The term $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ however is only defined implicitly through running the diffusion model, making the above integral difficult to evaluate. One way around this is to train a neural network to directly approximate $\nabla \log p_{t|\mathbf{y}}(\mathbf{x}_t|\mathbf{y})$ (Batzolis et al., 2021). Alternatively, if one already has access to an approximation of $\nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$, one can train an auxiliary network to approximate the term $\nabla \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t)$ in equation 4, (Song et al., 2020; Dhariwal & Nichol, 2021). However, these methods can be time and training-data intensive, as one needs to retrain networks for each conditional task as well have access to paired training data from the joint distribution $(\mathbf{x}_0, \mathbf{y})$. Alternatively, one could indeed try to do a Monte-Carlo approximation of the score corresponding to equation 5. But this needs evaluating the probability flow ODE together with its derivative (Song et al., 2020, Section D.2) for each sample, which is prohibitive, also suffers from high variance (Mohamed et al., 2020, Section 3).

3 Tweedie Moment Projected Diffusions

In this section, we first introduce *Tweedie moment projections* in Section 3.1 below. Our method relies on the approximation $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) \approx \mathcal{N}(\mathbf{x}_0; \mathbf{m}_{0|t}(\mathbf{x}_t), \mathbf{C}_{0|t}(\mathbf{x}_t))$ to make the sampling process tractable. In

that case, since the conditional distribution of \mathbf{y} given \mathbf{x}_0 is also Gaussian, we can *compute the integral in equation 5 analytically* — $p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t)$ will just be another Gaussian in that case, its mean and covariance being determined through $\mathbf{m}_{0|t}$, $\mathbf{C}_{0|t}$, \mathbf{H} and σ_y . In particular, we can then use this Gaussian to approximate $\nabla \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t)$, since the score of a Gaussian is available in closed form.

3.1 Tweedie moment projections

Instead of just approximating the variance of $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ heuristically, we approximate it by projecting on to the closest Gaussian distribution using Tweedie’s formula for the second moment. Our approximation at this stage consists of two main steps: (i) Find the mean and covariance of $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ using Tweedie’s formula, and (ii) approximate this density with a Gaussian using the mean and covariance of $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ (moment projection). Due to this approximation, we will refer to resulting methods as *Tweedie Moment Projected Diffusions* (TMPD). We will first introduce Tweedie’s formula for the mean and covariance and then describe the moment projection.

Proposition 1 (Tweedie’s formula). *Let $\mathbf{m}_{0|t}$ and $\mathbf{C}_{0|t}$ be the mean and the covariance of $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$, respectively. Then given the marginal density $p_t(\mathbf{x}_t)$, the mean is given as*

$$\mathbf{m}_{0|t} = \mathbb{E}[\mathbf{x}_0|\mathbf{x}_t] = \frac{1}{\sqrt{\alpha_t}}(\mathbf{x}_t + v_t \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)).$$

Then the covariance $\mathbf{C}_{0|t}$ is given by

$$\begin{aligned} \mathbf{C}_{0|t} &= \mathbb{E}[(\mathbf{x}_0 - \mathbf{m}_{0|t})(\mathbf{x}_0 - \mathbf{m}_{0|t})^\top | \mathbf{x}_t] \\ &= \frac{v_t}{\alpha_t}(\mathbf{I}_{d_x} + v_t \nabla^2 \log p_t(\mathbf{x}_t)) = \frac{v_t}{\sqrt{\alpha_t}} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}. \end{aligned} \quad (6)$$

The proof is an adaptation of Meng et al. (2021, Theorem 1), see Appendix A.1. While $\mathbf{m}_{0|t}$ and $\mathbf{C}_{0|t}$ gives us the moments of the density $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$, we do not have the exact form of this density. At this stage, we employ *moment projection*, i.e., we choose the closest Gaussian in Kullback-Leibler (KL) divergence which is a distribution with same first and second moments, as formalised next.

Proposition 2 (Moment projection). *Let $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ be a distribution with mean $\mathbf{m}_{0|t}$ and covariance $\mathbf{C}_{0|t}$. Let $\hat{p}_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$ be the the closest Gaussian in KL divergence to $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$, i.e.,*

$$\hat{p}_{0|t}(\mathbf{x}_0|\mathbf{x}_t) = \arg \min_{q \in \mathcal{Q}} D_{\text{KL}}(p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) || q),$$

where \mathcal{Q} is the family of multivariate Gaussian distributions. Then

$$\hat{p}_{0|t}(\mathbf{x}_0|\mathbf{x}_t) = \mathcal{N}(\mathbf{x}_0; \mathbf{m}_{0|t}, \mathbf{C}_{0|t}).$$

This is a well-known moment matching result, see, e.g., Bishop (2006, Section 10.7). Merging Propositions 1 and 2 leads to the following *Tweedie moment projection*:

$$p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) \approx \mathcal{N}\left(\mathbf{x}_0; \mathbf{m}_{0|t}, \frac{v_t}{\sqrt{\alpha_t}} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}\right), \quad (7)$$

where $\mathbf{m}_{0|t}$ is given in Proposition 1. In the next section, we demonstrate how to use this approximation to obtain approximate likelihoods.

3.2 Tweedie Moment Projected Likelihood Approximation

We next use the approximation in equation 7 to compute the following integral *analytically*

$$\begin{aligned} p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) &= \int p_{\mathbf{y}|0}(\mathbf{y}|\mathbf{x}_0) p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) d\mathbf{x}_t \\ &= \mathcal{N}\left(\mathbf{y}; \mathbf{H} \mathbf{m}_{0|t}, \mathbf{H} \frac{v_t}{\sqrt{\alpha_t}} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y}\right) \end{aligned} \quad (8)$$

which leads to the approximation

$$\begin{aligned} f^{\mathbf{y}}(\mathbf{x}_t) &:= \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top (\mathbf{H} \frac{v_t}{\sqrt{\alpha_t}} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H} \mathbf{m}_{0|t}) \\ &\approx \nabla_{\mathbf{x}_t} \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t), \end{aligned} \quad (9)$$

where $\nabla_{\mathbf{x}_t}$ only operates on $\mathbf{m}_{0|t}$ in the above equation. We note that this approximation is exact for the case p_{data} is a Gaussian. However, otherwise, there are several approximations behind equation 9. In particular, the approximation in equation 9 treats $\mathbf{C}_{0|t}$ *constant* w.r.t. \mathbf{x}_t when computing the gradient (which is the case if p_{data} is Gaussian). For non-Gaussian p_{data} , this results in computationally efficient sampler, as otherwise the resulting terms can be expensive to compute.

3.3 Algorithms

Plugging the approximation in equation 9 into the reverse SDE in equation 2 together with the prior score as described in Section 2.1 results in a TMPD for conditional sampling to solve inverse problems. The SDE we will approximate numerically to sample from the conditional distribution is given by

$$d\mathbf{z}_t = \frac{1}{2}\beta(T-t)\mathbf{z}_t dt + \beta(T-t)(\nabla_{\mathbf{z}_t} \log p_{T-t}(\mathbf{z}_t) + f_{T-t}^{\mathbf{y}}(\mathbf{z}_t))dt + \sqrt{\beta(T-t)}d\bar{\mathbf{w}}_t \quad (10)$$

where $\mathbf{z}_0 \sim p_T$ and $f_t^{\mathbf{y}}(\mathbf{z}_t) \approx \nabla \log p_{\mathbf{y}|T-t}(\mathbf{y}|\mathbf{z}_t)$ is our approximation to the data likelihood, given by equation 9. We call this SDE the *TMPD SDE*.

We have two options to convert TMPD SDE into implementable methods: (1) score-based samplers which we abbreviate also as TMPD as they are Euler-Maruyama discretizations of the TMPD SDE akin to Song et al. (2020), (2) denoising diffusion models (TMPD-D). The denoising diffusion approach is derived from approximate reverse Markov chains and is the approach of DDPM and DPS methods (Ho et al., 2020; Chung et al., 2022a). We note that the Gaussian projection can be used in this discrete setting, assuming that the conditional density is available analytically as in Ho et al. (2020), and can be written as $p_{n|0}(\mathbf{x}_n|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_n; \sqrt{\alpha_n}\mathbf{x}_0, v_n\mathbf{I}_{d_x})$. The idea is to update the unconditional mean $\mathbf{m}_{0|n}(\mathbf{x}_n)$ of the density $p_{0|n}(\mathbf{x}_0|\mathbf{x}_n)$ with a Bayesian update: $p(\mathbf{x}_0|\mathbf{x}_n, \mathbf{y}) \propto p(\mathbf{y}|\mathbf{x}_n)p_{0|n}(\mathbf{x}_0|\mathbf{x}_n)$. Given a similar formulation as above, assuming we have a readily available approximation $p_{0|n}(\mathbf{x}_0|\mathbf{x}_n) \approx \mathcal{N}(\mathbf{x}_0; \mathbf{m}_{0|n}, \mathbf{C}_{0|n})$ and a likelihood similar to equation 8 where t can be replaced by n , we can compute the moments of $p(\mathbf{x}_0|\mathbf{x}_n, \mathbf{y})$ analytically, which we denote $\mathbf{m}_{0|n}^{\mathbf{y}}$ and $\mathbf{C}_{0|n}^{\mathbf{y}}$. The Bayes update for Gaussians gives (Bishop, 2006)

$$\mathbf{m}_{0|n}^{\mathbf{y}} = \mathbf{m}_{0|n} + \mathbf{C}_{0|n} \mathbf{H}^\top (\mathbf{H} \mathbf{C}_{0|n} \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_x})^{-1} (\mathbf{y} - \mathbf{H} \mathbf{m}_{0|n}). \quad (11)$$

Incorporating equation 11 for $n = N-1, \dots, 0$ into the usual Ancestral sampling (Ho et al., 2020) steps leads to Algorithm 1, termed TMPD-Denoising (TPMD-D). The update in equation 11 can be used in any discrete sampler such as denoising diffusion implicit models (DDIM) (Song et al., 2021a).

3.4 Computationally cheaper approximation of Moment Projection

We show in our experiments promising results for TMPD that motivate the exploration of less computationally expensive approximations to the full Jacobian. In particular, we empirically study a computationally inexpensive method that is applicable to inpainting and super-resolution, below.

To make the computational cost of TMPD smaller, we can make an approximation of the Gaussian Projection that requires fewer Jacobian-vector products and does not require linear solves. One approximation that we found useful for sampling from high dimensional diffusion models, e.g., high resolution images, is denoted here as diagonal Tweedie Moment Projection (DTMPD). Instead of the full second moment, DTMPD uses the diagonal of the second moment $\nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \approx \text{diag}(\nabla_{\mathbf{x}_t} \mathbf{m}_{0|t})$. Intuitively, this approximation will perform well empirically since it is a similar approximation to Π GDM that assumes dimensional independence of the distribution $p(\mathbf{x}_0|\mathbf{x}_t)$, but unlike Π GDM, this diagonal approximation is the same as using the closest dimensionally independent Gaussian in KL divergence to $p_{0|t}(\mathbf{x}_0|\mathbf{x}_t)$.

The biggest drawback of our method is that without further approximation, it doesn't scale up to the high dimensions of image data. This is because even calculating the diagonal of a Jacobian requires computing d_x vector-Jacobian products since in general every element of the Jacobian at a location \mathbf{x}_t , $\nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}$ depends on every element of \mathbf{x}_t . Therefore we must resort to a further approximation that exploits knowledge of the observation operator \mathbf{H} .

For the cases of super-resolution and inpainting, a further approximation that allows scaling up to the dimensions of image data is approximating the diagonal of the Jacobian by the row sum of the Jacobian which only requires a single vector jacobian product and brings the memory and time complexity of DTMPD down to that of IIGDM. We exploit the sparsity of \mathbf{H} to make the rowsum approximation of the diagonal more accurate by masking out (zeroing) the values in the vector Jacobian product that will not contribute to the diagonal of $\mathbf{H}\mathbf{C}_{0|n}\mathbf{H}^\top$. We discuss a justification of this approximation in Ap. E.1. We use this approximation in the image experiments and find that in practice it is only $(1.5 \pm 0.1) \times$ slower than IIGDM and DPS across all of our experiments (Sec. 6), with competitive sample quality for noisy inverse problems and without the need for expensive hyperparameter tuning.

4 Theoretical Guarantees

Because of the approximations, our method, as well as IIGDM Song et al. (2023) and DPS (Chung et al., 2022a) do not sample the exact posterior for general prior distributions. Therefore, one cannot hope for these methods to sample the true posterior and a priori it is not even clear how the sampled distribution relates to the true posterior. Without further justification, such methods should only be interpreted as *guidance methods*, where paths are guided to regions where a given observation \mathbf{y} is more likely, not as posterior sampling methods.

We justify our approximation by showing that the TMPD-SDE in equation 10 is able to sample the exact posterior in the Gaussian case. One can see that this is in contrast to IIGDM and DPS in our numerical experiments or by explicitly evaluating their approximations on simple one-dimensional examples.

Proposition 3 (Gaussian data distribution). *Assume that p_{data} is Gaussian. Then, the posterior score expression using equation 9 is exact, i.e., if there are no errors in the initial condition and drift approximation $s_\theta(\mathbf{x}_t, t) = \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$, the TMPD will sample $p_{\text{data}}(\cdot|\mathbf{y})$ at its final time.*

The proof is given in Appendix B.1. However, most distributions will not be Gaussian. The following theorem generalizes the above proposition to *non-Gaussian distributions*, as long as they have a density with respect to a Gaussian. We study how close our sampled measure will be to the true posterior distribution and give explicit bounds on the total variation distance in terms of the regularity properties of the density:

Theorem 1 (General data distribution). *Assume that the data distribution p_{data} can be written as*

$$p_{\text{data}}(\mathbf{x}_0) = \exp(\Phi(\mathbf{x}_0))\mathcal{N}(\mathbf{x}_0; \boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0), \quad (12)$$

for some $\boldsymbol{\mu}_0$ and $\boldsymbol{\Sigma}_0$. We furthermore assume that for some $M \geq 1$, it holds that $1/M \leq \exp(\Phi(\mathbf{x})) \leq M$ and $\|\nabla_{\mathbf{x}}\Phi(\mathbf{x})\| \leq L$ for all $\mathbf{x} \in \mathbb{R}^{d_x}$. Then

$$\|p_{\text{data}}(\cdot|\mathbf{y}) - q_T(\cdot)\|_{\text{TV}} \leq C(1 + T^{1/2}) \left((M^{5/2} - 1)(L^{1/2} + 1) + L^{1/2} \right),$$

where q_t denotes the law of the corresponding reverse-time process for equation 10 at time t and the constant C that only depends on $\mathbf{y}, \mathbf{H}, \sigma_y, \boldsymbol{\mu}_0$ and $\boldsymbol{\Sigma}_0$.

See Appendix B.2 for a proof.

Remark 1. We note that this analysis can be extended to account for (i) discretization error, (ii) the error in the initial distribution of the reverse SDE, (iii) the score approximation error using the techniques from Chen et al. (2023). We think this is a fruitful direction for our work and we leave it for future work.

In the limit, when p_{data} becomes more similar to a Gaussian, the Φ in Equation 12 converges to one, and therefore $M \rightarrow 1$ and $L \rightarrow 0$. In particular, the right hand side in equation 1 converges to 0 and we recover the result of Proposition 3. When p_{data} is not Gaussian, the right hand side of equation 1 gives us an upper bound of the our sample distribution to the true posterior.

5 Related Work

In this section, we review two closely related methods which we use as benchmarks and summarise the relationship between methods.

The first work is by Chung et al. (2022a), abbreviated DPS-D here² where $p_t(\mathbf{x}_0|\mathbf{x}_t)$ is approximated by a Dirac delta (point) distribution centred at $\mathbf{m}_{0|t}$ computed by Tweedie’s formula, within the integral given in equation 5. The mean is chosen the same way as our method. Authors choose a Dirac centred at this mean, in our framework this corresponds to choosing a zero covariance, i.e.,

$$\mathbf{m}_{0|t}^{\text{DPS-D}} = \mathbf{m}_{0|t} \quad \text{and} \quad \mathbf{C}_{0|t}^{\text{DPS-D}} = 0. \quad (13)$$

In the work Song et al. (2023), abbreviated IIGDM-D here³ the same estimator for the mean is chosen. However, the variance is set to a multiple of the identity, corresponding to choices

$$\mathbf{m}_{0|t}^{\text{IIG}} = \mathbf{m}_{0|t} \quad \text{and} \quad \mathbf{C}_{0|t}^{\text{IIG}} = r_t^2 \mathbf{I}_{d_x} \quad (14)$$

The choice of r_t is such that it matches the variance of the reverse SDE if p_{data} would be a standard normal distribution. Since they employ a different forward SDE to ours (variance exploding SDE), r_t is set, when the variance of the data is 1, to be equal to $v_t/(v_t + 1)$. In our case, with the OU/variance preserving SDE as a forward process, r_t would be equal to $v_t/(v_t + \alpha_t)$, when the variance of the data is 1.

Another relevant very recent work by Finzi et al. (2023) arrives at the approximation matching ours in the context of modelling physical constraints. However, we focus on general linear inverse problems outside physical domain in this paper together with a novel theoretical result. Also we note the work of Stevens et al. (2023) who consider maximum-a-posteriori (MAP) approach to find the moments of $p_{0|t}$.

6 Experiments

In this section, we demonstrate our results as well as the performance of other approximations to the likelihood provided in Chung et al. (2022a); Song et al. (2023). In particular, we perform comparisons for two of our methods TMPD (an SGM using our approximation) and TMPD-D (a DDPM sampler using equation 11). We compare these to DPS (an SGM sampler using the posterior approximation in equation 13), DPS-D (Chung et al., 2022a) (a DDPM-type sampler using equation 13), IIGDM (Song et al., 2023) (an SGM sampler using the posterior approximation in equation 14), and finally IIGDM-D (a DDIM-type sampler using equation 14, but in our experiments a DDPM-type sampler since we set the DDIM hyperparameter $\eta = 1.0$ which is defined in Algorithm 1 by Song et al. (2021a) who show that this is equivalent to a DDPM-type sampler).

We will make available the code to run all of our experiments.

6.1 Gaussian Mixture Model

We now demonstrate a non-linear SDE example and follow the Gaussian mixture model example of Cardoso et al. (2023) where the data distribution $p_0(\mathbf{x}_0)$ is a mixture of 25 Gaussian distributions. The means and variances of the components of the mixture are given in Appendix E.3. In this case, for each choice of observation \mathbf{y} , observation map \mathbf{H} and measurement noise standard deviation σ_y , the target posterior can be computed explicitly (see Appendix E.3).

To investigate the performance of posterior sampling methods, for each pair of dimensions and observation noise $(d_x, d_y, \sigma_y) \in \{8, 80, 800\} \times \{1, 2, 4\} \times \{10^{-2}, 10^{-1}, 10^0\}$ we randomly generate multiple measurement models $(\mathbf{y}, \mathbf{H}) \in \mathbb{R}^{d_y} \times \mathbb{R}^{d_y \times d_x}$, and equally weight each component of the Gaussian mixture. Further details are given in Appendix E.3. We chose to control the dimension to gain insight into the performance of

²Since the authors run DDPM-type sampler, we re-abbreviate DPS as DPS-D in this work (as DPS approximation itself can be also run with Euler-Maruyama schemes).

³Since the authors run DDIM-type sampler, we re-abbreviate IIGDM as IIGDM-D in this work (as IIGDM approximation itself can also be run with Euler-Maruyama schemes).




Figure 2: We display the first two dimensions of the GMM inverse problem for one of the measurement models tested ($\mathbf{H}, \sigma_y = 0.1, (d_x, d_y) = (80, 1)$). The blue dots represent samples from the target posterior, while the red dots correspond to samples generated by each of the algorithms used (the names of the algorithms are given at the bottom of each column).

posterior sampling methods under varying dimensions. We chose to control and the noise level since the different posterior sampling methods have accuracy that depends on the signal to noise ratio. Through randomly varying the observation model, we gain an insight into the performance of the posterior sampling methods with different levels of posterior multimodality. This example is interesting because it allows us to study the behaviour of our methods on non-linear problems in high dimensions whilst having access to the target posterior with which to compare (usually, obtaining a ‘ground-truth’ posterior is not feasible for non-linear problems).

To compare the posterior distribution estimated by each algorithm with the target posterior distribution, we use the sliced Wasserstein (SW) distance defined in Appendix E.3. We use 10^4 slices for the SW distance and compare 1000 samples of TMPD-D, IIGDM-D and DPS-D in Tables 1 obtained using 1000 denoising steps and 1000 samples of the true posterior distribution.

Table 1 indicates the Central Limit Theorem (CLT) 95% confidence intervals obtained

Table 1: Sliced Wasserstein for the GMM case. The full table is in Ap. E.3.

d_x	8	8	8	80	80	80	800	800	800
d_y	1	2	4	1	2	4	1	2	4
TMPD-D	1.6	0.7	0.3	2.7	1.0	0.3	3.1	1.4	0.4
DTMPD-D	1.8	3.3	0.4	2.8	3.2	0.7	3.7	3.5	0.7
DPS-D	4.7	1.8	0.7	5.6	3.2	1.2	5.8	3.5	1.4
IIGDM-D	2.6	2.1	3.8	3.2	2.8	0.6	3.5	3.1	0.4
TMPD-D	1.4	0.9	0.3	2.3	1.2	0.4	2.9	1.3	0.4
DTMPD-D	1.8	2.7	0.5	2.6	3.2	0.8	3.4	3.4	0.8
DPS-D	4.7	1.5	0.8	5.1	3.1	1.0	5.7	3.1	1.3
IIGDM-D	2.2	1.6	3.8	2.9	2.7	0.6	3.3	2.7	0.4
TMPD-D	0.9	0.9	0.6	1.5	1.1	0.9	1.5	1.2	0.9
DTMPD-D	0.9	1.7	0.9	1.4	2.1	0.9	1.4	2.0	1.1
DPS-D	5.2	3.5	2.5	6.9	3.9	1.7	6.8	4.7	0.9
IIGDM-D	1.5	2.3	1.8	1.6	1.4	0.9	2.0	2.0	0.6

by considering 20 randomly selected measurement models (\mathbf{H}) for each setting (d_x, d_y, σ_y) . Figure 2 shows the first two dimensions of the estimated posterior distributions corresponding to the configurations (80, 1) from Table 1 for one of the randomly generated measurement model ($\mathbf{H}, \sigma_y = 0.1$). These illustrations give us insight into the behaviour of the algorithms and their ability to accurately estimate the posterior distribution. We see that TMPD-D estimates the target posterior well compared to IIGDM-D and DPS-D. TMPD-D covers all of the modes, whereas IIGDM-D and DPS-D do not.

We perform the same experiment using 1000 samples of TMPD, DTMPD, DPS and IIGDM, obtained using 1000 Euler-Maruyama time-steps, and results are shown in Appendix E.3.

A direct comparison to Cardoso et al. (2023) using their original experimental setup is shown in Table 5 in Appendix E.3, which shows competitive performance for posterior sampling compared to Sequential Monte-Carlo, an exact sampling method.

6.2 Noisy observation inpainting and super-resolution

We consider inpainting and super-resolution problems on the FFHQ 256×256 (Karras et al., 2019) and CIFAR-10 32×32 (Krizhevsky et al., 2009) datasets. We compare TMPD to IIDGM and DPS. We also compare score-based diffusion models with their denoising-diffusion counterparts (denoted with suffix, -D).

Firstly, we follow the benchmark used by Chung et al. (2022a) and test on a Variance Preserving (VP) SDE, using a DDPM sampler, on FFHQ 256×256 using 1k validation images. The pre-trained diffusion model for FFHQ was taken from Chung et al. (2022a) and was used directly without any finetuning. We follow Chung et al. (2022a) and use various forward operators. For super-resolution, we use a downsampling ratio of 4 ($256 \times 256 \rightarrow 64 \times 64$) and bicubic interpolation; for ‘box’ mask inpainting we mask out 128×128 region and for ‘random’ mask inpainting we choose a random mask for each image masking between 30% and 70% of the pixels. Images are normalized to the range $[0, 1]$ and it is on this scale that we add Gaussian measurement noise with standard deviation $\sigma_y \in \{0.01, 0.05, 0.1, 0.2\}$. For quantitative comparison, we focus on two widely used perception distances, Fréchet Inception Distance (FID) and Learned Perceptual Image Patch Similarity (LPIPS) distance. FID evaluates consistency with the whole dataset by using summary statistics from the FFHQ-50k dataset. We also evaluate observation data similarity using various distances between a sampled image and ground truth image: LPIPS, mean-squared-error (MSE), peak signal-to-noise-ratio (PSNR) and structural similarity index measure (SSIM). For IIDGM-D we use the algorithm and default hyperparameters as described in Song et al. (2023). For DPS-D we use the algorithm in the codebase provided by the authors Chung et al. (2022a) and we use their suggested step-size hyperparameter for this task, and we use their default hyperparameters, such as using static-thresholding (clipping the denoised image at each step to a range $[-1, 1]$) whereas TMPD-D does not require static-thresholding or any hyperparameter tuning. The results for FFHQ sampled using VP DDPM are shown in the appendix Table 6. We observe that TMPD-D is competitive with DPD-D over a range of noise levels, however PiGDM-D is not able to produce high quality reconstructions for larger noise levels.

We note that the heuristics in the DPS and IIDGM implementations have been designed to work with the VP SDE (DDPM sampler), and therefore the performance may not be robust to the choice of SDE. Fig 14 shows an illustration of this when each method is applied to the VE SDE on a sample from the FFHQ validation dataset. We next compare performance to TMPD across VP and VE SDE samplers and a range of noise levels on CIFAR-10 64×64 using 1k validation images.

We use pretrained denoising networks for CIFAR-10 that are available here. For inpainting, we use ‘box’ and ‘half’ mask. For ‘half’-type inpainting, we mask out a 16×16 right half region of the image; for box-type inpainting, we mask out an 8×8 box region following Cardoso et al. (2023). For super-resolution, we use a downsampling ratio of 2 on each axis ($32 \times 32 \rightarrow 16 \times 16$) with a nearest-neighbour downsampling method; and a downsampling ratio of 4 ($32 \times 32 \rightarrow 8 \times 8$) with bicubic downsampling. Images are normalized to the range $[0, 1]$ and it is on this scale that we add Gaussian measurement noise with standard deviation $\sigma_y \in \{0.01, 0.05, 0.1\}$. Whereas no hyperparameters are required for our method, we chose the DPS scale hyperparameter by optimising LPIPS, MSE, PSNR and SSIM on a validation set of 128 images (see Fig. 9 for an example). We found that static thresholding (clipping the denoised image estimate to a range $[0, 1]$ at each sampling step) is critical for the stability and performance of both DPS-D and IIDGM-D. Whilst stability was noted as a limitation in Chung et al. (2022a), and they suggest that devising methods to stabilize the samplers would be a promising direction of research. We find that our TMPD-D method is stable across SDE, noise-level and observation maps, without the need for static thresholding. Whilst for VE IIDGM-D we found the original algorithm in Song et al. (2023) to be stable, for VP IIDGM-D, the original algorithm, whilst stable for FFHQ, was not stable, even with static-thresholding, for CIFAR-10. We chose to bring IIDGM-D a step closer to our algorithm by substituting their likelihood score into an Ancestral sampling algorithm, instead of a DDIM algorithm as suggested in Song et al. (2023), which produced stable samples.

The methods TMPD, IIDGM and DPS all have the same numerical solver of their respective reverse-SDE, and TMPD-D, IIDGM-D and DPS-D all use DDPM since DDIM and DDPM are equivalent algorithms with our chosen DDIM hyperparameter $\eta = 1.0$. Therefore, the sampling methods being compared only differ in the $\nabla_{\mathbf{x}_t} \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t)$ term in their reverse-SDE, and so this experiment allows us to study the

Table 2: Summary of results using the VE and VP-SDE for increasingly noisy $\sigma_y \in \{0.01, 0.05, 0.1\}$ observation inpainting and super-resolution problems on FFHQ 1k validation set. The full results are in Ap. E.4.

SDE		VE SDE		VP SDE	
PROBLEM	METHOD	FID ↓	LPIPS ↓	FID ↓	LPIPS ↓
$\sigma_y = 0.01$	TMPD-D	32.3	0.203 ± 0.039	29.6	0.230 ± 0.034
4× ‘BICUBIC’	DPS-D	47.0	0.273 ± 0.031	31.4	0.234 ± 0.048
SUPER-RESOLUTION	IIGDM-D	37.4	0.244 ± 0.030	29.7	0.198 ± 0.037
$\sigma_y = 0.05$	TMPD-D	32.1	0.268 ± 0.048	32.7	0.304 ± 0.043
4× ‘BICUBIC’	DPS-D	105.9	0.590 ± 0.036	29.3	0.280 ± 0.051
SUPER-RESOLUTION	IIGDM-D	106.8	0.592 ± 0.041	45.1	0.311 ± 0.047
$\sigma_y = 0.1$	TMPD-D	32.7	0.310 ± 0.053	38.0	0.348 ± 0.048
4× ‘BICUBIC’	DPS-D	114.0	0.569 ± 0.044	30.9	0.318 ± 0.051
SUPER-RESOLUTION	IIGDM-D	206.0	0.724 ± 0.034	119.6	0.589 ± 0.047
$\sigma_y = 0.01$	TMPD-D	30.2	0.114 ± 0.029	25.7	0.153 ± 0.033
‘BOX’ MASK	DPS-D	23.9	0.093 ± 0.019	31.5	0.175 ± 0.038
INPAINTING	IIGDM-D	27.1	0.108 ± 0.025	143.8	0.247 ± 0.024
$\sigma_y = 0.05$	TMPD-D	33.6	0.186 ± 0.036	27.0	0.240 ± 0.038
‘BOX’ MASK	DPS-D	39.7	0.318 ± 0.044	30.7	0.228 ± 0.046
INPAINTING	IIGDM-D	49.5	0.354 ± 0.044	159.3	0.448 ± 0.046
$\sigma_y = 0.1$	TMPD-D	34.0	0.223 ± 0.041	29.6	0.292 ± 0.049
‘BOX’ MASK	DPS-D	59.1	0.467 ± 0.053	29.3	0.259 ± 0.049
INPAINTING	IIGDM-D	72.6	0.529 ± 0.047	165.7	0.539 ± 0.083

behaviour of our method on inpainting and super-resolution compared to the different approximations of the smoothed likelihood. A summary of the results for CIFAR-10 sampled using VE DDPM are shown in Table 2. The complete results are in Tables 7 and 10 for VP and VE DDPM respectively, and Tables 8 and 11 for score-based VP and VE SDE, respectively. For more experimental details including illustration of samples used to generate the tables can be found in Appendix E.4. Our method is the only method able to provide high-quality reconstructions independently of the SDE, time discretization or noise level used. On the other hand, we see that DPS-D and IIGDM-D are not able to provide high-quality reconstructions for the VE SDE. For the continuous time methods, IIGDM and DPS are outperformed by TMPD for both VE and VP SDEs in the majority of tasks.

7 Discussions, limitations and future work

In this paper, we introduced TMPD, a diffusion modelling approach to solve inverse problems and sample from conditional distributions using unconditional diffusion models. On various tasks on the VP SDE, TMPD achieves competitive quality with other methods that aim to solve the noisy, linear inverse problem while avoiding the expensive, problem-specific training of conditional models. Our method is more versatile in that it can also be used for the VE SDE, and for large noise and different time discretizations.

TMPD is slower, as each iteration costs more memory and compute due to the Jacobian over the score model. Even with a diagonal and row-sum approximation to the Jacobian, the method is around $1.5\times$ slower than DPS and IIGDM. The row-sum approximation may not be suitable for inverse problems with more complicated or non-linear observation operators, therefore, it would be helpful to explore more efficient sampling techniques, such as ensemble methods or low rank approximations to the Jacobian.

Many state of the art diffusion models operate in a latent space which would make any observation nonlinear, and thus linear observations maps do not suffice in the latent diffusion setting. Furthermore, for inverse problems in general it may be excessive to train a diffusion model for the full data distribution, and it could be difficult to outperform a diffusion model trained for a specific inverse problem. For example, using diffusion models trained specifically for the super-resolution problem in a cascade together with noise conditioning augmentation (Ho et al., 2022) has been extremely effective in progressively generating high-fidelity images, see e.g., Saharia et al. (2022).

A limitation of our method is that without approximations, the full method may be costly to implement in high dimensions of image data. On the positive side, our approach does not require any hyperparameter tuning, does not require static-thresholding of the denoised image, and is more principled compared to existing approaches, as shown in Section 4. We show that our method, unlike DPS and IIGDM, does not fail for cases where the additive Gaussian noise is large. We provided a way to analyse similar methods and our moment approximations can be analysed more rigorously to provide deeper theoretical results for these kinds of methods. Our future work plans include expanding the analysis we provided in this work.

Impact Statement

Our results do not increase or introduce new specific risks, but as a novel way to sample images and generate new data, usual risks of image generation models apply to our work. In particular, the methods in this paper can be used to generate faces of people who do not exist as well as edit existing images in certain ways that creates misinformation risk. However, our method as it is proposed here do not integrate text-based editing, therefore we think these risks are very low for the particular version of this method.

References

- Georgios Batzolis, Jan Stanczuk, Carola-Bibiane Schönlieb, and Christian Etmann. Conditional image generation with score-based diffusion models. *arXiv preprint arXiv:2111.13606*, 2021.
- Christopher M Bishop. *Pattern recognition and machine learning*, volume 4. Springer, 2006.
- Ashish Bora, Ajil Jalal, Eric Price, and Alexandros G Dimakis. Compressed sensing using generative models. In *International conference on machine learning*, pp. 537–546. PMLR, 2017.
- Gabriel Cardoso, Yazid Janati El Idrissi, Sylvain Le Corff, and Eric Moulines. Monte carlo guided diffusion for bayesian linear inverse problems, 2023.
- Sitan Chen, Sinho Chewi, Jerry Li, Yuanzhi Li, Adil Salim, and Anru R Zhang. Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. In *International Conference on Learning Representations*, 2023.
- Hyungjin Chung, Jeongsol Kim, Michael Thompson Mccann, Marc Louis Klasky, and Jong Chul Ye. Diffusion posterior sampling for general noisy inverse problems. In *The Eleventh International Conference on Learning Representations*, 2022a.
- Hyungjin Chung, Byeongsu Sim, Dohoon Ryu, and Jong Chul Ye. Improving diffusion models for inverse problems using manifold constraints. *Advances in Neural Information Processing Systems*, 35:25683–25696, 2022b.
- Hyungjin Chung, Jeongsol Kim, Sehui Kim, and Jong Chul Ye. Parallel diffusion models of operator and image for blind inverse problems. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 6059–6069, 2023.
- Giannis Daras, Yuval Dagan, Alex Dimakis, et al. Score-guided intermediate level optimization: Fast langevin mixing for inverse problems. In *Proceedings of the 39th International Conference on Machine Learning (ICML)*, 2022.
- Prafulla Dhariwal and Alexander Nichol. Diffusion models beat gans on image synthesis. *Advances in neural information processing systems*, 34:8780–8794, 2021.
- Bradley Efron. Tweedie’s formula and selection bias. *Journal of the American Statistical Association*, 106(496):1602–1614, 2011. doi: 10.1198/jasa.2011.tm11181. URL <https://doi.org/10.1198/jasa.2011.tm11181>. PMID: 22505788.
- Berthy T Feng and Katherine L Bouman. Efficient bayesian computational imaging with a surrogate score-based prior. *arXiv preprint arXiv:2309.01949*, 2023.

- Berthy T Feng, Jamie Smith, Michael Rubinstein, Huiwen Chang, Katherine L Bouman, and William T Freeman. Score-based diffusion models as principled priors for inverse imaging. *arXiv preprint arXiv:2304.11751*, 2023.
- Marc Anton Finzi, Anudhyan Boral, Andrew Gordon Wilson, Fei Sha, and Leonardo Zepeda-Núñez. User-defined event sampling and uncertainty quantification in diffusion models for physical dynamical systems. In *International Conference on Machine Learning*, pp. 10136–10152. PMLR, 2023.
- Clark R. Givens and Rae Michael Shortt. A class of Wasserstein metrics for probability distributions. *Michigan Mathematical Journal*, 31(2):231 – 240, 1984. doi: 10.1307/mmj/1029003026. URL <https://doi.org/10.1307/mmj/1029003026>.
- Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models, 2020.
- Jonathan Ho, Chitwan Saharia, William Chan, David J Fleet, Mohammad Norouzi, and Tim Salimans. Cascaded diffusion models for high fidelity image generation. *The Journal of Machine Learning Research*, 23(1):2249–2281, 2022.
- Aapo Hyvärinen. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(24):695–709, 2005. URL <http://jmlr.org/papers/v6/hyvarinen05a.html>.
- Ajil Jalal, Marius Arvinte, Giannis Daras, Eric Price, Alexandros G Dimakis, and Jon Tamir. Robust compressed sensing mri with deep generative priors. *Advances in Neural Information Processing Systems*, 34:14938–14954, 2021.
- Zahra Kadkhodaie and Eero Simoncelli. Stochastic solutions for linear inverse problems using the prior implicit in a denoiser. *Advances in Neural Information Processing Systems*, 34:13242–13254, 2021.
- Ioannis Karatzas and Steven Shreve. *Brownian motion and stochastic calculus*, volume 113. Springer Science & Business Media, 2012.
- Tero Karras, Samuli Laine, and Timo Aila. A style-based generator architecture for generative adversarial networks. In *Proceedings of the IEEE/CVF conference on computer vision and pattern recognition*, pp. 4401–4410, 2019.
- Bahjat Kawar, Gregory Vaksman, and Michael Elad. Snips: Solving noisy inverse problems stochastically. *Advances in Neural Information Processing Systems*, 34:21757–21769, 2021.
- Bahjat Kawar, Michael Elad, Stefano Ermon, and Jiaming Song. Denoising diffusion restoration models. *Advances in Neural Information Processing Systems*, 35:23593–23606, 2022.
- Alex Krizhevsky, Geoffrey Hinton, et al. Learning multiple layers of features from tiny images. 2009.
- Rémi Laumont, Valentin De Bortoli, Andrés Almansa, Julie Delon, Alain Durmus, and Marcelo Pereyra. Bayesian imaging using plug & play priors: when Langevin meets Tweedie. *SIAM Journal on Imaging Sciences*, 15(2):701–737, 2022.
- Morteza Mardani, Jiaming Song, Jan Kautz, and Arash Vahdat. A variational perspective on solving inverse problems with diffusion models. *arXiv preprint arXiv:2305.04391*, 2023.
- Chenlin Meng, Yang Song, Wenzhe Li, and Stefano Ermon. Estimating high order gradients of the data distribution by denoising. *Advances in Neural Information Processing Systems*, 34:25359–25369, 2021.
- Shakir Mohamed, Mihaela Rosca, Michael Figurnov, and Andriy Mnih. Monte carlo gradient estimation in machine learning. *The Journal of Machine Learning Research*, 21(1):5183–5244, 2020.
- Jakiw Pidstrigach, Youssef Marzouk, Sebastian Reich, and Sven Wang. Infinite-dimensional diffusion models for function spaces. *arXiv preprint arXiv:2302.10130*, 2023.

Herbert E. Robbins. *An Empirical Bayes Approach to Statistics*, pp. 388–394. Springer New York, New York, NY, 1992. ISBN 978-1-4612-0919-5. doi: 10.1007/978-1-4612-0919-5_26. URL https://doi.org/10.1007/978-1-4612-0919-5_26.

Chitwan Saharia, William Chan, Saurabh Saxena, Lala Li, Jay Whang, Emily L Denton, Kamyar Ghasemipour, Raphael Gontijo Lopes, Burcu Karagol Ayan, Tim Salimans, et al. Photorealistic text-to-image diffusion models with deep language understanding. *Advances in Neural Information Processing Systems*, 35:36479–36494, 2022.

Jiaming Song, Chenlin Meng, and Stefano Ermon. Denoising diffusion implicit models. 2021a. URL <https://openreview.net/forum?id=St1giarCHLP>.

Jiaming Song, Arash Vahdat, Morteza Mardani, and Jan Kautz. Pseudoinverse-guided diffusion models for inverse problems. In *International Conference on Learning Representations*, 2023. URL https://openreview.net/forum?id=9_gsMA8MRKQ.

Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. *arXiv preprint arXiv:2011.13456*, 2020.

Yang Song, Liye Shen, Lei Xing, and Stefano Ermon. Solving inverse problems in medical imaging with score-based generative models. *arXiv preprint arXiv:2111.08005*, 2021b.

Tristan SW Stevens, Jean-Luc Robert, Faik C Yu, Jun Seob Shin, and Ruud JG van Sloun. Removing structured noise with diffusion models. *arXiv preprint arXiv:2302.05290*, 2023.

Andrew M Stuart. Inverse problems: a bayesian perspective. *Acta numerica*, 19:451–559, 2010.

Albert Tarantola. *Inverse problem theory and methods for model parameter estimation*. SIAM, 2005.

A Proofs for Section 3

A.1 Proof of Proposition 1

In order to prove this result, we adapt Theorem 1 of Meng et al. (2021). We write the proof for generic exponential family which can be adapted to our case easily. Let us consider

$$p_{t|0}(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_t; \sqrt{\alpha_t}\mathbf{x}_0, v_t\mathbf{I}_{d_x}).$$

and write $p_t(\mathbf{x}_t) = \int p_{t|0}(\mathbf{x}_t|\mathbf{x}_0)p_0(\mathbf{x}_0)d\mathbf{x}_0$. We are interested in finding the moments of the posterior

$$p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) = \frac{p_{t|0}(\mathbf{x}_t|\mathbf{x}_0)p_0(\mathbf{x}_0)}{p_t(\mathbf{x}_t)}.$$

Let us redefine the right handside in this Bayes’ rule using the exponential family parameterisation of the Gaussian $p_{t|0}(\mathbf{x}_t|\mathbf{x}_0)$

$$p(\boldsymbol{\eta}_0|\mathbf{x}_t) = \frac{p_{t|0}(\mathbf{x}_t|\boldsymbol{\eta}_0)q_0(\boldsymbol{\eta}_0)}{p_t(\mathbf{x}_t)},$$

where $\boldsymbol{\eta}_0 = \mathbf{x}_0\sqrt{\alpha_t}/v_t$ and

$$p_{t|0}(\mathbf{x}_t|\boldsymbol{\eta}_0) = e^{\boldsymbol{\eta}_0^\top \mathbf{x}_t - \psi(\boldsymbol{\eta}_0)} q(\mathbf{x}_t),$$

where

$$q(\mathbf{x}_t) = ((2\pi)^d v^{2d})^{-1/2} e^{-\mathbf{x}_t^\top \mathbf{x}_t / 2v_t}$$

Now let

$$\lambda(\mathbf{x}_t) = \log \frac{p_t(\mathbf{x}_t)}{q(\mathbf{x}_t)},$$

we can rewrite

$$p(\boldsymbol{\eta}_0|\mathbf{x}_t) = e^{\boldsymbol{\eta}_0^\top \mathbf{x}_t - \psi(\mathbf{x}_t) - \lambda(\mathbf{x}_t)} q_0(\boldsymbol{\eta}_0).$$

It is then easy to show that the moments of $\boldsymbol{\eta}_0$ are given by (Meng et al., 2021)

$$\begin{aligned}\mathbb{E}[\boldsymbol{\eta}_0|\mathbf{x}_t] &= \mathbf{J}_\lambda(\mathbf{x}_t), \\ \mathbb{E}[\boldsymbol{\eta}_0^\top|\mathbf{x}_t] &= \mathbf{J}_\lambda(\mathbf{x}_t)^\top, \\ \mathbb{E}[\boldsymbol{\eta}_0\boldsymbol{\eta}_0^\top|\mathbf{x}_t] &= \mathbf{S}_\lambda(\mathbf{x}_t) + \mathbf{J}_\lambda(\mathbf{x}_t)\mathbf{J}_\lambda(\mathbf{x}_t)^\top,\end{aligned}$$

where \mathbf{J}_λ is the Jacobian of λ w.r.t. \mathbf{x}_t and \mathbf{S}_λ is the Hessian of λ w.r.t. \mathbf{x}_t . Recall now that $\boldsymbol{\eta}_0 = \mathbf{x}_0\sqrt{\alpha_t}/v_t$ and

$$\begin{aligned}\lambda(\mathbf{x}_t) &= \log p_t(\mathbf{x}_t) + \frac{\mathbf{x}_t^\top \mathbf{x}_t}{2v_t} + C \\ \mathbf{J}_\lambda(\mathbf{x}_t) &= \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) + \frac{\mathbf{x}_t}{v_t} \\ \mathbf{S}_\lambda(\mathbf{x}_t) &= \nabla_{\mathbf{x}_t}^2 \log p_t(\mathbf{x}_t) + \frac{1}{v_t} \mathbf{I}_{d_x}.\end{aligned}$$

This implies that

$$\mathbb{E}[\mathbf{x}_0|\mathbf{x}_t] = \frac{1}{\sqrt{\alpha_t}}(\mathbf{x}_t + v_t \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)),$$

which proves the Tweedie's formula for the mean. For the covariance, note that

$$\begin{aligned}\text{Cov}(\boldsymbol{\eta}_0|\mathbf{x}_t) &= \mathbb{E}[\boldsymbol{\eta}_0\boldsymbol{\eta}_0^\top|\mathbf{x}_t] - \mathbb{E}[\boldsymbol{\eta}_0|\mathbf{x}_t]\mathbb{E}[\boldsymbol{\eta}_0|\mathbf{x}_t]^\top \\ &= \nabla^2 \log p_t(\mathbf{x}_t) + \frac{1}{v_t} \mathbf{I}_{d_x}.\end{aligned}$$

Since

$$\text{Cov}(\boldsymbol{\eta}_0|\mathbf{x}_t) = \frac{\alpha_t}{v_t^2} \text{Cov}(\mathbf{x}_0|\mathbf{x}_t),$$

we conclude

$$\text{Cov}(\mathbf{x}_0|\mathbf{x}_t) = \frac{v_t}{\alpha_t} (\mathbf{I}_{d_x} + v_t \nabla^2 \log p_t(\mathbf{x}_t)),$$

which concludes the proof. \square

B Proofs for Section 4

B.1 Proof of Proposition 3

If p_{data} is Gaussian, then the full process \mathbf{x}_t is a Gaussian process. In particular, $(\mathbf{x}_0, \mathbf{x}_t)$ are jointly Gaussian:

$$\mathbf{x}_0|\mathbf{x}_t \propto \mathcal{N}(\mathbb{E}[\mathbf{x}_0|\mathbf{x}_t], \mathbb{V}[\mathbf{x}_0|\mathbf{x}_t]),$$

where \mathbb{V} denotes the covariance. However, the right-hand side is precisely the approximation we make, due to Proposition 1. Therefore, the approximation we make in equation 9 is correct. Adding this to the learned drift $\mathbf{s}_\theta(\mathbf{x}_t, t) = \nabla \log p_t(\mathbf{x}_t)$, we get an expression for $\nabla \log p_t(\mathbf{x}_t|\mathbf{y})$ by equation 4.

B.2 Proof of Theorem 1

We first introduce three SDEs.

Conditioned SDEs : The first pair of SDEs is given by an OU process, but started in the conditional distribution:

$$d\mathbf{x}_t^c = -\frac{1}{2}\mathbf{x}_t^c dt + d\mathbf{w}_t, \quad \mathbf{x}_0^c \sim p_0^c := p_{\text{data}}(\mathbf{x}_0|\mathbf{y}). \quad (15)$$

We denote the marginals of \mathbf{x}_t^c by p_t^c to differentiate them from the marginals p_t of the OU process started in the correct distribution, defined in equation 2. Note that $p_t^c = p_t(\mathbf{x}_t|\mathbf{y})$. The time reversal $\mathbf{z}_t = \mathbf{x}_{T-t}$ then satisfies the SDE

$$\begin{aligned} d\mathbf{z}_t &= \frac{1}{2}\mathbf{z}_t dt + \nabla_{\mathbf{z}_t} \log p_{T-t}^c(\mathbf{z}_t) dt + d\mathbf{w}_t \\ &= \frac{1}{2}\mathbf{z}_t dt + \nabla_{\mathbf{z}_t} \log p_{T-t}(\mathbf{z}_t) dt + \nabla_{\mathbf{z}_t} \log p_{\mathbf{y}|T-t}(\mathbf{y}|\mathbf{z}_t) dt + d\mathbf{w}_t \\ \mathbf{z}_0 &\sim p_T^c, \end{aligned} \quad (16)$$

where we used equation 4.

Solutions to the above reverse SDE will sample our target measure $p_0^c = p_{\text{data}}(\mathbf{x}_0|\mathbf{y})$ at final time. Therefore, we want to study how our algorithm approximates solutions of \mathbf{x}^c .

Intermediate Gaussian SDE: Instead of bounding the distance of solutions to the above conditioned reverse SDE to our algorithm, we will instead introduce an intermediate process, which we will later use in a triangle inequality. The process will be a Gaussian process. Therefore, we denote it with a superscript G .

The process \mathbf{x}_t^G will be defined analogous to equation 2, but assuming that it is started in $\mathcal{N}(\mathbf{x}_0; \boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0)$ instead of p_{data} . Since the forward SDE in equation 2 is linear, all of the marginals of \mathbf{x}_t^G , which we denote by p_t^G , will also be Gaussian.

Again, we define a conditioned version of \mathbf{x} , called $\mathbf{x}^{G,c}$ analogous to equation 15, i.e. $\mathbf{x}_0^{G,c}$ will be distributed as \mathbf{x}_0^G conditioned on $\mathbf{y} = y$, i.e. $p^G(\mathbf{x}_0|\mathbf{y} = y)$. Since we have a linear observation model, $p^G(\mathbf{x}_0|\mathbf{y} = y)$ is still a Gaussian, and therefore $\mathbf{x}^{G,c}$ will also be a Gaussian process. Its reverse SDE $\mathbf{z}^{G,c}$ is defined through equation 16, just that every appearance of p will also have a superscript G ,

$$\begin{aligned} d\mathbf{z}_t &= \frac{1}{2}\mathbf{z}_t dt + \nabla_{\mathbf{z}_t} \log p_{T-t}^G(\mathbf{z}_t) dt + \nabla_{\mathbf{z}_t} \log p_{T-t}^G(\mathbf{y}|\mathbf{z}_t) dt + d\mathbf{w}_t \\ \mathbf{z}_0 &\sim p_T^c, \end{aligned} \quad (17)$$

Algorithm SDE: Finally, we define a third reverse SDE, which is the SDE that we are discretizing when implementing our algorithm. It is given by

$$\begin{aligned} d\mathbf{z}_t &= \frac{1}{2}\mathbf{z}_t dt + \nabla_{\mathbf{z}_t} \log p_{T-t}(\mathbf{z}_t) dt + f_{T-t}(\mathbf{z}_t) dt + d\mathbf{w}_t \\ \mathbf{z}_0 &\sim p_T^c, \end{aligned} \quad (18)$$

where

$$f_t(\mathbf{z}_t) = \nabla_{\mathbf{z}_t} \mathbb{E}[\mathbf{x}_0|\mathbf{x}_t = \mathbf{z}_t] \mathbf{H}^T (\mathbf{H} \mathbb{V}[\mathbf{x}_0|\mathbf{x}_t = \mathbf{z}_t] \mathbf{H}^T + \sigma_y^2 \mathbf{I})^{-1} (\mathbf{y} - \mathbf{H} \mathbb{E}[\mathbf{x}_0|\mathbf{x}_t = \mathbf{z}_t]) \quad (19)$$

and \mathbb{V} denotes the conditional covariance. Except for approximation errors due to time discretization and in the initial conditions, this is the SDE we are sampling from in our algorithm. We denote the marginal of \mathbf{z}_t by q_t . Now we are ready to write our proof.

Proof of Theorem 1. We have that $p_{\text{data}}(\cdot|\mathbf{y}) = p_0^c$

$$\|p_0^c - q_T\|_{\text{TV}} \leq \|p_0^c - p_0^{G,c}\|_{\text{TV}} + \|p_0^{G,c} - q_T\|_{\text{TV}}$$

We will bound the second term using Pinsker's inequality to get a KL term on the path space, as done in Chen et al. (2023). The proof then consists in bounding the first total variation term and the resulting KL term.

Bounding the first term: The first term is the total variation distance between $p_0(\mathbf{x}_0|\mathbf{y} = y)$ and $p_0^G(\mathbf{x}_0^G|\mathbf{y} = y)$. Now,

$$\begin{aligned} p_0(\mathbf{x}_0 = x_0|\mathbf{y} = y) &= \frac{p_{\mathbf{y}|0}(\mathbf{y} = y|\mathbf{x}_0 = x_0)p_0(\mathbf{x}_0 = x_0)}{p_{\mathbf{y}}(\mathbf{y} = y)} \\ &= \frac{p_{\mathbf{y}|0}^G(\mathbf{y} = y|\mathbf{x}_0 = x_0)p_0^G(\mathbf{x}_0 = x_0)}{p_{\mathbf{y}}^G(\mathbf{y} = y)} \frac{p_0(\mathbf{x}_0 = x_0)}{p_0^G(\mathbf{x}_0^G = x_0^G)} \frac{p_{\mathbf{y}}(\mathbf{y} = y)}{p_{\mathbf{y}}(\mathbf{y} = y)} \\ &= p_{\mathbf{y}|0}^G(\mathbf{x}_0 = x_0|\mathbf{y} = y) \exp(\Phi(x_0)) \frac{p_{\mathbf{y}}^G(\mathbf{y} = y)}{p_{\mathbf{y}}(\mathbf{y} = y)}, \end{aligned}$$

where we used that the conditional distribution of \mathbf{y} given \mathbf{x}_0 does not depend on the distribution of \mathbf{x}_0 . For the last term we get

$$\begin{aligned} p_{\mathbf{y}}(\mathbf{y} = y) &= \int p_{\mathbf{y}|0}(\mathbf{y} = y|\mathbf{x}_0 = x_0)p_0(\mathbf{x}_0 = x_0)dx_0 \\ &= \int p_{\mathbf{y}|0}^G(\mathbf{y} = y|\mathbf{x}_0 = x_0)p_0^G(\mathbf{x}_0 = x_0)\exp(\Phi(x_0))dx_0 \\ &= Np_{\mathbf{y}}^G(\mathbf{y} = y), \end{aligned}$$

with $N \in [1/M, M]$. Therefore,

$$\|p_0^c - p_0^{G,c}\|_{\text{TV}} \leq \int \left| \frac{p_0^c}{p_0^{G,c}}(x_0) - 1 \right| p_0^{G,c}(x_0) dx_0 \leq M^2 - 1,$$

where we used that $M - 1$ is always greater or equal to $1 - \frac{1}{M}$, since $M \geq 1$.

Bounding the second term: For the second term we get that $p_0^{G,c}$ is the final marginal of equation 17, while q_T is the final marginal of equation 18. We now use Pinsker's inequality,

$$\|p_0^{G,c} - q_T\|_{\text{TV}} \leq \sqrt{\text{KL}(p_0^{G,c} \| q_T)}.$$

We denote the path measures induced by equation 17 and equation 18 by $\mathbb{P}^{G,c}$ and \mathbb{Q} respectively. They have final time marginals $p_0^{G,c}$ and q_T . Therefore, we can bound the KL-Divergence of the marginals $p_0^{G,c}$ and q_T by the KL-Divergence on the full path space:

$$\text{KL}(p_0^{G,c} \| q_T) \leq \text{KL}(\mathbb{P}^{G,c} \| \mathbb{Q}).$$

We will assume that we can apply Girsanov's theorem and show later that this is justified. Using Girsanov's theorem, we can evaluate the Radon-Nikodym derivative $\frac{d\mathbb{Q}}{d\mathbb{P}^{G,c}}$ on the path space and therefore calculate the KL-divergence:

$$\begin{aligned} \text{KL}(\mathbb{P}^{G,c} \| \mathbb{Q}) &= \mathbb{E}_{\mathbb{P}^{G,c}} \left[\log \frac{d\mathbb{P}^{G,c}}{d\mathbb{Q}} \right] \\ &= \mathbb{E}_{\mathbb{P}^{G,c}} \left[- \int_0^t \nabla \log p_t(x_t) + f_{T-t}(\mathbf{x}_t) - \nabla \log p_{T-t}^{G,c}(\mathbf{x}_t) dw_t \right] \\ &\quad + \mathbb{E}_{\mathbb{P}^{G,c}} \left[\int_0^t \|\nabla \log p_t(x_t) + f_{T-t}(\mathbf{x}_t) - \nabla \log p_{T-t}^{G,c}(\mathbf{x}_t)\|^2 dt \right] \\ &= \int_0^t \mathbb{E}_{p_t^{G,c}} [\|\nabla \log p_t(x_t) + f_{T-t}(\mathbf{x}_t) - \nabla \log p_{T-t}^{G,c}(\mathbf{x}_t)\|^2] dt, \end{aligned}$$

where the term on the second line drops out because stochastic integrals have expectation 0. Now we have the drift of the Gaussian SDE given by

$$\begin{aligned}
& \nabla \log p_t^{G,c}(x_t) \\
&= \nabla \log p_t^G(x_t) + \nabla \log p^G(\mathbf{y}|\mathbf{x}_t^G = x_t) \\
&= \nabla \log p_t^G(x_t) \\
&+ \nabla_{x_t} \mathbb{E}[\mathbf{x}_0^G|\mathbf{x}_t^G = x_t] \mathbf{H}^\top (\mathbf{H} \mathbb{V}[\mathbf{x}_0^G|\mathbf{x}_t^G = x_t] \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H} \mathbb{E}[\mathbf{x}_0^G|\mathbf{x}_t^G = x_t]) \\
&=: \nabla \log p_t^G(x_t) + \tilde{f}_t(x_t),
\end{aligned} \tag{20}$$

see equation 43, while the drift of the algorithm SDE is given by

$$\nabla \log p_t(x_t) + f_t(x_t), \tag{21}$$

where f_t is given by equation 19. We see that the difference in the SDE drifts mainly consists of differences between conditional moments of p_t and p_t^G , as well as the derivatives of the conditional expectations. Therefore, the main difficulty of the proof is to bound these.

The density of the conditional distribution of $p_{0|t}$ is given by

$$\begin{aligned}
p_{0|t}(x_0|x_t) &= \frac{p_{0,t}(x_0, x_t)}{p_t(x_t)} = \frac{p_{0,t}^G(x_0, x_t) \exp(\Phi(x_0))}{p(x_t)} = \frac{p_{0|t}^G(x_0|x_t) p^G(x_t) \exp(\Phi(x_0))}{p(x_t)} \\
&= p_{0|t}^G \exp(\Phi(x_0) - \Phi(x_t)),
\end{aligned} \tag{22}$$

where

$$\exp(\Phi_t(x_t)) = \frac{p_t(x_t)}{p_t^G(x_t)} = \mathbb{E}[\exp(\Phi(\mathbf{x}_0))|\mathbf{x}_t = x_t]. \tag{23}$$

In the last equality, we used that

$$\frac{d\mathbb{P}}{d\mathbb{P}^G}(x_{[0,t]}) = \exp(\Phi(x_0)),$$

where we denoted by \mathbb{P} the path measure induced by equation 15. Therefore, also their marginals $p_t = \mathbb{P}_t^G$ and $p_t = \mathbb{P}_t$ have relative densities, which are given by integrating out the density to time t , as we did in equation 23.

By assumption $\exp(\Phi)$ is bounded from above and below by M and $1/M$ respectively, and by equation 23 the same holds for $\exp(\Phi_t)$. Therefore, by equation 22, $p_{t|0}$ is absolutely continuous with respect to $p_{t|0}^G$ with a density that is bounded from above and below by M^2 and $1/M^2$ respectively. We now obtain

$$\begin{aligned}
& |\mathbb{E}[\mathbf{x}_0|\mathbf{x}_t = x_t] - \mathbb{E}[\mathbf{x}_0^G|\mathbf{x}_t^G = x_t]| \\
&= \left| \int x_0 [p(x_0|x_t) - p^G(x_0|x_t)] dx_0 \right| = \left| \int x_0 p^G(x_0|x_t) (\exp(\Phi(x_0) - \Phi(x_t)) - 1) dx_0 \right| \\
&\leq (M^2 - 1) \mathbb{E}[|\mathbf{x}_0^G| | \mathbf{x}_t^G = x_t^G]
\end{aligned}$$

The same holds for every entry of the covariance matrix. We denote

$$\begin{aligned}
V_{ij} &:= \mathbb{V}[\mathbf{x}_0|\mathbf{x}_t]_{ij}, \\
V_{ij}^G &:= \mathbb{V}[\mathbf{x}_0^G|\mathbf{x}_t^G]_{ij} = N_{ij}, \\
|V_{ij} - V_{ij}^G| &\leq (M^2 - 1) N_{ij} \\
N_{ij} &:= \mathbb{E}[|\mathbf{x}_{0,i}^G \mathbf{x}_{0,j}^G| | \mathbf{x}_t^G]
\end{aligned}$$

Now we get that

$$\|V - V^G\|_F = \left(\sum_{ij} (V_{ij} - V_{ij}^G)^2 \right)^{1/2} \leq (M^2 - 1) \|V^G\|_F \max_{ij} (N_{ij}).$$

We can bound N_{ij} by

$$N_{ij} \leq \mathbb{E}[(\mathbf{x}_{0,i}^G)^2 + (\mathbf{x}_{0,j}^G)^2 | \mathbf{x}_t^G] \leq \text{Tr}(\mathbb{V}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t]). \quad (24)$$

The latter term does not actually depend on x_t , but only on t . By using the formulas in equation 39 we see that it can be bounded independently of t (only depending on m_0 and Σ_0). Therefore, we can put it into a constant. We get that

$$\|V - V^G\|_F \lesssim (M^2 - 1)\|V^G\|_F,$$

where

$$a \lesssim b \quad \Leftrightarrow \quad a \leq Cb,$$

with a constant C only depending on m_0 , Σ_0 , H , σ^y and the observation y . Now let v be a vector with $\|v\| = 1$. It holds that

$$\begin{aligned} v^T V_{ij} v &= \int (v^T (x_0 - \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]))^2 p_{0|t}^G(x_0 | x_t) \exp(\Phi(x_0) - \Phi_t(x_t)) dx_0 \\ &= \frac{1}{M} v^T V_{ij}^G v + N \int (v^T (\mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t] - \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]))^2 p_{0|t}(x_0 | x_t) dx_0 \\ &\geq v^T V_{ij}^G v \frac{1}{M} \end{aligned} \quad (25)$$

with $N \geq 1/M$. Since V and V^G are positive semidefinite, symmetric matrices, this implies that all eigenvalues of V are bounded by the lowest eigenvalue of V^G times $1/M$. We define

$$T_1 = HV^G H^T + \sigma_y^2 I_{d_y} \quad (26)$$

$$T_2 = HVH^T + \sigma_y^2 I_{d_y}. \quad (27)$$

We then need to bound

$$\|T_1^{-1} - T_2^{-1}\| \leq \|T_1 - T_2\|(\|T_1^{-1}\| + \|T_2^{-1}\|).$$

We start with

$$\|T_1 - T_2\|_{\text{op}} \leq \|H\|^2 \|V - V^G\|_{\text{op}} \lesssim \|V - V^G\|_F \lesssim (M^2 - 1)\|V^G\|_F.$$

We also used the equivalence of the Frobenius and operator norm here. Due to our eigenvalue bound (equation 25) on V , we also get an analogous bound on $V + \sigma_y I_{d_y}$. Therefore,

$$\|T_1^{-1} - T_2^{-1}\|_{\text{op}} \lesssim (M^2 - 1)\|V^G\|_F \|V^G\|_F^{-1} (1 + M) \leq (M^2 - 1)(M + 1) \leq M^3 - 1$$

where we used that the operator norm of the inverse, is the inverse of the operator norm and the equivalence of the operator norm to the Frobenius norm.

Finally, we need to bound

$$\begin{aligned} &\nabla_{x_t} \int x_0 p(x_0 | x_t) dx_0 \\ &= \int x_0 \nabla_{x_t} p^G(x_0 | x_t) \exp(\Phi(x_0) - \Phi(x_t)) dx_0 \\ &= \int x_0 \exp(\Phi(x_0) - \Phi(x_t)) \nabla_{x_t} p^G(x_0 | x_t) dx_0 + \int x_0 \nabla_{x_t} \Phi(x_t) \exp(\Phi(x_0) - \Phi(x_t)) p^G(x_0 | x_t) dx_0 \\ &= \int x_0 \exp(\Phi(x_0) - \Phi(x_t)) \nabla_{x_t} p^G(x_0 | x_t) dx_0 + \nabla_{x_t} \Phi(x_t) \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]. \end{aligned} \quad (28)$$

Also

$$\begin{aligned}
& \left\| \int x_0 \exp(\Phi(x_0) - \Phi(x_t)) \nabla_{x_t} p^G(x_0|x_t) dx_0 - \nabla_{x_t} \mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t] \right\| \\
&= \left\| \int \nabla_{x_t} p^G(x_0|x_t) x_0^T (\exp(\Phi(x_0) - \Phi(x_t)) - 1) dx_0 \right\| \\
&\leq (M^2 - 1) \int p^G(x_0|x_t) \|\nabla \log p^G(x_0|x_t) x_0^T\| dx_0 \\
&\leq (M^2 - 1) \left(\int p^G(x_0|x_t) \|\nabla \log p^G(x_0|x_t) x_0^T\|^2 dx_0 \right)^{1/2} \\
&\leq (M^2 - 1) \left(\int p^G(x_0|x_t) x_0^T \Sigma_{0|t}^{-1} (x_0 - m_{0|t}^{x_t}) dx_0 \right)^{1/2} \\
&\lesssim (M^2 - 1) \left(\int p^G(x_0|x_t) (x_0 - m_{0|t}^{x_t})^T \Sigma_{0|t}^{-1} (x_0 - m_{0|t}^{x_t}) dx_0 \right)^{1/2} \lesssim \|x_t\| (M^2 - 1) \sqrt{d_x} \\
&\lesssim \|x_t\| (M^2 - 1),
\end{aligned} \tag{29}$$

here we used that we can upper bound the operator norm of a positive semidefinite matrix by its trace from the third to the fourth line. We denoted by $m_{0|t}^{x_t}$ and $\Sigma_{0|t}$ the mean and covariance of $p^G(\mathbf{x}_0^G | \mathbf{x}_t^G = x_t)$. From the second to last to the last line we used that the $m_{0|t}^{x_t}$ depends on x_t linearly, and the magnitude of the linear dependence can be bounded uniformly in t (see equation 39). The integral in the last line is the variance of a standard normal random variable, which evaluates to d_x . Furthermore,

$$\nabla \Phi_t(x_t) = \mathbb{E}[\nabla \Phi_0(\mathbf{x}_0) | \mathbf{x}_t = x_t] \leq L,$$

see for example the Proof of Theorem 1 in Pidstrigach et al. (2023). Therefore,

$$\begin{aligned}
& \|\nabla_{x_t} \mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t] - \nabla_{x_t} \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]\| \\
&\lesssim \|x_t\| (M^2 - 1) + L \|\mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]\| \\
&\leq \|x_t\| (M^2 - 1) + L (\|\mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]\| + \|\mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t]\|) \\
&\lesssim \|x_t\| (M^2 - 1) + L (M^2 - 1 + \|x_t\|) = \|x_t\| (M^2 - 1 + L) + (M^2 - 1)L =: B_{x_t}
\end{aligned}$$

Furthermore, from equation 28 we see that

$$\|\nabla_{x_t} \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = x_t]\| \leq \|x_t\| (M^2 - 1) + L (M^2 - 1 + \|x_t\|) = B_{x_t}$$

too. We now subtract the full drifts in equation 20 and equation 21 and arrive at

$$\begin{aligned}
& \|\nabla \tilde{f}(x_t) - f_t(x_t)\| \\
&\leq B_{x_t} \times \|T_1^{-1}(y - H \mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t])\| \\
&\quad + B_{x_t} \|T_1^{-1} - T_2^{-1}\|_{\text{op}} \|y - H \mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = x_t]\| \\
&\quad + B_{x_t} \|T_2^{-1}\|_{\text{op}} \|H \mathbb{E}[\mathbf{x}_0^G | \mathbf{x}_t^G = z_t] - H \mathbb{E}[\mathbf{x}_0 | \mathbf{x}_t = z_t]\| \\
&\lesssim B_{x_t} + B_{x_t} (M^3 - 1) + B_{x_t} M (M^2 - 1) \\
&\leq B_{x_t} (M^3 - 1) = (M^5 - 1)(\|x_t\| + L) + (M^3 - 1)L \|x_t\|.
\end{aligned}$$

Furthermore, we get that

$$\|\nabla \log p_t^G(x_t) - \nabla \log p_t(x_t)\| \leq \|\nabla \log \frac{p_t^G(x_t)}{p_t(x_t)}\| \leq \|\nabla_{x_t} \Phi_t(x_t)\| \leq L.$$

Plugging in equation 20, we get that

$$\|\nabla \log p_t(x_t) + f_t(x_t) - p_t^{G,c}(x_t)\| \leq (M^5 - 1)(\|x_t\| + L) + (M^3 - 1)L \|x_t\| + L.$$

Using this expression, we see that the drift change is linear in $\|x_t\|$. We now want to apply an iterated version of Novikov's condition to show that our application of Girsanov's Theorem is justified. To that end,

we follow the argument Karatzas & Shreve (2012, Corollary 3.5.16). We repeat it here for completeness. We see that

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}^{G,c}} \left[\exp \left(\frac{1}{2} \int_{t_i}^{t_{i+1}} \|\nabla \log p_t(x_t) + f_t(x_t) - \nabla \log p_t^{G,c}(x_t)\|^2 dt \right) \right] \\ & \leq \mathbb{E}_{\mathbb{P}^{G,c}} \left[\exp \left(c \int_{t_i}^{t_{i+1}} 1 + \|x_t\|^2 dt \right) \right]. \end{aligned}$$

Since the expectation is regarding a Gaussian random variable x_t , we can make it finite as long as we pick $\Delta_i = t_{i+1} - t_i$ small enough. By setting $t_0 = 0$ and $t_1 > 0$, we can show equivalence on $[t_0, t_1]$. We can then iterate this procedure, to get equivalence on $[t_1, t_2]$ and so on. Since the lowest and highest eigenvalues of Σ_t are bounded from below and above respectively, and m_t is also bounded, the Δ_i can be bounded from below. Therefore, we get equivalence on $[0, T]$ this way in at most $\lceil T/\Delta \rceil$ steps.

Furthermore, we see that the likelihood KL-divergence can be bounded by

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}^{G,c}} \left[\frac{1}{2} \int_0^T \|\nabla \log p_t(x_t) + f_t(x_t) - \nabla \log p_t^{G,c}(x_t)\|^2 dt \right] \\ & \lesssim T((M^5 - 1)(L + 1) + (M^3 - 1)L + L) \leq T((M^5 - 1)(L + 1) + L), \end{aligned}$$

where we used that the mean and covariance of the Gaussian process x_t (under $\mathbb{P}^{G,c}$) can be bounded by a constant only depending on y , Σ_t and m_t . Taking the square root proves our theorem. \square

C Variance Exploding SDE (time-rescaled Brownian motion)

The second main type of SDE used in denoising-diffusion is the Variance Exploding (Song et al., 2020) or time-rescaled Brownian motion, is

$$d\mathbf{x}_t = \sqrt{\frac{dv(t)}{dt}} d\mathbf{w}_t, \quad \mathbf{x}_0 \sim p_0 = p_{\text{data}} \quad (30)$$

where $(\mathbf{w}_t)_{t \in [0, T]}$ is a Brownian motion and $v : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is an increasing function where $v(0) = 0$. The transition kernel of the time-rescaled Brownian motion is

$$p_{t|0}(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_0, v_t \mathbf{I}_{d_x}),$$

denoting $v(t) = v_t$. The signal \mathbf{x}_0 is sampled from a second SDE, the reverse process, given in forward time as

$$d\mathbf{x}_t = -\nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) dt - \sqrt{\frac{dv_t}{dt}} d\bar{\mathbf{w}}_t, \quad \mathbf{x}_T \sim p_T.$$

D Tweedie's formula

In this section, we give an alternative derivation of the Tweedie's formula for VPSDEs and VESDEs.

Tweedie's formula (Robbins, 1992; Efron, 2011) gives the minimum mean squared error (MMSE) estimator of $\mathbf{x}_0 | \mathbf{x}_t$ when $\mathbf{x}_t | \mathbf{x}_0$ is Gaussian and the score $\nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$ is available. Tweedie's formula applied to the time-rescaled SDEs is given below.

D.1 Variance Preserving SDE (time-rescaled Ornstein-Uhlenbeck process)

Using the Variance Preserving SDE or time-rescaled Ornstein-Uhlenbeck process in equation 2, given the random variable

$$\sqrt{\frac{\alpha_t}{v_t}} \mathbf{x}_0 := \mathbf{v}_0 \sim p_{\mathbf{v}_0},$$

where, we now define $p_{\mathbf{v}_0}(\mathbf{v}_0)$ as the probability density of the random variable \mathbf{v}_0 evaluated at the realization \mathbf{v}_0 , then the random variable

$$\frac{\mathbf{x}_t}{\sqrt{v_t}} := \mathbf{v}_t \sim \mathcal{N}(\mathbf{v}_0, \mathbf{I}_{d_x}),$$

has a marginal probability density evaluated at the realization \mathbf{v}_t is the convolution of the random variable \mathbf{v}_0 with the Gaussian kernel

$$p_{\mathbf{v}_t}(\mathbf{v}_t) = \int \phi(\mathbf{v}_t - \mathbf{v}_0) p_{\mathbf{v}_0}(\mathbf{v}_0) d\mathbf{v}_0.$$

Then, Tweedie's formula (Robbins, 1992; Efron, 2011) gives

$$\mathbb{E}_{\mathbf{v}_0 \sim p_{\mathbf{v}_0} | \mathbf{v}_t}[\mathbf{v}_0] = \mathbf{v}_t + \nabla_{\mathbf{v}_t} \log p_{\mathbf{v}_t}(\mathbf{v}_t)$$

From continuous change of random variables, $p_{\mathbf{v}_t}(\mathbf{v}_t) = p_{\mathbf{x}_t}(\mathbf{v}_t \sqrt{v_t}) \left| \frac{d\mathbf{x}_t}{d\mathbf{v}_t} \right| = p_{\mathbf{x}_t}(\mathbf{x}_t) \sqrt{v_t}$, giving $\log p_{\mathbf{v}_t}(\mathbf{v}_t) = \log p_{\mathbf{x}_t}(\mathbf{x}_t) + \log \sqrt{v_t}$, which yields

$$\nabla_{\mathbf{v}_t} \log p_{\mathbf{v}_t}(\mathbf{v}_t) = \sqrt{v_t} \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t).$$

Now, substituting into Tweedie's formula,

$$\mathbb{E}_{\mathbf{v}_0 \sim p_{\mathbf{v}_0} | \mathbf{v}_t}[\mathbf{v}_0] = \frac{\mathbf{x}_t}{\sqrt{v_t}} + \sqrt{v_t} \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t),$$

giving

$$\mathbb{E}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0} | \mathbf{x}_t}[\mathbf{x}_0] = \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t + \frac{1}{\sqrt{\alpha_t}} v_t \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t).$$

The covariance is given as (Robbins, 1992; Efron, 2011), $\mathbb{V}_{\mathbf{v}_0 \sim p_{\mathbf{v}_0} | \mathbf{v}_t}[\mathbf{v}_0 | \mathbf{v}_t] = I + \nabla_{\mathbf{v}_t} \nabla_{\mathbf{v}_t} \log p_{\mathbf{v}_t}(\mathbf{v}_t)$, where

$$\nabla_{\mathbf{v}_t} \nabla_{\mathbf{v}_t} \log p_{\mathbf{v}_t}(\mathbf{v}_t) = \nabla_{\mathbf{v}_t} (\sqrt{v_t} \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t)) \quad (31)$$

$$= v_t \nabla_{\mathbf{x}_t}^2 \log p_{\mathbf{x}_t}(\mathbf{x}_t) \quad (32)$$

$$(33)$$

and so

$$\mathbb{V}_{\mathbf{v}_0 \sim p_{\mathbf{v}_0} | \mathbf{v}_t}[\mathbf{v}_0] = \mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t}^2 \log p_{\mathbf{x}_t}(\mathbf{x}_t)$$

giving

$$\mathbb{V}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0} | \mathbf{x}_t}[\mathbf{x}_0] = \frac{v_t}{\alpha_t} (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t)).$$

In practice, we make the approximation $\nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t) \approx \mathbf{s}_\theta(\mathbf{x}_t, t)$, giving

$$\mathbb{E}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0} | \mathbf{x}_t}[\mathbf{x}_0] \approx \mathbf{m}_{0|t} := \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t + \frac{1}{\sqrt{\alpha_t}} v_t \mathbf{s}_\theta(\mathbf{x}_t, t)$$

and

$$\mathbb{V}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0} | \mathbf{x}_t}[\mathbf{x}_0] \approx \frac{v_t}{\alpha_t} (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \mathbf{s}_\theta(\mathbf{x}_t, t)),$$

which yields a Gaussian approximation that matches the first and second moments,

$$p_{\mathbf{x}_0 | \mathbf{x}_t}(\mathbf{x}_0 | \mathbf{x}_t) \approx \mathcal{N} \left(\mathbf{m}_{0|t}, \frac{v_t}{\alpha_t} (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \mathbf{s}_\theta(\mathbf{x}_t, t)) \right) \quad (34)$$

$$= \mathcal{N} \left(\mathbf{m}_{0|t}, \frac{v_t}{\alpha_t} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \right). \quad (35)$$

The matrix $\frac{v_t}{\alpha_t} (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \mathbf{s}_\theta(\mathbf{x}_t, t))$ needs to be inverted to calculate the log likelihood, and therefore must be both symmetric and positive definite for all time, which puts the requirement that $\mathbf{s}_\theta(\mathbf{x}_t, t)$ can be written as the negative gradient of a potential, however it is only approximated as such $\mathbf{s}_\theta(\mathbf{x}_t, t) \approx \nabla \log p_{\mathbf{x}_t}(\mathbf{x}_t)$.

D.2 Variance Exploding SDE (time-rescaled Brownian motion)

Using the Variance Exploding SDE (Song et al., 2020) or time-rescaled Brownian motion, given the random variable

$$\sqrt{\frac{1}{v_t}} \mathbf{x}_0 := \mathbf{v}_0 \sim p_{\mathbf{z}_0}, \quad (36)$$

a similar calculation as in D.1 gives

$$\mathbb{E}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0 | \mathbf{x}_t}}[\mathbf{x}_0] = \mathbf{x}_t + v_t \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t),$$

and

$$\mathbb{V}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0 | \mathbf{x}_t}}[\mathbf{x}_0] = v_t (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t)).$$

Again, in practice, we make the approximation $\nabla_{\mathbf{x}_t} \log p_{\mathbf{x}_t}(\mathbf{x}_t) = \mathbf{s}_\theta(\mathbf{x}_t, t)$, giving

$$\mathbb{E}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0 | \mathbf{x}_t}}[\mathbf{x}_0] \approx \mathbf{m}_{0|t} := \mathbf{x}_t + v_t \mathbf{s}_\theta(\mathbf{x}_t, t)$$

and

$$\mathbb{V}_{\mathbf{x}_0 \sim p_{\mathbf{x}_0 | \mathbf{x}_t}}[\mathbf{x}_0] \approx v_t (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \mathbf{s}_\theta(\mathbf{x}_t, t)),$$

which yields a Gaussian approximation that matches the first and second moments,

$$p_{\mathbf{x}_0 | \mathbf{x}_t}(\mathbf{x}_0 | \mathbf{x}_t) \approx \mathcal{N}(\mathbf{m}_{0|t}, v_t (\mathbf{I}_{d_x} + v_t \nabla_{\mathbf{x}_t} \mathbf{s}_\theta(\mathbf{x}_t, t))) \quad (37)$$

$$= \mathcal{N}(\mathbf{m}_{0|t}, v_t \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}). \quad (38)$$

E Algorithmic details and numerics

The code for all of the experiments and instructions to run them are available at X.

Algorithm 1 TMPD-D (Ancestral sampling, VP)

```

input  $\mathbf{y}, \sigma_y$ 
 $\mathbf{x}_N \sim \mathcal{N}(0, \mathbf{I}_{d_x})$ 
for  $n = N - 1, \dots, 0$  do
   $\mathbf{m}_{0|t} = \frac{1}{\sqrt{\alpha_n}}(\mathbf{x}_n + v_n \mathbf{s}_\theta(\hat{\mathbf{x}}_n, t_n))$ 
   $\mathbf{m}_{0|t}^y = \mathbf{m}_{0|t} + \frac{v_n}{\sqrt{\alpha_n}} \nabla_{\mathbf{x}_n} \mathbf{m}_{0|t} \mathbf{H}^\top (\mathbf{H} \frac{v_n}{\sqrt{\alpha_n}} \nabla_{\mathbf{x}_n} \mathbf{m}_{0|t} \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H} \mathbf{m}_{0|t})$ 
   $\mathbf{z}_n \sim \mathcal{N}(0, \mathbf{I}_{d_x})$ 
   $\sigma_n = \sqrt{(1 - \alpha_{n-1}) \beta_n / (1 - \alpha_n)}$ 
   $\mathbf{x}_{n-1} \leftarrow \frac{\sqrt{1 - \beta_n (1 - \alpha_{n-1})}}{1 - \alpha_n} \mathbf{x}_n + \frac{\sqrt{\alpha_{n-1} \beta_n}}{1 - \alpha_n} \mathbf{m}_{0|t}^y + \sigma_n \mathbf{z}_n$ 
end for
output  $\mathbf{x}_0$ 

```

E.1 Computational Complexity

Let N is the number of noise scales, and let d_y is the dimensions of the observation, d_x are the dimension of the image and T_s is the time complexity of evaluating the score network. Computing a jacobian-vector product has time complexity T_s . Then, the time complexity of TMPD (not including fast matrix-vector products) is $\mathcal{O}(N(d_y^3 + T_s d_y + T_s + T_s))$. IIGDM comes at a smaller computational cost due to needing only $\mathcal{O}(1)$ vector-jacobian-products, instead of $\mathcal{O}(d_y)$, resulting in a time complexity of $\mathcal{O}(N(T_s + T_s))$. DPS comes at the same complexity as IIGDM.

Whereas TMPD requires calculating the Jacobian which has memory complexity of at least $\mathcal{O}(d_x d_y)$. This is too large for high resolution image problems where the dimension of the observation is large. In comparison,

the memory complexity of IIGDM depends on the observation operator, but for the class of problems that are explored in Song et al. (2023) is $\mathcal{O}(d_x)$. DPS comes at the same memory complexity as IIGDM.

As mentioned in the text, we make the following approximation

$$\text{DTMPD} \quad \nabla_{\mathbf{x}_t} \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) \approx \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top (\mathbf{H} \frac{v_t}{\sqrt{\alpha_t}} \text{diag}(\nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}) \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H} \mathbf{m}_{0|t}).$$

Whilst this approximation does not require a linear solve, taking out the $\mathcal{O}(Nd_y^3)$ time complexity term, we would still like to take out the $\mathcal{O}(NT_s d_y)$ time complexity and $\mathcal{O}(d_x d_y)$ memory complexity term from calculating and storing the Jacobian, respectively, since this is too large for solving high resolution image applications. A further approximation approximates the diagonal of the Jacobian by the row sum of the Jacobian which only requires one vector jacobian product and brings the memory and time complexity of DTMPD down to that of IIGDM and DPS. We use this approximation in the image experiments and find that in practice it is only $(1.5 \pm 0.1) \times$ slower than IIGDM and DPS across all of our experiments. The row sum will be a good approximation of the diagonal when the Jacobian is approximately diagonal, which happens when there is small linear correlation between observation pixels, which we found to work well for super-resolution and inpainting. In inpainting, we have further improved the accuracy of the rowsum by instead of calculating the vector Jacobian product evaluated at $\mathbf{H}\mathbf{1}$, masking out values of the vector in the vector Jacobian product using the inpainting observation operator since they won't contribute to the diagonal values of the of the variance. A fruitful direction of research would be to explore low rank approximations of the Jacobian matrix.

E.2 Gaussian

When the data distribution $p_0(\mathbf{x}_0)$ is a (multivariate) Gaussian, then the reverse SDE is a linear SDE and we can calculate all of the terms needed to sample from the target posterior using diffusion explicitly. Moreover, we can sample from the target posterior using a direct or implicit method such as Cholesky decomposition. In this simple example, we compare samples from the direct method to various conditional diffusion methods (TMPD, IIGDM, DPM), by plotting a sample estimate of the L^2 Wasserstein distance between the sample and the target Gaussian measures (Givens & Shortt, 1984) by using the analytical mean and covariance of the target distribution and empirical estimate of the mean and covariance of the sample distribution. To generate $p_0(\mathbf{x}_0)$, we use an equally spaced grid of vectors $\mathbf{u}_i \in [-5.0, 5.0]^2$ for $i \in 1, 2, \dots, 32^2$, pick a Matern 5/2 kernel for the covariance function $k(\mathbf{u}_i, \mathbf{u}_j) = (1 + \sqrt{5}|\mathbf{u}_i - \mathbf{u}_j| + \frac{5}{3}|\mathbf{u}_i - \mathbf{u}_j|^2) \exp(-\sqrt{3}|\mathbf{u}_i - \mathbf{u}_j|)$ which defines the prior $p_0(\mathbf{x}_0) = \mathcal{N}(\mathbf{m}_0, \mathbf{C}_0)$ covariance $\mathbf{C}_0 \in \mathbb{R}^{1024 \times 1024}$ where $\mathbf{C}_{0ij} = k(\mathbf{u}_i, \mathbf{u}_j)$ and we define the mean as a zero vector $\mathbf{m}_0 = \mathbf{0} \in \mathbb{R}^{1024}$. To compute analytically the distribution of $p_{0|\mathbf{y}}(\mathbf{x}_0|\mathbf{y})$, we sample $\mathbf{y} = \mathbf{H}\mathbf{x}_0 + \mathbf{z}$, $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \sigma_y^2 \mathbf{I}_{d_y})$ and use the standard Gaussian formula to calculate the mean and covariance of $\mathbf{x}_0|\mathbf{y}$ which in this case are a complete description of $p_{0|\mathbf{y}}(\mathbf{x}_0|\mathbf{y})$. The L^2 Wasserstein estimate is plotted over an increasing sample size $N \in [9, 1500]$ and for $\sigma_y = 0.1$ in Figure 1.

We provide an illustration of the mean and uncertainty captured by the diffusion samples using 1500 samples of each diffusion model to produce a Monte-Carlo estimate of the mean and diagonal variance vector, and compare these to the exact mean and diagonal variance.

Below, we provide a calculation comparing the exact diffusion posterior for the linear diffusion posterior sde to the approximations used in IIGDM (Song et al., 2023) and TMPD. Let us assume that the target

distribution be known $p_0(\mathbf{x}_0) = \mathcal{N}(\mathbf{m}_0, \mathbf{C}_0)$. Then, via Bayes' rule,

$$\begin{aligned}
\log p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) &= \log p_{t|0}(\mathbf{x}_t|\mathbf{x}_0) + \log p_0(\mathbf{x}_0) + \text{constant} \\
&= -\frac{1}{2}(\mathbf{x}_t - \sqrt{\alpha_t}\mathbf{x}_0)^\top (v_t\mathbf{I}_{d_x})^{-1}(\mathbf{x}_t - \sqrt{\alpha_t}\mathbf{x}_0) \\
&\quad -\frac{1}{2}(\mathbf{x}_0 - \mathbf{m}_0)^\top \mathbf{C}_0^{-1}(\mathbf{x}_0 - \mathbf{m}_0) + \text{constant} \\
&= -\frac{1}{2}(\sqrt{\alpha_t}\mathbf{x}_0)^\top (v_t\mathbf{I}_{d_x})^{-1}(\sqrt{\alpha_t}\mathbf{x}_0) + \mathbf{x}_t^\top (v_t\mathbf{I}_{d_x})^{-1}(\sqrt{\alpha_t}\mathbf{x}_0) \\
&\quad -\frac{1}{2}\mathbf{x}_0^\top \mathbf{C}_0^{-1}\mathbf{x}_0 + \mathbf{m}_0^\top \mathbf{C}_0^{-1}\mathbf{x}_0 + \text{constant} \\
&= -\frac{1}{2}(\mathbf{x}_0 - \mathbf{m}_t)^\top \boldsymbol{\Sigma}_t^{-1}(\mathbf{x}_0 - \mathbf{m}_t),
\end{aligned}$$

so

$$p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) = \mathcal{N}(\mathbf{m}_t, \boldsymbol{\Sigma}_t) \quad (39)$$

where

$$\boldsymbol{\Sigma}_t = \left(\left(\frac{v_t}{\alpha_t} \mathbf{I}_{d_x} \right)^{-1} + \mathbf{C}_0^{-1} \right)^{-1} \quad (40)$$

and

$$\mathbf{m}_t = \boldsymbol{\Sigma}_t \left(\frac{\sqrt{\alpha_t}}{v_t} \mathbf{x}_t + \mathbf{C}_0^{-1} \mathbf{m}_0 \right).$$

We also have that,

$$p_t(\mathbf{x}_t) = \mathcal{N}(\sqrt{\alpha_t}\mathbf{m}_0, \mathbf{C}_t)$$

where $\mathbf{C}_t = \alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x}$, and thus

$$\log p_t(\mathbf{x}_t) = -\frac{1}{2}(\mathbf{x}_t - \mathbf{m}_0\sqrt{\alpha_t})^\top \mathbf{C}_t^{-1}(\mathbf{x}_t - \mathbf{m}_0\sqrt{\alpha_t}).$$

From

$$p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) = \int p_{\mathbf{y}|0}(\mathbf{y}|\mathbf{x}_0) p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) d\mathbf{x}_0 = \mathcal{N}(\mathbf{H}\mathbf{m}_t, \mathbf{H}\boldsymbol{\Sigma}_t\mathbf{H}^\top + \sigma_y^2\mathbf{I}_{d_y}),$$

we have that

$$\log p_t(\mathbf{y}|\mathbf{x}_t) = -\frac{1}{2}(\mathbf{y} - \mathbf{H}\mathbf{m}_t)^\top (\mathbf{H}\boldsymbol{\Sigma}_t\mathbf{H}^\top + \sigma_y^2\mathbf{I}_{d_y})^{-1}(\mathbf{y} - \mathbf{H}\mathbf{m}_t).$$

The posterior score can be calculated (see Cardoso et al. (2023)) from applying Bayes' theorem,

$$\log p_{t|\mathbf{y}}(\mathbf{x}_t|\mathbf{y}) = \log p_t(\mathbf{x}_t) + \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) + \text{Constant},$$

and taking gradients with respect to the state \mathbf{x}_t gives

$$\nabla_{\mathbf{x}_t} \log p_{t|\mathbf{y}}(\mathbf{x}_t|\mathbf{y}) = \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) + \nabla_{\mathbf{x}_t} \log p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) \quad (41)$$

$$= -(\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}(\mathbf{x}_t - \mathbf{m}_0\sqrt{\alpha_t}) \quad (42)$$

$$+ \frac{\sqrt{\alpha_t}}{v_t} \boldsymbol{\Sigma}_t^\top \mathbf{H}^\top (\mathbf{H}\boldsymbol{\Sigma}_t\mathbf{H}^\top + \sigma_y^2\mathbf{I}_{d_y})^{-1}(\mathbf{y} - \mathbf{H}\boldsymbol{\Sigma}_t(\frac{\sqrt{\alpha_t}}{v_t}\mathbf{x}_t + \mathbf{C}_0^{-1}\mathbf{m}_0)). \quad (43)$$

Setting $\mathbf{m}_0 = \mathbf{0}$ in (43) gives,

$$\nabla_{\mathbf{x}_t} \log p_{t|\mathbf{y}}(\mathbf{x}_t|\mathbf{y}) = -(\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}\mathbf{x}_t \quad (44)$$

$$+ \frac{\sqrt{\alpha_t}}{v_t} \boldsymbol{\Sigma}_t^\top \mathbf{H}^\top (\mathbf{H}\boldsymbol{\Sigma}_t\mathbf{H}^\top + \sigma_y^2\mathbf{I}_{d_y})^{-1}(\mathbf{y} - \frac{\sqrt{\alpha_t}}{v_t}\mathbf{H}\boldsymbol{\Sigma}_t\mathbf{x}_t) \quad (45)$$

E.2.1 Comparison of the exact posterior to the approximation made in Pseudo-Inverse-Guidance

We aim to compare (45), to the approximation used in Π GDM (Song et al., 2023);

$$p_{0|t}(\mathbf{x}_0|\mathbf{x}_t) \approx \mathcal{N}(\mathbf{m}_{0|t}, r_t^2 \mathbf{I}_{d_x}),$$

where $\mathbf{m}_{0|t} = \mathbf{x}_t + v_t \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$, which is the MMSE estimate of $\mathbf{x}_0|\mathbf{x}_t$ and r_t is chosen empirically. This gives $p_{\mathbf{y}|t}(\mathbf{y}|\mathbf{x}_t) \approx \mathcal{N}(\mathbf{H}\mathbf{m}_{0|t}, r_t^2 \mathbf{H}\mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})$, which in turn gives

$$\nabla_{\mathbf{x}_t} \log p_{t|y}(\mathbf{x}_t|\mathbf{y}) \approx \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) \quad (46)$$

$$+ \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top (r_t^2 \mathbf{H}\mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H}\mathbf{m}_{0|t}) \quad (47)$$

which is computationally tractable in the standard diffusion model setting where the score is non-linear and approximated as $\nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) \approx \mathbf{s}_\theta(\mathbf{x}_t, t)$. Substituting in the known score and again setting $\mathbf{m}_0 = \mathbf{0}$ score to compare to the linear case (45) gives,

$$\nabla_{\mathbf{x}_t} \log p_{t|y}(\mathbf{x}_t|\mathbf{y}) \approx \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) \quad (48)$$

$$+ \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top (r_t^2 \mathbf{H}\mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H}\mathbf{m}_{0|t}) \quad (49)$$

$$= -(\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1} \mathbf{x}_t \quad (50)$$

$$+ \frac{1}{\sqrt{\alpha_t}} (\mathbf{I}_{d_x} - v_t (\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}) \mathbf{H}^\top (r_t^2 \mathbf{H}\mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} \quad (51)$$

$$(\mathbf{y} - \frac{1}{\sqrt{\alpha_t}} \mathbf{H} (\mathbf{I}_{d_x} - v_t (\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}) \mathbf{x}_t) \quad (52)$$

comparing terms, Π GDM is making the approximation

$$\nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} = \frac{1}{\sqrt{\alpha_t}} \mathbf{I}_{d_x} - \frac{v_t}{\sqrt{\alpha_t}} (\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1} \approx \frac{\sqrt{\alpha_t}}{v_t} \boldsymbol{\Sigma}_t.$$

Note that since,

$$\frac{\sqrt{\alpha_t}}{v_t} \boldsymbol{\Sigma}_t = \frac{\sqrt{\alpha_t}}{v_t} (\mathbf{C}_0^{-1} + (\frac{v_t}{\alpha_t} \mathbf{I}_{d_x})^{-1})^{-1} \quad (53)$$

$$= \frac{1}{v_t \sqrt{\alpha_t}} ((\alpha_t \mathbf{C}_0)^{-1} + (v_t \mathbf{I}_{d_x})^{-1})^{-1} \quad (54)$$

$$= \frac{1}{v_t \sqrt{\alpha_t}} (v_t \mathbf{I} - v_t^2 (\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}) \quad \text{Woodbury identity} \quad (55)$$

$$= \frac{1}{\sqrt{\alpha_t}} (\mathbf{I} - v_t (\alpha_t \mathbf{C}_0 + v_t \mathbf{I}_{d_x})^{-1}) \quad (56)$$

$$= \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t}, \quad (57)$$

this is exact. The other approximation Π GDM is making is $r_t^2 \mathbf{I} \approx \boldsymbol{\Sigma}_t$, and note that this approximation is accurate with $r_t^2 = v_t$ as $t \rightarrow 0$ but inaccurate as $t \rightarrow 1$. But the exact linear SDE can be recovered by instead using TMPD,

$$\nabla_{\mathbf{x}_t} \log p_{t|y}(\mathbf{x}_t|\mathbf{y}) = \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) \quad (58)$$

$$+ \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top (\mathbf{H} \frac{v_t}{\sqrt{\alpha_t}} \nabla_{\mathbf{x}_t} \mathbf{m}_{0|t} \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \mathbf{H}\mathbf{m}_{0|t}) \quad (59)$$

$$= \nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t) \quad (60)$$

$$+ \frac{\sqrt{\alpha_t}}{v_t} \boldsymbol{\Sigma}_t \mathbf{H}^\top (\mathbf{H} \boldsymbol{\Sigma}_t \mathbf{H}^\top + \sigma_y^2 \mathbf{I}_{d_y})^{-1} (\mathbf{y} - \frac{\sqrt{\alpha_t}}{v_t} \mathbf{H} \boldsymbol{\Sigma}_t \mathbf{x}_t) \quad (61)$$

Table 3: Sliced Wasserstein for the GMM example using the reverse VP SDEs discretized with Euler-Maruyama.

d_x	d_y	$\sigma_y = 0.01$				$\sigma_y = 0.1$				$\sigma_y = 1.0$			
		TMPD	DTMPD	IGDM	DPS	TMPD	DTMPD	IGDM	DPS	TMPD	DTMPD	IGDM	DPS
8	1	1.5 \pm 0.5	1.5 \pm 0.5	1.5 \pm 0.4	5.7 \pm 2.2	1.4 \pm 0.5	1.4 \pm 0.5	1.2 \pm 0.4	5.6 \pm 2.1	0.9 \pm 0.3	0.9 \pm 0.3	0.9 \pm 0.3	0.9 \pm 0.3
8	2	0.7 \pm 0.3	3.2 \pm 1.4	0.4 \pm 0.3	6.2 \pm 0.8	0.9 \pm 0.3	2.7 \pm 1.1	0.5 \pm 0.3	6.2 \pm 2.4	0.9 \pm 0.2	1.8 \pm 0.8	1.0 \pm 0.3	1.2 \pm 0.4
8	4	0.3 \pm 0.3	0.6 \pm 0.4	0.1 \pm 0.1	-	0.3 \pm 0.2	0.7 \pm 0.4	0.1 \pm 0.0	8.4 \pm 3.1	0.6 \pm 0.2	0.9 \pm 0.5	0.2 \pm 0.1	0.3 \pm 0.2
80	1	2.7 \pm 0.7	2.7 \pm 0.7	2.9 \pm 1.4	9.1 \pm 1.3	2.3 \pm 0.7	2.3 \pm 0.7	2.1 \pm 1.1	4.7 \pm 1.8	1.5 \pm 0.7	1.5 \pm 0.7	1.8 \pm 0.8	1.9 \pm 0.9
80	2	1.0 \pm 0.5	3.3 \pm 1.0	0.8 \pm 0.7	2.2 \pm 0.9	1.2 \pm 0.5	3.3 \pm 1.0	0.8 \pm 0.7	6.0 \pm 2.1	1.1 \pm 0.2	2.2 \pm 1.0	1.3 \pm 0.5	1.5 \pm 0.5
80	4	0.3 \pm 0.1	0.9 \pm 0.5	0.1 \pm 0.0	-	0.4 \pm 0.2	1.0 \pm 0.5	0.1 \pm 0.1	4.4 \pm 1.6	0.9 \pm 0.2	1.0 \pm 0.4	0.4 \pm 0.2	0.5 \pm 0.3
800	1	3.1 \pm 0.7	3.1 \pm 0.7	3.2 \pm 1.0	6.8 \pm 1.2	2.9 \pm 0.6	2.9 \pm 0.6	2.8 \pm 0.7	6.4 \pm 1.5	1.5 \pm 0.4	1.5 \pm 0.4	1.3 \pm 0.3	1.3 \pm 0.3
800	2	1.3 \pm 0.4	3.6 \pm 1.2	0.8 \pm 0.5	7.4 \pm 0.9	1.3 \pm 0.3	3.2 \pm 1.1	0.8 \pm 0.4	6.4 \pm 1.9	1.2 \pm 0.3	1.9 \pm 0.5	1.1 \pm 0.3	1.1 \pm 0.3
800	4	0.3 \pm 0.2	0.9 \pm 0.6	0.6 \pm 0.5	-	0.4 \pm 0.2	0.9 \pm 0.6	0.1 \pm 0.0	5.8 \pm 1.4	0.9 \pm 0.2	1.1 \pm 0.5	0.4 \pm 0.2	0.4 \pm 0.2

E.3 GMM

For a given dimension d_x , we consider p_0 a mixture of 25 Gaussian random variables. The components have mean $\mu_{i,j} := (8i, 8j, \dots, 8i, 8j) \in \mathbb{R}^{d_x}$ for $(i, j) \in -2, -1, 0, 1, 2^2$ and unit variance. We have set the associated unnormalized weights $\omega_{i,j} = 1.0$. We have set $\sigma_\delta^2 = 10^{-4}$.

Note that $p_t(\mathbf{x}_t) = \int p_{t|0}(\mathbf{x}_t|\mathbf{x}_0)p_0(\mathbf{x}_0)d\mathbf{x}_0$. As $p_0(\mathbf{x}_0)$ is a mixture of Gaussians, $p_t(\mathbf{x}_t)$ is also a mixture of Gaussians with means $\sqrt{\alpha_t}\mu_{i,j}$ and unitary variances. Therefore, using automatic differentiation libraries, we can calculate $\nabla_{\mathbf{x}_t} \log p_t(\mathbf{x}_t)$. We chose $\beta_{\max} = 500.0$ and $\beta_{\min} = 0.1$. We use 1000 timesteps for the time-discretization. For the pair of dimensions and chosen observation noise standard deviation (d_x, d_y, σ_y) the measurement model (y, \mathbf{H}) is drawn as follows:

- \mathbf{H} : We first draw $\tilde{\mathbf{H}} \sim \mathcal{N}(\mathbf{0}_{d_y \times d_x}, \mathbf{I}_{d_y \times d_x})$ and compute the SVD decomposition of $\tilde{\mathbf{H}} = \mathbf{U}\mathbf{S}\mathbf{V}^\top$. Then, we sample for $(i, j) \in -2, -1, 0, 1, 2^2$, $s_{i,j}$ according to a uniform in $[0, 1]$. Finally, we set $\mathbf{H} = \mathbf{U}\text{diag}(s_{i,j})_{(i,j) \in -2, -1, 0, 1, 2^2} \mathbf{V}^\top$.
- \mathbf{y} : We then draw $\mathbf{x}_* \sim p_0$ and set $\mathbf{y} := \mathbf{H}\mathbf{x}_* + \mathbf{z}$ where $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \sigma_y^2 \mathbf{I}_{d_y})$.

Once we have drawn both $\mathbf{x}_* \sim p_0$ and $(\mathbf{y}, \mathbf{H}, \sigma_y)$, the posterior can be exactly calculated using Bayes formula and gives a mixture of Gaussians with mixture components $c_{i,j}$ and associated weights $\tilde{\omega}_{i,j}$,

$$c_{i,j} := \mathcal{N}(\Sigma(\mathbf{H}^\top \mathbf{y} / \sigma_y^2 + \mu_{i,j}), \Sigma), \quad (62)$$

$$\tilde{\omega}_i := \omega_i \mathcal{N}(\mathbf{y}; \mathbf{H}\mu_{i,j}, \sigma_\delta^2 \mathbf{I}_{d_x} + \mathbf{H}\mathbf{H}^\top), \quad (63)$$

where $\Sigma = (\mathbf{I}_{d_x} + \frac{1}{\sigma_\delta^2} \mathbf{H}^\top \mathbf{H})^{-1}$.

Euler-Maruyama solver To compare the posterior distribution estimated by each algorithm with the target posterior distribution, we use 10^4 slices for the SW distance and compare 1000 samples of the continuous SDEs defined by the TMPD, DTMPD, Song et al. (2023) and Chung et al. (2022a) approximations obtained using 1000 Euler-Maruyama time-steps with 1000 samples of the true posterior distribution. Table 3 indicates the Central Limit Theorem (CLT) 95% confidence intervals obtained by considering 20 randomly selected measurement models (\mathbf{H}) for each setting (d_x, d_x, σ_y) .

DDPM Table 4 compares 1000 samples of TMPD-D, IGDM-D and DPS-D which are obtained using 1000 denoising steps and is the extended version of Table 1. Figure 2 shows the first two dimensions of the estimated posterior distributions corresponding to the configurations (80, 1) and (800, 1) from Table 4 for one of the randomly generated measurement model (\mathbf{H}) . These illustrations give us insight into the behaviour of the algorithms and their ability to accurately estimate the posterior distribution. We observe that TMPD-D (and the Euler-Maruyama method TMPD) is the only method that covers the modes of the posterior distribution. Finally, a direct comparison to Cardoso et al. (2023) using their original experimental setup is shown in Table 5, which shows competitive performance for posterior sampling compared to Sequential Monte-Carlo, an exact sampling method.

Table 4: Sliced Wasserstein for the GMM example using VP DDPM.

d_x	d_y	$\sigma_{\mathbf{y}} = 0.01$				$\sigma_{\mathbf{y}} = 0.1$				$\sigma_{\mathbf{y}} = 1.0$			
		TMPD-D	DTMPD-D	IIGDM-D	DPS-D	TMPD-D	DTMPD-D	IIGDM-D	DPS-D	TMPD-D	DTMPD-D	IIGDM-D	DPS-D
8	1	1.6 ± 0.5	1.8 ± 0.6	2.6 ± 0.9	4.7 ± 1.5	1.4 ± 0.5	1.8 ± 0.7	2.2 ± 0.9	4.7 ± 1.6	0.9 ± 0.3	0.9 ± 0.2	1.5 ± 0.4	5.2 ± 1.3
8	2	0.7 ± 0.3	3.3 ± 1.5	2.1 ± 1.0	1.8 ± 1.5	0.9 ± 0.3	2.7 ± 1.1	1.6 ± 0.6	1.5 ± 0.9	0.9 ± 0.2	1.7 ± 0.8	2.3 ± 0.4	3.5 ± 1.2
8	4	0.3 ± 0.3	0.4 ± 0.2	3.8 ± 2.3	0.7 ± 0.6	0.3 ± 0.2	0.5 ± 0.2	3.8 ± 2.2	0.8 ± 0.6	0.6 ± 0.2	0.9 ± 0.5	1.8 ± 0.3	2.5 ± 0.9
80	1	2.7 ± 0.7	2.8 ± 0.9	3.2 ± 1.0	5.6 ± 1.8	2.3 ± 0.7	2.6 ± 0.9	2.9 ± 0.8	5.1 ± 1.8	1.5 ± 0.7	1.4 ± 0.6	1.6 ± 0.5	6.9 ± 1.8
80	2	1.0 ± 0.5	3.2 ± 1.1	2.8 ± 1.3	3.2 ± 1.9	1.2 ± 0.5	3.2 ± 1.1	2.7 ± 1.2	3.1 ± 1.9	1.1 ± 0.2	2.1 ± 1.0	1.4 ± 0.2	3.9 ± 1.2
80	4	0.3 ± 0.1	0.7 ± 0.4	0.6 ± 0.4	1.2 ± 1.1	0.4 ± 0.2	0.8 ± 0.4	0.6 ± 0.4	1.0 ± 1.1	0.9 ± 0.3	0.9 ± 0.4	0.9 ± 0.2	1.7 ± 0.6
800	1	3.1 ± 0.7	3.7 ± 0.7	3.5 ± 1.1	5.8 ± 1.6	2.9 ± 0.6	3.4 ± 0.7	3.3 ± 0.9	5.7 ± 1.6	1.5 ± 0.4	1.4 ± 0.4	2.0 ± 0.4	6.8 ± 1.0
800	2	1.4 ± 0.4	3.5 ± 0.7	3.1 ± 1.1	3.5 ± 1.7	1.3 ± 0.3	3.4 ± 0.7	2.7 ± 0.9	3.1 ± 1.4	1.2 ± 0.3	2.0 ± 0.4	2.0 ± 0.5	4.7 ± 1.3
800	4	0.4 ± 0.2	0.7 ± 0.5	0.4 ± 0.2	1.4 ± 1.0	0.4 ± 0.2	0.8 ± 0.5	0.4 ± 0.2	1.3 ± 0.9	0.9 ± 0.2	1.1 ± 0.5	0.6 ± 0.2	0.9 ± 0.4

Table 5: Comparison to Cardoso et al. (2023) using their original experimental setup. TMPD-D and DTMPD-D use 1000 steps of DDPM.

d_x	d_y	MCGDIFF	TMPD-D	DTMPD-D
8	1	1.43 ± 0.55	1.83 ± 0.5	1.82 ± 0.5
8	2	0.49 ± 0.24	0.95 ± 0.3	2.27 ± 0.9
8	4	0.38 ± 0.25	0.61 ± 0.3	0.72 ± 0.4
80	1	1.39 ± 0.45	2.81 ± 0.8	2.81 ± 0.8
80	2	0.67 ± 0.24	1.14 ± 0.4	2.82 ± 0.9
80	4	0.28 ± 0.14	0.95 ± 0.5	0.95 ± 0.5
800	1	2.40 ± 1.00	2.96 ± 0.6	2.96 ± 0.6
800	2	1.31 ± 0.60	1.60 ± 0.5	3.07 ± 1.1
800	4	0.47 ± 0.19	0.60 ± 0.2	0.84 ± 0.5

E.4 Inpainting and super-resolution

Since the DPS-D method was derived and tuned specifically for VP SDE, we look at the VP SDE experiments in Section E.4.1 separately from the VE SDE experiments in Section E.4.2. In each comparison, we use the same score network for each method and the same sampling or discretization numerical method. All methods are discretized using 1000 denoising steps. For the Markov chain methods we use DDPM and for the SDE methods we use an Euler-Maruyama discretization. In contrast to DPS(-D) IIGDM(-D), we observe the robustness of our method across both SDEs and inpainting and super-resolution observation maps.

E.4.1 VP SDE

FFHQ 256×256 For VP SDE, our FFHQ 256×256 experiment compares diffusion methods (DDPM) TMPD-D to DPS-D and IIGDM-D, and results are shown in Table 6. Fig 3, Fig 4 and Fig 5 are a visual summary of Table 6, plotting the LPIPS, SSIM and FID metrics against increasing noise for different observation maps. Some uncured samples that were used to generate Table 6 are shown in Fig 6, 7 and 8. We observe that all methods can successfully produce high quality reconstructions in the low noise regime but, visually, only DPS-D and TMPD-D successfully produce high quality reconstructions in the high noise regime.

Whereas DPS-D requires a hyperparameter search, there are no hyperparameters for TMPD-D. For IIGDM-D we use the algorithm and default hyperparameters as described in Song et al. (2023). For DPS-D we use the algorithm in the codebase provided by the authors Chung et al. (2022a) and we use their suggested step-size hyperparameter for this task, and we use their default hyperparameters, such as using static-thresholding (clipping the denoised image at each step to a range $[-1, 1]$) whereas TMPD-D does not require static-thresholding or any hyperparameter tuning.

CIFAR10 64×64 Our CIFAR-10 64×64 experiment compares TMPD to DPS and IIGDM, and also compares diffusion methods (DDPM) in Table 7 and score-based methods (discretized with Euler-Maruyama) in Table 8. Various samples used to produce the figures in Tables 7 and 8 are shown in Fig 10 and 11.

Table 6: Quantitative evaluation of solving linear inverse problems for VP DDPM with increasing noise on FFHQ 256×256-1k validation dataset.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD-D	29.6	0.230 ± 0.034	1.60E-03 ± 7.74E-04	28.4 ± 1.9	0.784 ± 0.046
4× ‘BICUBIC’	DPS-D	31.4	0.234 ± 0.048	1.90E-03 ± 1.07E-03	27.8 ± 2.2	0.776 ± 0.062
SUPER-RESOLUTION	PIGDM-D	29.7	0.198 ± 0.037	1.56E-03 ± 8.72E-04	28.6 ± 2.1	0.809 ± 0.051
$\sigma_y = 0.05$	TMPD-D	32.7	0.304 ± 0.043	2.90E-03 ± 5.64E-03	26.0 ± 1.7	0.699 ± 0.060
4× ‘BICUBIC’	DPS-D	29.3	0.280 ± 0.051	2.90E-03 ± 5.73E-03	26.0 ± 1.8	0.719 ± 0.066
SUPER-RESOLUTION	PIGDM-D	45.1	0.311 ± 0.047	3.08E-03 ± 5.79E-03	25.7 ± 1.7	0.682 ± 0.062
$\sigma_y = 0.1$	TMPD-D	38.0	0.348 ± 0.048	4.33E-03 ± 4.72E-03	24.0 ± 1.6	0.635 ± 0.066
4× ‘BICUBIC’	DPS-D	30.9	0.318 ± 0.051	4.06E-03 ± 5.38E-03	24.4 ± 1.6	0.664 ± 0.069
SUPER-RESOLUTION	PIGDM-D	119.6	0.589 ± 0.047	1.10E-02 ± 5.56E-03	19.7 ± 1.0	0.376 ± 0.055
$\sigma_y = 0.2$	TMPD-D	45.6	0.401 ± 0.049	7.03E-03 ± 2.56E-03	21.8 ± 1.5	0.559 ± 0.071
4× ‘BICUBIC’	DPS-D	38.1	0.385 ± 0.061	7.59E-03 ± 3.50E-03	21.6 ± 1.8	0.570 ± 0.081
SUPER-RESOLUTION	PIGDM-D	295.7	0.780 ± 0.033	5.65E-02 ± 5.20E-03	12.5 ± 0.4	0.117 ± 0.035
$\sigma_y = 0.01$	TMPD-D	25.7	0.153 ± 0.033	9.04E-03 ± 7.25E-03	21.4 ± 2.9	0.829 ± 0.031
‘BOX’ MASK	DPS-D	31.5	0.175 ± 0.038	7.79E-03 ± 6.89E-03	22.4 ± 3.3	0.833 ± 0.035
INPAINTING	PIGDM-D	143.8	0.247 ± 0.024	2.58E-02 ± 7.11E-03	16.1 ± 1.3	0.759 ± 0.017
$\sigma_y = 0.05$	TMPD-D	27.0	0.240 ± 0.038	9.68E-03 ± 6.62E-03	20.9 ± 2.6	0.760 ± 0.036
‘BOX’ MASK	DPS-D	30.7	0.228 ± 0.046	8.28E-03 ± 7.93E-03	22.0 ± 3.1	0.782 ± 0.047
INPAINTING	PIGDM-D	159.3	0.448 ± 0.046	2.71E-02 ± 7.91E-03	15.8 ± 1.2	0.504 ± 0.080
$\sigma_y = 0.1$	TMPD-D	29.6	0.292 ± 0.049	1.06E-02 ± 1.15E-02	20.6 ± 2.5	0.709 ± 0.053
‘BOX’ MASK	DPS-D	29.3	0.259 ± 0.049	8.03E-03 ± 6.97E-03	22.0 ± 2.9	0.746 ± 0.051
INPAINTING	PIGDM-D	165.7	0.539 ± 0.083	2.84E-02 ± 6.57E-03	15.6 ± 1.0	0.418 ± 0.185
$\sigma_y = 0.2$	TMPD-D	33.8	0.346 ± 0.061	1.13E-02 ± 8.65E-03	20.2 ± 2.4	0.649 ± 0.070
‘BOX’ MASK	DPS-D	34.9	0.337 ± 0.056	8.01E-03 ± 4.22E-03	21.5 ± 2.0	0.662 ± 0.067
INPAINTING	PIGDM-D	199.7	0.590 ± 0.099	3.17E-02 ± 1.01E-02	15.2 ± 1.4	0.461 ± 0.202
$\sigma_y = 0.01$	TMPD-D	24.6	0.090 ± 0.033	4.87E-04 ± 6.27E-04	34.3 ± 2.8	0.931 ± 0.036
‘RANDOM’ MASK	DPS-D	32.7	0.137 ± 0.033	5.57E-04 ± 5.02E-04	33.4 ± 2.6	0.913 ± 0.031
INPAINTING	PIGDM-D	24.7	0.069 ± 0.023	4.71E-04 ± 5.18E-04	34.5 ± 3.0	0.940 ± 0.026
$\sigma_y = 0.05$	TMPD-D	27.1	0.187 ± 0.032	8.86E-04 ± 4.84E-04	30.9 ± 1.8	0.851 ± 0.033
‘RANDOM’ MASK	DPS-D	38.0	0.256 ± 0.049	1.74E-03 ± 1.04E-03	28.1 ± 2.0	0.794 ± 0.056
INPAINTING	PIGDM-D	44.8	0.315 ± 0.043	2.15E-03 ± 6.65E-04	26.8 ± 1.2	0.647 ± 0.068
$\sigma_y = 0.1$	TMPD-D	29.9	0.247 ± 0.038	1.74E-03 ± 8.94E-03	28.7 ± 1.8	0.787 ± 0.049
‘RANDOM’ MASK	DPS-D	34.8	0.274 ± 0.054	2.21E-03 ± 1.17E-03	27.1 ± 2.1	0.763 ± 0.065
INPAINTING	PIGDM-D	62.7	0.490 ± 0.059	6.71E-03 ± 1.80E-03	22.0 ± 1.7	0.410 ± 0.129
$\sigma_y = 0.2$	TMPD-D	34.6	0.306 ± 0.046	2.71E-03 ± 2.89E-03	26.1 ± 1.7	0.714 ± 0.060
‘RANDOM’ MASK	DPS-D	35.9	0.317 ± 0.059	3.59E-03 ± 1.73E-03	24.9 ± 1.9	0.701 ± 0.073
INPAINTING	PIGDM-D	104.1	0.623 ± 0.109	1.77E-02 ± 8.62E-03	18.6 ± 3.8	0.316 ± 0.218

We found the hyperparameter suggested for the DDIM method for the VP SDE in Song et al. (2023) $r_t^2 = v_t/(\alpha_t + v_t)$, which is calculated by assuming the data distribution $p_0(\mathbf{x}_0)$ is a standard normal and then calculating the posterior variance, to give unstable solutions for the algorithm given in Song et al. (2023). To make the method stable, we instead plug the IIGDM posterior score approximation into a DDIM sampler in a similar way to Algorithm 1, which, for the VPSDE, brings the algorithm IIGDM-D closer to our method; we are then able to choose $r_t^2 = v_t/(\alpha_t + v_t)$ for both VP DDIM and VP SDE methods.

For DDPM we use the step-size constant suggested in Chung et al. (2022a) for inpainting, $\zeta_i = \zeta'/\|\mathbf{y} - \mathbf{H}\mathbf{m}_{0|t}\|$, where we tune ζ' over the suggested range of $\zeta' \in [0.1, 1.0]$ in Chung et al. (2022a) across LPIPS, MSE, PSNR and SSIM, as shown in Fig. 9 for each individual inverse problem (each line in the Tables 7 and 8).

Table 7: Noisy observation inpainting and super-resolution for VP DDPM on CIFAR-10 1k validation set.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD-D	33.7	0.090 \pm 0.048	0.007 \pm 0.034	23.8 \pm 3.7	0.784 \pm 0.073
‘BOX’ MASK	DPS-D	31.5	0.064 \pm 0.033	0.004 \pm 0.003	25.8 \pm 3.6	0.841 \pm 0.068
INPAINTING	IGDM-D	37.1	0.316 \pm 0.108	0.012 \pm 0.029	19.9 \pm 1.9	0.546 \pm 0.143
$\sigma_y = 0.01$	TMPD-D	40.4	0.272 \pm 0.071	0.028 \pm 0.020	16.4 \pm 3.0	0.584 \pm 0.071
‘HALF’ MASK	DPS-D	33.1	0.221 \pm 0.072	0.028 \pm 0.021	16.8 \pm 3.6	0.637 \pm 0.093
INPAINTING	IGDM-D	35.8	0.397 \pm 0.103	0.031 \pm 0.020	15.9 \pm 2.8	0.419 \pm 0.132
$\sigma_y = 0.05$	TMPD-D	35.9	0.128 \pm 0.061	0.006 \pm 0.011	23.5 \pm 3.3	0.763 \pm 0.079
‘BOX’ MASK	DPS-D	31.1	0.078 \pm 0.037	0.004 \pm 0.003	25.5 \pm 3.3	0.830 \pm 0.070
INPAINTING	IGDM-D	36.3	0.319 \pm 0.110	0.012 \pm 0.031	20.0 \pm 1.9	0.545 \pm 0.142
$\sigma_y = 0.05$	TMPD-D	40.4	0.292 \pm 0.075	0.029 \pm 0.020	16.4 \pm 3.1	0.572 \pm 0.076
‘HALF’ MASK	DPS-D	32.5	0.230 \pm 0.075	0.028 \pm 0.023	16.7 \pm 3.5	0.626 \pm 0.096
INPAINTING	IGDM-D	35.6	0.398 \pm 0.109	0.032 \pm 0.021	15.9 \pm 2.8	0.421 \pm 0.133
$\sigma_y = 0.1$	TMPD-D	38.9	0.168 \pm 0.072	0.007 \pm 0.012	22.5 \pm 2.9	0.728 \pm 0.081
‘BOX’ MASK	DPS-D	31.6	0.101 \pm 0.043	0.004 \pm 0.003	24.5 \pm 2.9	0.807 \pm 0.078
INPAINTING	IGDM-D	36.3	0.318 \pm 0.109	0.012 \pm 8.553	19.7 \pm 1.8	0.546 \pm 0.140
$\sigma_y = 0.1$	TMPD-D	43.8	0.350 \pm 0.088	0.030 \pm 0.028	16.2 \pm 2.9	0.547 \pm 0.076
‘HALF’ MASK	DPS-D	33.0	0.522 \pm 0.097	0.031 \pm 0.022	16.3 \pm 3.3	0.602 \pm 0.097
INPAINTING	IGDM-D	36.2	0.276 \pm 0.085	0.034 \pm 0.023	15.5 \pm 2.7	0.412 \pm 0.128
$\sigma_y = 0.01$	TMPD-D	33.2	0.117 \pm 0.051	0.004 \pm 0.004	24.7 \pm 3.0	0.835 \pm 0.071
2× ‘NEAREST’	DPS-D	32.5	0.099 \pm 0.044	0.004 \pm 0.003	25.1 \pm 3.1	0.847 \pm 0.073
SUPER-RESOLUTION	IGDM-D	35.6	0.407 \pm 0.118	0.016 \pm 0.006	18.2 \pm 1.7	0.442 \pm 0.152
$\sigma_y = 0.01$	TMPD-D	41.5	0.278 \pm 0.084	0.011 \pm 0.006	20.4 \pm 2.7	0.563 \pm 0.114
4× ‘BICUBIC’	DPS-D	33.9	0.220 \pm 0.079	0.010 \pm 0.006	20.8 \pm 3.0	0.609 \pm 0.135
SUPER-RESOLUTION	IGDM-D	39.4	0.279 \pm 0.081	0.011 \pm 0.006	20.1 \pm 2.5	0.546 \pm 0.111
$\sigma_y = 0.05$	TMPD-D	33.9	0.156 \pm 0.065	0.005 \pm 0.025	24.1 \pm 2.8	0.810 \pm 0.079
2× ‘NEAREST’	DPS-D	34.4	0.127 \pm 0.048	0.004 \pm 0.003	24.4 \pm 2.7	0.825 \pm 0.070
SUPER-RESOLUTION	IGDM-D	35.1	0.407 \pm 0.118	0.016 \pm 0.006	18.2 \pm 1.7	0.440 \pm 0.153
$\sigma_y = 0.05$	TMPD-D	38.3	0.332 \pm 0.093	0.013 \pm 0.032	19.6 \pm 2.4	0.501 \pm 0.116
4× ‘BICUBIC’	DPS-D	38.2	0.265 \pm 0.087	0.011 \pm 0.006	20.2 \pm 2.4	0.567 \pm 0.128
SUPER-RESOLUTION	IGDM-D	23.2	0.522 \pm 0.101	0.034 \pm 0.012	15.0 \pm 1.7	0.215 \pm 0.114
$\sigma_y = 0.1$	TMPD-D	35.0	0.208 \pm 0.085	0.006 \pm 0.005	23.1 \pm 2.5	0.760 \pm 0.091
2× ‘NEAREST’	DPS-D	32.1	0.152 \pm 0.058	0.005 \pm 0.003	23.8 \pm 2.5	0.802 \pm 0.074
SUPER-RESOLUTION	IGDM-D	35.0	0.407 \pm 0.122	0.017 \pm 0.006	18.1 \pm 1.8	0.435 \pm 0.150
$\sigma_y = 0.1$	TMPD-D	36.8	0.378 \pm 0.102	0.015 \pm 0.007	18.7 \pm 2.1	0.435 \pm 0.121
4× ‘BICUBIC’	DPS-D	34.8	0.308 \pm 0.095	0.013 \pm 0.006	19.4 \pm 2.2	0.513 \pm 0.132
SUPER-RESOLUTION	IGDM-D	23.0	0.521 \pm 0.104	0.035 \pm 0.013	14.9 \pm 1.7	0.212 \pm 0.117

Table 8: Noisy observation inpainting and super-resolution for the reverse VP SDEs on CIFAR-10 1k validation set.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD	33.2	0.088 \pm 0.050	0.007 \pm 0.019	23.8 \pm 3.7	0.785 \pm 0.071
‘BOX’ MASK	DPS	61.8	0.646 \pm 0.078	0.111 \pm 0.049	9.9 \pm 1.9	0.050 \pm 0.068
INPAINTING	IGDM	34.8	0.073 \pm 0.035	0.004 \pm 0.004	25.1 \pm 3.5	0.816 \pm 0.068
$\sigma_y = 0.01$	TMPD	37.4	0.267 \pm 0.068	0.030 \pm 0.036	16.3 \pm 3.0	0.581 \pm 0.070
‘HALF’ MASK	DPS	65.8	0.645 \pm 0.079	0.112 \pm 0.050	9.9 \pm 1.9	0.052 \pm 0.069
INPAINTING	IGDM	33.3	0.236 \pm 0.068	0.027 \pm 0.021	17.0 \pm 3.6	0.620 \pm 0.087
$\sigma_y = 0.05$	TMPD	33.5	0.117 \pm 0.052	0.006 \pm 0.017	23.5 \pm 3.2	0.763 \pm 0.074
‘BOX’ MASK	DPS	62.2	0.647 \pm 0.078	0.110 \pm 0.048	10.0 \pm 1.9	0.055 \pm 0.067
INPAINTING	IGDM	35.2	0.103 \pm 0.047	0.004 \pm 0.003	24.8 \pm 2.9	0.798 \pm 0.074
$\sigma_y = 0.05$	TMPD	37.9	0.286 \pm 0.073	0.030 \pm 0.023	16.2 \pm 3.1	0.567 \pm 0.075
‘HALF’ MASK	DPS	65.1	0.647 \pm 0.080	0.117 \pm 0.054	9.7 \pm 2.0	0.052 \pm 0.063
INPAINTING	IGDM	34.1	0.256 \pm 0.078	0.027 \pm 0.022	16.8 \pm 3.3	0.605 \pm 0.094
$\sigma_y = 0.1$	TMPD	35.1	0.159 \pm 0.065	0.007 \pm 0.006	22.6 \pm 2.8	0.731 \pm 0.078
‘BOX’ MASK	DPS	61.9	0.647 \pm 0.074	0.112 \pm 0.052	9.9 \pm 1.9	0.054 \pm 0.071
INPAINTING	IGDM	36.2	0.140 \pm 0.059	0.005 \pm 0.003	23.7 \pm 2.6	0.763 \pm 0.083
$\sigma_y = 0.1$	TMPD	41.9	0.310 \pm 0.077	0.031 \pm 0.020	16.0 \pm 2.9	0.541 \pm 0.078
‘HALF’ MASK	DPS	81.0	0.646 \pm 0.082	0.112 \pm 0.052	10.0 \pm 2.0	0.052 \pm 0.069
INPAINTING	IGDM	38.2	0.279 \pm 0.074	0.029 \pm 0.021	16.4 \pm 3.2	0.577 \pm 0.095
$\sigma_y = 0.01$	TMPD	32.4	0.123 \pm 0.055	0.005 \pm 0.008	24.5 \pm 3.0	0.826 \pm 0.077
2× ‘NEAREST’	DPS	71.6	0.645 \pm 0.079	0.118 \pm 0.054	9.7 \pm 2.0	0.049 \pm 0.068
SUPER-RESOLUTION	IGDM	41.7	0.192 \pm 0.079	0.005 \pm 0.003	23.3 \pm 2.4	0.773 \pm 0.079
$\sigma_y = 0.01$	TMPD	22.9	0.525 \pm 0.098	0.035 \pm 0.013	14.9 \pm 1.7	0.206 \pm 0.109
4× ‘BICUBIC’	DPS	77.9	0.640 \pm 0.076	0.120 \pm 0.060	9.7 \pm 2.1	0.046 \pm 0.070
SUPER-RESOLUTION	IGDM	36.7	0.461 \pm 0.093	0.023 \pm 0.009	16.7 \pm 1.7	0.305 \pm 0.106
$\sigma_y = 0.05$	TMPD	32.7	0.158 \pm 0.060	0.005 \pm 0.003	23.9 \pm 2.6	0.800 \pm 0.075
2× ‘NEAREST’	DPS	72.5	0.648 \pm 0.078	0.118 \pm 0.053	9.7 \pm 1.9	0.050 \pm 0.069
SUPER-RESOLUTION	IGDM	37.3	0.229 \pm 0.086	0.006 \pm 0.003	22.6 \pm 2.1	0.735 \pm 0.088
$\sigma_y = 0.05$	TMPD	35.7	0.328 \pm 0.091	0.013 \pm 0.011	19.5 \pm 2.3	0.495 \pm 0.117
4× ‘BICUBIC’	DPS	77.9	0.641 \pm 0.071	0.120 \pm 0.061	9.7 \pm 2.1	0.047 \pm 0.070
SUPER-RESOLUTION	IGDM	34.8	0.479 \pm 0.094	0.026 \pm 0.010	16.2 \pm 1.7	0.276 \pm 0.112
$\sigma_y = 0.1$	TMPD	32.9	0.207 \pm 0.081	0.006 \pm 0.005	23.0 \pm 2.4	0.753 \pm 0.088
2× ‘NEAREST’	DPS	72.2	0.644 \pm 0.077	0.118 \pm 0.056	9.7 \pm 2.0	0.051 \pm 0.067
SUPER-RESOLUTION	IGDM	35.9	0.273 \pm 0.098	0.007 \pm 0.003	21.7 \pm 2.1	0.682 \pm 0.105
$\sigma_y = 0.1$	TMPD	34.7	0.379 \pm 0.096	0.016 \pm 0.007	18.5 \pm 2.0	0.429 \pm 0.120
4× ‘BICUBIC’	DPS	78.3	0.639 \pm 0.074	0.120 \pm 0.060	9.7 \pm 2.1	0.050 \pm 0.068
SUPER-RESOLUTION	IGDM	34.4	0.492 \pm 0.097	0.028 \pm 0.010	15.8 \pm 1.7	0.258 \pm 0.112




Figure 3: 4 \times bicubic super-resolution FID vs LPIPS (left) and SSIM (right) using the VP SDE on FFHQ-1k validation dataset for increasing observation noise.




Figure 4: ‘box’ mask inpainting FID vs LPIPS (left) and SSIM (right) using the VP SDE on FFHQ-1k validation dataset for increasing observation noise.

E.4.2 VE SDE

FFHQ 256 \times 256 For VE SDE, our FFHQ 256 \times 256 experiment compares diffusion methods (DDPM) TMPD-D to DPS-D and IIGDM-D, and results are shown in Table 9. Fig 12 and Fig 13 are a visual summary of Table 9, plotting the LPIPS, SSIM and FID metrics against increasing noise. Some uncured samples that were used to generate Table 9 are shown in Fig 14 and 15. Since the DPS-D and IIGDM-D algorithms were developed for the VP SDE, the methods are not performant for the VE SDE, whereas TMPD-D is performant across a wide range of noise levels.

CIFAR10 64 \times 64 For VE SDE, our CIFAR-10 64 \times 64 experiments compare TMPD to DPS and IIGDM, and also compare diffusion methods (DDPM) in Table 10 and score-based methods (discretized with Euler-Maruyama) in Table 11. Various samples used to produce the figures in Tables 10 and 11 are shown in Fig 16 and 17.

For IIGDM(-D), we are able to use the hyperparameter $r_t^2 = v_t/(1 + v_t)$ as suggested by Song et al. (2023), which is calculated by assuming the data distribution $p_0(\mathbf{x}_0)$ is a standard normal and then calculating the




Figure 5: ‘random’ mask inpainting FID vs LPIPS (left) and SSIM (right) using the VP SDE on FFHQ-1k validation dataset for increasing observation noise.



Figure 6: 4 \times bicubic super-resolution samples from the VP SDE on FFHQ across a range of observation noise.

posterior variance, for both VE DDIM and VE SDE methods, but note some instability for the Π GDM VE SDE method for small noise, as shown in Fig 10.

For DDPM we use the step-size constant suggested in Chung et al. (2022a) for inpainting, $\zeta_i = \zeta' / \|\mathbf{y} - \mathbf{H}\mathbf{m}_{0|t}\|$, where we tune ζ' over the suggested range of $\zeta' \in [0.1, 1.0]$ in Chung et al. (2022a) across LPIPS, MSE, PSNR and SSIM, as shown in Fig. 9 for each individual inverse problem (each line in the Tables 10 and 11).



Figure 7: ‘Random’ mask inpainting samples from the VP SDE on FFHQ across a range of observation noise.

Table 9: Quantitative evaluation of solving linear inverse problems for VE DDPM with increasing noise on FFHQ 256×256-1k validation dataset.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD-D	32.3	0.203 ± 0.039	1.57E-03 $\pm 8.89\text{E-}04$	28.6 ± 2.1	0.810 ± 0.052
4× ‘BICUBIC’	DPS-D	47.0	0.273 ± 0.031	1.70E-03 $\pm 8.36\text{E-}04$	28.1 ± 1.8	0.747 ± 0.037
SUPER-RESOLUTION	IIGDM-D	37.4	0.244 ± 0.030	1.74E-03 $\pm 9.56\text{E-}04$	28.1 ± 2.0	0.755 ± 0.042
$\sigma_y = 0.05$	TMPD-D	32.1	0.268 ± 0.048	2.62E-03 $\pm 1.17\text{E-}03$	26.2 ± 1.8	0.733 ± 0.066
4× ‘BICUBIC’	DPS-D	105.9	0.590 ± 0.036	6.18E-03 $\pm 8.50\text{E-}04$	22.1 ± 0.6	0.404 ± 0.040
SUPER-RESOLUTION	IIGDM-D	106.8	0.592 ± 0.041	7.92E-03 $\pm 1.07\text{E-}03$	21.0 ± 0.6	0.353 ± 0.042
$\sigma_y = 0.1$	TMPD-D	32.7	0.310 ± 0.053	3.99E-03 $\pm 1.84\text{E-}03$	24.3 ± 1.7	0.679 ± 0.071
4× ‘BICUBIC’	DPS-D	114.0	0.569 ± 0.044	8.47E-03 $\pm 4.95\text{E-}03$	21.1 ± 1.7	0.483 ± 0.046
SUPER-RESOLUTION	IIGDM-D	206.0	0.724 ± 0.034	2.46E-02 $\pm 2.16\text{E-}03$	16.1 ± 0.4	0.176 ± 0.034
$\sigma_y = 0.01$	TMPD-D	30.2	0.114 ± 0.029	2.84E-03 $\pm 2.56\text{E-}03$	26.5 ± 2.8	0.907 ± 0.020
‘BOX’ MASK	DPS-D	23.9	0.093 ± 0.019	2.53E-03 $\pm 1.77\text{E-}03$	26.9 ± 2.8	0.899 ± 0.017
INPAINTING	IIGDM-D	27.1	0.108 ± 0.025	2.50E-03 $\pm 1.55\text{E-}03$	26.7 ± 2.3	0.877 ± 0.015
$\sigma_y = 0.05$	TMPD-D	33.6	0.186 ± 0.036	3.24E-03 $\pm 3.19\text{E-}03$	25.8 ± 2.5	0.847 ± 0.030
‘BOX’ MASK	DPS-D	39.7	0.318 ± 0.044	3.81E-03 $\pm 1.92\text{E-}03$	24.7 ± 2.0	0.677 ± 0.086
INPAINTING	IIGDM-D	49.5	0.354 ± 0.044	4.67E-03 $\pm 1.50\text{E-}03$	23.5 ± 1.3	0.555 ± 0.091
$\sigma_y = 0.1$	TMPD-D	34.0	0.223 ± 0.041	3.42E-03 $\pm 2.24\text{E-}03$	25.3 ± 2.3	0.801 ± 0.045
‘BOX’ MASK	DPS-D	59.1	0.467 ± 0.053	7.78E-03 $\pm 3.19\text{E-}03$	21.4 ± 1.7	0.476 ± 0.110
INPAINTING	IIGDM-D	72.6	0.529 ± 0.047	9.88E-03 $\pm 2.48\text{E-}03$	20.2 ± 1.2	0.356 ± 0.118




Figure 8: ‘Box’ mask inpainting samples from the VP SDE on FFHQ across a range of observation noise.




Figure 9: DPS scale hyperparameter search across LPIPS, MSE, PSNR and SSIM for CIFAR-10 $4\times$ bicubic interpolation super-resolution with $\sigma = 0.05$. Plotted are the mean values ± 2 standard deviations over a 128 sample/batch size, calculated over 10 values of ζ' . We chose an optimal value of 0.15 for the DPS scale hyperparameter in this case.



Figure 10: Inpainting samples from the VP SDE on CIFAR-10. The observation model was ‘box’ mask with Gaussian ($\sigma_y = 0.05$) noise.



Figure 11: 2× nearest-neighbour super-resolution samples from the VP SDE on CIFAR-10. The observation model was Gaussian ($\sigma_y = 0.05$) noise.




Figure 12: $4\times$ bicubic super-resolution FID vs LPIPS (left) and SSIM (right) using the VE SDE on FFHQ-1k validation dataset for increasing observation noise.




Figure 13: ‘box’ mask inpainting FID vs LPIPS (left) and SSIM (right) using the VE SDE on FFHQ-1k validation dataset for increasing observation noise.




Figure 14: Whilst TMPD-D with the VE SDE remains robust to increasing noise, DPS-D and IIGDM-D do not. This is illustrated by samples from the different guidance methods for a $4\times$ super-resolution ($256 \times 256 \rightarrow 64 \times 64$) problem with increasing Gaussian observation noise distorting a ground truth image. The top row measurement has $\sigma_y = 0.01$, the middle row $\sigma_y = 0.05$ and the bottom row $\sigma_y = 0.1$. For the full results, see Table 9.




Figure 15: $4\times$ bicubic super-resolution samples from the VE SDE on FFHQ. The observation model was Gaussian ($\sigma_y = 0.05$) noise.

Table 10: Noisy observation inpainting and super-resolution for VE DDPM on CIFAR-10 1k validation set.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD-D	34.1	0.094 ± 0.044	0.006 ± 0.005	23.6 ± 3.3	0.782 ± 0.067
‘BOX’ MASK	DPS-D	40.6	0.085 ± 0.036	0.005 ± 0.004	24.2 ± 3.1	0.802 ± 0.064
INPAINTING	IGDM-D	40.0	0.076 ± 0.035	0.004 ± 0.003	25.0 ± 3.3	0.824 ± 0.066
$\sigma_y = 0.01$	TMPD-D	39.0	0.271 ± 0.069	0.028 ± 0.020	16.5 ± 3.1	0.582 ± 0.072
‘HALF’ MASK	DPS-D	42.8	0.247 ± 0.065	0.029 ± 0.020	16.5 ± 3.2	0.595 ± 0.077
INPAINTING	IGDM-D	42.2	0.237 ± 0.067	0.029 ± 0.020	16.4 ± 3.1	0.610 ± 0.079
$\sigma_y = 0.05$	TMPD-D	36.7	0.163 ± 0.070	0.007 ± 0.011	22.6 ± 2.8	0.724 ± 0.082
‘BOX’ MASK	DPS-D	101.5	0.224 ± 0.077	0.008 ± 0.004	21.8 ± 2.1	0.681 ± 0.087
INPAINTING	IGDM-D	99.9	0.225 ± 0.076	0.007 ± 0.003	21.8 ± 1.7	0.682 ± 0.093
$\sigma_y = 0.05$	TMPD-D	40.9	0.320 ± 0.078	0.029 ± 0.020	16.4 ± 2.9	0.544 ± 0.073
‘HALF’ MASK	DPS-D	105.5	0.362 ± 0.078	0.032 ± 0.021	15.7 ± 2.7	0.500 ± 0.075
INPAINTING	IGDM-D	97.1	0.366 ± 0.079	0.033 ± 0.020	15.4 ± 2.5	0.503 ± 0.079
$\sigma_y = 0.1$	TMPD-D	38.1	0.233 ± 0.086	0.009 ± 0.009	21.4 ± 2.5	0.652 ± 0.102
‘BOX’ MASK	DPS-D	80.4	0.313 ± 0.078	0.012 ± 0.006	19.7 ± 2.1	0.596 ± 0.088
INPAINTING	IGDM-D	140.5	0.378 ± 0.086	0.013 ± 0.003	18.8 ± 0.9	0.550 ± 0.113
$\sigma_y = 0.1$	TMPD-D	40.8	0.359 ± 0.086	0.030 ± 0.019	16.1 ± 2.8	0.495 ± 0.087
‘HALF’ MASK	DPS-D	99.5	0.434 ± 0.072	0.038 ± 0.023	14.8 ± 2.4	0.400 ± 0.083
INPAINTING	IGDM-D	138.8	0.476 ± 0.078	0.039 ± 0.019	14.5 ± 1.9	0.402 ± 0.095
$\sigma_y = 0.01$	TMPD-D	32.7	0.126 ± 0.058	0.004 ± 0.003	24.4 ± 2.9	0.828 ± 0.073
2× ‘NEAREST’	DPS-D	42.3	0.134 ± 0.053	0.004 ± 0.002	24.8 ± 2.5	0.839 ± 0.063
SUPER-RESOLUTION	IGDM-D	34.9	0.110 ± 0.042	0.004 ± 0.002	24.8 ± 2.8	0.839 ± 0.066
$\sigma_y = 0.01$	TMPD-D	38.6	0.295 ± 0.082	0.011 ± 0.006	20.2 ± 2.4	0.544 ± 0.109
4× ‘BICUBIC’	DPS-D	53.4	0.296 ± 0.074	0.011 ± 0.006	20.1 ± 2.3	0.547 ± 0.101
SUPER-RESOLUTION	IGDM-D	37.8	0.250 ± 0.077	0.011 ± 0.006	20.4 ± 2.5	0.577 ± 0.119
$\sigma_y = 0.05$	TMPD-D	34.0	0.204 ± 0.082	0.006 ± 0.004	23.1 ± 2.3	0.762 ± 0.089
2× ‘NEAREST’	DPS-D	98.8	0.283 ± 0.077	0.007 ± 0.003	21.9 ± 1.6	0.711 ± 0.081
SUPER-RESOLUTION	IGDM-D	99.6	0.298 ± 0.081	0.008 ± 0.003	21.1 ± 1.4	0.677 ± 0.092
$\sigma_y = 0.05$	TMPD-D	36.8	0.380 ± 0.097	0.015 ± 0.006	18.8 ± 2.0	0.440 ± 0.113
4× ‘BICUBIC’	DPS-D	126.1	0.467 ± 0.072	0.017 ± 0.007	18.2 ± 1.7	0.434 ± 0.096
SUPER-RESOLUTION	IGDM-D	104.2	0.419 ± 0.088	0.016 ± 0.006	18.1 ± 1.6	0.454 ± 0.117
$\sigma_y = 0.1$	TMPD-D	37.1	0.285 ± 0.100	0.008 ± 0.006	21.5 ± 2.1	0.657 ± 0.112
2× ‘NEAREST’	DPS-D	144.6	0.421 ± 0.082	0.012 ± 0.003	19.4 ± 1.1	0.571 ± 0.101
SUPER-RESOLUTION	IGDM-D	175.8	0.456 ± 0.085	0.016 ± 0.003	17.9 ± 0.9	0.514 ± 0.115
$\sigma_y = 0.1$	TMPD-D	36.1	0.439 ± 0.099	0.019 ± 0.008	17.5 ± 1.9	0.353 ± 0.115
4× ‘BICUBIC’	DPS-D	155.1	0.535 ± 0.067	0.022 ± 0.007	16.8 ± 1.3	0.360 ± 0.092
SUPER-RESOLUTION	IGDM-D	196.2	0.539 ± 0.078	0.030 ± 0.007	15.3 ± 1.1	0.327 ± 0.106

Table 11: Noisy observation inpainting and super-resolution for the reverse VE SDEs on CIFAR-10 1k validation set.

PROBLEM	METHOD	FID ↓	LPIPS ↓	MSE ↓	PSNR ↑	SSIM ↑
$\sigma_y = 0.01$	TMPD	40.0	0.102 \pm 0.047	0.005 \pm 0.004	23.7 \pm 3.1	0.773 \pm 0.069
‘BOX’ MASK	DPS	103.6	0.637 \pm 0.074	0.114 \pm 0.052	9.9 \pm 1.9	0.050 \pm 0.068
INPAINTING	HGDM	78.9	0.094 \pm 0.039	0.005 \pm 0.004	24.1 \pm 3.1	0.787 \pm 0.071
$\sigma_y = 0.01$	TMPD	45.8	0.279 \pm 0.066	0.030 \pm 0.029	16.3 \pm 3.0	0.569 \pm 0.069
‘HALF’ MASK	DPS	110.8	0.638 \pm 0.072	0.117 \pm 0.056	9.8 \pm 2.0	0.045 \pm 0.067
INPAINTING	HGDM	50.5	0.264 \pm 0.068	0.027 \pm 0.020	16.7 \pm 3.1	0.585 \pm 0.077
$\sigma_y = 0.05$	TMPD	45.3	0.167 \pm 0.065	0.007 \pm 0.024	22.6 \pm 2.7	0.718 \pm 0.078
‘BOX’ MASK	DPS	103.4	0.638 \pm 0.070	0.115 \pm 0.055	9.8 \pm 1.9	0.047 \pm 0.067
INPAINTING	HGDM	81.9	0.160 \pm 0.056	0.006 \pm 0.004	22.8 \pm 2.4	0.720 \pm 0.078
$\sigma_y = 0.05$	TMPD	51.1	0.319 \pm 0.071	0.029 \pm 0.019	16.2 \pm 2.8	0.532 \pm 0.071
‘HALF’ MASK	DPS	109.3	0.638 \pm 0.070	0.116 \pm 0.057	9.8 \pm 2.0	0.044 \pm 0.067
INPAINTING	HGDM	56.0	0.311 \pm 0.074	0.029 \pm 0.021	16.4 \pm 3.0	0.540 \pm 0.078
$\sigma_y = 0.1$	TMPD	47.6	0.231 \pm 0.079	0.008 \pm 0.005	21.5 \pm 2.3	0.650 \pm 0.097
‘BOX’ MASK	DPS	104.3	0.639 \pm 0.068	0.113 \pm 0.049	9.9 \pm 1.9	0.048 \pm 0.065
INPAINTING	HGDM	84.6	0.220 \pm 0.079	0.008 \pm 0.004	21.6 \pm 2.1	0.665 \pm 0.105
$\sigma_y = 0.1$	TMPD	54.1	0.366 \pm 0.080	0.032 \pm 0.020	15.8 \pm 2.7	0.480 \pm 0.085
‘HALF’ MASK	DPS	110.4	0.639 \pm 0.068	0.117 \pm 0.054	9.7 \pm 2.0	0.046 \pm 0.067
INPAINTING	HGDM	58.5	0.349 \pm 0.081	0.031 \pm 0.021	16.1 \pm 2.9	0.499 \pm 0.090
$\sigma_y = 0.01$	TMPD	43.5	0.141 \pm 0.070	0.007 \pm 0.034	24.1 \pm 3.1	0.810 \pm 0.092
2 \times ‘NEAREST’	DPS	118.7	0.641 \pm 0.066	0.117 \pm 0.054	9.7 \pm 1.9	0.048 \pm 0.065
SUPER-RESOLUTION	HGDM	59.4	0.258 \pm 0.083	0.007 \pm 0.003	22.0 \pm 2.2	0.716 \pm 0.075
$\sigma_y = 0.01$	TMPD	49.6	0.439 \pm 0.093	0.020 \pm 0.008	17.3 \pm 1.8	0.345 \pm 0.115
4 \times ‘BICUBIC’	DPS	128.3	0.642 \pm 0.071	0.122 \pm 0.059	9.6 \pm 2.0	0.046 \pm 0.068
SUPER-RESOLUTION	HGDM	52.9	0.561 \pm 0.083	0.042 \pm 0.015	14.1 \pm 1.7	0.167 \pm 0.085
$\sigma_y = 0.05$	TMPD	47.5	0.213 \pm 0.087	0.007 \pm 0.012	22.8 \pm 2.5	0.744 \pm 0.100
2 \times ‘NEAREST’	DPS	119.1	0.638 \pm 0.070	0.118 \pm 0.059	9.7 \pm 2.0	0.049 \pm 0.068
SUPER-RESOLUTION	HGDM	60.0	0.326 \pm 0.097	0.009 \pm 0.004	20.7 \pm 1.9	0.619 \pm 0.104
$\sigma_y = 0.05$	TMPD	51.2	0.379 \pm 0.094	0.016 \pm 0.029	18.6 \pm 2.1	0.428 \pm 0.117
4 \times ‘BICUBIC’	DPS	128.0	0.642 \pm 0.068	0.122 \pm 0.061	9.6 \pm 2.0	0.043 \pm 0.067
SUPER-RESOLUTION	HGDM	53.8	0.548 \pm 0.084	0.038 \pm 0.014	14.5 \pm 1.7	0.186 \pm 0.087
$\sigma_y = 0.1$	TMPD	51.6	0.292 \pm 0.097	0.009 \pm 0.020	21.3 \pm 2.2	0.646 \pm 0.116
2 \times ‘NEAREST’	DPS	120.5	0.644 \pm 0.074	0.121 \pm 0.061	9.7 \pm 2.0	0.048 \pm 0.068
SUPER-RESOLUTION	HGDM	61.9	0.386 \pm 0.100	0.012 \pm 0.005	19.5 \pm 1.9	0.523 \pm 0.117
$\sigma_y = 0.1$	TMPD	49.6	0.439 \pm 0.093	0.020 \pm 0.008	17.3 \pm 1.8	0.345 \pm 0.115
4 \times ‘BICUBIC’	DPS	128.3	0.642 \pm 0.071	0.122 \pm 0.059	9.6 \pm 2.0	0.046 \pm 0.068
SUPER-RESOLUTION	HGDM	52.9	0.561 \pm 0.083	0.042 \pm 0.015	14.1 \pm 1.7	0.167 \pm 0.085




Figure 16: Inpainting samples from the VE SDE on CIFAR-10. The observation model was ‘box’ mask with Gaussian ($\sigma_y = 0.05$) noise.




Figure 17: $2\times$ nearest-neighbour super-resolution samples from the VE SDE on CIFAR-10. The observation model was Gaussian ($\sigma_y = 0.05$) noise.