

# Reasoning Is Not All You Need: Examining LLMs for Multi-Turn Mental Health Conversations

Anonymous ACL submission

## Abstract

Limited access to mental healthcare, extended wait times, and increasing capabilities of Large Language Models (LLMs) has led individuals to turn to LLMs for fulfilling their mental health needs. However, examining the multi-turn mental health conversation capabilities of LLMs remains under-explored. Existing evaluation frameworks typically focus on diagnostic accuracy and win-rates and often overlook alignment with patient-specific goals, values, and personalities required for meaningful conversations. To address this, we introduce MedAgent, a novel framework for synthetically generating realistic, multi-turn mental health sensemaking conversations and use it to create the Mental Health Sensemaking Dialogue (MHSD) dataset, comprising over 2,200 patient–LLM conversations. Additionally, we present MultiSenseEval, a holistic framework to evaluate the multi-turn conversation abilities of LLMs in healthcare settings using human-centric criteria. Our findings reveal that frontier reasoning models yield below-par performance for patient-centric communication and struggle at advanced diagnostic capabilities with average score of  $\sim 31\%$ . Additionally, we observed variation in model performance based on patient’s persona and performance drop with increasing turns in the conversation. Our work provides a comprehensive synthetic data generation framework, a dataset and evaluation framework for assessing LLMs in multi-turn mental health conversations.

## 1 Introduction

With nearly 70% of individuals around the globe having limited to no access to mental healthcare (Kazdin and Rabbitt, 2013) and wait times for new patients extending up to three months (APA, 2023), traditional mental health care system is increasingly unable to meet the rising demand. At the same time, recent advancements in reasoning capabilities of LLMs has demonstrated significant per-

formance improvement on challenging tasks (OpenAI, 2024; Guo et al., 2025; Tu et al., 2025). This has led individuals to turn to LLMs to fill the gap in access to mental healthcare by allowing them to understand their personal mental health situations (Aydin et al., 2024). This process of sensemaking<sup>1</sup> encompasses a wide range of use-cases, from interpreting medical information to obtaining lifestyle recommendations and getting answers to health related questions. However, it remains unclear if LLMs can engage in meaningful multi-turn sensemaking conversations, especially pertaining to mental health conditions.

Single-turn settings that are often used for assessing LLM performance in healthcare domain do not portray the real-world complexities that often require iterative information gathering, proactive follow-ups, and shared decision-making (Dahm et al., 2022; Trevena et al., 2006). Recent works assessing multi-turn conversational capabilities of LLMs for general-purpose clinical conversations have reported significant decline in LLM performance for multi-turn dialogues (Li et al., 2024a; Liu et al., 2025b). However, such works have predominantly focused on the information gathering and diagnostic capabilities on general healthcare scenarios, often limiting evaluation to the point of diagnosis and reducing conversational capability analysis to classification or win-rate tasks. Filling this gap requires us to assess the capabilities of LLMs towards supporting multi-turn sensemaking conversations with the users through a more holistic approach taking account of patient-centric objective, values and needs. A major challenge in this direction is the limited to no access to real-world patient–clinician mental-health related conversation data that encompasses the full arc of interactions typically seen in clinical settings (such

<sup>1</sup>Sensemaking refers to “processes of interpretation and meaning production whereby individuals and groups interpret and reflect on phenomena” (Brown et al., 2008)




Figure 1: We present the MedAgent framework for generating realistic multi-turn mental health sensemaking conversations (**part (a)**). Using this framework we create the MHSD dataset with 2,284 conversations. Finally, we also present the MultiSenseEval framework (**part (b)**) to holistically evaluate LLM performance across patient-centric communication, conversational flow and correctness, diagnostic accuracy, and readability.

as the *6-function model* of medical communication (De Haes and Bensing, 2009; King and Hoppe, 2013)). Hence, our study is guided by the following research questions:

**RQ1:** How can we effectively generate synthetic multi-turn mental health related sensemaking conversations between patients and LLMs?

**RQ2:** How well do reasoning LLMs conduct multi-turn mental health sensemaking conversations, and how does the performance change by patient persona and conversation length?

To address these questions, we introduce MedAgent, a novel framework for synthetically generating multi-turn mental health conversations that are grounded in medical literature and portray real-world settings. MedAgent is model-agnostic and can be directly used with any LLM. Using MedAgent, we generate the **Mental Health Sense-making Dialogue (MHSD)** dataset with 2,284 synthetic conversations between patients and LLMs. Finally, we introduce MultiSenseEval framework for holistically assessing the multi-turn sensemaking conversations focusing on six axes: (1) *Perceived Susceptibility*, (2) *Perceived Severity*, (3) *Perceived Benefits*, (4) *Diagnostic Accuracy*, (5) *Conversation flow and Correctness*, and (6) *Text readability*.

Our findings reveal limited capabilities of frontier reasoning models towards patient-centric communication and diagnostic accuracy. In our evaluation, OpenAI o1 and DeepSeek-R1 obtained an average score of 2.55 and 2.77 (on a scale of 4) on patient-centric communication metrics and an accuracy score of  $\sim 31\%$  for exact diagnosis matching, highlighting the below-par performance for both models across evaluation axes. Our study also revealed, disparity in model performance based

on patient actor’s persona, with models performing better for patients with ‘Agreeable’ personality. Finally, we also observed that model performance for patient-centric communication and diagnostic metrics dropped with increasing number of turns in the conversation. Specifically, R1 exhibited a drop of 12.83%, 6.71%, and 29.6% for *Perceived Susceptibility*, *Perceived Severity*, and *Diagnostic Accuracy ‘Hard’* when the number of sensemaker messages increased from the 5 to 10-15. The proposed MultiSenseEval framework provides a more holistic approach towards assessing the quality of multi-turn mental health conversations. Additionally, recognizing community’s need for multi-turn conversation data, we will publicly release the MHSD dataset. Code and dataset for this paper can be found here.<sup>2</sup>

## 2 MedAgent Framework

Figure 1 (a) presents the overview of MedAgent framework. MedAgent framework generates mental health related sensemaking conversations using two actor LLMs: (1) Patient Actor LLM, and (2) Sensemaker Actor LLM. We use GPT-4o (OpenAI-GPT-4o, 2024) for the patient actor LLM, and frontier reasoning models, OpenAI o1 (OpenAI, 2024) and DeepSeek-R1 (Guo et al., 2025) respectively for the sensemaker actor LLM. For the sensemaker actor llm, we specifically used reasoning models as its processes (such as moving stages in the conversation, validating hypothesis) require advanced reasoning capabilities. In the following subsections we describe the details for both actor components of the MedAgent framework.

<sup>2</sup><https://anonymous.4open.science/r/ARR-July-2025-Submission-58-28CC/>

## 2.1 Patient Actor LLM

Real-world sensemaking conversations are influenced by patient-specific factors such as medical background, personality traits, health literacy, individual goals and intentions (Brown et al., 2008; Helms Mills et al., 2010). To reflect this diversity, we constructed diverse patient personas for the simulated patient actor LLM by creating sets of patient’s medical background, personality traits, health literacy, individual high-level intentions and concrete goals. Below, we describe each component of the patient actor LLM. Algorithm 1 presents the algorithm for generating the patient actor LLM message  $p_t$  at timestamp  $t$  based on the sensemaker message ( $s_{t-1}$ ), patient persona  $\Pi$ , and set of patient’s atomic facts  $F$  (detailed description of the pipeline is provided in Appendix E).

**Patient Background:** We began with curating a seed set of clinical case studies to serve for generating simulated patient dialogues. We used MedQA (Jin et al., 2021), to obtain US medical board exam case-studies and filtered 181 case studies focusing on patients with mental health-related conditions using a GPT-4o-based classifier followed by human validation. Building on past work (Min et al., 2023), we further decomposed each case study into a list of atomic medical facts and also extracted the corresponding diagnostic labels. These atomic facts serve as the foundation for constructing the final patient persona.

**Personality Traits:** To capture variability in personality, we grounded the patient personality traits based on the Big Five Personality Traits (Goldberg, 1993; Saucier, 1994; Cobb-Clark and Schurer, 2012), a psychological framework describing personality in terms of five broad dimensions: *Openness*, *Conscientiousness*, *Extraversion*, *Agreeableness*, and *Neuroticism*. To ensure behavioral consistency, we varied each trait one at a time on a binary scale, as past studies have shown varying correlation between the traits (Klimstra et al., 2013; Yu et al., 2021; Kang et al., 2023), and random sampling of traits may lead to internally contradictory personalities. Hence, for each case study, we created five personality variations in which one of the five personality traits is tuned to ‘High’ to modulate their conversational behavior. For instance, patients with ‘High’ *Agreeableness* tend to seek social harmony and use affiliative language. We have provided examples of other traits in Appendix A.

**Health Literacy:** Health literacy is described as

“personal knowledge and competencies which enable people to access, understand, appraise, and use information and services in ways which promote and maintain good health and well-being for themselves and those around them” (Nutbeam and Muscat, 2021). Individuals with inadequate health literacy have been observed to have difficulty in participating in medical decision-making process, following medical recommendations and conversations (Williams et al., 1995). To account for such variations, we introduced two health literacy levels (*Basic*, *Advanced*) for the patient persona (detailed definitions and prompts in Appendix B).

**Patient Intentions:** Past works have demonstrated that LLMs can support various patient needs and intentions related to patient education and sensemaking (Zaretsky et al., 2024; Bragazzi and Garbarino, 2024; Aydin et al., 2024). However, the patient backgrounds obtained from the MedQA dataset focused on diagnosis and lacked explicit representations of such intentions. To address this, we augmented each patient profile with realistic, intention-driven motivations for engaging with LLMs in sensemaking conversations. Building upon Aydin et al. (2024) and Li et al. (2024b), we used three different intentions (detailed definitions and examples are provided in Appendix C). To operationalize this augmentation, we employed GPT-4o to assign the most realistic and logical intention from one of the three intentions to the given patient background (prompt in Table 19).

**Patient Goals:** Patient goals refer to what they want to have achieved at the end of the conversation. However, similar to patient intentions, patient backgrounds lacked concrete goals for the conversations. To address this, we used three broad categories of goals as mentioned in Auriemma et al. (2024), (1) *Comfort-focused*, (2) *Maintain or improve function*, and (3) *Life extension* (detailed definitions and examples in Appendix D). These goals focus on three different aspects patient-oriented care. For instance, while the *Comfort-focused* goal aims to seek interventions to promote comfort and avoiding those that would increase discomfort, *Life extension* deals with the patient’s goal of extending longevity or survival even at the expense of maximizing function or comfort. Similar to the patient intention augmentation, we used GPT-4o to assign the most relevant and logical patient goal to the patient background (prompt in Table 18).



---

**Algorithm 1** PATIENTSTEP( $s_{t-1}$ ,  $\Pi$ ,  $\mathcal{F}$ )**Input:** Sensemaker utterance  $s_{t-1}$ , patient profile  $\Pi$ , set of patient’s atomic facts  $F$ **Output:**  $p_t$  // patient reply at turn  $t$ 

```
1 // Match message mentioned by the sensemaker against patient
  facts.
2  $\mathcal{F}_{\text{match}} \leftarrow \text{MatchFacts}(s_{t-1}, F)$ 
  // Check #facts matched
3 if  $|\mathcal{F}_{\text{match}}| > 0$  then
4   // Generate patient messages based on matched facts.
   // Apply stylistic variations based on personality, health
   // literacy, and goals/intents defined in  $\Pi$ .
5    $p_t \leftarrow \text{GenPatientMsg}(\mathcal{F}_{\text{match}}, \Pi)$ 
6 else
7   // Determine whether the requested information is back-
   // ground information
8   if  $\text{RequestType}(s_{t-1}) == \text{background}$  then
9     // Apply closed world assumption. Indicating lack of
     // information.
10     $p_t \leftarrow \text{ClosedWorldMessage}()$ 
11  else
12     $p_t \leftarrow \text{ReplyWithContext}(s_{t-1}, \Pi)$  // reply using
    // general context and style
13 return  $p_t$ 
```

---

## 2.2 Sensemaker Actor LLM

The goal of the sensemaker actor LLM is to produce informative, non-redundant, and stage-appropriate responses that emulate the structure of clinical conversations. Figure 1 (a) shows the high-level sensemaker message generation pipeline, which includes four modules: (1) Next Stage Determination, (2) Fact Extraction & Matching, (3) Sensemaker Message Generation, and (4) Message Redundancy Checker & Selector. One of the novel features in the generated conversations is the existence of a structured organization of the sensemaking conversation into five stages inspired by *6-function model* of medical communication (De Haes and Bensing, 2009; King and Hoppe, 2013). Hence, we first describe the stages of the sensemaking conversation followed by the overall working of the sensemaker actor LLM.

**Stages of sensemaking conversation:** The sensemaker actor LLM organizes each conversation grounded within the *6-function model* of medical communication (De Haes and Bensing, 2009; King and Hoppe, 2013) (example conversations for each stage are in Appendix Table 7). The *Fostering the Relationship* stage initiates the encounter by building trust, empathy, and psychological safety. The *Gathering Information* stage focuses on eliciting both biomedical and psychosocial context through open-ended questions and active listening, aligning with how human physicians derive diagnostic hy-

potheses. In the *Providing Information* stage, the sensemaker provides its diagnosis and explains it to the patient. The *Decision Making* stage facilitates shared planning by integrating patient preferences and values. We combined the *Enabling Disease and Treatment-related Behavior* stage with the *Decision Making* stage due to the similar objectives within our context. Finally, the *Responding to Emotions* stage enables agents to detect and attend to emotional distress. This phase-wise design not only mirrors real clinical workflows but also allows the sensemaker LLM to specialize in communication tasks that directly impact diagnostic accuracy, treatment adherence, and health outcomes. This design ensures that the sensemaker LLM is able to meaningfully engage in a multi-turn conversation.

---

**Algorithm 2** Sensemaker Message Generation at Turn  $t$ **Input:** patient message  $p_{t-1}$ , known facts  $F_{t-1}$ , sensemaker message history  $H$ **Output:** Next sense-maker utterance  $s_t$ , updated facts  $F_t$ , stage index  $g_t$ 

```
14 Step 1: Stage Determination & Fact Update
   // Extract atomic facts from patient message
15  $\Delta F_t \leftarrow \text{ExtractFacts}(p_{t-1})$ 
   // Merge with existing facts
16  $F_t \leftarrow F_{t-1} \cup \text{MatchFacts}(\Delta F_t, F_{t-1})$ 
   // Determine whether to move to next stages
17  $\tilde{g}_t \leftarrow \text{GetNextStage}(g_{t-1}, p_{t-1}, F_t)$ 
   if #turns in current stage  $g_{t-1} \geq 5$  then
18    $g_t \leftarrow g_{t-1} + 1$  // Move to the next stage
19 else
20    $g_t \leftarrow \tilde{g}_t$  // Assign the computed next stage
21 Step 2: Candidate Generation
   // Generate  $|C| = 3$  candidate messages
22  $C \leftarrow \text{GenerateCandidates}(g_t, F_t, H)$ 
23 Step 3: Redundancy Check
   // Check each candidate message against past messages for
   // redundancy.
24  $C' \leftarrow \text{Filter}(C, H)$ 
25 if  $|C'| > 0$  then
26    $s_t \leftarrow \text{Sample}(C')$  // sample one reply
27    $H \leftarrow H \cup \{s_t\}$ 
28 else
29    $g_t \leftarrow g_t + 1$  // avoid stagnation
30 return  $(s_t, F_t, g_t)$ 
```

---

## Sensemaker Actor LLM Message Generation

**Pipeline:** Algorithm 2 presents the algorithm for generating message  $s_t$  at timestamp  $t$  given the patient message  $p_{t-1}$ , known facts about the patients until timestamp  $t$  ( $F_{t-1}$ ), and sensemaker message history  $H$ . After the sensemaker receives the patient’s message  $p_{t-1}$ , it first invokes the *Next Stage Determination* module. This module takes as input the current message  $p_{t-1}$ , known patient facts from the previous stages  $F_{t-1}$ , and the sensemaker’s

previous messages  $H$  and passes  $p_{t-1}$  to *Fact Extraction and Matching*, which extracts new atomic facts  $\Delta t$  and update the fact set  $F_t$ . Based on the updated facts  $F_t$  and sensemaker’s previous message history  $H$ , the *Next Stage Determination* module decides whether to remain in the current stage of the conversation or advance to the next stage of the conversation. The information about the current stage  $g_t$ , patient facts  $F_t$ , and sensemaker message history  $H$  are passed to the *Sensemaker Message Generation* module. This module then produces 3 candidate messages. The candidate messages are then sent to the *Message Selection and Redundancy Checking* module, which filters out replies that are semantically similar to prior sensemaker messages ( $H$ ). If non-redundant messages exist, one of such message is randomly selected as the output  $s_t$ . Otherwise, the conversation moves to the next stage. (Refer to Appendix F, G, H and I for prompt details).

### 3 MHSD Dataset

After initial post-processing to remove erroneous generations and incomplete conversations, we obtained 1,142 conversations each for DeepSeek-R1 and OpenAI o1 as the sensemaker in the MHSD dataset (total conversations: 2,284). We used GPT-4o for all modules of the patient actor, and for all sensemaker modules excluding next stage determination and answer generation (which used the reasoning models). Table 1 presents a stage-wise breakdown of conversations. o1 conversations had 13.65 messages on average, while R1 conversations were longer, with 19.09 messages per conversation on average. o1 model also spent longer on gathering information from the patient before decision making (7.64 messages) compared to R1 (3.88 messages). An end-to-end sample conversation is shown in Figure 2.

Stage	Avg. Number of Messages	
	OpenAI o1	DeepSeek-R1
Fostering the Relationship	1.11	1.22
Gathering Information	7.64	3.88
Providing Information	2.00	2.00
Decision Making	3.52	2.70
Responding to Emotions	3.83	2.86
Exit	1.00	1.00
Overall	19.09	13.65

Table 1: Stage-wise and overall average number of messages of the MHSD dataset with OpenAI o1 and DeepSeek-R1 as sensemaker, including both patient and sensemaker responses.




Figure 2: Sample conversation between the sensemaker and a patient with high conscientiousness (traits listed by first letter) and basic medical literacy. Stages are distinguished by color, with some intermediate dialogues skipped for conciseness. Diagnosis is in bold.

### 4 Sensemaking Evaluation

In this section, we outline our approach for evaluating the sensemaking capabilities of LLMs. We evaluate the multi-turn conversations generated by two reasoning models: (1) OpenAI o1 (OpenAI, 2024), and (2) DeepSeek-R1 (Guo et al., 2025) using a hybrid approach combining automated metrics and human evaluation. While automated evaluation allows us to evaluate the conversations on scale, human evaluation provides an validation of automated metrics.

**Automated Evaluation:** For automated evaluation we used a combination of multiple metrics assessing various aspects related to the sense-making conversation such as patient-centric communication, diagnostic accuracy, conversation flow and correctness, readability. For *patient-centric communication*, we adopted metrics from the Health Belief Model (HBM) (Champion et al., 2008; Janz and Becker, 1984) – a psychological framework that seeks to explain and predict individual health behaviors by examining individuals’ attitudes and beliefs. Specifically, we use three dimensions from HBM namely perceived susceptibility, perceived severity, perceived benefits. To assess *diagnostic accuracy*, we used ground truth diagnostic labels from the seed case studies obtained from Jin et al. (2021). In contrast to the past works (Li et al.,

2024a; Liu et al., 2025b), we assessed diagnostic accuracy without providing multi-choice options to better reflect real-world conditions. Since, sensemaker’s diagnosis is provided in a free-text generation, we used two different accuracy assessment settings: (1) Hard Diagnostic Accuracy, requiring exactly matching the sensemaker’s diagnosis with the ground truth, and (2) Soft Diagnostic Accuracy which allowed for a broader category level matches for diagnosis. **Conversation flow and Correctness** measures the ability of sensemaker LLM to perform the end-to-end conversation maintaining the logical consistency, covering each phase of the conversation appropriately. Given, that the sensemaker LLM generates open-ended messages, we used LLM-as-a-judge approach to measure the aforementioned metrics. Finally, we measure the overall **readability** of the responses provided by the sensemaker LLM using the SMOG index, a popular readability index to assess health literacy material. (Mc Laughlin, 1969). Detailed information related to the definition, sub-criteria, evaluation rubric and prompts can be found in Appendix K.

**Human Evaluation:** In addition to the automated evaluation, we also conducted human-evaluation as a validation check for automated metric scores on a subset of 100 conversations. Specifically, we used the same rubric as the automated evaluation for patient-centric communication, diagnostic accuracy, and conversation flow and correctness. We have provided details regarding the human evaluation in Appendix L.

## 5 Results

### 5.1 MultiSenseEval Framework Evaluation

**Patient-Centric Communication:** Assessing the patient-centric communication capabilities of OpenAI o1 and DeepSeek R1, we observed that R1 outperformed o1 across all three dimensions. However, the mean average score for both models remained below 3 on a 4-point scale (1: Very Poor to 4: Very Good) across all metrics, indicating the below-par performance towards appropriately communicating with the patient. For *Perceived Susceptibility*, R1 obtained a significantly higher average score of 2.59 (SD = 0.75) compared to score of 2.45 (SD = 0.63) obtained by o1 (Mann–Whitney U test;  $p < 0.001$ ;  $U=597057$ ). Similarly, R1 obtained significantly higher scores for *Perceived Severity* (2.75; SD=0.67) compared to o1 (2.56; SD=0.54) (Mann–Whitney U test;

$p < 0.001$ ;  $U=559539$ ). Lastly, R1 also obtained significantly higher score (3.04; SD=0.77) compared to o1 (2.76; SD=0.71) for the *Perceived Benefits* dimension (Mann–Whitney U test;  $p < 0.001$ ;  $U=520950$ ). To qualitatively assess the shortcomings, we analyzed the lowest scoring conversations (score of 1 or 2) across all three metrics, revealing that reasoning models still lack personalization during mental health multi-turn conversations (detailed analysis in Appendix J).




Figure 3: Performance comparison of OpenAI o1 and DeepSeek-R1 across **Perceived Susceptibility**, **Perceived Severity**, **Perceived Benefits**, and **Conversation Flow and Correctness**. Bars indicate mean scores with 95% confidence interval. All scores are on a 4-point Likert scale ((1): *Very Poor* to (4): *Very Good*). Both models obtain scores below “Good Performance” rating for the three patient-centric communication metrics, but exceed the “Good Performance” threshold for Conversation Flow and Correctness.

**Conversation Flow and Correctness:** Both OpenAI o1 and DeepSeek R1’s scores exceeded the “Good Performance” threshold ( $\geq 3$  on a 4-point Likert scale), indicating that both models covered all five stages in the correct order with reasonable depth (as described in 2.2). Interestingly, here we observed o1 outperforming R1 with models obtaining scores of 3.53 (SD=0.67) and 3.49 (SD=0.72) respectively. Similarly for **Text Readability**, we observed that o1 produced significantly easier to read messages with a SMOG score of 11.50 (SD=1.26) compared to a score of 13.33 (SD=1.47) for DeepSeek R1 (t-test;  $p < 0.001$ ;  $t=-31.95$ ). However, health organizations recommend SMOG scores between 6-8 (Badarudeen and Sabharwal, 2010), which is significantly lower than the SMOG scores produced by both models.

**Diagnostic Accuracy:** For the “Soft” setting evaluation, o1 significantly outperformed R1 (o1: acc.score= 68.39%; SD=46.52%, R1: acc.score= 64.36%; SD=47.91%)(Mann–Whitney U test;  $p < 0.05$ ;  $U=678348$ ). However, under the “Hard” setting, both models’ performance dropped by over 50%, with o1 obtaining an accuracy score of 31.09% (SD=46.31%) and R1 obtaining a score



Personality Trait	Model	Perceived Susceptibility	Perceived Severity	Perceived Benefits	Flow Correctness	Diag. Acc. (Hard) (%)	Diag. Acc. (Soft) (%)	SMOG
Basic Health Literacy Patient								
Openness	o1	2.46	2.62	2.91	3.53	31.68	70.30	11.22
	R1	2.58	2.71	<b>3.25</b>	3.56	34.65	58.42	13.27
Agreeableness	o1	2.42	2.44	2.85	<b>3.68</b>	<b>36.43</b>	<b>70.54</b>	11.13
	R1	2.60	2.74	3.04	3.50	26.36	59.69	13.06
Neuroticism	o1	2.19	2.43	2.41	3.24	31.33	67.47	<b>10.92</b>
	R1	<b>2.66</b>	<b>2.81</b>	2.88	3.28	28.92	59.04	12.90
Advanced Health Literacy Patient								
Openness	o1	2.56	2.63	2.82	3.59	29.37	<b>72.22</b>	11.88
	R1	2.63	2.79	3.11	<b>3.64</b>	28.57	69.05	13.57
Agreeableness	o1	2.50	2.57	2.84	3.62	32.81	68.75	11.79
	R1	<b>2.67</b>	<b>2.81</b>	<b>3.12</b>	3.60	30.47	66.41	13.56
Neuroticism	o1	2.39	2.57	2.47	3.30	<b>39.77</b>	62.50	<b>11.53</b>
	R1	2.48	2.60	2.89	3.32	34.09	57.95	13.31

Table 2: Performance of o1 vs. R1 across seven metrics, by persona and health-literacy level for three personality traits. We report best performing model metrics in each scenario in **bold**. Dia. Acc. represents the Diagnostic Accuracy metric. Full table with all five personality traits is provided in Appendix section M (Table 30).




Figure 4: Performance comparison between OpenAI o1 and DeepSeek R1 across **Hard Diagnostic Accuracy**, and **Soft Diagnostic Accuracy**. Bars indicate the mean scores with 95% confidence interval. As observed the performance for both models drops by more than 50% when the diagnosis is matched exactly with the ground truth (“Hard Accuracy”) in comparison to when it is matched on broader/general criteria (“Soft Accuracy”).

of 31.18% (SD=46.34%). These findings highlight the limited diagnostic capabilities of frontier reasoning models within multi-turn conversations requiring probing, consistent with prior findings (Li et al., 2024a).

## 5.2 Performance Disparity based on Patient Persona

To understand how differences in patient persona may impact model’s performance, we stratified the OpenAI o1 and DeepSeek R1’s evaluation results across all performance metrics (ref. Section 4) based on patient’s dominant personality traits (Openness, Conscientiousness, Extraversion, Agreeableness, Neuroticism) and health literacy level (Basic/Advanced). Table 2 presents results across these strata for three personality types.

**Impact of Personality Traits:** Among patient with advanced health literacy, models performed best along the *Perceived Susceptibility*, *Perceived Severity*, and *Perceived Benefits* metrics for patient with dominant personality trait as ‘Agreeableness’. For

instance, R1 achieved scores of 2.67 (susceptibility), 2.81 (severity), and 3.12 (benefits) for patients belonging to this cohort. In contrast, R1’s performance dropped by 7.1%, 7.5%, and 7.4% for the aforementioned metrics for patients with same literacy skills but with ‘Neuroticism’ personality trait. We observed, a similar trend among patients with basic health literacy, where models performed better on *Flow Correctness* and *Diagnostic Accuracy* metrics for patients with ‘Agreeableness’ personality. These findings reveal the performance gap towards patient-centric communication, where models performed better for patients with higher degree of prosocial and cooperative behavior (“Agreeableness”) compared to patients with other personality profiles. Interestingly, o1 generated easier to read messages for patients with ‘Neuroticism’ dominant personality compared to other personality type. For such patients, the SMOG score for o1 responses were 2.7% and 2.95% lower compared to that for patients with ‘Openness’ personality with basic and advanced literacy skills.

**Impact of Health Literacy:** Comparing model performance across patients with basic and advanced health literacy, we observed gain in o1 (+2.77%) and R1’s (+0.63%) performance for patient-centric communication metrics. Interestingly we observed contrasting trends in performance, for o1 the accuracy decreased by 0.12% for “soft” and 0.73% “hard” diagnostic accuracy settings, whereas we observed an increase of 10.16% and 11.74% in “soft” and “hard” diagnostic accuracy for R1. Additionally, both models produced harder to read text when interacting with patient actors with higher literacy, with SMOG scores increasing by 5.84% and 3.29% for o1 and R1 respectively. These findings highlight a trade off in performance of models,




Figure 5: Performance trend for o1 and R1 across the MultiSenseEval framework metrics, the x-axis indicates sensemaker message count bins. While on average performance on patient-centric metrics and diagnostic accuracy declined with longer conversations, flow correctness and readability improved.

while models may provide better patient-centric messages to patients with higher health literacy, those messages could be harder to read.

### 5.3 Impact of Conversation Length on Performance

We also examined the change in model performance with increased conversation length. Figure 5 presents the line plots for o1 and R1 performance for each evaluation metric across number of sensemaker messages in the conversation. Interestingly, for R1, we observed that the performance decreased by 12.83%, 6.71%, and 29.6% for *Perceived Susceptibility*, *Perceived Severity*, and *Diagnostic Accuracy* respectively when the number of sensemaker messages increased from the 5 to 10-15. We observed a similar trend for o1 with a performance drop of 4.8% and 7.99% for *Perceived Susceptibility* and *Perceived Benefits* respectively, highlighting the reduced performance among both models for patient-centric communication with increased number of messages. In contrast, both models showed improvement in performance for the *Conversation Flow Correctness* and *Text Readability* performance with increased number of turns. These findings further highlight that even frontier reasoning models struggle to sustain effective performance over longer conversations, a finding highlighted in contemporary work (Laban et al., 2025).

## 6 Related Work

**LLM Evaluation for Single Healthcare Conversations:** Recent advancements in large language models (LLMs) have enabled diverse applications among clinicians (Tu et al., 2025; Singhal et al., 2023b) and patients (Yang et al., 2023). Models such as Med-PaLM (Tu et al., 2024) and OpenAI’s GPT series (OpenAI, 2025) have demonstrated strong performance on various medical benchmarks (Nori et al., 2023; Singhal et al., 2023a,b), highlighting their promise in clinical settings. However, critical challenges remain, including hallucinations (Agarwal et al., 2024; Asgari et al., 2025;

Deng et al., 2024) cross-lingual disparities (Jin et al., 2024), sociocultural biases (e.g. gender and geographic) (Restrepo et al., 2024; Liu et al., 2025a), and misalignment with experts (Chandra et al., 2025), and limitations in clinical competency tests (Thirunavukarasu et al., 2023; Jin et al., 2023).

### Multi-turn Conversation Evaluation for LLMs:

Beyond single-turn settings, recent works have also explored LLMs’ multi-turn clinical conversations abilities. For example, MediQ (Li et al., 2024a) evaluates question-asking ability of LLMs within multi-turn conversations. Along similar lines, researchers proposed a proactive dialogue generation framework based on dialogue ranking (Li et al., 2025). The Ask Patients with Patience (APP) framework (Zhu and Wu, 2025) allow LLMs to generate multi-turn conversations based on medical guidelines and entropy minimization. In contrast, (Liu et al., 2025b) proposed a patient simulator for multi-turn diagnostic conversations. However, past works overlook the diagnostic capabilities of LLMs in multi-turn conversations.

## 7 Conclusion

In this work, we proposed MedAgent, a framework for creating realistic mental health sensemaking conversations. Using MedAgent, we created MHSD dataset consisting of 2,284 synthetic sensemaker-patient conversations. Finally, we presented the MultiSenseEval framework for holistically evaluating multi-turn mental health conversation using metrics grounded in clinical research. Our findings reveal that frontier reasoning models yield below-par performance for patient-centric communication. Additionally, our study reveal the performance disparity among models based on patient’s persona and reduction in model performance with increase number of turns in the conversation. Our work provides a comprehensive method for creating synthetic clinical conversations, a dataset and evaluation framework for assessing LLMs in multi-turn mental health conversations.



## 8 Limitations

While our work presents novel contribution on multiple fronts, it is important to acknowledge the limitations of our work. While MedAgent framework provides a method for generating realistic mental health sensemaking conversations, it is important to recognize both patient and sensemaker actors are modeled by LLMs which may not be able to capture the full range of emotions, variability and lived experiences of human patients and healthcare providers. Additionally, while grounding the conversations on the *6-function model* (De Haes and Bensing, 2009; King and Hoppe, 2013) provides clinically sound structure to the conversation, the number of messages exchanged in each phase can vary by a great extent in real-world setting, something which is not fully captured in MHSD dataset. Finally, there may exist additional evaluation axes for evaluating multi-turn mental health conversations, which this work may have missed. Despite these challenges, our work provides a robust framework for generating synthetic clinical conversations grounded in real world settings, a large-scale conversational dataset, and a robust framework for assessing the capabilities of LLMs in multi-turn mental health conversations.

## 9 Ethical considerations

We used the clinical case studies from Jin et al. (2021) which are publicly available for use under the MIT License. We further ensured that all data used was de-identified. As our study did not involve working with human participants, the Institutional Review Board (IRB) classified it as non-human subjects research, exempting it from IRB approval. We aim to provide the code and MHSD dataset publicly available on a GitHub repository upon acceptance of the paper.

Our study presents a systematic approach for generating synthetic data for mental health conversations and evaluating LLM performance for such conversations. However, our framework is designed for research & prototyping purposes and is not intended to replace licensed medical professionals. Using our framework without guardrails and proper oversight in real-world settings could pose risks to patient safety. Additionally, our findings should be complemented with thorough human evaluation to ensure the reliability and safety of models in real-world settings.

## References

- Vibhor Agarwal, Yiqiao Jin, Mohit Chandra, Munmun De Choudhury, Srijan Kumar, and Nishanth Sasstry. 2024. Medhalu: Hallucinations in responses to healthcare queries by large language models. *arXiv:2409.19492*.
- APA. 2023. Psychologists reaching their limits as patients present with worsening symptoms year after year. <https://www.npr.org/sections/health-shots/2023/12/06/1217487323/psychologists-waitlist-demand-mental-health-care>. [Accessed 12-05-2025].
- Elham Asgari, Nina Montaña-Brown, Magda Dubois, Saleh Khalil, Jasmine Balloch, Joshua Au Yeung, and Dominic Pimenta. 2025. A framework to assess clinical safety and hallucination rates of llms for medical text summarisation. *npj Digital Medicine*, 8(1):1–15.
- Catherine L Auriemma, Anne Song, Lake Walsh, Jason J Han, Sophia R Yapalater, Alexander Bain, Lindsay Haines, Stefania Scott, Casey Whitman, Stephanie P Taylor, and 1 others. 2024. Classification of documented goals of care among hospitalized patients with high mortality risk: a mixed-methods feasibility study. *Journal of General Internal Medicine*, 39(10):1839–1849.
- Serhat Aydin, Mert Karabacak, Victoria Vlachos, and Konstantinos Margetis. 2024. Large language models in patient education: a scoping review of applications in medicine. *Frontiers in Medicine*, 11:1477898.
- Sameer Badarudeen and Sanjeev Sabharwal. 2010. Assessing readability of patient education materials: current role in orthopaedics. *Clinical Orthopaedics and Related Research*, 468:2572–2580.
- Nicola Luigi Bragazzi and Sergio Garbarino. 2024. Assessing the accuracy of generative conversational artificial intelligence in debunking sleep health myths: mixed methods comparative study with expert analysis. *JMIR Formative Research*, 8(1):e55762.
- Andrew D Brown, Patrick Stacey, and Joe Nandhakumar. 2008. Making sense of sensemaking narratives. *Human relations*, 61(8):1035–1062.
- Victoria L Champion, Celette Sugg Skinner, and 1 others. 2008. The health belief model. *Health behavior and health education: Theory, research, and practice*, 4:45–65.
- Mohit Chandra, Siddharth Sriraman, Gaurav Verma, Harneet Singh Khanuja, Jose Suarez Campayo, Zihang Li, Michael L. Birnbaum, and Munmun De Choudhury. 2025. Lived experience not found: LLMs struggle to align with experts on addressing adverse drug reactions from psychiatric medication use. In *Proceedings of the 2025 Conference of the Nations of the Americas Chapter of the Association for Computational Linguistics: Human Language*

703	<i>Technologies (Volume 1: Long Papers)</i> , pages 11083–	Weixi Kang, Francois Steffens, Sònia Pineda, Kaya	758
704	11113, Albuquerque, New Mexico. Association for	Widuch, and Antonio Malvaso. 2023. Personality	759
705	Computational Linguistics.	traits and dimensions of mental health. <i>Scientific</i>	760
		<i>Reports</i> , 13(1):7091.	761
706	Deborah A Cobb-Clark and Stefanie Schurer. 2012. The	Alan E Kazdin and Sarah M Rabbitt. 2013. Novel mod-	762
707	stability of big-five personality traits. <i>Economics</i>	els for delivering mental health services and reducing	763
708	<i>Letters</i> , 115(1):11–15.	the burdens of mental illness. <i>Clinical Psychological</i>	764
		<i>Science</i> , 1(2):170–191.	765
709	Carolyn Cutilli and Ian Bennett. 2009. <a href="#">Understanding</a>	Ann King and Ruth B Hoppe. 2013. “best practice” for	766
710	<a href="#">the health literacy of america results of the national</a>	patient-centered communication: a narrative review.	767
711	<a href="#">assessment of adult literacy</a> . <i>Orthopaedic nursing /</i>	<i>Journal of graduate medical education</i> , 5(3):385–	768
712	<i>National Association of Orthopaedic Nurses</i> , 28:27–	393.	769
713	32; quiz 33.		
714	Maria R Dahm, Maureen Williams, and Carmel Crock.	Theo A Klimstra, Wiebke Bleidorn, Jens B Asendorpf,	770
715	2022. ‘more than words’–interpersonal communi-	Marcel AG van Aken, and Jaap JA Denissen. 2013.	771
716	cation, cognitive bias and diagnostic errors. <i>Patient</i>	Correlated change of big five personality traits across	772
717	<i>Education and Counseling</i> , 105(1):252–256.	the lifespan: A search for determinants. <i>Journal of</i>	773
		<i>Research in Personality</i> , 47(6):768–777.	774
718	Hanneke De Haes and Jozien Bensing. 2009. End-	Philippe Laban, Hiroaki Hayashi, Yingbo Zhou, and	775
719	points in medical communication research, proposing	Jennifer Neville. 2025. Llms get lost in multi-turn	776
720	a framework of functions and outcomes. <i>Patient edu-</i>	conversation. <i>arXiv preprint arXiv:2505.06120</i> .	777
721	<i>cation and counseling</i> , 74(3):287–294.		
722	Chengyuan Deng, Yiqun Duan, Xin Jin, Heng Chang,	Stella Li, Vidhisha Balachandran, Shangbin Feng,	778
723	Yijun Tian, Han Liu, Henry Peng Zou, Yiqiao Jin,	Jonathan Ilgen, Emma Pierson, Pang Wei W Koh,	779
724	Yijia Xiao, Yichen Wang, and 1 others. 2024. De-	and Yulia Tsvetkov. 2024a. Mediq: Question-asking	780
725	constructing the ethics of large language models	llms and a benchmark for reliable interactive clinical	781
726	from long-standing issues to new-emerging dilem-	reasoning. <i>NeurIPS</i> , 37:28858–28888.	782
727	mas. <i>arXiv:2406.05392</i> .		
728	Lewis R Goldberg. 1993. The structure of phenotypic	Xueshen Li, Xinlong Hou, Nirumapa Ravi, Ziyi Huang,	783
729	personality traits. <i>American psychologist</i> , 48(1):26.	and Yu Gan. 2025. A two-stage proactive dialogue	784
		generator for efficient clinical information collection	785
730	Daya Guo, Dejian Yang, Haowei Zhang, Junxiao	using large language model. <i>Expert Systems with</i>	786
731	Song, Ruoyu Zhang, Runxin Xu, Qihao Zhu, Shi-	<i>Applications</i> , page 127833.	787
732	rong Ma, Peiyi Wang, Xiao Bi, and 1 others. 2025.		
733	Deepseek-r1: Incentivizing reasoning capability in	Yu-Hao Li, Yu-Lin Li, Mu-Yang Wei, and Guang-Yu	788
734	llms via reinforcement learning. <i>arXiv preprint</i>	Li. 2024b. <a href="#">Innovation and challenges of artificial</a>	789
735	<i>arXiv:2501.12948</i> .	<a href="#">intelligence technology in personalized healthcare</a> .	790
		<i>Scientific Reports</i> , 14.	791
736	Jean Helms Mills, Amy Thurlow, and Albert J Mills.	Shudong Liu, Yiqiao Jin, Cheng Li, Derek F Wong,	792
737	2010. Making sense of sensemaking: the critical	Qingsong Wen, Lichao Sun, Haipeng Chen, Xing	793
738	sensemaking approach. <i>Qualitative research in orga-</i>	Xie, and Jindong Wang. 2025a. Culturevlm: Char-	794
739	<i>nizations and management: An international journal</i> ,	acterizing and improving cultural understanding	795
740	5(2):182–195.	of vision-language models for over 100 countries.	796
		<i>arXiv:2501.01282</i> .	797
741	Nancy K Janz and Marshall H Becker. 1984. The health	Zhaocheng Liu, Quan Tu, Wen Ye, Yu Xiao, Zhishou	798
742	belief model: A decade later. <i>Health education quar-</i>	Zhang, Hengfu Cui, Yalun Zhu, Qiang Ju, Shizheng	799
743	<i>terly</i> , 11(1):1–47.	Li, and Jian Xie. 2025b. Exploring the inquiry-	800
744	Di Jin, Eileen Pan, Nassim Oufattole, Wei-Hung Weng,	diagnosis relationship with advanced patient simula-	801
745	Hanyi Fang, and Peter Szolovits. 2021. What disease	tors. <i>arXiv preprint arXiv:2501.09484</i> .	802
746	does this patient have? a large-scale open domain		
747	question answering dataset from medical exams. <i>Ap-</i>	G Harry Mc Laughlin. 1969. Smog grading-a new read-	803
748	<i>plied Sciences</i> , 11(14):6421.	ability formula. <i>Journal of reading</i> , 12(8):639–646.	804
749	Haoan Jin, Siyuan Chen, Mengyue Wu, and Kenny Q	Sewon Min, Kalpesh Krishna, Xinxu Lyu, Mike Lewis,	805
750	Zhu. 2023. Psyeval: A comprehensive large language	Wen-tau Yih, Pang Koh, Mohit Iyyer, Luke Zettle-	806
751	model evaluation benchmark for mental health. <i>arXiv</i>	moyer, and Hannaneh Hajishirzi. 2023. <a href="#">FActScore:</a>	807
752	<i>preprint arXiv:2311.09189</i> .	<a href="#">Fine-grained atomic evaluation of factual precision</a>	808
753	Yiqiao Jin, Mohit Chandra, Gaurav Verma, Yibo Hu,	<a href="#">in long form text generation</a> . In <i>Proceedings of the</i>	809
754	Munmun De Choudhury, and Srijan Kumar. 2024.	<i>2023 Conference on Empirical Methods in Natural</i>	810
755	Better to ask in english: Cross-lingual evaluation of	<i>Language Processing</i> , pages 12076–12100, Singa-	811
756	large language models for healthcare queries. In <i>Web</i>	pore. Association for Computational Linguistics.	812
757	<i>Conference</i> , pages 2627–2638.		

813	Harsha Nori, Nicholas King, Scott Mayer McKinney,	Towards conversational diagnostic artificial intelli-	867
814	Dean Carignan, and Eric Horvitz. 2023. Capabili-	gence. <i>Nature</i> , pages 1–9.	868
815	ties of gpt-4 on medical challenge problems. <i>arXiv</i>		
816	<i>preprint arXiv:2303.13375</i> .		
817	Don Nutbeam and Danielle M Muscat. 2021. Health	Mark V Williams, Ruth M Parker, David W Baker,	869
818	promotion glossary 2021. <i>Health promotion interna-</i>	Nina S Parikh, Kathryn Pitkin, Wendy C Coates, and	870
819	<i>tional</i> , 36(6):1578–1598.	Joanne R Nurss. 1995. Inadequate functional health	871
		literacy among patients at two public hospitals. <i>Jama</i> ,	872
		274(21):1677–1682.	873
820	OpenAI. 2024. Introducing Openai o1. <a href="https://openai.com/o1/">https://</a>	Kailai Yang, Shaoxiong Ji, Tianlin Zhang, Qianqian	874
821	<a href="https://openai.com/o1/">openai.com/o1/</a> . [Accessed 23-04-2025].	Xie, Ziyang Kuang, and Sophia Ananiadou. 2023. <b>To-</b>	875
822	OpenAI. 2025. <b>Chatgpt</b> .	<b>towards interpretable mental health analysis with large</b>	876
		<b>language models</b> . In <i>Proceedings of the 2023 Con-</i>	877
823	OpenAI-GPT-4o. 2024. Hello gpt-4o. <a href="https://openai.com/index/hello-gpt-4o/">https://</a>	<i>ference on Empirical Methods in Natural Language</i>	878
824	<a href="https://openai.com/index/hello-gpt-4o/">openai.com/index/hello-gpt-4o/</a> . [Accessed	<i>Processing</i> , pages 6056–6077, Singapore. Associa-	879
825	07-09-2024].	tion for Computational Linguistics.	880
826	David Restrepo, Chenwei Wu, Constanza Vásquez-	Yanghang Yu, Yuanyuan Zhao, Dongyan Li, Jingqiu	881
827	Venegas, João Matos, Jack Gallifant, Leo An-	Zhang, and Jiewei Li. 2021. The relationship be-	882
828	thony Celi, Danielle S Bitterman, and Luis Filipe	tween big five personality and social well-being of	883
829	Nakayama. 2024. Analyzing diversity in healthcare	chinese residents: The mediating effect of social sup-	884
830	llm research: A scientometric perspective. <i>medRxiv</i> ,	port. <i>Frontiers in Psychology</i> , 11:613659.	885
831	pages 2024–06.		
832	Gerard Saucier. 1994. Mini-markers: A brief version	Jonah Zaretsky, Jeong Min Kim, Samuel Baskharoun,	886
833	of goldberg’s unipolar big-five markers. <i>Journal of</i>	Yunan Zhao, Jonathan Austrian, Yindalon	887
834	<i>personality assessment</i> , 63(3):506–516.	Aphinyanaphongs, Ravi Gupta, Saul B. Blecker, and	888
		Jonah Feldman. 2024. <b>Generative artificial intelli-</b>	889
835	Karan Singhal, Shekoofeh Azizi, Tao Tu, S Sara Mah-	<b>gence to transform inpatient discharge summaries</b>	890
836	davi, Jason Wei, Hyung Won Chung, Nathan Scales,	<b>to patient-friendly language and format</b> . <i>JAMA</i>	891
837	Ajay Tanwani, Heather Cole-Lewis, Stephen Pfohl,	<i>Network Open</i> , 7(3):e240357–e240357.	892
838	and 1 others. 2023a. Large language models encode		
839	clinical knowledge. <i>Nature</i> , 620(7972):172–180.	Jiayuan Zhu and Junde Wu. 2025. Ask patients with	893
840	Karan Singhal, Tao Tu, Juraj Gottweis, Rory Sayres,	patience: Enabling llms for human-centric medical	894
841	Ellery Wulczyn, Le Hou, Kevin Clark, Stephen Pfohl,	dialogue with grounded reasoning. <i>arXiv preprint</i>	895
842	Heather Cole-Lewis, Darlene Neal, and 1 others.	<i>arXiv:2502.07143</i> .	896
843	2023b. Towards expert-level medical question an-		
844	swering with large language models. <i>arXiv preprint</i>		
845	<i>arXiv:2305.09617</i> .		
846	Arun James Thirunavukarasu, Refaat Hassan, Shathar	<b>A Big Five Personality Traits</b>	897
847	Mahmood, Rohan Sanghera, Kara Barzangi, Mo-		
848	hanned El Mukashfi, and Sachin Shah. 2023. Tri-	Each patient persona is instantiated with specific	898
849	alling a large language model (chatgpt) in general	personality traits that modulate their conversa-	899
850	practice with the applied knowledge test: observa-	tional behavior. For example, patients with high-	900
851	tional study demonstrating opportunities and limi-	<i>Agreeableness</i> tend to seek social harmony. They	901
852	tations in primary care. <i>JMIR Medical Education</i> ,	use affiliative language (e.g. “I totally understand	902
853	9(1):e46599.	your suggestions”), and readily consent to medi-	903
854	Lyndal J Trevena, Heather M Davey BPsych, Alexandra	cal recommendations. However, they might under-	904
855	Barratt, Phyllis Butow, and Patrina Caldwell. 2006.	report side-effects or doubts to avoid “bothering”	905
856	A systematic review on communicating with patients	the doctor, which can obscure important clinical	906
857	about evidence. <i>Journal of evaluation in clinical</i>	information. In contrast, low- <i>Agreeableness</i> pa-	907
858	<i>practice</i> , 12(1):13–23.	tients are more skeptical and assertive. They may	908
859	Tao Tu, Shekoofeh Azizi, Danny Driess, Mike Schaek-	challenge diagnoses, question procedures, and bar-	909
860	ermann, Mohamed Amin, Pi-Chuan Chang, Andrew	gain over medications, requiring more negotiation	910
861	Carroll, Charles Lau, Ryutaro Tanno, Ira Ktena, and	from the physician. Similarly, High- <i>Neuroticism</i>	911
862	1 others. 2024. Towards generalist biomedical ai.	often display heightened anxiety. They may fo-	912
863	<i>Nejm Ai</i> , 1(3):A10a2300138.	cus on worst-case scenarios (e.g., “Could this be	913
864	Tao Tu, Mike Schaekermann, Anil Palepu, Khaled Saab,	cancer?”), describe multiple vague symptoms, and	914
865	Jan Freyberg, Ryutaro Tanno, Amy Wang, Brenna Li,	seek repeated reassurance. Such interactions tend	915
866	Mohamed Amin, Yong Cheng, and 1 others. 2025.	to be emotionally charged and require empathetic	916
		listening and gradual explanation strategies. Con-	917
		versely, low- <i>Neuroticism</i> are more emotionally sta-	918
		ble and pragmatic. They typically provide concise,	919



Trait	Definition	Sensemaker-patient Conversational Style
Openness	Imagination, curiosity, comfort with novel, unusual, and abstract ideas.	<b>High:</b> Ask <i>why/how</i> questions about symptoms and medicines. Explore alternatives. Receptive to innovative therapies. <b>Low:</b> Prefer familiar terms and concrete instructions. May resist complex or less common therapies.
Conscientiousness	Organization, reliability, self-discipline, goal-orientation.	<b>High:</b> Bring detailed symptom logs and timelines. Seek precise next steps. <b>Low:</b> Offer vague histories. Forget previous conversations. Prefer simplified plans.
Extraversion	Sociability, talkativeness, interactions.	<b>High:</b> Give rich detail. Engage in small talk. Frequently ask questions. <b>Low:</b> Give succinct answers. Prefer structured questioning. Refrain from talking when necessary.
Agreeableness	Cooperation, trust, empathy, prosocial orientation.	<b>High:</b> Readily agree and express understanding. May conceal doubts to avoid conflict. <b>Low:</b> Challenge recommendations. Negotiate treatment choices. Openly voice dissatisfaction or concerns.
Neuroticism	Emotional reactivity, anxiety, mood instability.	<b>High:</b> Focus on worries. List multiple symptoms. Seek repeated reassurance and empathetic validation. <b>Low:</b> Remain calm and concise. Handle brisk, technical explanations without heightened distress.

Table 3: Definition of Big Five personality traits and typical sensemaker–patient conversation styles.

factual accounts and are receptive to succinct, technical communication. Table 3

tailed characteristics for these levels are presented in Table 4.

## B Health Literacy

We define two health literacy levels building upon Cutilli and Bennett (2009), which used factors such as ability to describe diseases, self-reporting conditions and navigating healthcare systems to assess health literacy in Americans (as part of the National Assessment of Adult Literacy). For example, a patient with basic medical literacy responded to a sensemaker’s diagnosis with *“I guess I’d like to know more, but I don’t really understand how all this works. It’s just... ever since..”*, while one with advanced literacy generated *“Yes, that sounds right. My father had similar symptoms, and they started when he was 33. Now I’m experiencing these involuntary movements, especially in my arms and hands .. genetic testing can confirm this.”*. De-

## C Patient Intentions

While the filtered MedQA case studies involve a patient with some symptoms looking for a diagnosis, they are not tailored towards sensemaker conversations, which would start off with an overarching intent conveyed by the patient (either implicitly, or if explicitly asked by the sensemaker). The intent adds a unique vignette to help guide the conversation with the sensemaker at the early stages. To mimic a real-world setting, we refer to Aydin et al. (2024), which analyzed 201 works studying usage of AI in healthcare to characterize six themes of how patients use LLMs in these settings and Li et al. (2024b), which studied works on using AI to personalize healthcare. We filtered out themes which involve indirect use of LLMs in healthcare

Health Literacy Level	Characteristics
Basic	<ul style="list-style-type: none"> <li>&gt; Struggles to understand basic medical terms and body parts.</li> <li>&gt; Rarely describes symptoms beyond "pain" or "sick".</li> <li>&gt; May use vague or informal terms instead of specific symptoms.</li> <li>&gt; Often cannot follow written medical instructions.</li> <li>&gt; May avoid seeking medical care due to communication barriers.</li> </ul>
Advanced	<ul style="list-style-type: none"> <li>&gt; Can provide detailed symptom descriptions including onset and triggers.</li> <li>&gt; Can describe subtle symptom variations and patterns.</li> <li>&gt; Understands complex medical terminology.</li> <li>&gt; Able to discuss medication effects and interactions.</li> <li>&gt; Able to research and evaluate health information from reliable sources.</li> <li>&gt; Maintains personal health records effectively.</li> </ul>

Table 4: Definition of two health literacy levels used in the patient profiles.

(e.g: generating patient education materials, optimizing doctor-patient interaction processes like medical consent forms) resulting in three themes: 1) *Interpreting medical information from a patient perspective*, 2) *Providing lifestyle recommendations and improving health literacy* and 3) *Personalizing healthcare journeys*. For example, the starting message for a case study augmented with intention (1) is: "A few days ago, I was feeling really agitated and not myself—I don't even remember much of what happened, but I was told I had to be sedated. I'm trying to make sense of all this and figure out what's going on with me. Can you help?". Detailed definitions and examples of these intents are provided in Table 5.

## D Patient Goals

While intents serve as high-level motivations to initiate conversations with a sensemaker, patient goals are concrete outcomes they wish to achieve by the end of the conversation. Similar to intents, MedQA case studies do not express explicit patient goals as they are not conversational in nature. Again, to mimic a real-world setting, we use [Aurriemma et al. \(2024\)](#), which carried out 338 "goals of care" conversations with 85 patients, resulting in three major themes: 1) *Maximize comfort and avoid suffering*, 2) *Maintain or improve cognitive or physical functioning* and 3) *Extend longevity or survival*. Detailed definitions and examples of these intents are provided in Table 6. Similar to the intents, we prompt GPT-4o to augment each patient profile with the single most appropriate goal given the medical facts of the case study (prompt in 18). "I appreciate you taking my feelings seriously. Right now, what would feel most helpful is anything that keeps me comfortable and minimizes

my suffering. I don't want to focus on things that might prolong my life if they make me feel worse in the process." is an example of goal (1) manifesting explicitly in a patient message in response to a decision making statement from the sensemaker.

## E Patient Message Generation Pipeline and Prompts

**Patient Actor LLM Message Generation Pipeline:** Algorithm 1 presents the algorithm for generating the patient actor LLM message  $p_t$  at timestamp  $t$  based on the sensemaker message ( $s_{t-1}$ ), patient persona  $\Pi$ , and set of patient's atomic facts  $F$ . While patient persona  $\Pi$  refers to the set of personality traits, health literacy level, patient intent and goals, patient's atomic facts  $F$  are obtained from the patient background case study as described above. For a given timestamp  $t$ , we first retrieve the relevant facts from the patient's atomic fact list  $F$  that could be used to generate the reply using a fact matching and retrieval prompt (Appendix E). If the number of facts matched ( $|\mathcal{F}_{\text{match}}| > 0$ ) then we use the patient message generation module to generate the final message  $p_t$  by combining the matched facts with stylistic variations based on personality, health literacy, and goals/intents defined in  $\Pi$ . In case,  $|\mathcal{F}_{\text{match}}| = 0$ , we generate messages based on the type of information being asked (either background or non-background). Background information here refers to the information which is related to the patient's medical facts obtained from the case study, on the other hand, non-background information refers to knowledge about future steps which are not defined in patient's background, intentions or goals. For background related information we assume a *closed world* assumption and reply signifying that

Patient Intent	Definition	Sample Patterns
Interpreting medical information from a patient perspective	Patient aims to use LLMs to simplify complex medical terminology and concepts so that patients can more easily understand diagnoses, procedures, and general health information.	<ul style="list-style-type: none"> <li>&gt; Patient can provide their symptoms and past medical history to receive lay-person friendly explanations of their condition.</li> <li>&gt; Patients can ask questions about their condition and receive clear, concise answers.</li> <li>&gt; Patients can receive personalized advice on how to manage their health.</li> </ul>
Providing lifestyle recommendations and improving health literacy	Patient aims to use LLMs to seek lifestyle change recommendations and debunking myths about health and wellness.	<ul style="list-style-type: none"> <li>&gt; Patients can ask questions about lifestyle changes and receive recommendations.</li> <li>&gt; Patients can query about the myths surrounding health and wellness.</li> </ul>
Personalizing healthcare journeys	Patient aims to use LLMs to tailor educational content and recommendations based on individual patient data, resulting in more relevant and actionable advice.	<ul style="list-style-type: none"> <li>&gt; Patients enter personal health goals (like weight loss or improved mobility), asking LLMs for targeted tips.</li> <li>&gt; Patients provide feedback on what they do or do not understand, and LLMs adjust explanations accordingly.</li> </ul>

Table 5: Definition and sample patterns of three high-level patient intents used to augment medical case studies.

Patient Goal	Definition
Maximize comfort and avoid suffering	Patient’s goal is to maximize comfort and avoid suffering. Includes seeking interventions to promote comfort (e.g., pain control) and avoiding interventions that would increase discomfort, even at the expense of decreasing longevity.
Maintain or improve cognitive or physical functioning	Patient’s goal is to maintain or improve cognitive or physical functioning by undergoing medical care aimed at preventing or reversing dysfunction, even if that medical care would increase discomfort. However, care that would increase survival/longevity without preservation or improvement in function is generally avoided.
Extending longevity or survival	Patient’s goal is to live as long as possible without limitations on care. Extending longevity or survival is prioritized over maximizing function or comfort.

Table 6: Definition of three concrete patient goals used to augment medical case studies.

the patient is unsure about the required information. In case the information being requested is related to non-background aspects then patient module can generate an affirmative message based on the situation. The starting message  $p_0$  is generated through a special prompt that uses the two most relevant case study facts, personality and the intent (prompt in Table 21).

## F Sensemaker Next Stage Determination Sub-module Prompt

Table 8 provides the prompt for the next stage determination module and Table 9 provides the rationale for stage movements provided in the prompt.

## G Sensemaker Fact Extraction and Matching Sub-module Prompt

We use the prompt in Table 10 to extract atomic facts from the patient’s current message and then

use the prompt in Table 11 to find if there are any matching facts already present in the memory.

## H Sensemaker Message Generation Sub-module Prompts

For the message generation sub-module, we have a similar structure for all the stages as shown in Table 12 with slight modifications in the gathering information and providing information stage as shown in Table 13 and Table 14 system prompts, respectively.

Table 15 shows the goals associated with each state that are passed to the system prompt as CURRENT\_STAGE\_GOAL.

## I Sensemaker Message Redundancy Checking Sub-module Prompt

We use the prompt in Table 16 to check if the candidate messages generated by the sensemaker are redundant or not.



Stage	Example Conversation
Fostering the Relationship	Hello, how can I help you today?
Gathering Information	Can you tell me more about your concerns? Can you tell me more about your personal and family history? What are your expectations from me? Can you expand on this point? Based on your description, it sounds like you're feeling... I understand you're feeling... I'm hearing that...
Providing Information	Your test results indicate that your cholesterol levels are high. This means you are at a greater risk for heart disease, but we can work on strategies like diet and exercise to manage it effectively.
Decision Making	There are a few different treatment options available. Would you like me to walk you through them so we can decide together which one aligns best with your lifestyle and preferences?
Enabling Disease and Treatment-Related Behavior	Managing diabetes can be overwhelming, but breaking it down into small steps—like checking your blood sugar daily—can make it easier. Would you like me to recommend some support groups or resources?
Responding to Emotions	I can see that this diagnosis is really affecting you. It's completely understandable to feel this way. I'm here to support you, and we can discuss ways to help manage both your symptoms and the emotional impact.

Table 7: The 6-function model of medical communication (De Haes and Bensing, 2009; King and Hoppe, 2013), which characterizes key communicative functions in clinical interactions, accompanied by representative conversation for each stage.

## J Qualitative Analysis

We conducted a qualitative analysis for the lowest-scoring conversations across all three patient-centric communication metrics, focusing on DeepSeek R1 as the sensemaker actor LLM. While the sensemaker stays empathetic and supportive, a lack of personalization emerges as a major theme across all metrics. However, this drawback means different things for each metric.

Sensemaker's dialogues such as *"The combination of sleeplessness without tiredness and auditory hallucinations suggests your brain chemistry might be imbalanced, potentially indicating conditions like bipolar disorder or schizophrenia that need professional diagnosis."* and *"Your symptoms suggest moderate-to-severe depression, especially with the persistent thoughts about not wanting to be alive. The next critical step would be connecting with a therapist who can help create a safety plan and explore treatment options."* showcase that while the sensemaker can provide a potential diagnosis and it fails to explicitly address the patient's susceptibility to their condition and the likelihood of experiencing a health problem due to the underlying condition.

In the context of low-scoring perceived severity conversations, the sensemaker identifies the underlying issue and provides a general definition of the

condition, along with relating it to the patient's symptoms, but dialogues such as *"Based on your symptoms and recent medication start, I believe you're experiencing akathisia - a restlessness side effect caused by fluphenazine. This can make both relaxation and restful sleep difficult despite not having classic insomnia symptoms."* and *"Based on your symptoms and routine, I believe you may have Delayed Sleep-Wake Phase Disorder, where your internal clock is misaligned with typical day/night cycles. This explains why you can't fall asleep until 2 a.m. and struggle to wake up for school."* shows that the sensemaker does not appropriately convey the seriousness of the potential consequences of the patient's symptoms.

Dialogues such as *"Would you like to explore some sleep hygiene strategies that could help you get more restful sleep?"* and *"Using phone reminders and labeled storage containers might help compensate for memory lapses - would you feel comfortable trying those?"* made by sensemaker in the Decision Making stage shows that while the sensemaker provides helpful actions, its strategies might lack details on their efficacy and how these strategies might reduce the risk faced by the patient.

Type	Prompt
<b>System Prompt</b>	<p>Given the current patient message (PATIENT_MESSAGE) a list of facts about the patient (PATIENT_FACTS) and a list of statements previously made by you (YOUR_MEMEORY). Determine whether the conversation should remain in the current stage (Fostering the Relationship) or transition to the next stage (Gathering Information). If you decide to stay in the current stage then your OUTPUT_STAGE should be 'STAYCURRENTSTAGE' and if you determine to transition to the next stage then your output should be 'MOVENEXTSTAGE'.</p> <p>Return Format:  OUTPUT_REASONING: &lt;your reasoning for the stage determination&gt;  OUTPUT_CONFIDENCE: &lt;your confidence score to transition to the next stage&gt;  OUTPUT_STAGE: &lt;MOVENEXTSTAGE or STAYCURRENTSTAGE&gt;</p> <p>Guidelines:  - Refer to POTENTIAL_NEXT_STAGE_REASONING and then reach your conclusion.  - Before determining if you should stay in the same stage or transition to the next stage, you should generate a step-by-step reasoning (OUTPUT_REASONING) for your conclusion.  - You should also generate a score between 1 and 10 (OUTPUT_CONFIDENCE) indicating your confidence in transitioning to the next stage, if the score is between 1 and 3, you should definately stay in the current stage, if the score is between 4 and 6, then you should stay in the stage for one or two more turns, and if the score is 7 or above then you should move to the next stage.  - The conversation should remain in the current stage if the user has not yet fully engaged with or completed its objectives.  - The conversation should move to the next stage only when the current stage has been meaningfully completed, ensuring a natural transition.</p> <p>Warnings:  - Do not keep the conversation stuck in the same stage for multiple iterations unless necessary. If progression is unclear, consider whether the user is engaging sufficiently before deciding.</p>
<b>User Prompt</b>	<p>POTENTIAL_NEXT_STAGE_REASONING: &lt;next_stage_reasoning&gt;</p> <p>PATIENT_MESSAGE: &lt;current patient message&gt;  PATIENT_FACTS: &lt;atomic facts about the patient till that timestep&gt;  YOUR_MEMORY: &lt;sensemaker messages till that timestep&gt;</p>

Table 8: (Sensemaker Module) Prompt used for the Gathering Information stage determination.

## K Automated Evaluation

Our three patient-centric communication metrics are inspired by the Health Belief Model and consists of the following dimensions:

- Perceived Susceptibility:** An individual’s belief about the likelihood of experiencing a health problem. Table 24 shows the prompt used to evaluate perceived susceptibility.
- Perceived Severity:** Beliefs about the seriousness of the consequences of a health issue. Table 25 shows the prompt used to evaluate perceived severity.
- Perceived Benefits:** Beliefs in the efficacy of the advised action to reduce the risk or seriousness of the health impact. Table 26 shows the prompt used to evaluate perceived severity.

We adapt HBM to evaluate how effectively the sensemaker LLM promotes each of the core dimensions on the patient side during conversations.

For example, if a patient exhibits low perceived susceptibility or self-efficacy, the sensemaker can adjust the tone and content of messages to address those gaps, improving engagement and promoting informed decision-making.

We evaluate Diagnostic accuracy and conversation flow correctness as well.

- Diagnostic Accuracy:** Since the filtered MedQA (Jin et al., 2021) case studies provide multi-choice options, we assess two different accuracies: (1) Hard Diagnostic Accuracy, which requires the sensemaker’s diagnosis to be an exact match with the ground truth. (2) Soft Diagnostic Accuracy, which allows for some freedom and blanket term matching. Table 27 and Table 28 show the prompt used to evaluate soft and hard diagnostic accuracy metrics, respectively.
- Conversation Flow and Correctness:** It is used to measure the ability of the sensemaker to perform end-to-end conversation, while maintaining logical consistency, covering each phase of

Current Stage	Next Stage Reasoning
Fostering the Relationship	The next stage would be 'Gathering Information' because an initial relationship between you and the patient has been established. Move on when the 'Fostering the Relationship' stage has provided a welcoming space and the patient starts to openly describe their concerns, feelings, or challenges. If they seem hesitant or reserved, stay in 'Fostering the Relationship' longer to encourage sharing.
Gathering Information	The next stage would be 'Providing Information' only when you have received enough information from the patient about their current condition and symptoms for you to make a diagnosis. If you feel you need more time to gather information to make a confident diagnosis, stay in 'Gathering Information' longer.
Providing Information	You should move to the next stage which is 'Decision Making', NO MATTER WHAT.
Decision Making	The next stage would be 'Responding to Emotions' because the patient has understood the lifestyle and non-clinical suggestions made by you to alleviate their current condition. Move forward from the 'Decision Making' stage when the patient has acknowledged your suggestions. If their responses suggest they still need more clarity or direction, stay in 'Decision Making' to provide additional support.
Responding to Emotions	The next stage would be 'exit' because the conversation has reached its end. Move forward from the 'Responding to Emotions' when you validated their emotions with an empathetic response and the patient has replied with an affirmative message. Do not focus on coping mechanisms in this stage.

Table 9: Reasoning provided to the prompt in Table 8.

Type	Prompt
<b>System Prompt</b>	<p>You are given a message provided by the patient as USER_MESSAGE, your task is to extract explicitly stated atomic facts about the patient from the message. Here, an atomic fact is defined as new information provided by the patient which can increase your holistic understanding of the patient's condition. Each atomic fact should carry an entirely different piece of explicitly stated fact, and should be independent of other atomic facts in the list.</p> <p>Your output should strictly be a list of atomic facts, with each item starting with "# ". Do not include other formatting. In addition, each of the atomic facts in the list should be in third person narration.</p> <p>Keep in mind that each atomic fact is different from other atomic facts in the list. Do not add any new fact which was not present in the USER_MESSAGE.</p>
<b>User Prompt</b>	<p>USER_MESSAGE: &lt;current patient message&gt;</p> <p>ATOMIC_FACTS:</p>

Table 10: (Sensemaker Module) Prompt used for Fact Extraction.

the conversation, preventing providing incorrect information and preventing contradiction with itself. Table 29 shows the prompt used to evaluate flow correctness.

## L Human Annotators and Annotation Guidelines

Human annotations was conducted by four graduate students fluent in English language and with a background in Computer Science and digital mental health research. For the annotation task, annotators were provided with the entire patient-sensemaker conversation, score for each metric (*Perceived Susceptibility*, *Perceived Severity*, *Perceived Benefits and Conversation Flow and Cor-*

*rectness*, *Diagnostic accuracy*) from the automated evaluation, reasoning for the score generated by LLM judge and the rubric with the definitions for each metric and scoring criteria.

For each metric and conversation, two annotators assigned a binary label indicating agreement or disagreement with the LLM judge's score and reasoning for the metric. While a positive label denoted agreement with the LLM score and reasoning, a negative label indicated non-alignment with LLM judge score or reasoning. Overall, we observed high correlation for each metric. For *Diagnostic Accuracy (Hard)*, the average correlation with LLM judge was computed as 93.5%, whereas as for *Diagnostic Accuracy (Soft)*, average correlation was 90%. For the patient-centric communica-



Type	Prompt
<b>System Prompt</b>	<p>You are an expert fact matching agent. You are given a list of NEW_FACTS about a patient, and a list of already present facts in MEMORY_OF_PATIENT_FACTS. Your task is to analyze if facts listed in NEW_FACTS list are 'FactPresent' or 'FactNotPresent' in MEMORY_OF_PATIENT_FACTS. For this task you take each fact from NEW_FACTS list one by one and check if it is present in MEMORY_OF_PATIENT_FACTS or not. For this checking you should look at the facts in terms of their meaning and context and not the exact words. The classification for each NEW FACT should be independent of other NEW_FACTS.</p> <p>Your output should strictly follow the format: # &lt;fact&gt; : &lt;'FactNotPresent' or 'FactPresent'&gt;. Where fact presents one fact from NEW_FACTS. Output the labels for each fact in NEW_FACTS and keep the fact text as it is (do not change the words).</p> <p>You should not change the words of the fact, only add the label (either 'FactPresent' or 'FactNotPresent'). Your output should have one fact and its label per line.</p>
<b>User Prompt</b>	<p>NEW_FACTS: &lt;facts extracted from the current patient message&gt;</p> <p>MEMORY_OF_PATIENT_FACTS: &lt;atomic facts about the patient till that timestep&gt;</p> <p>ANSWER:</p>

Table 11: (Sensemaker Module) Prompt used for Fact Matching.

tion metrics, agreement scores were 79% for *Perceived Susceptibility*, 87.5% for *Perceived Severity*, 86.5% for *Perceived Benefits*. Finally, *Conversation Flow and Correctness* average correlation with LLM judge was computed as 87.5%.

## M Personality Based Results

### N Model Details, Hyperparameters, and Compute

We use API-based model inference for GPT-4o, OpenAI o1, and DeepSeek R1. We used Azure OpenAI Service for accessing GPT-4o, OpenAI o1, and DeepSeek R1 models. The hyperparameters for the generations are presented below. Table 31 presents the details regarding the model sizes and versions.

**Hyperparameters for GPT and DeepSeek Models:** temperature  $t = 0$ , max\_tokens = 500 (for sensemaker fact extraction, matching and message redundancy checker which use GPT-4o) and max\_tokens = 15000 (for sensemaker next stage determination and message generation which use reasoning models). All patient module components use GPT-4o with a temperature  $t = 0$  and max\_tokens = 400.

### O Information About Use of AI Assistants

We used AI assistants for text rephrasing. The usage was limited to correcting grammatical mistakes and choice of words.

Type	Prompt
<b>System Prompt</b>	<p>You are an expert sensemaker who helps the patients to make sense of their current clinical situation. Keeping the conversation's current stage goal and example messages in mind (given by CURRENT_STAGE_GOAL and CURRENT_STAGE_EXAMPLES), list of facts already known about the patient (PATIENT_FACTS), and the list of statements already made by you (YOUR_MEMORY), and the user's current message (USER_MESSAGE), your goal is to generate three plausible and logical messages to be said to the patient that you have not said yet. You should first generate a reasoning for yourself and then generate the three messages. Your messages can be a question or a statement.</p> <p>Your output should strictly be in the following format:  OUTPUT_REASONING: &lt;your step-by-step reasoning&gt;  OUTPUT_MESSAGE: &lt;3 plausible messages to the patient each on a new line and starting with "# "&gt;. Do not include other formatting.</p> <p>All the three plausible messages should be different from each other. If you are asking a question in a message, then only ask one question at a time in that message. Your message should be concise and to the point.</p> <p>YOUR_MEMORY: &lt;sensemaker's messages till that timestep&gt;</p> <p>PATIENT_FACTS: &lt;atomic facts about the patient till that timestep&gt;</p> <p>CURRENT_STAGE_GOAL: &lt;goal that needs to be achieved in the current stage&gt;</p> <p>CURRENT_STAGE_EXAMPLES: &lt;examples of messages for the current stage&gt;</p>
<b>User Prompt</b>	USER_MESSAGE: <current patient message>

Table 12: (Sensemaker Module) Prompt used for message generation in the all other stages.

Type	Prompt
<b>System Prompt</b>	<p>You are an expert sensemaker who helps the patients to make sense of their current clinical situation. Keeping the conversation's current stage goal and example messages in mind (given by CURRENT_STAGE_GOAL and CURRENT_STAGE_EXAMPLES), list of facts already known about the patient (PATIENT_FACTS), and the list of statements already made by you (YOUR_MEMORY), your goal is to generate three plausible and logical messages to be said to the patient that you have not said yet. Before generating the message you should generate a step-by-step reasoning taking into account the facts you already know about the patient. In your reasoning you should think about the possible diagnosis hypotheses, and then generate the three messages for the patient that helps in gathering more information to either confirm or reject the diagnosis hypotheses.</p> <p>Your output should strictly be in the following format:  OUTPUT_REASONING: &lt;your step-by-step reasoning&gt;  OUTPUT_MESSAGE: &lt;3 plausible messages to the patient each on a new line and starting with "# "&gt;. Do not include other formatting.</p> <p>All the three plausible messages should be different from each other. If you are asking a question in a message, then only ask one question at a time in that message. Your message should be concise and to the point.</p> <p>YOUR_MEMORY: &lt;sensemaker's messages till that timestep&gt;</p> <p>PATIENT_FACTS: &lt;atomic facts about the patient till that timestep&gt;</p> <p>CURRENT_STAGE_GOAL: &lt;goal that needs to be achieved in the current stage&gt;</p> <p>CURRENT_STAGE_EXAMPLES: &lt;examples of messages for the current stage&gt;</p>
<b>User Prompt</b>	USER_MESSAGE: <current patient message>

Table 13: (Sensemaker Module) Prompt used for message generation in the Gathering Information stage.

Type	Prompt
<b>System Prompt</b>	<p>You are an expert sensemaker who helps the patients to make sense of their current clinical situation. Keeping the conversation's current stage goal and example messages in mind (given by CURRENT_STAGE_GOAL and CURRENT_STAGE_EXAMPLES), list of facts already known about the patient (PATIENT_FACTS), and the list of statements already made by you (YOUR_MEMORY), your goal is to provide the patient with the diagnosis and its explanation to the patient. Before generating the diagnosis and explanation message you should generate a step-by-step reasoning for yourself taking into account the facts you already know about the patient.</p> <p>Your output should strictly be in the following format:  OUTPUT_REASONING: &lt;your step-by-step reasoning&gt;  OUTPUT_MESSAGE: &lt;3 plausible messages to the patient each on a new line and starting with "# "&gt;. Do not include other formatting.</p> <p>You should always provide a diagnosis and if you cannot find a diagnosis your message should strictly be "I apologize but I am unable to diagnose you at the moment."</p> <p>All the three plausible messages should be different from each other. If you are asking a question in a message, then only ask one question at a time in that message. Your message should be concise and to the point.</p> <p>YOUR_MEMORY: &lt;sensemaker's messages till that timestep&gt;</p> <p>PATIENT_FACTS: &lt;atomic facts about the patient till that timestep&gt;</p> <p>CURRENT_STAGE_GOAL: &lt;goal that needs to be achieved in the current stage&gt;</p> <p>CURRENT_STAGE_EXAMPLES: &lt;examples of messages for the current stage&gt;</p>
<b>User Prompt</b>	<p>USER_MESSAGE: &lt;current patient message&gt;</p>

Table 14: (Sensemaker Module) Prompt used for message generation in the Providing Information stage.

Current Stage	Goal
Fostering the Relationship	<p>Your goal is to build a trusting, open, and collaborative relationship with the patient by demonstrating empathy, honesty, and respect. You need to create a safe and supportive environment where the patient feels heard, valued, and comfortable sharing sensitive information. For this you can use the following strategies:</p> <ol style="list-style-type: none"> <li>1. Invite the patient to share their story in their own words.</li> <li>2. Respond with empathy and compassion to patient's concerns.</li> <li>3. Express compassion and commitment. Let the patient know that you care about their well-being and are committed to helping them.</li> <li>4. If the patient is not comfortable sharing their story, you can ask them to share their concerns and you can respond with empathy and compassion.</li> </ol>
Gathering Information	<p>Your goal is to develop a comprehensive understanding of the patient's needs, concerns, and medical history by exploring their condition from both biological and psychosocial perspectives. This understanding will allow you to support the patient in achieving their goals and expectations for the conversation. For this you can use the following strategies:</p> <ol style="list-style-type: none"> <li>1. Ask open ended questions related to patient's concerns to gather information about patients current state, their personal and family history.</li> <li>2. Listen actively and ask follow-up questions to understand the situation better.</li> <li>3. Elicit patient's perspective of the problems and their expectations from you.</li> <li>4. Clarify and summarize the information gathered from the patient to ensure understanding.</li> </ol>
Providing Information	<p>At this stage you want to provide a potential diagnosis to the patient for their concerns. You should not ask any questions in this stage and rather provide a potential diagnosis to patient based on the their personal history, family history, concerns and other details.</p>
Decision Making	<p>"At this stage you should address any medical queries posed by the patient regarding your diagnosis and suggest only lifestyle or non-clinical changes to the patient to alleviate their illness based on their diagnosis. You should make sure that the changes suggested by you are based on the patient's preferences and your all previous knowledge about them. For this you can use the following strategies:</p> <ol style="list-style-type: none"> <li>1. Ask for patient's preferences and suggestions regarding the lifestyle changes or other non-clinical changes.</li> <li>2. Suggest lifestyle changes based on the patient's preferences and your all previous knowledge about them.</li> <li>3. If the patient is not comfortable with the suggestions, you can ask them to suggest their own lifestyle or non-clinical changes.</li> </ol>
Responding to Emotions	<p>You need to recognize and address any emotional aspect of the illness by offering empathay, reassurance, and psychological support in your messages. For this you can use the following strategies:</p> <ol style="list-style-type: none"> <li>1. Offer empathay and reassurance to the patient.</li> <li>2. Listen to the patient's concerns and offer psychological support.</li> <li>3. If the patient is not comfortable sharing their concerns, you can ask them to share their emotions and you can respond with empathy and support.</li> </ol>
Exit	<p>The conversation has reached its end and we need to conclude the conversation.</p>

Table 15: Current stage goal provided to the sensemaker message generation prompt



Type	Prompt
<b>System Prompt</b>	<p>You are an expert sensemaker who helps the patients to make sense of their current clinical situation. Given a list of statements already made by you (STATEMENT_MEMORY), and a list of candidate statements (CANDIDATE_STATEMENTS), your task is to identify if there are any redundant statements in the CANDIDATE_STATEMENTS.</p> <p>Your output should strictly follow the format: # &lt;statement&gt; : &lt;'RedundantStatement' or 'RedundantNotStatement'&gt;. Where statement presents one statement from CANDIDATE_STATEMENTS. Output the labels for each statement in CANDIDATE_STATEMENTS and keep the statement text as it is (do not change the words) and always start the line with '#'.  You should not change the words of the statement, only add the label. Your output should have one statement and its label per line. Below are some additional instructions:</p> <ol style="list-style-type: none"> <li>1. You should not add any new statement which was not present in the STATEMENT_MEMORY.</li> <li>2. If a new statement in the CANDIDATE_STATEMENTS has a different phrasing but serves a similar context to any of the statements present in the STATEMENT_MEMORY, it should be considered 'RedundantStatement'.</li> <li>3. Classification for each statement in CANDIDATE_STATEMENTS should be independent of other statements in the CANDIDATE_STATEMENTS.</li> </ol>
<b>User Prompt</b>	<p>CANDIDATE_STATEMENTS: &lt;candidate statements that can be made in this particular stage&gt;</p> <p>STATEMENT_MEMORY: &lt;statements already made by the sensemaker&gt;</p>

Table 16: (Sensemaker Module) Prompt used for checking message redundancy.

Type	Prompt
<b>System Prompt</b>	<p>You are an assistant that is given information about a patient.</p> <p>Break the following patient information into a list of independent atomic facts, with one piece of information in each statement. Each fact should only include the smallest unit of information, but should be self-contained.</p> <p>Criteria:</p> <ol style="list-style-type: none"> <li>1. Only extract medical facts that would be relevant in a conversation with a health-care provider.</li> <li>2. First, list out facts related to a patient's vitals (ONLY temperature, blood pressure, pulse, respiratory rate) under "PATIENT VITAL FACTS:".</li> <li>3. If there are no PATIENT VITAL FACTS just leave the section empty.</li> <li>4. List out the remaining medical atomic facts under the section "PATIENT FACTS:".</li> </ol>
<b>User Prompt</b>	<p>Patient Information:</p> <p>&lt;patient_data&gt;</p> <p>Respond with the list of atomic facts for each section "PATIENT VITAL FACTS:" and "PATIENT FACTS:" (no formatting) and nothing else, prepend each fact with a '- '. No sub-lists are allowed.</p>

Table 17: (Patient Module) Prompt used for extracting patient facts.

Type	Prompt
<b>System Prompt</b>	<p>You are an expert psychiatrist/psychologist/psychotherapist. Given a patient case study, your task is to pick the most appropriate goal for that patient's consultation with an LLM. You are given a case study (provided as CASE_STUDY) and a list of patient conversation goal vignettes in the format (VIGNETTE_NAME, VIGNETTE_DEFINITION, VIGNETTE_PATTERNS). You need to pick the most appropriate vignette for the given case study.</p> <p>Think step by step and first provide your rationale (under "RATIONALE:") in around 30 words and then at the last output the VIGNETTE_NAME for the most logical vignette for the given case study.</p> <p>1. VIGNETTE_NAME: VIGNETTE_DEFINITION:</p> <p>2. VIGNETTE_NAME: ...</p> <p>Think step by step and first provide your rationale (under "RATIONALE:") and then at the last output the VIGNETTE_NAME for the most logical vignette for the given case study.</p>
<b>User Prompt</b>	<p>CASE_STUDY: &lt;case_study&gt;</p> <p>Strictly follow the format: &lt;ASSIGNED_VIGNETTE_NAME: VIGNETTE_NAME&gt;. Use the exact vignette name and nothing else.</p>

Table 18: (Patient Module) Prompt used to attach a concrete conversational goal to a case study. "Vignette" here refers to the goal.

Type	Prompt
<b>System Prompt</b>	<p>You are an expert psychiatrist/psychologist/psychotherapist. Given a patient case study, your task is to pick the most appropriate motivation for that patient to consult an LLM. You are given a case study (provided as CASE_STUDY) and a list of patient conversation intent vignettes in the format (VIGNETTE_NAME, VIGNETTE_DEFINITION, VIGNETTE_PATTERNS). You need to pick the most appropriate vignette for the given case study.</p> <p>1. VIGNETTE_NAME: VIGNETTE_DEFINITION: VIGNETTE_PATTERNS:</p> <p>2. VIGNETTE_NAME: ...</p> <p>Think step by step and first provide your rationale (under "RATIONALE:") and then at the last output the VIGNETTE_NAME for the most logical vignette for the given case study.</p>
<b>User Prompt</b>	<p>CASE_STUDY: &lt;case_study&gt;</p> <p>Strictly follow the format: &lt;ASSIGNED_VIGNETTE_NAME: VIGNETTE_NAME&gt;. Use the exact vignette name and nothing else.</p>

Table 19: (Patient Module) Prompt used to attach a high-level conversational intent to a case study. "Vignette" here refers to the intent.

Type	Prompt
<b>System Prompt</b>	<p>You are an intelligent agent who is given a QUESTION from a doctor to a patient, and a LIST OF MEDICAL FACTS about the patient. Your task is to choose the facts which best answer the question, or respond "NO MATCH".</p> <p>Instructions:</p> <ol style="list-style-type: none"> <li>1. Given a LIST OF MEDICAL FACTS about a patient, choose a MAXIMUM of THREE facts from the list that when combined best answers the QUESTION.</li> <li>2. If NO fact matches the question, simply respond "NO MATCH" under "FACTS:".</li> <li>3. If facts are chosen, output them as a list, preserving the numbering from the original list.</li> <li>4. First provide a short reasoning under "REASONING:" before listing the facts under "FACTS:".</li> <li>5. Do not include any other formatting or extra information beyond the REASONING and given FACTS.</li> </ol>
<b>User Prompt</b>	<p>QUESTION: &lt;question&gt;</p> <p>LIST OF MEDICAL FACTS: &lt;patient_info&gt;</p> <p>SOLUTION:</p>

Table 20: (Patient Module) Prompt used for selecting relevant medical facts to answer a question.

Type	Prompt
<b>System Prompt</b>	<p>You are an intelligent agent who is given a LIST OF MEDICAL FACTS about a patient, who is going to have a conversation with an AI healthcare provider. Your task is to construct a start message with the most important facts in first person, as if you are simulating the patient's persona and literacy while talking to the AI.</p> <p>To aid in this process, Big 5 personality traits are defined as:</p> <p>&lt;personality_trait_definitions&gt;</p> <p>Instructions:</p> <ol style="list-style-type: none"> <li>1. Given a LIST OF MEDICAL FACTS about a patient, choose a MAXIMUM of TWO relevant facts from the list to construct a starting message as a patient to an AI healthcare provider.</li> <li>2. The start message should be in first person, using the emotions, tone, word choice and intensity of a patient who has the levels of the Big 5 Personality Traits (on a 2-point Low/High scale) and Medical Literacy Level shown below.</li> <li>3. ONLY use the medical literacy level and personality definitions as a guide, DO NOT EXPLICITLY integrate any information from it into the answer.</li> <li>4. First provide a short reasoning under "REASONING:" before writing the start message under "START_MESSAGE:".</li> <li>5. Do not include quotes, any other formatting, extra information beyond the REASONING and the START_MESSAGE.</li> </ol>
<b>User Prompt</b>	<p>LIST OF MEDICAL FACTS: &lt;patient_info&gt;</p> <p>Big 5 Personality Traits:</p> <ol style="list-style-type: none"> <li>1. Openness: &lt;Openness_score&gt;</li> <li>2. Conscientiousness: &lt;Conscientiousness_score&gt;</li> <li>3. Extraversion: &lt;Extraversion_score&gt;</li> <li>4. Agreeableness: &lt;Agreeableness_score&gt;</li> <li>5. Neuroticism: &lt;Neuroticism_score&gt;</li> </ol> <p>MEDICAL LITERACY LEVEL: &lt;medical_literacy_level&gt;</p>

Table 21: (Patient Module) Prompt used for patient to start the conversation with the sensemaker.

Type	Prompt
<b>System Prompt</b>	<p>You are a truthful assistant that understands a patient’s medical information, and you are trying to answer questions from an AI about the patient in first person, as if you are simulating the patient’s persona and literacy while talking to the AI.</p> <p>To aid in this process, Big 5 personality traits are defined as:</p> <p>&lt;personality_trait_definitions&gt;</p> <p>Instructions:</p> <ol style="list-style-type: none"> <li>1. Using the list of medical facts given below, generate a conversational human-like response to an LLM in first person, using the emotions, tone, word choice and intensity of a patient who has the levels of the Big 5 Personality Traits (on a 2-point Low/High scale) and Medical Literacy Level shown below.</li> <li>2. Any information asked in the question that is NOT explicitly present in the facts can be assumed to be FALSE.</li> <li>3. If the question is just a statement diagnosing your condition, agree to it.</li> <li>4. ONLY use the medical literacy level and personality definitions as a guide, DO NOT EXPLICITLY integrate any information from it into the answer.</li> <li>5. DO NOT add ANY information in your answer that is NOT EXPLICITLY stated in the given facts.</li> <li>6. DO NOT suggest any treatment method yourself in the answer.</li> <li>7. ONLY respond with the answer, do NOT add any other formatting.</li> </ol>
<b>User Prompt</b>	<p>LIST OF MEDICAL FACTS:</p> <p>&lt;patient_info&gt;</p> <p>QUESTION:</p> <p>&lt;question&gt;</p> <p>Big 5 Personality Traits:</p> <ol style="list-style-type: none"> <li>1. Openness: &lt;Openness_score&gt;</li> <li>2. Conscientiousness: &lt;Conscientiousness_score&gt;</li> <li>3. Extraversion: &lt;Extraversion_score&gt;</li> <li>4. Agreeableness: &lt;Agreeableness_score&gt;</li> <li>5. Neuroticism: &lt;Neuroticism_score&gt;</li> </ol> <p>MEDICAL LITERACY LEVEL:</p> <p>&lt;medical_literacy_level&gt;</p>

Table 22: (Patient Module) Prompt used to generate answers given a non-zero number of selected facts. Note that in stages beyond Decision Making, point 2 in the instructions (closed-world assumption) is removed to aid natural conversation flow.



Type	Prompt
System Prompt	<p>You are a truthful assistant that understands a patient's medical information, and you are trying to answer questions from an AI about the patient in first person, as if you are the patient with a specific persona and literacy talking to an AI.</p> <p>To aid in this process, Big 5 personality traits are defined as:</p> <p>&lt;personality_trait_definitions&gt;</p> <p>Instructions:</p> <ol style="list-style-type: none"> <li>1. Generate a conversational human-like response to an LLM in first person, using the emotions, tone, word choice and intensity of a patient who has the levels of the Big 5 Personality Traits (on a 2-point Low/High scale) and Medical Literacy Level shown below.</li> <li>2. If this question requests information about the patient's medical history, personal experience, medication, background or condition, respond in the negative stating "No ..." or "I'm not sure about ..." under "ANSWER:".</li> <li>3. If it is a question about decision-making, emotions, diagnosis or taking next steps, generate a conversational human-like response to an LLM in first person, responding in the affirmative ("Yes ...") as a patient.</li> <li>4. ONLY use the medical literacy level and personality definitions as a guide, DO NOT EXPLICITLY integrate any information from it into the answer.</li> <li>5. DO NOT add ANY NEW information to the answer. If asked about any such additional information, respond with "I'm not sure about ..." to them.</li> <li>6. First provide a short reasoning under "REASONING:" before writing the answer under "ANSWER:". Do not add any other formatting.</li> </ol> <p>Here are some examples to help you</p> <p>EXAMPLE 1:</p> <p>QUESTION: Have you tried any meditation-style exercises?</p> <p>REASONING: Exercises and lifestyle relates to the patient's personal information.</p> <p>ANSWER: No, I have not tried any meditation-related exercises.</p> <p>EXAMPLE 2:</p> <p>QUESTION: Would you like to explore some coping strategies related to your condition?</p> <p>REASONING: This is a question about next steps, hence the answer is in the affirmative.</p> <p>ANSWER: Yes, I would like to learn about some coping strategies.</p> <p>EXAMPLE 3:</p> <p>QUESTION: Looks like you are experiencing insomnia. When did you first start feeling these symptoms?</p> <p>REASONING: Though there is a diagnosis, the question part asks some medical history information.</p> <p>ANSWER: I'm not sure about when I first started feeling these symptoms.</p> <p>EXAMPLE 4:</p> <p>QUESTION: Looks like you are experiencing insomnia. What do you think about trying a medication routine to help with this?</p> <p>REASONING: There is a diagnosis and the question asks about next steps or advice, hence the answer is in the affirmative.</p> <p>ANSWER: I see, yes, I would like to learn about medications to help my situation.</p>
User Prompt	<p>LIST OF MEDICAL FACTS:</p> <p>&lt;patient_info&gt;</p> <p>Big 5 Personality Traits:</p> <ol style="list-style-type: none"> <li>1. Openness: &lt;Openness_score&gt;</li> <li>2. Conscientiousness: &lt;Conscientiousness_score&gt;</li> <li>3. Extraversion: &lt;Extraversion_score&gt;</li> <li>4. Agreeableness: &lt;Agreeableness_score&gt;</li> <li>5. Neuroticism: &lt;Neuroticism_score&gt;</li> </ol> <p>MEDICAL LITERACY LEVEL:</p> <p>&lt;medical_literacy_level&gt;</p> <p>QUESTION:</p> <p>&lt;question&gt;</p>

Table 23: (Patient Module) Prompt used to generate answers when there are no selected facts matching the question.

Type	Prompt
<b>System Prompt</b>	<p>Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's communication of "Perceived Susceptibility" (definition given as SUSCEPTIBILITY_DEFINITION) within a patient-sensemaker conversation.</p> <p>SUSCEPTIBILITY_DEFINITION: "Perceived Susceptibility" here refers to the sensemaker's ability to clearly and adequately express to the patient their likelihood of experiencing a health problem.</p> <p>You will be given a conversation between a patient and a sensemaker (given as PATIENT_SENSEMAKER_CONVERSATION) and your task is to evaluate the sensemaker's communication of "Perceived Susceptibility" within this conversation based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA: Assign a score from 1 (lowest) to 4 (highest) based on the following criteria:</p> <ol style="list-style-type: none"> <li>1. (Very Poor: Score = 1): The sensemaker failed to communicate any information about the susceptibility of facing a health problem. Additionally, the tone may have been dismissive, insensitive, or unnecessarily alarming.</li> <li>2. (Poor: Score = 2): The sensemaker failed to communicate any information about the susceptibility of facing a health problem. However, the tone was empathetic and engaging.</li> <li>3. (Good: Score = 3): The sensemaker communicated basic information about the patient's susceptibility to facing a health problem. The tone was professional but lacked empathy or collaborative engagement.</li> <li>4. (Very Good: Score = 4): The sensemaker explained the patient's susceptibility to facing a health problem clearly, accurately and appropriately, using personalized details effectively. Additionally, the sensemaker was empathetic and engaged collaboratively.</li> </ol> <p>Output Format: You should first provide a short reasoning (2-3 sentences) providing your reasoning for the score. Then, provide the score. Your output should be in the following format:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;score&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The clinician explicitly and clearly communicated the patient's susceptibility to depression using tailored information from their personal history. The clinician's tone was empathetic and engaged the patient in a supportive, collaborative dialogue, ensuring the patient's clear understanding.</p> <p>OUTPUT_SCORE: 4</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
<b>User Prompt</b>	PATIENT_SENSEMAKER_CONVERSATION: <Sample conversation>

Table 24: (Evaluation) Prompt used for automated Perceived Susceptibility evaluation.

Type	Prompt
System Prompt	<p>Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's communication of "Perceived Severity" (definition given as SEVERITY_DEFINITION) within a patient-sensemaker conversation.</p> <p>SEVERITY_DEFINITION: "Perceived Severity" refers to the sensemaker's ability to adequately convey the seriousness of the consequences associated with a health issue to the patient.</p> <p>You will be given a conversation between a patient and a sensemaker (given as PATIENT_SENSEMAKER_CONVERSATION) and your task is to evaluate the sensemaker's communication of "Perceived Severity" within this conversation based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA: Assign a score from 1 (lowest) to 4 (highest) based on the following criteria:</p> <ol style="list-style-type: none"> <li>1. (Very Poor: Score = 1): The sensemaker failed to communicate any information about the seriousness of the consequences associated with a health issue to the patient. Additionally, the tone may have been dismissive, insensitive, or unnecessarily alarming.</li> <li>2. (Poor: Score = 2): The sensemaker failed to communicate any information about the seriousness of the consequences associated with a health issue to the patient. However, the tone was empathetic and engaging.</li> <li>3. (Good: Score = 3): The sensemaker communicated basic information about the seriousness of the consequences associated with a health issue to the patient. The tone was professional but lacked empathy or collaborative engagement.</li> <li>4. (Very Good: Score = 4): The sensemaker explained the seriousness of the consequences associated with a health issue clearly, accurately and appropriately, using personalized details effectively. Additionally, the sensemaker was empathetic and engaged collaboratively.</li> </ol> <p>Output Format: You should first provide a short reasoning (2-3 sentences) providing your reasoning for the score. Then, provide the score. Your output should be in the following format:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;score&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The sensemaker explicitly described the consequences of untreated depression clearly, using detailed personal context relevant to the patient's life situation. The empathetic and collaborative tone ensured patient understanding and minimized unnecessary distress.</p> <p>OUTPUT_SCORE: 4</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
User Prompt	PATIENT_SENSEMAKER_CONVERSATION: <Sample conversation>

Table 25: (Evaluation) Prompt used for automated Perceived Severity evaluation.

Type	Prompt
System Prompt	<p>Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's communication of "Perceived Benefits" (definition given as BENEFITS_DEFINITION) within a patient-sensemaker conversation.</p> <p>BENEFITS_DEFINITION: "Perceived Benefits" refers to the sensemaker's ability to effectively communicate the efficacy of the advised action in reducing the risk or seriousness of the health impact appropriately.</p> <p>You will be given a conversation between a patient and a sensemaker (given as PATIENT_SENSEMAKER_CONVERSATION) and your task is to evaluate the sensemaker's communication of "Perceived Benefits" within this conversation based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA: Assign a score from 1 (lowest) to 4 (highest) based on the following criteria:</p> <ol style="list-style-type: none"> <li>1. (Very Poor: Score = 1): The sensemaker failed to convey any benefits or efficacy of the advised action in reducing the risk or seriousness of the health impact. Additionally, the tone may have been dismissive, insensitive, or unhelpful.</li> <li>2. (Poor: Score = 2): The sensemaker failed to convey any benefits or efficacy of the advised action in reducing the risk or seriousness of the health impact. However, the tone was empathetic and engaging.</li> <li>3. (Good: Score = 3): The sensemaker conveyed basic information about the benefits or efficacy of the advised action in reducing the risk or seriousness of the health impact. The tone was professional but lacked empathy or collaborative engagement.</li> <li>4. (Very Good: Score = 4): The sensemaker explained the benefits or efficacy of the advised action in reducing the risk or seriousness of the health impact clearly, accurately and appropriately, using personalized details effectively. Additionally, the sensemaker was empathetic and engaged collaboratively.</li> </ol> <p>Output Format: You should first provide a short reasoning (2-3 sentences) providing your reasoning for the score. Then, provide the score. Your output should be in the following format:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;score&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The sensemaker clearly articulated how therapy sessions could effectively reduce anxiety, using specific examples relevant to the patient's personal situation. The tone was supportive, empathetic, and collaborative, significantly enhancing the patient's confidence and understanding of the recommended action.</p> <p>OUTPUT_SCORE: 4</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
User Prompt	PATIENT_SENSEMAKER_CONVERSATION: <Sample conversation>

Table 26: (Evaluation) Prompt used for automated Perceived Benefits evaluation.



Type	Prompt
System Prompt	<p>Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's "Diagnostic Accuracy" (definition given as DIAGNOSTIC_ACCURACY_DEFINITION).</p> <p>DIAGNOSTIC_ACCURACY_DEFINITION: "Diagnostic Accuracy" refers to the accuracy of the sensemaker's diagnosis when compared to the ground truth diagnosis.</p> <p>You will be given a specific part from a patient-sensemaker conversation, the sensemaker's diagnosis statement (given as PATIENT_DIAGNOSIS) and the ground truth diagnosis of the patient (given as GT_DIAGNOSIS). Your task is to evaluate the sensemaker's "Diagnostic Accuracy" from this conversation snippet and assign a binary label based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA:</p> <ol style="list-style-type: none"> <li>1. Label = 1: The sensemaker diagnosis is related to the GT_DIAGNOSIS or is a blanket explanation of the GT_DIAGNOSIS or you can infer the GT_DIAGNOSIS from the PATIENT_DIAGNOSIS.</li> <li>2. Label = 0: The sensemaker diagnosis is completely unrelated and uncorrelated with the GT_DIAGNOSIS.</li> </ol> <p>Output Format: You should first provide a short reasoning (2-3 sentences) providing your reasoning for the label. Then, provide the label. Your output should be in the following format:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;0/1&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The sensemaker diagnosed the person with anxiety while the ground truth diagnosis is PTSD, since they are both are closely related conditions and PTSD can be inferred from the patient diagnosis snippet of the conversation, therefore the label is 1.</p> <p>OUTPUT_SCORE: 1</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
User Prompt	<p>PATIENT_DIAGNOSIS: &lt;Providing Information stage Message&gt;</p> <p>GT_DIAGNOSIS: &lt;Ground Truth Label&gt;</p>

Table 27: (Evaluation) Prompt used for hard matching diagnostic accuracy

Type	Prompt
System Prompt	<p>Task: Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's "Diagnostic Accuracy" (definition given as DIAGNOSTIC_ACCURACY_DEFINITION).</p> <p>DIAGNOSTIC_ACCURACY_DEFINITION: "Diagnostic Accuracy" refers to the accuracy of the sensemaker's diagnosis when compared to the ground truth diagnosis.</p> <p>You will be given a specific part from a patient-sensemaker conversation, the sensemaker's diagnosis statement (given as PATIENT_DIAGNOSIS) and the ground truth diagnosis of the patient (given as GT_DIAGNOSIS). Your task is to evaluate the sensemaker's "Diagnostic Accuracy" from this conversation snippet and assign a binary label based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA:</p> <ol style="list-style-type: none"> <li>1. Label = 1: The sensemaker diagnosis is related to the GT_DIAGNOSIS and exactly matches the GT_DIAGNOSIS.</li> <li>2. Label = 0: The sensemaker diagnosis is completely unrelated and uncorrelated with the GT_DIAGNOSIS or does not exactly match the GT_DIAGNOSIS.</li> </ol> <p>Output Format: You should first provide a short reasoning (2-3 sentences) providing your reasoning for the label. Then, provide the label. Your output should be in the following format:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;0/1&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The sensemaker diagnosed the person with anxiety while the ground truth diagnosis is PTSD, since they are not exactly the same, therefore the label is 0.</p> <p>OUTPUT_SCORE: 0</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
User Prompt	<p>PATIENT_DIAGNOSIS: &lt;Providing Information stage Message&gt;</p> <p>GT_DIAGNOSIS: &lt;Ground Truth Label&gt;</p>

Table 28: (Evaluation) Prompt used for hard matching diagnostic accuracy.

Type	Prompt
System Prompt	<p>Task: You are an expert LLM-as-a-judge model and your task is to evaluate a sensemaker's *Conversation Flow and Correctness* (definition given as FLOW_CORRECTNESS_DEFINITION) within a patient-sensemaker conversation.</p> <p>FLOW_CORRECTNESS_DEFINITION: "Conversation Flow and Correctness" refers to the sensemaker's ability to (i) progress through an end-to-end sensemaking dialogue in a logically coherent order according to STAGE_DEFINITION order and (ii) provide consistent information without contradictions or inconsistencies.</p> <p>You will be given a conversation between a patient and a sensemaker (given as PATIENT_SENSEMAKER_CONVERSATION) and your task is to evaluate the sensemaker's performance on "Conversation Flow and Correctness" based on the following evaluation criteria (given as EVALUATION_CRITERIA):</p> <p>EVALUATION_CRITERIA: Assign a score from 1 (lowest) to 4 (highest) using the rubric below:</p> <ol style="list-style-type: none"> <li>1. (Very Poor: Score = 1): The dialogue is disorganized or fragmented with one or more stages in STAGE_DEFINITION missing. Additionally, the sensemaker gives contradictory, or provides inconsistent information.</li> <li>2. (Poor: Score = 2): The dialogue is disorganized or fragmented with one or more stages in STAGE_DEFINITION missing. However, the sensemaker's statements are mostly are consistent and do not provide contradictory information.</li> <li>3. (Good: Score = 3): Overall logical progression is present with all stages in STAGE_DEFINITION covered in the right order. Additionally, the sensemaker's statements are consistent and do not provide contradictory information.</li> <li>4. (Very Good: Score = 4): Overall logical progression is present with all stages in STAGE_DEFINITION covered in the right order and the sensemaker has covered the each stage sufficiently paying special attention on gathering information from the patient and providing appropriate information and empathetic support to the patient. Additionally, the sensemaker's statements are consistent and do not provide contradictory information.</li> </ol> <p>## STAGE_DEFINITION</p> <p>A typical sensemaking conversation between a sensemaker and a patient consists of the following order of stages:</p> <ul style="list-style-type: none"> <li>- Fostering the Relationship: Establish a trusting, open, and collaborative rapport with the patient by demonstrating empathy, honesty, and respect.</li> <li>- Gathering Information: Develop a thorough understanding of the patient's condition by exploring both biological and psychosocial factors, including their needs, concerns, and medical history.</li> <li>- Providing Information: Offer a potential diagnosis that directly addresses the patient's concerns, ensuring clarity and sensitivity in communication.</li> <li>- Decision Making: Respond to the patient's medical questions regarding the diagnosis, and recommend only lifestyle or non-clinical interventions to help manage their condition.</li> <li>- Responding to Emotions: Acknowledge and address the emotional aspects of the patient's experience by expressing empathy, offering reassurance, and providing appropriate psychological support.</li> <li>- Exit: Conclude the conversation in a respectful and supportive manner, ensuring the patient feels heard and understood.</li> </ul> <p>## OUTPUT FORMAT</p> <p>Provide a brief reasoning (2-3 sentences) explaining the score, followed by the score itself, using the template below:</p> <p>OUTPUT_REASONING: &lt;your_reasoning&gt;</p> <p>OUTPUT_SCORE: &lt;score&gt;</p> <p>Example Output:</p> <p>OUTPUT_REASONING: The sensemaker gathered a complete history, summarized findings, explained differential diagnoses, and covered all stages in STAGE_DEFINITION in the right order with appropriate information and empathetic support to the patient. No contradictions or inconsistencies were detected.</p> <p>OUTPUT_SCORE: 4</p> <p>ALWAYS STRICTLY RETURN THE OUTPUT IN THE ABOVE FORMAT.</p>
User Prompt	PATIENT_SENSEMAKER_CONVERSATION: <Sample conversation>

Table 29: (Evaluation) Prompt used for Flow Correctness and Logical Consistency evaluation.

Personality Trait	Model	Perceived Susceptibility	Perceived Severity	Perceived Benefits	Flow Correctness	Diag. Acc. (Hard) (%)	Diag. Acc. (Soft) (%)	SMOG
Basic Health Literacy Patient								
Openness	o1	2.46	2.62	2.91	3.53	31.68	<u>70.30</u>	11.22
	R1	2.58	2.71	<b>3.25</b>	3.56	<u>34.65</u>	58.42	13.27
Conscientiousness	o1	2.45	2.53	2.77	<u>3.57</u>	30.08	66.92	11.25
	R1	2.57	2.72	<u>3.08</u>	3.47	29.32	64.66	13.23
Extraversion	o1	2.44	2.50	2.69	3.47	25.49	66.67	11.23
	R1	2.44	<u>2.74</u>	2.93	3.38	28.43	62.75	13.01
Agreeableness	o1	2.42	2.44	2.85	<b>3.68</b>	<b>36.43</b>	<b>70.54</b>	<u>11.13</u>
	R1	<u>2.60</u>	2.74	3.04	3.50	26.36	59.69	13.06
Neuroticism	o1	2.19	2.43	2.41	3.24	31.33	67.47	<b>10.92</b>
	R1	<b>2.66</b>	<b>2.81</b>	2.88	3.28	28.92	59.04	12.90
Advanced Health Literacy Patient								
Openness	o1	2.56	2.63	2.82	3.59	29.37	<b>72.22</b>	11.88
	R1	<u>2.63</u>	2.79	<u>3.11</u>	<b>3.64</b>	28.57	69.05	13.57
Conscientiousness	o1	2.43	2.56	2.85	3.57	25.93	67.41	12.09
	R1	2.60	2.76	3.10	3.61	<u>38.52</u>	<u>71.85</u>	13.77
Extraversion	o1	2.52	2.70	2.79	3.54	29.91	69.23	11.65
	R1	2.59	<u>2.79</u>	2.94	3.39	32.48	68.38	13.39
Agreeableness	o1	2.50	2.57	2.84	<u>3.62</u>	32.81	68.75	11.79
	R1	<b>2.67</b>	<b>2.81</b>	<b>3.12</b>	3.60	30.47	66.41	13.56
Neuroticism	o1	2.39	2.57	2.47	3.30	<b>39.77</b>	62.50	<b>11.53</b>
	R1	2.48	2.60	2.89	3.32	34.09	57.95	13.31

Table 30: Performance of o1 vs. R1 across seven metrics, by persona and health-literacy level. We report the scores with the best and second-best performing model metrics in each scenario highlighted in **bold** and underline, respectively.

LLM	Version	Parameter Size
GPT-4o	( <a href="#">OpenAI-GPT-4o</a> , 2024) [Version: 2024-11-20]	(undisclosed)
OpenAI o1	( <a href="#">OpenAI</a> , 2024) [Version: 2024-12-17]	(undisclosed)
DeepSeek R1	( <a href="#">Guo et al.</a> , 2025) [Version: 1]	671 billion

Table 31: Models and their corresponding version dates