Stochastic Momentum Methods for Non-smooth Non-Convex Finite-Sum Coupled Compositional Optimization

Xingyu Chen* tamucxy2023@tamu.edu

Bokun Wang* bokun-wang@tamu.edu

Ming Yang*
myang@tamu.edu

Qihang Lin[†] qihang-lin@uiowa.edu

Tianbao Yang^{*‡} tianbao-yang@tamu.edu.

Abstract

Finite-sum Coupled Compositional Optimization (FCCO), characterized by its coupled compositional objective structure, emerges as an important optimization paradigm for addressing a wide range of machine learning problems. In this paper, we focus on a challenging class of non-convex non-smooth FCCO, where the outer functions are non-smooth weakly convex or convex and the inner functions are smooth or weakly convex. Existing state-of-the-art result face two key limitations: (1) a high iteration complexity of $O(1/\epsilon^6)$ under the assumption that the stochastic inner functions are Lipschitz continuous in expectation; (2) reliance on vanilla SGD-type updates, which are not suitable for deep learning applications. Our main contributions are two fold: (i) We propose stochastic momentum methods tailored for non-smooth FCCO that come with provable convergence guarantees; (ii) We establish a **new state-of-the-art** iteration complexity of $O(1/\epsilon^5)$. Moreover, we apply our algorithms to multiple inequality constrained non-convex optimization problems involving smooth or weakly convex functional inequality constraints. By optimizing a smoothed hinge penalty based formulation, we achieve a new **state-of-the-art** complexity of $O(1/\epsilon^5)$ for finding an (nearly) ϵ -level KKT solution. Experiments on three tasks demonstrate the effectiveness of the proposed algorithms.

1 Introduction

In this paper, we consider the finite-sum coupled compositional optimization (FCCO) problem

$$\min_{\mathbf{w} \in \mathbb{R}^d} F(\mathbf{w}) := \frac{1}{n} \sum_{i=1}^n f_i(g_i(\mathbf{w})), \tag{1}$$

where $f_i : \mathbb{R}^{d_1} \to \mathbb{R}$ is continuous, $g_i : \mathbb{R}^d \to \mathbb{R}^{d_1}$ is continuous and satisfies $g_i(\mathbf{w}) = \mathbb{E}g_i(\mathbf{w}, \xi_i)$, and the expectation is taken over the random variable ξ_i for $i = 1, \ldots, n$.

FCCO has been effectively applied to optimizing a wide range of risk functions known as X-risks [1, 2, 3, 4], and group distributionally robust optimization (GDRO) [5, 6]. Recently, it was also applied to solving non-convex inequality constrained optimization problems [7, 8]. Several optimization algorithms have been developed for non-convex FCCO (1) under different conditions.

^{*}Department of CSE, Texas A&M University, College Station, USA

[†]Tippie College of Business, The University of Iowa, Iowa City, USA.

[‡]Correspondence to: tianbao-yang@tamu.edu.

Table 1: Comparison between our algorithms and prior works for the **non-smooth non-convex FCCO** problem. The complexity of SONX is for finding a nearly ϵ -stationary solution (Definition 3.1), and that of SONEX and ALEXR2 for smooth inner functions are for finding an approximate ϵ -stationary solution (Definition 4.1), which implies a nearly ϵ -stationary solution under a verifiable condition of the inner functions. The complexity of ALEXR2 for weakly convex inner functions is for finding a nearly ϵ -stationary solution to the outer smoothed objective. In this table, "WC" means weakly convex, "C" means convex, "C" means monotonically non-decreasing, "SM" means smooth, "PMS" means that the proximal mapping can be easily computed, "MLC0" means that the function is mean Lipschitz continuous of zero-order (Assumption 4.6).

Algorithm	f_i	g_i	Complexity	Loop	Update type
SONX [6]	WC, ↗	WC, MLC0	$O(\epsilon^{-6})$	Single	SGD
SONEX (Ours)	WC, PMS	SM, MLC0	$O(\epsilon^{-5})$	Single	Momentum or Adam
ALEXR2 (Ours)	C, PMS	SM	$O(\epsilon^{-5})$	Double	Momentum or Adam
ALEXR2 (Ours)	C, PMS, ↗	WC	$O(\epsilon^{-5})$	Double	Momentum or Adam

Table 2: Comparison between our algorithms and prior works for solving **non-convex inequality constrained optimization** problem. g_0 is the objective and g_i are constraint functions.

Algorithm	g_0	g_i	Constraint Sampling	Complexity	Loop	Update type
OSS [9]	WC	WC	No	$O(\epsilon^{-6})$	Double	SGD
ICPPAC [10]	WC	WC	No	$O(\epsilon^{-6})$	Double	SGD
SSG [11]	WC	C	No	$O(\epsilon^{-8})$	Single	SGD
Li et al. [7]	SM	SM	Yes	$O(\epsilon^{-7})$	Single	Momentum or Adam
Yang et al. [8]	WC	WC, MLC0	Yes	$O(\epsilon^{-6})$	Single	SGD
Liu et al. [12]	WC	WC, MLC0	No	$O(\epsilon^{-6})$	Single	SGD
SONEX (Ours) ALEXR2 (Ours)	SM WC	SM, MLC0 WC	Yes Yes	$O(\epsilon^{-5}) \\ O(\epsilon^{-5})$	Single Double	Momentum or Adam Momentum or Adam

Most of them require the smoothness of f_i and g_i . Hu et al. [6] has initiated the study of non-smooth non-convex FCCO where both the inner and outer functions could be non-smooth. However, their results face two key limitations: (1) a high iteration complexity of $O(1/\epsilon^6)$ under the assumption that the stochastic inner functions are Lipschitz continuous in expectation; (2) reliance on vanilla SGD-type updates, which are not suitable for deep learning applications.

Novelty. This paper addresses these limitations by proposing stochastic momentum methods for non-smooth FCCO, where f_i is non-smooth and g_i could be smooth or weakly convex, and improving the convergence rate to $O(1/\epsilon^5)$. To the best of our knowledge, this is the first work to propose stochastic momentum methods for solving non-smooth FCCO problems. Unlike [6] that directly solves the original problem, the key to our methods is to smooth the outer non-smooth functions based on the Moreau envelope smoothing or equivalently the Nesterov smoothing when they are convex, which is referred to as *outer smoothing*. When the inner functions are non-smooth weakly convex, we further smooth the transformed objective using another layer of Moreau envelope smoothing, which is referred to as *nested smoothing*. Then we propose stochastic momentum algorithms to optimize these smoothed objectives.

Contributions and Significance. We establish two main results regarding non-smooth FCCO. In the first result, we consider non-smooth FCCO with non-smooth outer functions and smooth inner functions. With outer smoothing, we propose a single-loop stochastic momentum method named SONEX to solve the resulting smoothed objective. Our main contribution here lies at a theory that guarantees a convergence rate of $O(1/\epsilon^5)$ under a meaningful convergence measure. Specially, we show that when the outer smoothing parameter is small enough, the proposed algorithm is guaranteed to find a novel notion of approximate ϵ -stationary solution to the original problem, which implies the standard nearly ϵ -stationary solution to the original problem under a verifiable condition of inner functions. In the second result, we consider non-smooth FCCO with non-smooth convex outer functions and non-smooth weakly inner functions. With nested smoothing, we propose a novel double-loop stochastic momentum method named ALEXR2 for solving the resulting smoothed objective and establish a convergence rate of $O(1/\epsilon^5)$ for finding an ϵ -stationary solution of the smoothed objective. Table 1 compares our methods with those of the existing works for non-smooth FCCO problems from different aspects.

Then we consider novel applications of these two algorithms to **non-convex inequality constrained optimization**. By optimizing a *smoothed hinge penalty function* with the proposed stothastic momentum methods, we derive a new state-of-the-art rate of $O(1/\epsilon^5)$ for finding an ϵ -KKT solution when objective and constraint functions are smooth, and for finding a nearly ϵ -KKT solution when objective and the constraint functions are weakly convex. Table 2 summarizes the complexity of our methods and existing works for solving non-convex inequality constrained optimization problems.

2 Related work

FCCO. FCCO is a special case of stochastic composite optimization (SCO) [13, 14] and conditional stochastic optimization (CSO) [15] when the outer expectation is a finite sum. FCCO was first introduced in [4] as a model for optimizing the average precision. Later, [2] proposed the SOX algorithm and improved the convergence rate by adding gradient momentum and further being improved in [16] by replacing exponential moving average (EMA) with MSVR and STORM estimators for inner function and overall gradient, respectively. All these techniques have gain success in optimizing Deep X-Risks [1] such as smooth surrogate losses of AUC and contrastive loss [17, 18]. For convex FCCO problem, [2] reformulate FCCO as a saddle point problem and use restarting technique to boost convergence rate. For non-smooth weakly-convex FCCO (NSWC FCCO), [6] leverages MSVR technique and requires $O(\epsilon^{-6})$ iterations to achieve a nearly ϵ -stationary point.

Smoothing techniques: An effective approach for solving non-smooth optimization is to approximate the original problem by a smoothed one using different smoothing techniques, including Nesterov's smoothing [19], randomized smoothing [20], and Moreau envelope [21, 22]. Our method is related to the one based on Moreau envelope, which is a prevalent technique for weakly convex non-smooth problems. This method approximates the objective function by its smooth Moreau envelope and then computes an ϵ -stationary point of the Moreau envelope, which is also a nearly ϵ -stationary solution of the original problem [22, 23]. The similar technique has been extended to min-max problems [24, 25] and constrained problems [9, 10, 11, 26]. Our methods are different from these works because we consider the Moreau envelope of each outer function and design stochastic momentum methods. Similar outer smoothing techniques have been developed in [27] for compositional objectives. However, different from our work (1) their algorithm cannot handle multiple outer functions of a finite sum structure as (1); (2) their algorithm is not momentum method; (3) their convergence theory requires mean Lipchitz continuity of the gradient of the inner functions for finding a primal-dual ϵ -stationary (KKT) solution of the min-max formulation of the original problem.

Non-convex constrained optimization: Most existing works for stochastic non-convex constrained optimization focus on the smooth problems. For example, [28] proposes a double-loop inexact augmented Lagrangian method for stochastic smooth nonconvex equality constrained optimization under a regularity assumption on the gradients of the constraints. A quadratic penalty method is developed by [29] under similar assumptions but uses only a single loop. Both [29] and [28] consider only smooth problems while we also consider non-smooth problems. For non-smooth but weakly convex problems, the existing methods can compute a nearly ϵ -KKT point based on the Moreau envelopes of the original problem with iteration complexity $O(\epsilon^{-6})$ [9, 10, 11, 26, 8]. The penalty-based method by [12] reduces the number of evaluations of the subgradients to $O(\epsilon^{-4})$ but still needs to evaluate $O(\epsilon^{-6})$ function values. In contrast, our method only needs complexity $O(\epsilon^{-5})$ to compute a nearly ϵ -KKT point. For smooth constrained optimization with non-convex equality constraints, variance-reduced stochastic gradient methods have been developed based on quadratic penalty [29, 30]. One can convert inequality constraints into equality constraints by adding slack variables s^2 or s (requiring $s \ge 0$) to each inequality constraint [30]. However, adding s^2 will introduce spurious stationary points [31, 32], while adding $s \ge 0$ will make the domain or the constraint functions unbounded which violate the assumptions in [29, 30]. More importantly, we consider non-smooth constraint functions while [29, 30] only consider smooth cases.

3 Preliminaries

Let $\|\cdot\|$ be the Euclidean norm. For $g(\cdot): \mathbb{R}^d \to \mathbb{R}^{d_1}$, let $\nabla g(\cdot) \in \mathbb{R}^{d \times d_1}$ be its Jacobian and let $\|\nabla g(\cdot)\| = \max_{\mathbf{u} \in \mathbb{R}^d, \|\mathbf{u}\| = 1} \|\nabla g(\cdot)\mathbf{u}\|$ be the operator norm. Let $g_i(\mathbf{w}, \mathcal{B}) = \frac{1}{|\mathcal{B}|} \sum_{\xi \in \mathcal{B}} g_i(\mathbf{w}, \xi), i = 1, 2, \cdots, n$ be an estimator of g_i based on samples from a minibatch \mathcal{B} . A function $f(\cdot)$ is monotonically non-decreasing if for any $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^{d_1}$ satisfying that

 $\mathbf{x}_{1,k} \leq \mathbf{x}_{2,k}, k \in [d_1]$ where \mathbf{x}_k denote the k-th element of \mathbf{x} , it holds that $f(\mathbf{x}_1) \leq f(\mathbf{x}_2)$. A function $f(\cdot)$ is ρ -weakly convex for $\rho \geq 0$ if $f(\cdot) + \frac{\rho}{2} \|\cdot\|^2$ is convex. A function $g(\cdot)$ is L-smooth for $L \geq 0$ if g is differentiable and $\nabla g(\cdot)$ is L-Lipschitz continuous. A vector-valued mapping $g(\cdot)$ is L-smooth for $L \geq 0$ if g is differentiable at each component and $\nabla g(\cdot)$ is L-Lipschitz continuous w.r.t to the operator norm. Given a ρ -weakly convex function $f(\cdot)$, we denote its regular subdifferential as $\partial f(\cdot)$. For a ρ -weakly convex function $f(\cdot)$ and a constant $\lambda \in (0, \rho^{-1})$, the proximal mapping and the Moreau envelope of $f(\cdot)$ with parameter λ are defined as follows, respectively,

$$\operatorname{prox}_{\lambda f}(\mathbf{u}) := \operatorname*{arg\,min}_{\mathbf{v}} f(\mathbf{v}) + \frac{1}{2\lambda} \left\| \mathbf{u} - \mathbf{v} \right\|^2 \text{ and } f_{\lambda}(\mathbf{u}) := \min_{\mathbf{v}} f(\mathbf{v}) + \frac{1}{2\lambda} \left\| \mathbf{u} - \mathbf{v} \right\|^2.$$

It is known that, for $\lambda \in (0, \rho^{-1})$, $f_{\lambda}(\cdot)$ is $L_f := \max\{\frac{1}{\lambda}, \frac{\rho}{1-\lambda\rho}\}$ -smooth [33] and

$$\nabla f_{\lambda}(\mathbf{u}) = \frac{\mathbf{u} - \operatorname{prox}_{\lambda f}(\mathbf{u})}{\lambda} \in \partial f(\operatorname{prox}_{\lambda f}(\mathbf{u})). \tag{2}$$

For a non-smooth weakly convex objective, we follow existing works [21] [6] and aim to find a nearly ϵ -stationary solution defined below.

Definition 3.1. A solution \mathbf{w} is an ϵ -stationary solution of (1) if $\operatorname{dist}(0, \partial F(\mathbf{w})) \leq \epsilon$. A solution \mathbf{w} is a nearly ϵ -stationary solution of (1) if there exists \mathbf{w}' such that $\|\mathbf{w} - \mathbf{w}'\| \leq \epsilon$ and $\operatorname{dist}(0, \partial F(\mathbf{w}')) \leq \epsilon$. We make the following assumption on (1) throughout the paper.

Assumption 3.2. We assume that f_i is C_f -Lipschitz continuous, g_i is C_g -Lipschitz continuous for $i=1,\ldots,n$, $\operatorname{prox}_{\lambda f_i}(\mathbf{w})$ and its subgradient are easily computable, and one of the following conditions hold:

A1. f_i is ρ_f -weakly convex, and $g_i(\mathbf{w})$ is L_g -smooth for all i.

A2. f_i is convex, and $g_i(\mathbf{w})$ is L_g -smooth for all i.

A3. f_i is convex and monotonically non-decreasing, and $g_i(\mathbf{w})$ is ρ_q -weakly convex for all i.

Remark: The Lipschitz continuity of f and g in Assumption 3.2 are fairly standard for FCCO problems. Since we target at non-smooth f and g, Lipchitz continuity is a minimal assumption. In the considered applications of GDRO and learning with fairness constraints in section 6, f is hinge and hence Lipschitz continuous.

4 Smoothing of Non-smooth FCCO

We first describe the main idea of our methods and then present detailed algorithms and their convergence. Under the Assumption 3.2, we can show F is also weakly convex. To improve the convergence rates and accommodate deep learning applications, we develop stochastic momentum methods, which can be easily modified to incorporate with adaptive step sizes (cf. Appendix E). The challenges of developing provable stochastic momentum methods for solving (1) lie at the non-smoothness of f_i and potentially g_i .

Let us first consider smooth g_i and defer the discussion for weakly convex g_i to subsection 4.2. The key idea is to use a smoothed version of f_i , denoted by $f_{i,\lambda}$, and to approximate (1) by

$$\min_{\mathbf{w} \in \mathbb{R}^d} F_{\lambda}(\mathbf{w}) := \frac{1}{n} \sum_{i=1}^n f_{i,\lambda}(g_i(\mathbf{w})). \tag{3}$$

We can easily show that when g_i is L_g -smooth, then $F_{\lambda}(\mathbf{w})$ is L_F -smooth with $L_F = C_g^2 \max\{\frac{1}{\lambda}, \frac{\rho_f}{1-\lambda\rho_f}\} + C_f L_g$ (cf. Lemma A.1). As a result, the problem becomes a smooth FCCO where both inner functions and outer functions are smooth, which makes it possible to employ existing techniques to develop stochastic momentum methods to find an ϵ -stationary point of (3), which is a point \mathbf{w} satisfying $\mathbb{E}[\|\nabla F_{\lambda}(\mathbf{w})\|] \leq \epsilon$. However, what this implies for solving the original problem and what constitutes a good choice of λ remain unclear. Below, we present a theory to address these questions. We introduce the following notion of stationarity of the original problem.

Definition 4.1. A solution \mathbf{w} is an approximate ϵ -stationary solution of (1) if there exist $\mathbf{t}_1, \ldots, \mathbf{t}_m$ and $\mathbf{y}_i \in \partial f_i(\mathbf{t}_i)$ for $i = 1, \ldots, n$ such that $\|\mathbf{t}_i - g_i(\mathbf{w})\| \leq \epsilon, \ i = 1, \ldots, n$, and $\|\frac{1}{n} \sum_{i=1}^n \nabla g_i(\mathbf{w}) \mathbf{y}_i\| \leq \epsilon$.

This definition implies that when $\epsilon \to 0$, the solution converges to a stationary solution of $F(\cdot)$. Given this definition, our first theorem is stated below whose proof is given in Section A.3.

Algorithm 1 SONEX for solving (3)

```
1: Input: T, \lambda, \mathbf{w}_0 = \mathbf{w}_{-1}, \mathbf{v}_0, \eta, \gamma, \gamma', \beta, \forall t.
 2: Sample a batch of data \mathcal{B}_{i,2} from the distribution of \xi_i for i=1,\ldots,n.
 3: Set u_{i,0} = g_i(\mathbf{w}_0, \mathcal{B}_{i,2}) for i = 1, \dots, n.
4: for t = 0, 1, \dots, T - 1 do
            Sample \mathcal{B}_1^t \subset \{1,\ldots,n\} and a batch of data \mathcal{B}_{i,2}^t from the distribution of \xi_i for i \in \mathcal{B}_1^t.
            for i=1,\ldots,n do
 6:
 7:
                u_{i,t+1} = \begin{cases} (1-\gamma)u_{i,t} + \gamma g_i(\mathbf{w}_t, \mathcal{B}_{i,2}^t) + \gamma'(g_i(\mathbf{w}_t, \mathcal{B}_{i,2}^t) - g_i(\mathbf{w}_{t-1}, \mathcal{B}_{i,2}^t)) & \text{if } i \in \mathcal{B}_1^t \\ u_{i,t} & \text{if } i \notin \mathcal{B}_1^t \end{cases}
 8:
            Compute G_t = \frac{1}{|\mathcal{B}_i^t|} \sum_{i \in \mathcal{B}_1^t} \nabla f_{i,\lambda}(u_{t,i}) \nabla g_i(\mathbf{w}_t, \mathcal{B}_{i,2}^t)
 9:
             Update \mathbf{v}_{t+1} = (1 - \beta)\mathbf{v}_t + \beta G_t
10:
             Update \mathbf{w}_{t+1} = \mathbf{w}_t - \eta \mathbf{v}_{t+1} or Adam-type update
11:
12: end for
13: Output: \mathbf{w}_{\tau} with \tau randomly sampled from \{1, 2, \dots, T\}
```

Theorem 4.2. If w is an ϵ -stationary solution to $F_{\lambda}(\cdot)$ with $\lambda = \epsilon/C_f$ such that $\|\nabla F_{\lambda}(\mathbf{w})\| \le \epsilon$, then w is an approximate ϵ -stationary solution to the original objective $F(\cdot)$.

A remaining question is whether an approximate ϵ -stationary point is also a nearly ϵ -stationary solution of the original problem (1). We present a theorem below under the following assumption.

Assumption 4.3. There exist c>0 and $\delta>0$ such that, if \mathbf{w} is an ϵ -stationary point of F_{λ} with $\lambda=\epsilon/C_f$ and $\epsilon\leq c$, it holds that $\lambda_{\min}\left(\nabla\mathbf{g}(\mathbf{w})\nabla\mathbf{g}(\mathbf{w})^{\top}\right)\geq\delta$, where $\nabla\mathbf{g}(\mathbf{w})=[\nabla g_1(\mathbf{w})^{\top},\ldots,\nabla g_n(\mathbf{w})^{\top}]^{\top}$,

Remark: In Assumption 4.3, once ϵ is smaller than c, the lower bound δ does not depend on ϵ . The empirical justification for this assumption is provided in subsection F.5.

Theorem 4.4. Suppose Assumptions 3.2 (A1) and 4.3 hold. If \mathbf{w} is an approximate ϵ -stationary solution of (1) with $\epsilon \leq \min\left\{c, \frac{\delta^2}{16nC_g^2L_g}\right\}$, \mathbf{w} is also a nearly ϵ -stationary solution of (1).

Remark: The proof is given in Appendix A.4. It is notable that the Assumption 4.3 is easily verifiable (cf. Appendix F.5 for empirical evidence). In addition, when we consider the applications in non-convex constrained optimization, this condition is commonly used in existing analyses [7, 8].

4.1 Single-loop Methods for Smooth Inner Functions

Next, we present a single-loop algorithm for finding an ϵ -stationary point of (3). The key ingredients of the algorithm include two parts: (1) maintaining and updating n sequences $u_{i,t}$ for tracking each $g_i(\mathbf{w}_t), i=1,\ldots,n$, which are updated in a coordinate-wise manner; (2) a stochastic momentum update. We present the detailed updates in Algorithm 1, which is referred to as SONEX. We note that the algorithm is inspired by existing stochastic momentum methods for smooth FCCO. First, Step 7 for updating $u_{i,t+1}$ is from the MSVR algorithm [16]. Different from MSVR, we directly utilize the stochastic momentum update in Step 10, 11, which are similar to SOX [2]. In contrast, MSVR also leverages a variance reduction technique (STROM) to compute an estimate of the stochastic gradient, which requires a stronger assumption that $\nabla g_i(\mathbf{w};\xi)$ is Lipschitz continuous in expectation. Our analysis shows that this is not helpful for improving the convergence, as the complexity will be dominated by a term related to the smoothness of $f_{i,\lambda}$, which is in the order of $O(1/\epsilon)$.

We assume the following conditions of the stochastic estimators of g_i and their gradients.

Assumption 4.5. There exist constants $\sigma_0 \ge 0$ and $\sigma_1 \ge 0$ such that the following statements hold for $g_i(\mathbf{w})$ and $g_i(\mathbf{w}, \xi_i)$: for $i = 1, \ldots, n$ and any $\mathbf{w} \in \mathbb{R}^d$

$$\mathbb{E}\|g_i(\mathbf{w},\xi_i) - g_i(\mathbf{w})\|^2 \le \sigma_0^2, \mathbb{E}\|\partial g_i(\mathbf{w},\xi_i) - \partial g_i(\mathbf{w})\|^2 \le \sigma_1^2$$

Assumption 4.6. There exist a constant C_q such that $\mathbb{E}_{\xi} ||g_i(\mathbf{w}, \xi) - g_i(\mathbf{w}', \xi)||_2 \le C_q ||\mathbf{w} - \mathbf{w}'||_2$.

It is notable that these assumptions have been made in [6] which is important for analyzing variance reduction technique such as MSVR. We refer to Assumption 4.6 as mean Lipchitz continuity of zero-order (MLC0). Let B_1 and B_2 denote outer and inner batch sizes.

Theorem 4.7. Under Assumption 3.2 (A1), 4.5 and 4.6, by setting $\lambda = \Theta(\epsilon), \beta = \Theta(\min\{B_1, B_2\}\epsilon^2), \gamma = \Theta(B_2\epsilon^4), \eta = \Theta(\frac{B_1\sqrt{B_2}}{n}\epsilon^3)$, SONEX with $\gamma' = 1 - \gamma + \frac{n-B_1}{B_1(1-\gamma)}$ and $\gamma \leq \frac{1}{2}$ converges to an approximate ϵ -stationary solution of (1) within $T = O(\frac{n}{B_1\sqrt{B_2}}\epsilon^{-5})$ iterations.

Combining the above theorem with Theorem 4.4, we obtain the following guarantee:

Corollary 4.8. Under Assumption 3.2(A1), 4.3, 4.5 and 4.6, with the same setting as in Theorem 4.7, SONEX converges to a nearly ϵ -stationary solution of (1) within $T = O(\frac{n}{B_1\sqrt{B_2}}\epsilon^{-5})$ iterations.

We compare our proposed SONEX with SONX for solving (1). Under assumption 4.6, the rate of SONX is $O(1/\epsilon^6)$ which is worse than our result of $O(1/\epsilon^5)$ in Corollary 4.8.

4.2 A Double-loop Algorithm for Smooth or Weakly-Convex Inner Functions

In this subsection, we further improve our results to achieve the same convergence rate of $O(1/\epsilon^5)$ by (1) removing the MCL0 assumption instead assuming that f_i are convex; (2) designing a stochastic momentum method with a convergence guarantee for weakly convex g_i . It is worth mentioning that the convexity of f_i holds for a broad range of real applications such as group DRO and the application in non-convex constrained optimization as discussed in next section. Before introducing our new algorithm, we need to first reformulate the problem.

Let us first consider the scenario that satisfies Assumption 3.2 (A3). For simplicity of notation, we denote $f_{i,\lambda}$ by \bar{f}_i . When f_i is convex, we cast the problem (3) into a minimax formulation:

$$\min_{\mathbf{w} \in \mathbb{R}^d} F_{\lambda}(\mathbf{w}) := \frac{1}{n} \sum_{i=1}^n \max_{\mathbf{y}_i \in \mathbb{R}^{d_1}} \left\{ \mathbf{y}_i^{\mathsf{T}} g_i(\mathbf{w}) - \bar{f}_i^*(\mathbf{y}_i) \right\}, \tag{4}$$

where \bar{f}_i^* is the conjugate function of \bar{f}_i . In Appendix A.2, we show that this is also equivalent to the classical Nesterov's smoothing [19]. Under Assumption 3.2 (A3), \bar{f}_i is also C_f -Lipschitz continuous, convex, non-decreasing because of (2). As a consequence, $\mathrm{dom}(\bar{f}_i^*) \subset \{\mathbf{y}_i \in \mathbb{R}_+^{d_1} | \|\mathbf{y}_i\| \leq C_f\}$. This implies that, for any $\mathbf{y}_i \in \mathrm{dom}(\bar{f}_i^*)$, function $\mathbf{y}_i^\top g_i(\mathbf{w})$ is $\|\mathbf{y}_i\|_1 \rho_g$ -weakly convex in \mathbf{w} and also $\sqrt{d_1}C_f\rho_g$ -weakly convex because $\|\mathbf{y}_i\|_1 \leq \sqrt{d_1}\|\mathbf{y}_i\| \leq \sqrt{d_1}C_f$. This further implies that $F_{\lambda}(\cdot)$ is $\rho_{F_{\lambda}} := \sqrt{d_1}C_f\rho_g$ -weakly convex.

Different from last subsection, there is another challenge we need to deal with in order to develop a stochastic momentum method, which comes from that g_i are non-smooth. To tackle this challenge, we use another Moreau envelope smoothing of $F_{\lambda}(\cdot)$ and solve the following problem:

$$\min_{\mathbf{w} \in \mathbb{R}^d} \left\{ F_{\lambda,\nu}(\mathbf{w}) := \min_{\mathbf{z} \in \mathbb{R}^d} \left\{ F_{\lambda}(\mathbf{z}) + \frac{1}{2\nu} \|\mathbf{z} - \mathbf{w}\|^2 \right\} \right\}$$
 (5)

$$= \min_{\mathbf{w} \in \mathbb{R}^d} \left\{ \min_{\mathbf{z} \in \mathbb{R}^d} \max_{\mathbf{y} \in \mathbb{R}^{nd_1}} \frac{1}{n} \sum_{i=1}^n \mathbf{y}_i^\top g_i(\mathbf{z}) - \bar{f}_i^*(\mathbf{y}_i) + \frac{1}{2\nu} \|\mathbf{z} - \mathbf{w}\|^2 \right\},$$
(6)

where $\nu \in (0, \rho_{F_{\lambda}}^{-1})$ and $\mathbf{y} = [\mathbf{y}_{1}^{\top}, \dots, \mathbf{y}_{n}^{\top}]^{\top}$. The benefit of doing this is that (i) the resulting objective $F_{\lambda,\nu}(\mathbf{w})$ is smooth (cf. Lemma C.2), which allows us to employ stochastic momentum method; (ii) the inner $\min_{\mathbf{z}} \max_{\mathbf{y}}$ becomes a strongly-convex and strongly-concave problem due to that \bar{f}_{i} is smooth and its conjugate is strongly convex, and $\nu \in (0, \rho_{F_{\lambda}}^{-1})$ (cf. Lemma C.3).

Recall that $\nabla F_{\lambda,\nu}(\mathbf{w}) = \frac{1}{\nu}(\mathbf{w} - \operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}))$. Given an approximate solution of $\operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w})$, denoted by $\hat{\mathbf{z}}$, of the inner minimization problem in (5), we can use $\frac{1}{\nu}(\mathbf{w} - \hat{\mathbf{z}})$ as an inexact gradient $F_{\lambda,\nu}(\mathbf{w})$ to update \mathbf{w} with a momentum method in order to solve (5). To obtain an estimate $\hat{\mathbf{z}}_t$ at the t-th iteration, we employ a recent algorithm ALEXR [5] for solving the inner minimax problem. With an estimate $\hat{\mathbf{z}}_t$, we will update \mathbf{w}_{t+1} by a momentum update. We present the detailed steps of our double-loop algorithm named ALEXR2 in Algorithm 2, where $U_{\bar{f}_i^*}(\mathbf{y}_i, \mathbf{y}_i')$ is the Bregman divergence induced by \bar{f}_i^* , namely, $U_{\bar{f}_i^*}(\mathbf{y}_i, \mathbf{y}_i') := \bar{f}_i^*(\mathbf{y}_i) - \bar{f}_i^*(\mathbf{y}_i') - \langle \partial \bar{f}_i^*(\mathbf{y}_i'), \mathbf{y}_i - \mathbf{y}_i' \rangle$. For the sake of convergence analyses we need $U_{\bar{f}_i^*}$ to be bounded, as stated in the assumption below.

Algorithm 2 ALEXR2 for solving (6)

```
1: Input: T, \mathbf{w}_0 \in \mathbb{R}^d, \mathbf{v}_0 = 0, K_t, \alpha, \beta, \nu, \eta, \theta, \gamma > 0
  2: for t = 0, 1, \dots, T - 1 do
                 Set \mathbf{z}_{t,0} = \mathbf{z}_{t,-1} = \mathbf{w}_t and initialize \mathbf{y}_{t,0} = [\mathbf{y}_{t,0}^{(1)\top}, \dots, \mathbf{y}_{t,0}^{(n)\top}]^{\top} \in \mathbb{R}^{nd_1}.
                 for k=0,1,\ldots,K_t-1 do Sample \mathcal{B}_1^{t,k}\subset\{1,\ldots,n\} and two independent batches of data \mathcal{B}_{i,2}^{t,k} and \tilde{\mathcal{B}}_{i,2}^{t,k} from the
  4:
  5:
                        distribution of \xi_i for i \in \mathcal{B}_1^{t,k}.
                      Compute \tilde{g}_{t,k}^{(i)} = g_i(\mathbf{z}_{t,k}, \mathcal{B}_{i,2}^{t,k}) + \theta(g_i(\mathbf{z}_{t,k}, \mathcal{B}_{i,2}^{t,k}) - g_i(\mathbf{z}_{t,k-1}, \mathcal{B}_{i,2}^{t,k})) for i \in \mathcal{B}_1^{t,k}.

for i = 1, \dots, n do
\mathbf{y}_{t,k+1}^{(i)} = \begin{cases} \arg\max_{\mathbf{y}^{(i)}} \left\{ \mathbf{y}^{(i)^{\top}} \tilde{g}_{t,k}^{(i)} - \bar{f}_i^*(\mathbf{y}^{(i)}) - \frac{1}{\gamma} U_{\bar{f}_i^*}(\mathbf{y}^{(i)}, \mathbf{y}_{t,k}^{(i)}) \right\} & \text{if } i \in \mathcal{B}_1^{t,k} \\ \mathbf{y}_{t,k}^{(i)} & \text{if } i \notin \mathcal{B}_1^{t,k} \end{cases}
  6:
  7:
  8:
                       end for Compute G_{t,k} = \frac{1}{|\mathcal{B}_1^{t,k}|} \sum_{i \in \mathcal{B}_1^{t,k}} [\partial g_i(\mathbf{z}_{t,k}, \tilde{\mathcal{B}}_{i,2}^{t,k})]^{\top} \mathbf{y}_{t,k+1}^{(i)}
  9:
10:
                        \text{Update } \mathbf{z}_{t,k+1} = \arg\min_{\mathbf{z} \in \mathbb{R}^d} \left\{ \left\langle G_{t,k},\, \mathbf{z} \right\rangle + \tfrac{1}{2\nu} \left\| \mathbf{z} - \mathbf{w}_t \right\|^2 + \tfrac{1}{2\eta} \left\| \mathbf{z} - \mathbf{z}_{t,k} \right\|^2 \right\}
11:
12:
                 Let G_t = \frac{\beta}{\nu} (\mathbf{w}_t - \mathbf{z}_{t,K_t})
13:
                  Update \mathbf{v}_{t+1} = (1 - \beta)\mathbf{v}_t + \beta G_t
14:
                   Update \mathbf{w}_{t+1} = \mathbf{w}_t - \alpha \mathbf{v}_{t+1} or use Adam-type update
15:
16: end for
17: Output: \mathbf{w}_{\tau} with \tau randomly sampled from \{1, 2, \dots T\}
```

Besides, we would like to point out that the sequences of y_i can be updated similar to u_i sequences in SONEX similar to [5], i.e., $y_{t,k}^i = \nabla f_{i,\lambda}(u_{t,k}^i)$, where $u_{t,k+1}^i = (1-\hat{\gamma})u_{t,k}^i + \hat{\gamma}g_i(\mathbf{z}_{t,k};\mathcal{B}_{i,2}^{t,k}) + \hat{\gamma}\theta(g_i(\mathbf{z}_{t,k};\mathcal{B}_{i,2}^{t,k}) - g_i(\mathbf{z}_{t,k-1};\mathcal{B}_{i,2}^{t,k}))$, where $\hat{\gamma} = \gamma/(1+\gamma)$.

Assumption 4.9. We assume that f_i is a function s.t. $U_{\bar{f}^*}(y_1, y_2)$ is bounded for any $y_1, y_2 \in dom(\bar{f}_i^*) \subset \mathbb{R}^{d_1}$.

Remark: We point out that this condition is not strong under our setting where \bar{f}_i is lipschitz continuous and $dom(\bar{f}_i^*)$ is consequently bounded and it is satisfied by many practical convex lipschitz-continuous functions such as hinge function, smoothed hinge function(i.e. $f_{i,\lambda}$ or \bar{f}_i in this paper), etc.

The following theorem states the convergence of ALEXR2 with proof given in Section C.

Theorem 4.10. Suppose assumption 3.2 (A3), 4.5 and 4.9 hold and $\lambda = \epsilon/C_f$ in (4). For any $\epsilon > 0$, there exists $\theta \in (0,1)$ with $1-\theta = O(\epsilon^2)$ such that, by setting $\eta = \frac{1-\theta}{\theta} L_f$ and $\gamma = \frac{(1-\theta)n}{B_1}$, $\beta \leq \frac{1}{2}$, $\alpha = \frac{\beta}{2L_{F_{\lambda,\nu}}}$ and $K_t = \tilde{O}\left(\frac{n}{B_1B_2\epsilon^3} + \frac{1}{\epsilon^3}\right)$, ALEXR2 returns \mathbf{w}_{τ} as a nearly ϵ -stationary solution of (4) in expectation within $T = O(\epsilon^{-2})$ and a total iteration complexity of $\tilde{O}\left(\frac{n}{B_1B_2\epsilon^5} + \frac{1}{\epsilon^5}\right)$.

Remark: We can easily extend the above result to Assumption 3.2 (A2). When g_i are smooth, we do not need the monotonicity of f_i as the minimax problem $\min_{\mathbf{z}} \max_{\mathbf{y}}$ is still guaranteed to be strongly-convex and strongly-concave when $\nu \in (0, \rho_{F_{\lambda}}^{-1})$, where $\rho_{F_{\lambda}} := \sqrt{d_1} C_f L_g$ is the weak convexity parameter of F_{λ} .

Finally, we show that when g_i are smooth, we can also obtain a nearly ϵ -stationary solution to the original problem (1). Different from the result in Theorem 4.7, the above theorem only guarantees a nearly ϵ -stationary solution to F_{λ} . To address this gap, we can recover a nearly ϵ -stationary solution to (1) from \mathbf{w}_{τ} by running ALEXR starting with \mathbf{w}_{τ} with $K = \tilde{O}(\epsilon^{-5})$ iterations. This result is stated as the following corollary with the proof given in Section C.

Corollary 4.11. Suppose assumption 3.2 (A2), 4.3, 4.5, 4.9 hold. For any
$$\epsilon \leq \min\left\{c, \frac{\delta^2}{16nC_g^2L_g}\right\}$$
, let $\hat{\mathbf{w}}_{\tau} = ALEXR(\mathbf{w}_{\tau}, K)$ with $K = \tilde{O}\left(\frac{n}{B_1B_2\epsilon^5} + \frac{1}{\epsilon^5}\right)$, $\eta = \frac{1-\theta}{\theta}L_f$ and $\gamma = \frac{(1-\theta)n}{B_1}$ for $\theta \in (0, 1)$

with $1 - \theta = O(\epsilon^4)$. $\hat{\mathbf{w}}_{\tau}$ is a nearly ϵ -stationary solution of (1) and it is found with a total iteration complexity of $\tilde{O}(\frac{n}{B_1B_2\epsilon^5} + \frac{1}{\epsilon^5})$.

5 Smoothed Hinge Penalty Method for Constrained Optimization

In this section, we consider a constrained optimization problem with m > 1 inequality constraints:

$$\min_{\mathbf{w}} g_0(\mathbf{w}) \text{ s.t. } g_i(\mathbf{w}) \le 0, i = 1, \cdots, m \tag{7}$$

where $g_i : \mathbb{R}^d \to \mathbb{R}$ is continuous and satisfies $g_i(\mathbf{w}) = \mathbb{E}g_i(\mathbf{w}, \xi_i)$ and the expectation is taken over the random variable ξ_i for $i = 0, 1, \dots, m$. Following [8], we consider an exact penalty method for (7) by solving the following unconstrained problem:

$$\min_{\mathbf{w} \in \mathbb{R}^d} \Phi(\mathbf{w}) := g_0(\mathbf{w}) + \frac{\rho}{m} \sum_{i=1}^m [g_i(\mathbf{w})]_+$$
 (8)

where ρ is a sufficiently large number. Let $f(\cdot) := \rho[\cdot]_+$ so the penalty term $\frac{\rho}{m} \sum_{i=1}^m [g_i(\mathbf{w})]_+$ has the same structure as $F(\mathbf{w})$ in (1). Yang et al. [8] have employed SONX for solving the above problem and established a complexity of $O(1/\epsilon^6)$ for finding a nearly ϵ -KKT solution. Below, we apply the smoothing idea and ALEXR2 or SONEX for improving the convergence rate. Our key idea is to optimize an outer smoothed problem:

$$\min_{\mathbf{w} \in \mathbb{R}^d} \Phi_{\lambda}(\mathbf{w}) := g_0(\mathbf{w}) + \frac{1}{m} \sum_{i=1}^m f_{\lambda}(g_i(\mathbf{w})) = g_0(\mathbf{w}) + \frac{1}{m} \sum_{i=1}^m \max_{y_i \in [0,\rho]} \left\{ y_i g_i(\mathbf{w}) - \frac{\lambda}{2} y_i^2 \right\}, \quad (9)$$

where $f_{\lambda}(z) := \min_{z' \in \mathbb{R}} f(z') + \frac{1}{2\lambda} (z - z')^2$. Except for the term g_0 , the objective function of (9) has the same structure as (3) and (4) with $\bar{f}_i^*(y_i) = \frac{\lambda}{2} y_i^2 + \mathbf{1}_{[0,\rho]}(y_i)$.

Assumption 5.1. There exists a constant $\delta > 0$ such that $\sigma_{min}(\mathbf{J}(\mathbf{w})) \geq \delta$ for any matrix $\mathbf{J}(\mathbf{w}) = [\mathbf{h}_1(\mathbf{w}), \dots, \mathbf{h}_m(\mathbf{w})] \in \mathbb{R}^{d \times m}$ with $\mathbf{h}_i(\mathbf{w}) \in \partial g_i(\mathbf{w})$ and any \mathbf{w} satisfying $\max_{i=1,\dots,m} g_i(\mathbf{w}) > 0$.

Under this assumption, we have the following proposition whose proof is given in Section D.1.

Proposition 5.2. Suppose Assumption 5.1 hold. If $\rho > \frac{m(C_g+1)}{\delta}$ and $\lambda = \frac{\epsilon}{\rho}$, a nearly ϵ -stationary solution \mathbf{w} to (9) is also a nearly $O(\epsilon)$ -KKT solution to the original problem (7) in the sense that there exist $\hat{\mathbf{w}}$ and $\nu_i \geq 0$ for $i = 1, \ldots, m$ such that $\|\mathbf{w} - \hat{\mathbf{w}}\| \leq \epsilon$, $\mathrm{dist}(0, \partial g_0(\hat{\mathbf{w}}) + \sum_{i=1}^m \partial g_i(\hat{\mathbf{w}})\nu_i) \leq O(\epsilon)$, $\max_{i=1,\ldots,m} g_i(\hat{\mathbf{w}}) \leq O(\epsilon)$ and $|g_i(\hat{\mathbf{w}})\nu_i| \leq O(\epsilon)$, $\forall i = 1, 2, \cdots, m$.

To find a nearly ϵ -stationary solution to (9), we can adapt either SONEX if g_i are smooth or ALEXR2 if g_i are weakly convex with a minor change by including the stochastic gradient of g_0 in the calculation of G_t . The complexity of this approach is presented below with proofs given in Section D.2.

Theorem 5.3. Suppose assumption 4.5 and 5.1 hold, and the stochastic (sub)gradient of g_0 has bounded variance. Let $\rho > m(C_q + 1)/\delta$ and $\lambda = \epsilon/\rho$.

- if $\{g_i\}_{i=0}^m$ are weakly convex, then with $\tilde{\mathcal{O}}\left(\frac{m}{B_1B_2\delta^4\epsilon^5}\right)$ iterations ALEXR2 find a nearly ϵ -KKT solution \mathbf{w} for (7) satisfying that there exist $\hat{\mathbf{w}}$ and $\nu_i \geq 0$ for $i=1,\ldots,m$ such that $\mathbb{E}\|\mathbf{w}-\hat{\mathbf{w}}\| \leq \epsilon$ and $\mathbb{E} dist(0,\partial g_0(\hat{\mathbf{w}})+\sum_{i=1}^m \partial g_i(\hat{\mathbf{w}})\nu_i)) \leq O(\epsilon)$ and it holds with probability $1-O(\epsilon)$ that $\max_{i=1,\ldots,m} g_i(\hat{\mathbf{w}}) \leq O(\epsilon)$ and $|g_i(\hat{\mathbf{w}})\nu_i| \leq O(\epsilon)$, $\forall i=1,2,\cdots,m$.
- if $\{g_i\}_{i=0}^m$ are smooth, then with $\tilde{\mathcal{O}}\left(\frac{m}{B_1\sqrt{B_2}\delta^4\epsilon^5}\right)$ iterations SONEX can find an ϵ -KKT solution \mathbf{w} for (7) similar as above except for $\hat{\mathbf{w}} = \mathbf{w}$.

Notably, our method not only improve the rate but also enjoy an improved dependence on δ , compared to $O(\epsilon^{-6}\delta^{-6})$ in prior work [8].

 $[\]overline{\ \ }^4 ext{If } |g_i(\mathbf{w})| < +\infty ext{ for any } i ext{ and } \mathbf{w}, ext{ this high probability result can be replaced by } \mathbb{E}[\max_{i=1,\dots,m} g_i(\hat{\mathbf{w}})] \le O(\epsilon)$ and $\mathbb{E}[|q_i(\hat{\mathbf{w}})\nu_i|] < O(\epsilon), \forall i=1,2,\cdots,m.$




Figure 1: Training loss curves (left three) and testing accuracy (right one) of different methods for Group DRO with CVaR ratio r=0.15 on different datasets.




Figure 2: Training curves of AUC values (fig 1,3) and constraint violation (fig 2,4) of different methods. The format of legend is "Algorithm(penalty function, ρ)", and SH, H, smH mean square hinge, hinge and smoothed hinge, respectively.




Figure 3: Training curves of Target \triangle ACC values (fig 1,3) and constraint violation (fig 2,4) of different methods. The format of label is "Algorithm(penalty function, ρ)", and SH, smH mean square hinge and smoothed hinge, respectively.

6 Numerical Experiments

We conduct experiments to verify the effectiveness of the proposed algorithms. For non-smooth non-convex FCCO, we consider GDRO with CVaR divergence [5]. For non-convex constrained optimization, we follow [8] and consider two tasks, namely AUC maximization with ROC fairness constraints and continual learning with non-forgetting constraints.

GDRO with CVaR divergence. We consider the following GDRO with CVaR divergence, which minimizes top-k worst groups' losses [5]:

$$\min_{\theta} \min_{s} s + \frac{1}{n} \sum_{g=1}^{n} \frac{1}{r} [\mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{D}_g} \ell(\theta; (\mathbf{x}, y)) - s]_{+}$$
(10)

where n is the number of groups, r = k/n, and \mathcal{D}_q denotes the data of the g-th group.

We use 3 datasets: Camelyon17, Amazon [34], and CelebA [35]. The first two datasets are from WILDS Benchmark for evaluating methods tackling the distributional shift [36], where Camelyon17 has 30 groups and Amazon has 1252 groups. CelebA is a large-scale facial attribute dataset containing over 200,000 celebrity images annotated with 40 binary attributes. We select 4 binary attributes 'Attractive', 'Mouth_Slightly_Open', 'Male' and 'Blonde_Hair' and construct 16 groups, where 'Blonde_Hair' also serves as the label for classification. We compare SONEX with SONX and OOA, where OOA is a stochastic primal-dual algorithm ⁵. We use pretrained Densenet121 [38], Distilbert [39] and ResNet50 [40] for Camelyon17, Amazon and CelebA, respectively. We perform Adam-type update for SONEX on Amazon dataset and momentum-type update on Camelyon17 and CelebA datasets. We run each experiment for 3 random seeds and report their average performance. The hyperparameter tuning is discussed in Appendix F.1.

Results. We report our result under setting of r=0.15 in Figure 1, where the first 3 figures are the loss curves on the training dataset while the last one is the test accuracy. Our experiment results shows that SONEX performs better than SONX and OOA on GDRO tasks.

⁵We extend the original OOA [37] to GDRO with CVaR setting as in [5]

AUC maximization with ROC fairness constraints. The formulation is given in Appendix F.2, where the objective is AUC loss and constraints specify the tolerance of the gap between false positive rates (true positive rates) of two sensitive groups at different classification thresholds. For this experiment, we follow almost the same setting as in [8]. Two datasets are used, namely Adult [41] and COMPAS [42], which contain male/female, Caucasian/non-Caucasian groups, respectively. We set thresholds at $\Gamma = \{-3, -2, -1, \cdots, 3\}$ and the tolerance $\kappa = 0.005$, which gives us 14 constraints. We learn a simple neural network with 2 hidden layers. We compare ALEXR2 with the method in [7] that optimizes a squared-hinge penalty function with the SOX algorithm [2], the method in [8] that optimizes a hinge penalty function with the SONX algorithm [6], the double-loop method ICPPAC [10]. We perform Adam-type update for ALEXR2. We run each method for totally 60 epoches with a batch size of 128, repeat five times with different seeds and then report average of the AUC scores and constraint values. Hyperparameter tuning is presented in Appendix F.2.

Results. We compare the training curves of objective AUC values and the constraint violation measured by the worst constraint function value at each epoch for different methods in Figure 2 on different datasets. These results demonstrate that ALEXR2 optimizing smoothed hinge penalty function can better maximize AUC value on both datasets than the baseline methods while still having similar constraint satisfaction.

Continual Learning with non-forgetting constraints. We follow a similar experimental setup to [7] for fine-tuning a CLIP model for autonomous driving on the BDD100K dataset [43], a large-scale, multi-task driving image dataset. The formulation is given in Appendix F.3, where the objective is the contrastive loss on a target task (e.g., classifying foggy and overcast condition), and the constraints specify the logistic loss of the new model should not be larger than that of the old model for other different classes (e.g., clear, snowy, rainy, partly cloudy). Since the objective function is another FCCO, we modify SONEX such that the gradient estimator of the objective is computed similar as the smoothed square penalty function. We compare with SOX that optimizes the squared hinge loss use the Adam-type update for both methods. We do not compare with SONX that optimizes the hinge penalty function with SGD-type update as it fails for learning the Transformer network used in this experiment. Hyperparameter tuning is presented in Appendix F.3.

Results. We present training curves of accuracy improvement and constraint violation on different target tasks in Figure 3. It shows that our method can achieve higher accuracy on target tasks than the baseline method of using squared hinge function while retaining similar constraint satisfaction.

7 Conclusions and Discussion

In this paper, we have considered non-smooth non-convex finite-sum coupled compositional optimization, where the out functions are non-smooth and inner functions are smooth or weakly convex. We proposed stochastic momentum methods to improve the convergence rate and accelerate the training for deep neural networks. We also considered the applications of the proposed stochastic momentum methods to solving non-convex inequality constrained optimization problems and derived state-of-the-art results for finding an ϵ -KKT solution. Our experiments demonstrate the effectiveness of our methods. One limitation of our results is that our single-loop method SONEX still has a worse complexity than the double-loop method ALEXR2 in terms of dependence on the inner batch size.

Acknowledgments and Disclosure of Funding

X. Chen, B. Wang, M. Yang and T. Yang were partially supported by the National Science Foundation Career Award 2246753, the National Science Foundation Award 2246757 and 2306572. Q. Lin was partially supported by National Science Foundation Award 2147253.

References

- [1] Tianbao Yang. Algorithmic foundations of empirical X-risk minimization. arXiv preprint arXiv:2206.00439, 2022.
- [2] Bokun Wang and Tianbao Yang. Finite-sum coupled compositional stochastic optimization: Theory and applications. In *International Conference on Machine Learning*, pages 23292–23317. PMLR, 2022.
- [3] Dixian Zhu, Gang Li, Bokun Wang, Xiaodong Wu, and Tianbao Yang. When auc meets dro: Optimizing partial auc for deep learning with non-convex convergence guarantee. In *International Conference on Machine Learning*, pages 27548–27573. PMLR, 2022.
- [4] Qi Qi, Youzhi Luo, Zhao Xu, Shuiwang Ji, and Tianbao Yang. Stochastic optimization of areas under precision-recall curves with provable convergence. *Advances in neural information processing systems*, 34:1752–1765, 2021.
- [5] Bokun Wang and Tianbao Yang. Alexr: An optimal single-loop algorithm for convex finite-sum coupled compositional stochastic optimization. *arXiv* preprint arXiv:2312.02277, 2023.
- [6] Quanqi Hu, Dixian Zhu, and Tianbao Yang. Non-smooth weakly-convex finite-sum coupled compositional optimization. *Advances in Neural Information Processing Systems*, 36:5348–5403, 2023.
- [7] Gang Li, Wendi Yu, Yao Yao, Wei Tong, Yingbin Liang, Qihang Lin, and Tianbao Yang. Model developmental safety: A safety-centric method and applications in vision-language models. *arXiv* preprint *arXiv*:2410.03955, 2024.
- [8] Ming Yang, Gang Li, Quanqi Hu, Qihang Lin, and Tianbao Yang. Single-loop algorithms for stochastic non-convex optimization with weakly-convex constraints. *arXiv preprint arXiv:2504.15243*, 2025.
- [9] Runchao Ma, Qihang Lin, and Tianbao Yang. Quadratically regularized subgradient methods for weakly convex optimization with weakly convex constraints. In *International Conference on Machine Learning*, pages 6554–6564. PMLR, 2020.
- [10] Digvijay Boob, Qi Deng, and Guanghui Lan. Stochastic first-order methods for convex and nonconvex functional constrained optimization. *Mathematical Programming*, 197(1):215–279, 2023.
- [11] Yankun Huang and Qihang Lin. Oracle complexity of single-loop switching subgradient methods for non-smooth weakly convex functional constrained optimization. *Advances in Neural Information Processing Systems*, 36:61327–61340, 2023.
- [12] Wei Liu and Yangyang Xu. A single-loop spider-type stochastic subgradient method for expectation-constrained nonconvex nonsmooth optimization. *arXiv* preprint arXiv:2501.19214, 2025.
- [13] Mengdi Wang, Ethan X Fang, and Han Liu. Stochastic compositional gradient descent: algorithms for minimizing compositions of expected-value functions. *Mathematical Programming*, 161:419–449, 2017.
- [14] Mengdi Wang, Ji Liu, and Ethan X Fang. Accelerating stochastic composition optimization. *Journal of Machine Learning Research*, 18(105):1–23, 2017.
- [15] Yifan Hu, Siqi Zhang, Xin Chen, and Niao He. Biased stochastic first-order methods for conditional stochastic optimization and applications in meta learning. Advances in Neural Information Processing Systems, 33:2759–2770, 2020.
- [16] Wei Jiang, Gang Li, Yibo Wang, Lijun Zhang, and Tianbao Yang. Multi-block-single-probe variance reduced estimator for coupled compositional optimization. Advances in Neural Information Processing Systems, 35:32499–32511, 2022.
- [17] Zhuoning Yuan, Yuexin Wu, Zi-Hao Qiu, Xianzhi Du, Lijun Zhang, Denny Zhou, and Tianbao Yang. Provable stochastic optimization for global contrastive learning: Small batch does not harm performance. In *International Conference on Machine Learning*, pages 25760–25782. PMLR, 2022.
- [18] Zi-Hao Qiu, Quanqi Hu, Zhuoning Yuan, Denny Zhou, Lijun Zhang, and Tianbao Yang. Not all semantics are created equal: contrastive self-supervised learning with automatic temperature individualization. In Proceedings of the 40th International Conference on Machine Learning, pages 28389–28421, 2023.
- [19] Yu Nesterov. Smooth minimization of non-smooth functions. Math. Program., 103(1):127–152, May 2005.

- [20] Guy Kornowski and Ohad Shamir. Oracle complexity in nonsmooth nonconvex optimization. *Journal of Machine Learning Research*, 23(314):1–44, 2022.
- [21] Damek Davis and Dmitriy Drusvyatskiy. Stochastic model-based minimization of weakly convex functions. SIAM Journal on Optimization, 29(1):207–239, 2019.
- [22] Damek Davis and Benjamin Grimmer. Proximally guided stochastic subgradient method for nonsmooth, nonconvex problems. SIAM Journal on Optimization, 29(3):1908–1930, 2019.
- [23] Ahmet Alacaoglu, Yura Malitsky, and Volkan Cevher. Convergence of adaptive algorithms for constrained weakly convex optimization. Advances in Neural Information Processing Systems, 34:14214–14225, 2021.
- [24] Hassan Rafique, Mingrui Liu, Qihang Lin, and Tianbao Yang. Weakly-convex—concave min—max optimization: provable algorithms and applications in machine learning. *Optimization Methods and Software*, 37(3):1087–1121, 2022.
- [25] Renbo Zhao. A primal-dual smoothing framework for max-structured non-convex optimization. Mathematics of operations research, 49(3):1535–1565, 2024.
- [26] Zhichao Jia and Benjamin Grimmer. First-order methods for nonsmooth nonconvex functional constrained optimization with or without slater points. *arXiv* preprint arXiv:2212.00927, 2022.
- [27] Quoc Tran Dinh, Deyi Liu, and Lam Nguyen. Hybrid variance-reduced sgd algorithms for minimax problems with nonconvex-linear function. Advances in Neural Information Processing Systems, 33:11096– 11107, 2020.
- [28] Zichong Li, Pin-Yu Chen, Sijia Liu, Songtao Lu, and Yangyang Xu. Stochastic inexact augmented Lagrangian method for nonconvex expectation constrained optimization. *Computational Optimization and Applications*, 87(1):117–147, 2024.
- [29] Ahmet Alacaoglu and Stephen J Wright. Complexity of single loop algorithms for nonlinear programming with stochastic objective and constraints. In *International Conference on Artificial Intelligence and Statistics*, pages 4627–4635. PMLR, 2024.
- [30] Zichong Li, Pin-Yu Chen, Sijia Liu, Songtao Lu, and Yangyang Xu. Rate-improved inexact augmented lagrangian method for constrained nonconvex optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 2170–2178. PMLR, 2021.
- [31] Ellen Fukuda and Masao Fukushima. A note on the squared slack variables technique for nonlinear optimization. *Journal of the Operations Research Society of Japan*, 60:262–270, 07 2017.
- [32] Lijun Ding and Stephen J Wright. On squared-variable formulations. arXiv preprint arXiv:2310.01784, 2023.
- [33] Marien Renaud, Valentin De Bortoli, Arthur Leclaire, and Nicolas Papadakis. From stability of langevin diffusion to convergence of proximal mcmc for non-log-concave sampling. arXiv preprint arXiv:2505.14177, 2025.
- [34] Peter Bandi, Oscar Geessink, Quirine Manson, Marcory Van Dijk, Maschenka Balkenhol, Meyke Hermsen, Babak Ehteshami Bejnordi, Byungjae Lee, Kyunghyun Paeng, Aoxiao Zhong, et al. From detection of individual metastases to classification of lymph node status at the patient level: the camelyon17 challenge. IEEE Transactions on Medical Imaging, 2018.
- [35] Ziwei Liu, Ping Luo, Xiaogang Wang, and Xiaoou Tang. Deep learning face attributes in the wild. In *Proceedings of the IEEE international conference on computer vision*, pages 3730–3738, 2015.
- [36] Pang Wei Koh, Shiori Sagawa, Henrik Marklund, Sang Michael Xie, Marvin Zhang, Akshay Balsubramani, Weihua Hu, Michihiro Yasunaga, Richard Lanas Phillips, Irena Gao, et al. Wilds: A benchmark of in-the-wild distribution shifts. In *International conference on machine learning*, pages 5637–5664. PMLR, 2021.
- [37] Shiori Sagawa, Pang Wei Koh, Tatsunori B Hashimoto, and Percy Liang. Distributionally robust neural networks. In *International Conference on Learning Representations*, 2019.
- [38] Gao Huang, Zhuang Liu, Laurens Van Der Maaten, and Kilian Q Weinberger. Densely connected convolutional networks. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 4700–4708, 2017.

- [39] Victor Sanh, Lysandre Debut, Julien Chaumond, and Thomas Wolf. Distilbert, a distilled version of bert: smaller, faster, cheaper and lighter. *arXiv preprint arXiv:1910.01108*, 2019.
- [40] Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 770–778, 2016.
- [41] Barry Becker and Ronny Kohavi. Adult. UCI Machine Learning Repository, 1996. DOI: https://doi.org/10.24432/C5XW20.
- [42] Julia Angwin, Jeff Larson, Surya Mattu, and Lauren Kirchner. Machine bias. ProPublica, 2016.
- [43] Fisher Yu, Haofeng Chen, Xin Wang, Wenqi Xian, Yingying Chen, Fangchen Liu, Vashisht Madhavan, and Trevor Darrell. Bdd100k: A diverse driving dataset for heterogeneous multitask learning. In *Proceedings of the IEEE/CVF conference on computer vision and pattern recognition*, pages 2636–2645, 2020.
- [44] Junyu Zhang and Lin Xiao. Multilevel composite stochastic optimization via nested variance reduction. SIAM Journal on Optimization, 31(2):1131–1157, 2021.
- [45] Maurice Sion. On general minimax theorems. 1958.
- [46] Zhize Li, Hongyan Bao, Xiangliang Zhang, and Peter Richtarik. Page: A simple and optimal probabilistic gradient estimator for nonconvex optimization. In *Proceedings of the 38th International Conference on Machine Learning*, pages 6286–6295, 2021.
- [47] Heinz H Bauschke, Patrick L Combettes, Heinz H Bauschke, and Patrick L Combettes. *Correction to:* convex analysis and monotone operator theory in Hilbert spaces. Springer, 2017.
- [48] Zhishuai Guo, Yi Xu, Wotao Yin, Rong Jin, and Tianbao Yang. Unified convergence analysis for adaptive optimization with moving average estimator. *Machine Learning*, 2021.
- [49] R Tyrrell Rockafellar, Stanislav Uryasev, et al. Optimization of conditional value-at-risk. *Journal of risk*, 2:21–42, 2000.
- [50] Jianmo Ni, Jiacheng Li, and Julian McAuley. Justifying recommendations using distantly-labeled reviews and fine-grained aspects. In Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing and the 9th International Joint Conference on Natural Language Processing (EMNLP-IJCNLP), 2019.
- [51] Robin Vogel, Aurelien Bellet, and Stephan Clemencon. Learning fair scoring functions: Bipartite ranking under roc-based fairness constraints. In *International conference on artificial intelligence and statistics*, pages 784–792. PMLR, 2021.
- [52] Alec Radford, Jong Wook Kim, Chris Hallacy, Aditya Ramesh, Gabriel Goh, Sandhini Agarwal, Girish Sastry, Amanda Askell, Pamela Mishkin, Jack Clark, et al. Learning transferable visual models from natural language supervision. In *International conference on machine learning*, pages 8748–8763. PMLR, 2021.
- [53] Christoph Schuhmann, Richard Vencu, Romain Beaumont, Robert Kaczmarczyk, Clayton Mullis, Aarush Katta, Theo Coombes, Jenia Jitsev, and Aran Komatsuzaki. Laion-400m: Open dataset of clip-filtered 400 million image-text pairs. arXiv preprint arXiv:2111.02114, 2021.

NeurIPS Paper Checklist

1. Claims

Question: Do the main claims made in the abstract and introduction accurately reflect the paper's contributions and scope?

Answer: [Yes]

Justification: We have detailed explanation on all items described in the abstract presented in the paper.

Guidelines:

- The answer NA means that the abstract and introduction do not include the claims made in the paper.
- The abstract and/or introduction should clearly state the claims made, including the contributions made in the paper and important assumptions and limitations. A No or NA answer to this question will not be perceived well by the reviewers.
- The claims made should match theoretical and experimental results, and reflect how much the results can be expected to generalize to other settings.
- It is fine to include aspirational goals as motivation as long as it is clear that these goals are not attained by the paper.

2. Limitations

Question: Does the paper discuss the limitations of the work performed by the authors?

Answer: [Yes]

Justification: We discussed the limitation of this in the conclusion section.

Guidelines:

- The answer NA means that the paper has no limitation while the answer No means that the paper has limitations, but those are not discussed in the paper.
- The authors are encouraged to create a separate "Limitations" section in their paper.
- The paper should point out any strong assumptions and how robust the results are to violations of these assumptions (e.g., independence assumptions, noiseless settings, model well-specification, asymptotic approximations only holding locally). The authors should reflect on how these assumptions might be violated in practice and what the implications would be.
- The authors should reflect on the scope of the claims made, e.g., if the approach was only tested on a few datasets or with a few runs. In general, empirical results often depend on implicit assumptions, which should be articulated.
- The authors should reflect on the factors that influence the performance of the approach. For example, a facial recognition algorithm may perform poorly when image resolution is low or images are taken in low lighting. Or a speech-to-text system might not be used reliably to provide closed captions for online lectures because it fails to handle technical jargon.
- The authors should discuss the computational efficiency of the proposed algorithms and how they scale with dataset size.
- If applicable, the authors should discuss possible limitations of their approach to address problems of privacy and fairness.
- While the authors might fear that complete honesty about limitations might be used by reviewers as grounds for rejection, a worse outcome might be that reviewers discover limitations that aren't acknowledged in the paper. The authors should use their best judgment and recognize that individual actions in favor of transparency play an important role in developing norms that preserve the integrity of the community. Reviewers will be specifically instructed to not penalize honesty concerning limitations.

3. Theory assumptions and proofs

Question: For each theoretical result, does the paper provide the full set of assumptions and a complete (and correct) proof?

Answer: [Yes]

Justification: Assumptions are presented in the main paper. The detailed theorem statements and proofs are presented in the appendix.

Guidelines:

- The answer NA means that the paper does not include theoretical results.
- All the theorems, formulas, and proofs in the paper should be numbered and crossreferenced.
- All assumptions should be clearly stated or referenced in the statement of any theorems.
- The proofs can either appear in the main paper or the supplemental material, but if they appear in the supplemental material, the authors are encouraged to provide a short proof sketch to provide intuition.
- Inversely, any informal proof provided in the core of the paper should be complemented by formal proofs provided in appendix or supplemental material.
- Theorems and Lemmas that the proof relies upon should be properly referenced.

4. Experimental result reproducibility

Question: Does the paper fully disclose all the information needed to reproduce the main experimental results of the paper to the extent that it affects the main claims and/or conclusions of the paper (regardless of whether the code and data are provided or not)?

Answer: [Yes]

Justification: We provide all the information needed to reproduce the experimental results in the "Numerical Experiments" section and related sections in the Appendix.

Guidelines:

- The answer NA means that the paper does not include experiments.
- If the paper includes experiments, a No answer to this question will not be perceived
 well by the reviewers: Making the paper reproducible is important, regardless of
 whether the code and data are provided or not.
- If the contribution is a dataset and/or model, the authors should describe the steps taken to make their results reproducible or verifiable.
- Depending on the contribution, reproducibility can be accomplished in various ways. For example, if the contribution is a novel architecture, describing the architecture fully might suffice, or if the contribution is a specific model and empirical evaluation, it may be necessary to either make it possible for others to replicate the model with the same dataset, or provide access to the model. In general, releasing code and data is often one good way to accomplish this, but reproducibility can also be provided via detailed instructions for how to replicate the results, access to a hosted model (e.g., in the case of a large language model), releasing of a model checkpoint, or other means that are appropriate to the research performed.
- While NeurIPS does not require releasing code, the conference does require all submissions to provide some reasonable avenue for reproducibility, which may depend on the nature of the contribution. For example
 - (a) If the contribution is primarily a new algorithm, the paper should make it clear how to reproduce that algorithm.
- (b) If the contribution is primarily a new model architecture, the paper should describe the architecture clearly and fully.
- (c) If the contribution is a new model (e.g., a large language model), then there should either be a way to access this model for reproducing the results or a way to reproduce the model (e.g., with an open-source dataset or instructions for how to construct the dataset).
- (d) We recognize that reproducibility may be tricky in some cases, in which case authors are welcome to describe the particular way they provide for reproducibility. In the case of closed-source models, it may be that access to the model is limited in some way (e.g., to registered users), but it should be possible for other researchers to have some path to reproducing or verifying the results.

5. Open access to data and code

Question: Does the paper provide open access to the data and code, with sufficient instructions to faithfully reproduce the main experimental results, as described in supplemental material?

Answer: [No]

Justification: The data we used are public datasets. Code will be released after well-organized.

Guidelines:

- The answer NA means that paper does not include experiments requiring code.
- Please see the NeurIPS code and data submission guidelines (https://nips.cc/public/guides/CodeSubmissionPolicy) for more details.
- While we encourage the release of code and data, we understand that this might not be possible, so "No" is an acceptable answer. Papers cannot be rejected simply for not including code, unless this is central to the contribution (e.g., for a new open-source benchmark).
- The instructions should contain the exact command and environment needed to run to reproduce the results. See the NeurIPS code and data submission guidelines (https://nips.cc/public/guides/CodeSubmissionPolicy) for more details.
- The authors should provide instructions on data access and preparation, including how
 to access the raw data, preprocessed data, intermediate data, and generated data, etc.
- The authors should provide scripts to reproduce all experimental results for the new proposed method and baselines. If only a subset of experiments are reproducible, they should state which ones are omitted from the script and why.
- At submission time, to preserve anonymity, the authors should release anonymized versions (if applicable).
- Providing as much information as possible in supplemental material (appended to the paper) is recommended, but including URLs to data and code is permitted.

6. Experimental setting/details

Question: Does the paper specify all the training and test details (e.g., data splits, hyper-parameters, how they were chosen, type of optimizer, etc.) necessary to understand the results?

Answer: [Yes]

Justification: We provided all the implementation details in the "Numerical Experiments" section and related sections in Appendix.

Guidelines:

- The answer NA means that the paper does not include experiments.
- The experimental setting should be presented in the core of the paper to a level of detail that is necessary to appreciate the results and make sense of them.
- The full details can be provided either with the code, in appendix, or as supplemental material.

7. Experiment statistical significance

Question: Does the paper report error bars suitably and correctly defined or other appropriate information about the statistical significance of the experiments?

Answer: [Yes]

Justification: We ran multiple trails for each experiment setting and present the error bars in the plot or table. The error bar in this paper is 1-sigma standard deviation over the random seeds.

Guidelines:

- The answer NA means that the paper does not include experiments.
- The authors should answer "Yes" if the results are accompanied by error bars, confidence intervals, or statistical significance tests, at least for the experiments that support the main claims of the paper.

- The factors of variability that the error bars are capturing should be clearly stated (for example, train/test split, initialization, random drawing of some parameter, or overall run with given experimental conditions).
- The method for calculating the error bars should be explained (closed form formula, call to a library function, bootstrap, etc.)
- The assumptions made should be given (e.g., Normally distributed errors).
- It should be clear whether the error bar is the standard deviation or the standard error
 of the mean.
- It is OK to report 1-sigma error bars, but one should state it. The authors should preferably report a 2-sigma error bar than state that they have a 96% CI, if the hypothesis of Normality of errors is not verified.
- For asymmetric distributions, the authors should be careful not to show in tables or figures symmetric error bars that would yield results that are out of range (e.g. negative error rates).
- If error bars are reported in tables or plots, The authors should explain in the text how they were calculated and reference the corresponding figures or tables in the text.

8. Experiments compute resources

Question: For each experiment, does the paper provide sufficient information on the computer resources (type of compute workers, memory, time of execution) needed to reproduce the experiments?

Answer: [Yes]

Justification: Compute resources information is provided in the Appendix section.

Guidelines:

- The answer NA means that the paper does not include experiments.
- The paper should indicate the type of compute workers CPU or GPU, internal cluster, or cloud provider, including relevant memory and storage.
- The paper should provide the amount of compute required for each of the individual experimental runs as well as estimate the total compute.
- The paper should disclose whether the full research project required more compute than the experiments reported in the paper (e.g., preliminary or failed experiments that didn't make it into the paper).

9. Code of ethics

Question: Does the research conducted in the paper conform, in every respect, with the NeurIPS Code of Ethics https://neurips.cc/public/EthicsGuidelines?

Answer: [Yes]

Justification: The research conducted in the paper conform, in every respect, with the NeurIPS Code of Ethics.

Guidelines:

- The answer NA means that the authors have not reviewed the NeurIPS Code of Ethics.
- If the authors answer No, they should explain the special circumstances that require a deviation from the Code of Ethics.
- The authors should make sure to preserve anonymity (e.g., if there is a special consideration due to laws or regulations in their jurisdiction).

10. **Broader impacts**

Question: Does the paper discuss both potential positive societal impacts and negative societal impacts of the work performed?

Answer: [NA]

Justification: There is no societal impact of the work performed.

Guidelines:

• The answer NA means that there is no societal impact of the work performed.

- If the authors answer NA or No, they should explain why their work has no societal impact or why the paper does not address societal impact.
- Examples of negative societal impacts include potential malicious or unintended uses (e.g., disinformation, generating fake profiles, surveillance), fairness considerations (e.g., deployment of technologies that could make decisions that unfairly impact specific groups), privacy considerations, and security considerations.
- The conference expects that many papers will be foundational research and not tied to particular applications, let alone deployments. However, if there is a direct path to any negative applications, the authors should point it out. For example, it is legitimate to point out that an improvement in the quality of generative models could be used to generate deepfakes for disinformation. On the other hand, it is not needed to point out that a generic algorithm for optimizing neural networks could enable people to train models that generate Deepfakes faster.
- The authors should consider possible harms that could arise when the technology is being used as intended and functioning correctly, harms that could arise when the technology is being used as intended but gives incorrect results, and harms following from (intentional or unintentional) misuse of the technology.
- If there are negative societal impacts, the authors could also discuss possible mitigation strategies (e.g., gated release of models, providing defenses in addition to attacks, mechanisms for monitoring misuse, mechanisms to monitor how a system learns from feedback over time, improving the efficiency and accessibility of ML).

11. Safeguards

Question: Does the paper describe safeguards that have been put in place for responsible release of data or models that have a high risk for misuse (e.g., pretrained language models, image generators, or scraped datasets)?

Answer: [NA]

Justification: This paper poses no such risks.

Guidelines:

- The answer NA means that the paper poses no such risks.
- Released models that have a high risk for misuse or dual-use should be released with
 necessary safeguards to allow for controlled use of the model, for example by requiring
 that users adhere to usage guidelines or restrictions to access the model or implementing
 safety filters.
- Datasets that have been scraped from the Internet could pose safety risks. The authors should describe how they avoided releasing unsafe images.
- We recognize that providing effective safeguards is challenging, and many papers do
 not require this, but we encourage authors to take this into account and make a best
 faith effort.

12. Licenses for existing assets

Question: Are the creators or original owners of assets (e.g., code, data, models), used in the paper, properly credited and are the license and terms of use explicitly mentioned and properly respected?

Answer: [Yes]

Justification: For all assets we used in this paper, we cited their original source.

Guidelines:

- The answer NA means that the paper does not use existing assets.
- The authors should cite the original paper that produced the code package or dataset.
- The authors should state which version of the asset is used and, if possible, include a URL.
- The name of the license (e.g., CC-BY 4.0) should be included for each asset.
- For scraped data from a particular source (e.g., website), the copyright and terms of service of that source should be provided.

- If assets are released, the license, copyright information, and terms of use in the
 package should be provided. For popular datasets, paperswithcode.com/datasets
 has curated licenses for some datasets. Their licensing guide can help determine the
 license of a dataset.
- For existing datasets that are re-packaged, both the original license and the license of the derived asset (if it has changed) should be provided.
- If this information is not available online, the authors are encouraged to reach out to the asset's creators.

13. New assets

Question: Are new assets introduced in the paper well documented and is the documentation provided alongside the assets?

Answer: [NA]

Justification: This paper does not release new assets.

Guidelines:

- The answer NA means that the paper does not release new assets.
- Researchers should communicate the details of the dataset/code/model as part of their submissions via structured templates. This includes details about training, license, limitations, etc.
- The paper should discuss whether and how consent was obtained from people whose asset is used.
- At submission time, remember to anonymize your assets (if applicable). You can either create an anonymized URL or include an anonymized zip file.

14. Crowdsourcing and research with human subjects

Question: For crowdsourcing experiments and research with human subjects, does the paper include the full text of instructions given to participants and screenshots, if applicable, as well as details about compensation (if any)?

Answer: [NA]

Justification: This paper does not involve crowdsourcing nor research with human subjects.

Guidelines:

- The answer NA means that the paper does not involve crowdsourcing nor research with human subjects.
- Including this information in the supplemental material is fine, but if the main contribution of the paper involves human subjects, then as much detail as possible should be included in the main paper.
- According to the NeurIPS Code of Ethics, workers involved in data collection, curation, or other labor should be paid at least the minimum wage in the country of the data collector.

15. Institutional review board (IRB) approvals or equivalent for research with human subjects

Question: Does the paper describe potential risks incurred by study participants, whether such risks were disclosed to the subjects, and whether Institutional Review Board (IRB) approvals (or an equivalent approval/review based on the requirements of your country or institution) were obtained?

Answer: [NA]

Justification: This paper does not involve crowdsourcing nor research with human subjects. Guidelines:

- The answer NA means that the paper does not involve crowdsourcing nor research with human subjects.
- Depending on the country in which research is conducted, IRB approval (or equivalent) may be required for any human subjects research. If you obtained IRB approval, you should clearly state this in the paper.

- We recognize that the procedures for this may vary significantly between institutions and locations, and we expect authors to adhere to the NeurIPS Code of Ethics and the guidelines for their institution.
- For initial submissions, do not include any information that would break anonymity (if applicable), such as the institution conducting the review.

16. Declaration of LLM usage

Question: Does the paper describe the usage of LLMs if it is an important, original, or non-standard component of the core methods in this research? Note that if the LLM is used only for writing, editing, or formatting purposes and does not impact the core methodology, scientific rigorousness, or originality of the research, declaration is not required.

Answer: [NA]

Justification: The core method development in this research does not involve LLMs as any important, original, or non-standard components.

Guidelines:

- The answer NA means that the core method development in this research does not involve LLMs as any important, original, or non-standard components.
- Please refer to our LLM policy (https://neurips.cc/Conferences/2025/LLM) for what should or should not be described.

A Proofs of Technical Lemmas

A.1 Proof of Lemma A.1

Lemma A.1. Under Assumption 3.2(A1), $F(\cdot)$ in (1) is ρ_F -weakly convex with $\rho_F = \sqrt{d_1}L_gC_f + \rho_fC_g^2$. If g_i is L_g smooth, then $F_\lambda(\mathbf{w})$ is L_F -smooth with $L_F = C_g^2 \max\{\frac{1}{\lambda}, \frac{\rho_f}{1-\lambda\rho_f}\} + C_fL_g$.

Proof. Consider any \mathbf{w} and \mathbf{w}' in \mathbb{R}^d , any $i \in \{1, \dots, n\}$ and any $\mathbf{v}_i \in \partial f_i(\mathbf{t}_i)$ at $\mathbf{t}_i = g_i(\mathbf{w})$. By Assumption 3.2(A1), $\mathbf{v}_i^{\top} g_i(\mathbf{w})$ is $\|\mathbf{v}_i\|_1 L_q$ -smooth in \mathbf{w} and f_i is ρ_f -weakly convex, we have

$$f_{i}(g_{i}(\mathbf{w}')) \geq f_{i}(g_{i}(\mathbf{w})) + \mathbf{v}_{i}^{\top}(g_{i}(\mathbf{w}') - g_{i}(\mathbf{w})) - \frac{\rho_{f}}{2} \|g_{i}(\mathbf{w}') - g_{i}(\mathbf{w})\|^{2}$$

$$\geq f_{i}(g_{i}(\mathbf{w})) + \mathbf{v}_{i}^{\top}(g_{i}(\mathbf{w}') - g_{i}(\mathbf{w})) - \frac{\rho_{f}C_{g}^{2}}{2} \|\mathbf{w}' - \mathbf{w}\|^{2}$$

$$\geq f_{i}(g_{i}(\mathbf{w})) + \mathbf{v}_{i}^{\top}\nabla g_{i}(\mathbf{w}')(\mathbf{w}' - \mathbf{w}) - \frac{\|\mathbf{v}_{i}\|_{1}L_{g}}{2} \|\mathbf{w}' - \mathbf{w}\|^{2} - \frac{\rho_{f}C_{g}^{2}}{2} \|\mathbf{w}' - \mathbf{w}\|^{2}$$

$$\geq f_{i}(g_{i}(\mathbf{w})) + \mathbf{v}_{i}^{\top}\nabla g_{i}(\mathbf{w}')(\mathbf{w}' - \mathbf{w}) - \frac{\sqrt{d_{1}}C_{f}L_{g} + \rho_{f}C_{g}^{2}}{2} \|\mathbf{w}' - \mathbf{w}\|^{2},$$

where the second and the last inequalities hold because of lipschitzness of f_i and g_i from Assumption 3.2, the third because $\mathbf{v}_i^{\top} g_i(\mathbf{w})$ is $\|\mathbf{v}_i\|_1 L_g$ -smooth in \mathbf{w} .

By the ρ_f -weak convexity of f_i and lemma 13 in [33], we know that $f_{i,\lambda}$ is $L_{f_\lambda} := \max\{\frac{1}{\lambda}, \frac{\rho_f}{1-\lambda\rho_f}\}$ -smooth. Since $\nabla f_{i,\lambda}(\mathbf{w}) = (\mathbf{w} - \operatorname{prox}_{\lambda f_i}(\mathbf{w}))/\lambda \in \partial f_i(\operatorname{prox}_{\lambda f_i}(\mathbf{w}))$, we have $\|\nabla f_{i,\lambda}(\mathbf{w})\| \le C_f$ by Assumption 3.2. Therefore, by Assumption 3.2 again, we can easily show that $\nabla f_{i,\lambda}(g_i(\mathbf{w}))\nabla g_i(\mathbf{w})$ is $(C_g^2 L_{f_\lambda} + C_f L_g)$ -Lipschitz continuous (see Lemma 4.2 in [44]), so is $\nabla F_\lambda(\mathbf{w}) = \frac{1}{n} \sum_{i=1}^n \nabla f_{i,\lambda}(g_i(\mathbf{w}))\nabla g_i(\mathbf{w})$.

A.2 Proof of Proposition A.2

Proposition A.2. When f_i is convex, the smoothed problem (3) is equivalent to the Nesterov's smoothing, i.e.,

$$\min_{\mathbf{w}} \max_{\mathbf{y} \in \Omega^n} \frac{1}{n} \sum_{i=1}^n y_i^{\top} g_i(\mathbf{w}) - f_i^*(y_i) - \frac{\lambda}{2} ||y_i||_2^2$$
 (11)

where $\Omega = dom(f_i^*)$, $f_i^*(y_i) + \frac{\lambda}{2} ||y_i||_2^2 = f_{i,\lambda}^*(y_i)$. In addition, $f_{i,\lambda}(g) \leq f_i(g) \leq f_{i,\lambda}(g) + \frac{\lambda C_f^2}{2}$.

Proof. We prove by deriving (11) from (3).

$$F_{\lambda}(\mathbf{w}) = \frac{1}{n} \sum_{i=1}^{n} f_{i,\lambda}(g_{i}(\mathbf{w}))$$

$$= \min_{\mathbf{y} \in \Omega^{n}} \frac{1}{n} \sum_{i=1}^{n} f_{i}(\mathbf{y}_{i}) + \frac{1}{2\lambda} \|\mathbf{y}_{i} - g_{i}(\mathbf{w})\|^{2}$$

$$= \min_{\mathbf{y} \in \Omega^{n}} \max_{\mathbf{z} \in \mathcal{Z}^{n}} \frac{1}{n} \sum_{i=1}^{n} \mathbf{y}_{i}^{\top} \mathbf{z}_{i} - f_{i}^{*}(\mathbf{z}_{i}) + \frac{1}{2\lambda} \|\mathbf{y}_{i} - g_{i}(\mathbf{w})\|^{2}$$

$$= \max_{\mathbf{z} \in \mathcal{Z}^{n}} \min_{\mathbf{y} \in \Omega^{n}} \frac{1}{n} \sum_{i=1}^{n} \mathbf{y}_{i}^{\top} \mathbf{z}_{i} - f_{i}^{*}(\mathbf{z}_{i}) + \frac{1}{2\lambda} \|\mathbf{y}_{i} - g_{i}(\mathbf{w})\|^{2}$$

where the dual domian $\mathcal{Z} = \{\mathbf{z} : \mathbf{z} \in \partial f_i(\mathbf{t}), t \in \mathrm{dom} f_i\} \subseteq \{\mathbf{z} : \|\mathbf{z}\| \leq C_f\}$ is bounded; the second equality is from definition of $f_{i,\lambda}$, the third equality holds from convexity of f_i and the fourth equality holds from Sion's minimax theorem [45] and the convexity of f_i^* .

Note that the inner-level minimization problem can be solved exactly as $\mathbf{y}_i^* = g_i(\mathbf{w}) - \lambda \mathbf{z}_i$ since it's a quadratic function, we plug \mathbf{y}_i^* into the RHS, then we have:

$$F_{\lambda}(\mathbf{w}) = \max_{\mathbf{z} \in \mathcal{Z}^n} \frac{1}{n} \sum_{i=1}^n \mathbf{z}_i^{\top} g_i(\mathbf{w}) - f_i^*(\mathbf{z}_i) - \frac{\lambda}{2} \|\mathbf{z}_i\|^2$$
(12)

Besides, note that $f^+(\mathbf{x}, \mathbf{u}) = f_i(\mathbf{u}) + \frac{1}{2\lambda} \|\mathbf{u} - \mathbf{x}\|^2$ is jointly convex, $f_{i,\lambda}(\mathbf{x}) = \min_{\mathbf{u}} f_i(\mathbf{u}) + \frac{1}{2\lambda} \|\mathbf{u} - \mathbf{x}\|^2$ is also convex. Then we have

$$\frac{1}{n} \sum_{i=1}^{n} f_{i,\lambda}(g_i(\mathbf{w})) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{z}_i^{\top} g_i(\mathbf{w}) - f_{i,\lambda}^*(\mathbf{z}_i)$$

Compare this with 12 we have

$$f_{i,\lambda}^*(\mathbf{z}_i) = f_i^*(\mathbf{z}_i) + \frac{\lambda}{2} \|\mathbf{z}_i\|^2$$

Now we already have that

$$f_{i,\lambda}(g) = \max_{\mathbf{z}_i \in \mathcal{Z}} \mathbf{z}_i^\top g - f_i^*(\mathbf{z}_i) - \frac{\lambda}{2} \left\| \mathbf{z}_i \right\|^2 \text{ and } f_i(g) = \max_{\mathbf{z}_i \in \mathcal{Z}} \mathbf{z}_i^\top g - f_i^*(\mathbf{z}_i)$$

then $f_{i,\lambda}(g) \leq f_i(g)$ naturally holds. Besides,

$$f_i(g) = \max_{\mathbf{z}_i \in \mathcal{Z}} \mathbf{z}_i^\top g - f_i^*(\mathbf{z}_i) \le \max_{\mathbf{z}_i \in \mathcal{Z}} \mathbf{z}_i^\top g - f_i^*(\mathbf{z}_i) - \frac{\lambda}{2} \|\mathbf{z}_i\|^2 + \max_{\mathbf{z}_i \in \mathcal{Z}} \frac{\lambda}{2} \|\mathbf{z}_i\|^2 \le f_{i,\lambda}(g) + \frac{\lambda}{2} C_f^2$$

A.3 Proof of Theorem 4.2

Proof of Theorem 4.2. Suppose w is an ϵ -stationary solution to $F_{\lambda}(\cdot)$ with $\lambda = \epsilon/C_f$. It holds that

$$\|\nabla F_{\lambda}(\mathbf{w})\| = \left\| \frac{1}{n} \sum_{i=1}^{n} \nabla f_{i,\lambda}(g_i(\mathbf{w})) \nabla g_i(\mathbf{w}) \right\| = \left\| \frac{1}{n\lambda} \sum_{i=1}^{n} (g_i(\mathbf{w}) - \operatorname{prox}_{\lambda f_i}(g_i(\mathbf{w}))) \nabla g_i(\mathbf{w}) \right\| \le \epsilon.$$

By (2) and C_f -Lipschitz continuity of f_i , we have $||g_i(\mathbf{w}) - \operatorname{prox}_{\lambda f_i}(g_i(\mathbf{w}))|| \leq \lambda C_f \leq \epsilon$. Let $\mathbf{t}_i = \operatorname{prox}_{\lambda f_i}(g_i(\mathbf{w}))$ and $\mathbf{y}_i = (g_i(\mathbf{w}) - \operatorname{prox}_{\lambda f_i}(g_i(\mathbf{w})))/\lambda \in \partial f_i(\operatorname{prox}_{\lambda f_i}(g_i(\mathbf{w}))) = \partial f_i(\mathbf{t}_i)$. It holds that $||\mathbf{t}_i - g_i(\mathbf{w})|| \leq \epsilon$, $i = 1, \ldots, n$, and $||\frac{1}{n} \sum_{i=1}^n \nabla g_i(\mathbf{w}) \mathbf{y}_i|| \leq \epsilon$, so \mathbf{w} is an approximate ϵ -stationary solution to the original objective $F(\cdot)$ by definition.

A.4 Proof of Theorem 4.4

We need the following lemma.

Lemma A.3. Suppose Assumption 3.2 (A1) or (A2) holds. If $\lambda_{\min} \left(\nabla g_i(\mathbf{w}) \nabla g_i(\mathbf{w})^{\top} \right) \geq \delta$, it holds that $\lambda_{\min} \left(\nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \right) \geq \frac{\delta}{2}$ for any \mathbf{w}' satisfying $\|\mathbf{w}' - \mathbf{w}\| \leq \frac{\delta}{4C_qL_q}$.

Proof. Consider any \mathbf{w}' satisfying $\|\mathbf{w}' - \mathbf{w}\| \leq \frac{\delta}{4C_gL_g}$. Note that

$$\lambda_{\min} \left(\nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \right) = \min_{\mathbf{u} \in \mathbb{R}^{d_1}, \|\mathbf{u}\| = 1} \mathbf{u}^{\top} \nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \mathbf{u}$$

Let

$$\mathbf{u}_{\mathbf{w}'} \in \underset{\mathbf{u} \in \mathbb{R}^{d_1}, \|\mathbf{u}\| = 1}{\arg\min} \mathbf{u}^{\top} \nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \mathbf{u}.$$

It holds that

$$\lambda_{\min} \left(\nabla g_i(\mathbf{w}) \nabla g_i(\mathbf{w})^{\top} \right) - \lambda_{\min} \left(\nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \right)$$

$$\leq \mathbf{u}_{\mathbf{w}'}^{\top} \nabla g_i(\mathbf{w}) \nabla g_i(\mathbf{w})^{\top} \mathbf{u}_{\mathbf{w}'} - \mathbf{u}_{\mathbf{w}'}^{\top} \nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \mathbf{u}_{\mathbf{w}'}$$

$$= \mathbf{u}_{\mathbf{w}'}^{\top} \left(\nabla g_i(\mathbf{w}) \nabla g_i(\mathbf{w})^{\top} - \nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \right) \mathbf{u}_{\mathbf{w}'}$$

$$\leq \| \nabla g_i(\mathbf{w}) \nabla g_i(\mathbf{w})^{\top} - \nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^{\top} \|$$

$$\leq 2C_g L_g \| \mathbf{w} - \mathbf{w}' \|.$$

This implies

$$\lambda_{\min} \left(\nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^\top \right) \ge \lambda_{\min} \left(\nabla g_i(\mathbf{w}') \nabla g_i(\mathbf{w}')^\top \right) - 2C_g L_g \|\mathbf{w} - \mathbf{w}'\| \ge \delta - \frac{\delta}{2} = \frac{\delta}{2}.$$

Proof of Theorem 4.4. Suppose \mathbf{w} is an approximate ϵ -stationary solution of (1) with $\epsilon \leq \min\left\{c, \frac{\delta^2}{16nC_g^2L_g}\right\}$. There exist $\mathbf{t}_1, \ldots, \mathbf{t}_m$ and $\mathbf{y}_i \in \partial f_i(\mathbf{t}_i)$ for $i=1,\ldots,n$ such that $\|\mathbf{t}_i - g_i(\mathbf{w})\| \leq \epsilon, \ i=1,\ldots,n$, and $\|\frac{1}{n}\sum_{i=1}^n \nabla g_i(\mathbf{w})\mathbf{y}_i\| \leq \epsilon.$

Consider the following optimization problem

$$\min_{\mathbf{v}} \left\{ q(\mathbf{v}) := \frac{1}{2n} \sum_{i=1}^{n} \|\mathbf{t}_i - g_i(\mathbf{v})\|^2 \right\}. \tag{13}$$

We want to show that the optimal objective value of (13) is zero and there exists an optimal solution \mathbf{w}' such that $\|\mathbf{w}' - \mathbf{w}\| \leq \frac{4C_g n\epsilon}{\delta}$.

By the condition on ϵ , it holds that $\frac{4C_g n \epsilon}{\delta} \leq \frac{\delta}{4C_g L_g}$. Hence, by Lemma A.3, if $\|\mathbf{v} - \mathbf{w}\| \leq \frac{4C_g n \epsilon}{\delta}$, it holds that

$$\|\nabla q(\mathbf{v})\|^2 = \frac{1}{n^2} \left\| \sum_{i=1}^n \nabla g_i(\mathbf{v})^\top \left[g_i(\mathbf{v}) - \mathbf{t}_i \right] \right\|^2 \ge \frac{\delta}{2n^2} \sum_{i=1}^n \|g_i(\mathbf{v}) - \mathbf{t}_i\|^2 = \frac{\delta}{n} q(\mathbf{v}).$$

Moreover, it always holds that

$$\|\nabla q(\mathbf{v})\|^2 \le \frac{C_g^2}{n} \sum_{i=1}^n \|g_i(\mathbf{v}) - \mathbf{t}_i\|^2 = 2C_g^2 q(\mathbf{v}).$$

Let L_q be the smoothness parameter of $q(\mathbf{v})$ on the compact set $\{\mathbf{v} \in \mathbb{R}^d | \|\mathbf{v} - \mathbf{w}\| \leq \frac{4C_gn\epsilon}{\delta}\}$. Let \mathbf{v}_t for $t=0,1,\ldots$ be generated by the gradient descent method using $\mathbf{v}_0=\mathbf{w}$ and a step size of $\eta=\frac{1}{L_q}$, i.e., $\mathbf{v}_{t+1}=\mathbf{v}_t-\eta\nabla q(\mathbf{v}_t)$. Suppose $\|\mathbf{v}_s-\mathbf{w}\|\leq \frac{4C_gn\epsilon}{\delta}$ for $s=0,1,\ldots,t$ (which is true at least when t=0). By the standard convergence analysis of the gradient descent method for a L_q -smooth objective function, we have

$$q(\mathbf{v}_{t+1}) \leq q(\mathbf{v}_t) + \langle \nabla q(\mathbf{v}_t), \mathbf{v}_{t+1} - \mathbf{v}_t \rangle + \frac{L_q}{2} \|\mathbf{v}_{t+1} - \mathbf{v}_t\|^2$$

$$\leq q(\mathbf{v}_t) - \frac{1}{2L_q} \|\nabla q(\mathbf{v}_t)\|^2 \leq q(\mathbf{v}_t) - \frac{\delta}{2L_q n} q(\mathbf{v}_t).$$

Applying this inequality recursively gives

$$\frac{1}{2C_g^2} \|\nabla q(\mathbf{v}_t)\|^2 \le q(\mathbf{v}_t) \le \left(1 - \frac{\delta}{2L_q n}\right)^t q(\mathbf{v}_0).$$

Recall that $\|\mathbf{t}_i - g_i(\mathbf{w})\| \le \epsilon$ such that $q(\mathbf{v}_0) = \frac{1}{2n} \sum_{i=1}^n \|\mathbf{t}_i - g_i(\mathbf{w})\|^2 \le \frac{\epsilon^2}{2}$. By the triangular inequality, we have

$$\|\mathbf{v}_{t+1} - \mathbf{w}\| \le \sum_{s=0}^{t} \eta \|\nabla q(\mathbf{v}_s)\| \le \sum_{s=0}^{t} \eta C_g \left(1 - \frac{\delta}{2L_q n}\right)^{\frac{s}{2}} \epsilon \le \frac{4\eta C_g L_q n}{\delta} \epsilon = \frac{4C_g n\epsilon}{\delta}.$$

By induction, we have $\|\mathbf{v}_t - \mathbf{w}\| \le \frac{4C_g n\epsilon}{\delta}$ for $t = 0, 1, \dots$

Let \mathbf{w}' be any limiting point of $\{\mathbf{v}_t\}_{t\geq 0}$. It holds that $\|\mathbf{w}' - \mathbf{w}\| \leq \frac{4C_gn\epsilon}{\delta}$ and

$$q(\mathbf{w}') \le \lim_{t \to \infty} \left(1 - \frac{\delta}{2L_q}\right)^t q(\mathbf{v}_0) = 0,$$

meaning that \mathbf{w}' is the optimal solution to (13) with the objective value equal to zero, which implies $\mathbf{t}_i = g_i(\mathbf{w}')$ for $i = 1, \dots, n$.

Since w is an approximate ϵ -stationary point of (1) and $\mathbf{y}_i \in \partial f_i(\mathbf{t}_i)$, we have

$$\left\| \frac{1}{n} \sum_{i} \partial f_{i}(g_{i}(\mathbf{w}')) \nabla g_{i}(\mathbf{w}') \right\|$$

$$= \left\| \frac{1}{n} \sum_{i} \partial f_{i}(\mathbf{t}_{i}) \nabla g_{i}(\mathbf{w}') \right\|$$

$$\leq \left\| \frac{1}{n} \sum_{i} \partial f_{i}(\mathbf{t}_{i}) \nabla g_{i}(\mathbf{w}) \right\| + \left\| \frac{1}{n} \sum_{i} \partial f_{i}(\mathbf{t}_{i}) \left(\nabla g_{i}(\mathbf{w}') - \nabla g_{i}(\mathbf{w}) \right) \right\|$$

$$= \left\| \frac{1}{n} \sum_{i} \mathbf{y}_{i} \nabla g_{i}(\mathbf{w}) \right\| + \left\| \frac{1}{n} \sum_{i} \partial f_{i}(\mathbf{t}_{i}) \left(\nabla g_{i}(\mathbf{w}') - \nabla g_{i}(\mathbf{w}) \right) \right\|$$

$$\leq \epsilon + C_{f} L_{g} \left\| \mathbf{w} - \mathbf{w}' \right\| \leq \left(1 + \frac{4C_{g} C_{f} L_{g} n}{\delta} \right) \epsilon,$$

where the last inequality is by Assumptions 3.2. This means \mathbf{w}' is an $O(\epsilon)$ -stationary solution of (1) and thus \mathbf{w} is a nearly $O(\epsilon)$ -stationary solution of (1).

B Convergence Analysis of SONEX

In this section, we present the proofs for Theorems 4.7. Let \mathbb{E}_t be the conditional expectation conditioning on \mathcal{B}_1^s and $\mathcal{B}_{i,2}^s$ for $i=1,\ldots,n$ and $s=0,1,\ldots,t-1$. To facilitate the proofs, the following quantities are defined based on the notations in Algorithm 1 for $t=0,\ldots,T-1$,

$$V_{t+1} := \left\| \mathbf{v}_{t+1} - \nabla F_{\lambda}(\mathbf{w}_t) \right\|^2$$
(14)

$$U_{t+1} := \frac{1}{n} \left\| \mathbf{u}_{t+1} - \mathbf{g}(\mathbf{w}_t) \right\|^2$$
 (15)

$$\mathbf{u}_t \coloneqq [u_{1,t}, \dots, u_{n,t}]^\top \tag{16}$$

$$\mathbf{g}(\mathbf{w}_t) := [g_1(\mathbf{w}_t), \dots, g_n(\mathbf{w}_t)]^{\top}. \tag{17}$$

B.1 Proof of Theorem 4.7

Lemma B.1 (Lemma 2 in [46]). Suppose $\eta \leq \frac{1}{2L_F}$ in Algorithm 1 with L_F defined in lemma A.1. It holds that

$$F_{\lambda}(\mathbf{w}_{t+1}) \le F_{\lambda}(\mathbf{w}_{t}) + \frac{\eta}{2} V_{t+1} - \frac{\eta}{2} \|\nabla F_{\lambda}(\mathbf{w}_{t})\|^{2} - \frac{\eta}{4} \|\mathbf{v}_{t+1}\|^{2},$$
 (18)

where V_{t+1} is defined as in (14).

A recursion for V_{t+1} is provided below.

Lemma B.2 (Lemma 9 in [2]). Let V_{t+1} be defined as in (14). Suppose $\beta \leq \frac{2}{7}$ in Algorithm 1. It holds that

$$\mathbb{E}\left[V_{t+2}\right] \leq (1-\beta)\mathbb{E}\left[V_{t+1}\right] + \frac{2L_F^2\eta^2\mathbb{E}\left[\|\mathbf{v}_{t+1}\|^2\right]}{\beta} + \frac{3L_f^2C_1^2}{n}\mathbb{E}\left[\|\mathbf{u}_{t+2} - \mathbf{u}_{t+1}\|^2\right] + \frac{2\beta^2C_f^2(C_g^2 + \sigma_1^2)}{\min\{B_1, B_2\}} + 5\beta L_f^2C_1^2\mathbb{E}\left[U_{t+2}\right]$$
(19)

where $C_1^2 := C_q^2 + \frac{\sigma_1^2}{B_2}$, and U_{t+1} , \mathbf{u}_t and $\mathbf{g}(\mathbf{w}_t)$ are defined as in (15), (16) and (17), respectively.

The following lemma provides the recursion for the estimation error U_{t+1} of the MSVR estimator of $g(\mathbf{w})$.

Lemma B.3 (Lemma 1 in [16]). Suppose $\gamma' = \frac{n-B_1}{B_1(1-\gamma)} + (1-\gamma)$ and $\gamma \leq \frac{1}{2}$ and the MSVR update is used in Algorithm 1. It holds that

$$\mathbb{E}\left[U_{t+2}\right] \le \left(1 - \frac{\gamma B_1}{n}\right) \mathbb{E}\left[U_{t+1}\right] + \frac{8nC_g^2}{B_1} \mathbb{E}\left\|\mathbf{w}_{t+1} - \mathbf{w}_t\right\|^2 + \frac{2B_1\gamma^2\sigma_1^2}{nB_2}.$$
 (20)

The following lemma handles the term $\|\mathbf{u}_{t+1} - \mathbf{u}_t\|^2$.

Lemma B.4 (Lemma 6 in [16]). Under the same assumptions as Lemma B.3, it holds that

$$\mathbb{E}[\|\mathbf{u}_{t+2} - \mathbf{u}_{t+1}\|^2] \le \frac{4B_1\gamma^2}{n} \mathbb{E}[U_{t+1}] + \frac{9n^2C_g^2}{B_1} \mathbb{E}\|\mathbf{w}_{t+1} - \mathbf{w}_t\|^2 + \frac{2B_1\gamma^2\sigma_0^2}{B_2}.$$
 (21)

Now we are ready to proof Theorem 4.7

Proof of Theorem 4.7. Note that $\|\mathbf{w}_{t+1} - \mathbf{w}_t\|^2 = \eta^2 \|\mathbf{v}_{t+1}\|^2$. Let P be a non-negative constant to be determined later. Taking the weighted summation of both sides (18), (19), (20), and (21) as specified in the following formula

$$\frac{1}{\eta} \times (18) + \frac{1}{\beta} \times (19) + P \times (20) + \frac{3L_f^2 C_1^2}{n\beta} \times (21),$$

we have

$$\begin{split} &\frac{1}{\eta}\mathbb{E}F_{\lambda}(\mathbf{w}_{t+1}) + \frac{1}{\beta}\mathbb{E}V_{t+2} + \left(P - 5L_f^2C_1^2\right)\mathbb{E}U_{t+2} \\ \leq &\frac{1}{\eta}\mathbb{E}F_{\lambda}(\mathbf{w}_t) + \left(\frac{1}{\beta} - \frac{1}{2}\right)\mathbb{E}V_{t+1} + \left(P\left(1 - \frac{\gamma B_1}{n}\right) + \frac{12L_f^2C_1^2\gamma^2B_1}{\beta n^2}\right)\mathbb{E}U_{t+1} \\ &- \left(\frac{1}{4} - \frac{2L_f^2\eta^2}{\beta^2} - \frac{8nC_g^2P\eta^2}{B_1} - \frac{27nL_f^2C_1^2C_g^2\eta^2}{B_1\beta}\right)\mathbb{E}\left\|\mathbf{v}_{t+1}\right\|^2 \\ &+ \frac{2\beta C_f^2(C_g^2 + \sigma_1^2)}{\min\{B_1, B_2\}} + \frac{2B_1\gamma^2\sigma_1^2P}{nB_2} + \frac{6L_f^2C_1^2\gamma^2\sigma_0^2B_1}{\beta nB_2} - \frac{1}{2}\mathbb{E}\left\|\nabla F_{\lambda}(\mathbf{w}_t)\right\|^2 \end{split}$$

We choose η such that

$$\frac{1}{4} - \frac{2L_F^2 \eta^2}{\beta^2} - \frac{8nC_g^2 P \eta^2}{B_1} - \frac{27nL_f^2 C_1^2 C_g^2 \eta^2}{B_1 \beta} \ge 0.$$
 (22)

Moreover, we set P to be the solution of the following linear equation

$$P\left(1 - \frac{\gamma B_1}{n}\right) + \frac{12L_f^2 C_1^2 \gamma^2 B_1}{\beta n^2} = P - 5L_f^2 C_1^2,$$

which means

$$P = \frac{n}{\gamma B_1} \left(5 L_f^2 C_1^2 + \frac{12 L_f^2 C_1^2 \gamma^2 B_1}{\beta n^2} \right).$$

Combining the results above gives

$$\begin{split} &\frac{1}{\eta} \mathbb{E} F_{\lambda}(\mathbf{w}_{t+1}) + \frac{1}{\beta} \mathbb{E} V_{t+2} + \left(P - 5L_f^2 C_1^2\right) \mathbb{E} U_{t+2} \\ \leq &\frac{1}{\eta} \mathbb{E} F_{\lambda}(\mathbf{w}_t) + \frac{1}{\beta} \mathbb{E} V_{t+1} + \left(P - 5L_f^2 C_1^2\right) \mathbb{E} U_{t+1} \\ &+ \frac{2\beta C_f^2 (C_g^2 + \sigma_1^2)}{\min\{B_1, B_2\}} + \frac{2B_1 \gamma^2 \sigma_1^2 P}{n B_2} + \frac{6L_f^2 C_1^2 \gamma^2 \sigma_0^2 B_1}{\beta n B_2} - \frac{1}{2} \mathbb{E} \left\| \nabla F_{\lambda}(\mathbf{w}_t) \right\|^2. \end{split}$$

Summing both sides of the inequality above over $t = 0, 1, \dots, T-1$ and organizing term lead to

$$\frac{1}{T} \sum_{t=0}^{T-1} \|\nabla F_{\lambda}(\mathbf{w}_{t})\|^{2} \leq \frac{2}{T} \left(\frac{1}{\eta} (F_{\lambda}(\mathbf{w}_{0}) - F_{\lambda}(\mathbf{w}_{T})) + \frac{1}{\beta} \mathbb{E} V_{1} + \left(P - 5L_{f}^{2} C_{1}^{2} \right) \mathbb{E} U_{1} \right) \\
+ \frac{4\beta C_{f}^{2} (C_{g}^{2} + \sigma_{1}^{2})}{\min\{B_{1}, B_{2}\}} + \frac{4B_{1} \gamma^{2} \sigma_{1}^{2} P}{nB_{2}} + \frac{12L_{f}^{2} C_{1}^{2} \gamma^{2} \sigma_{0}^{2} B_{1}}{\beta n B_{2}} \tag{23}$$

Note that $L_F=O(\frac{1}{\epsilon}), L_f=O(\frac{1}{\epsilon})$ and $P=O(\frac{n}{B_1\gamma\epsilon^2}+\frac{\gamma}{n\beta\epsilon^2})$. We require that all the terms in the right-hand side to be in the order of $O(\epsilon^2)$. To ensure $\frac{4\beta C_f^2(C_g^2+\sigma_1^2)}{\min\{B_1,B_2\}}=O(\epsilon^2)$, we must have $\beta=O(\min\{B_1,B_2\}\epsilon^2)$. To ensure $\frac{4B_1\gamma^2\sigma_1^2P}{nB_2}=O(\frac{\gamma}{B_2\epsilon^2}+\frac{B_1\gamma^3}{n^2B_2\beta\epsilon^2})=O(\epsilon^2)$, we must have $\gamma=O(B_2\epsilon^4)$. With these orders of β and γ , we also have $P=O(\frac{n}{B_1B_2\epsilon^6})$ and $\frac{12L_f^2C_1^2\gamma^2\sigma_0^2B_1}{\beta nB_2}=O(\epsilon^4)$. Therefore the last three terms in (23) are $O(\epsilon^2)$.

To satisfy (22), we will need to set

$$\eta = O\left(\min\left\{\frac{\beta}{L_F}, \sqrt{\frac{B_1}{nP}}, \frac{1}{L_f}\sqrt{\frac{B_1\beta}{n}}\right\}\right)
= O\left(\min\left\{\min\{B_1, B_2\}\epsilon^3, \frac{B_1\sqrt{B_2}}{n}\epsilon^3, \sqrt{\frac{B_1\min\{B_1, B_2\}}{n}}\epsilon^2\right\}\right)
= O\left(\frac{B_1\sqrt{B_2}}{n}\epsilon^3\right).$$

We also set the sizes of $\mathcal{B}_{i,2}$ to be $O(\frac{1}{\epsilon^3})$ so that $\mathbb{E}U_1 = O(\epsilon^3)$. Also, it is easy to see that $F_{\lambda}(\mathbf{w}_0) - F_{\lambda}(\mathbf{w}_T) = O(1)$ and $\mathbb{E}V_1 = O(1)$. Then the first term in (23) becomes

$$\frac{1}{T}O\left(\frac{1}{\eta} + \frac{1}{\beta} + P\epsilon^3\right) = \frac{1}{T}O\left(\frac{n}{B_1\sqrt{B_2}\epsilon^3} + \frac{1}{\min\{B_1,B_2\}\epsilon^2} + \frac{n}{B_1B_2\epsilon^3}\right)$$

which will become $O(\epsilon^2)$ when $T = O(\frac{n}{B_1 \sqrt{B_2} \epsilon^5})$.

C Convergence Analysis of ALEXR2

To establish the complexity of Algorithm 2, we first present the convergence result of ALEXR [5] which is the inner loop of ALEXR2, and it shows the complexity of Algorithm 2 to produce $\hat{\mathbf{z}}_t$ such that $\|\hat{\mathbf{z}}_t - \text{prox}_{\nu F_{\lambda}}(\mathbf{w}_t)\|_2^2 \leq O(\epsilon')$.

Theorem C.1 (A Variant of Theorem 1 in [5] when g_i is non-smooth ρ_g -weakly convex and $\mu = \frac{1}{\nu} - \sqrt{d_1}C_f\rho_g$). Suppose Assumptions 3.2 (A3) and 4.5 hold. For any $\epsilon' > 0$, there exists $\theta \in (0,1)$ with $1-\theta = O(\epsilon\epsilon')$ such that, by setting $\eta = \frac{1-\theta}{(4\theta-1)\sqrt{d_1}C_f\rho_g}$ and $\gamma = (\frac{B_1}{n(1-\theta)}-1)^{-1}$, the inner loop of Algorithm 2 (i.e., ALEXR) guarantees $\frac{\mu}{2}\mathbb{E}\left\|\hat{\mathbf{z}}_t - \operatorname{prox}_{\nu F_\lambda}(\mathbf{w}_t)\right\|_2^2 \leq \epsilon'$ for any t after $K_t = \tilde{O}\left(\frac{n}{B_1} + C_g\sqrt{\frac{nL_f}{B_1\sqrt{d_1}\rho_gC_f}} + \frac{nL_f\sigma_0^2}{B_2B_1\epsilon'} + \frac{C_f\sigma_1^2}{B_2\sqrt{d_1}\rho_g\epsilon'} + \frac{C_fC_g^2}{B_1\sqrt{d_1}\rho_g\epsilon'} + \frac{C_fC_g^2}{\sqrt{d_1}\rho_g\epsilon'}\right) = \tilde{O}(\frac{n\sigma_0^2}{B_2B_1\epsilon\epsilon'})$ iterations.

In this section, we provide the convergence analysis for ALEXR2 in Algorithm 2.

Lemma C.2. $F_{\lambda,\nu}(\mathbf{w})$ is $L_{F_{\lambda,\nu}}$ -smooth with $L_{F_{\lambda,\nu}} = \max\{\frac{1}{\nu}, \frac{\rho_{F_{\lambda}}}{1-\nu\rho_{F_{\lambda}}}\}$ and $\rho_{F_{\lambda}} = \sqrt{d_1}C_f\rho_g$.

Lemma C.3. The inner min-max problem in 6 is a $(\frac{1}{\nu} - \rho_{F_{\lambda}})$ -strongly-convex and λ -strongly-concave problem when $\nu \in (0, \rho_{F_{\lambda}}^{-1})$.

Proof. As discussed in Section 4.2, the term $\mathbf{y}_i^{\top} g_i(\mathbf{w})$ is $\rho_{F_{\lambda}}$ -weakly convex with $\rho_{F_{\lambda}} = \sqrt{d_1} C_f \rho_g$. Therefore, when $\nu \in (0, \rho_{F_{\lambda}}^{-1})$, the objective in (6) becomes strongly convex in \mathbf{z} for any fixed $\mathbf{y} \in \mathbb{R}^{nd_1}$.

To establish strong concavity in y, we note that the term \bar{f}_i^* is the Fenchel conjugate of a $\frac{1}{\lambda}$ -smooth function \bar{f}_i . It is a standard result in convex analysis (see, e.g., Theorem 18.15 in [47]) that the conjugate of an L-smooth function is $\frac{1}{L}$ -strongly concave. Thus, \bar{f}_i^* is λ -strongly convex.

The remaining terms in the objective are either independent of \mathbf{y} or linear in \mathbf{y} , and hence do not affect strong concavity. Therefore, the objective is strongly convex in \mathbf{z} and strongly concave in \mathbf{y} when $\nu \in (0, \rho_F^{-1})$.

C.1 Proof of Technical Lemma

Note that our proof for theorem C.1 is almost the same as that of theorem 1 in [5], we only show the difference in this and the following subsections.

Lemma C.4 (Generalization of Lemma 7 in ALEXR [5] to weakly-convex g_i). Suppose that Assumptions 3.2(A2), 4.5 hold. Then, the following holds for Algorithm 2.

$$\frac{1}{n} \mathbb{E} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{*}), \, \bar{y}_{k+1}^{(i)} \right\rangle - \mathbb{E} \left\langle G_{k}, \, \mathbf{w}_{k+1} - \mathbf{w}_{*} \right\rangle
\leq \frac{\frac{C_{f}^{2} \sigma_{1}^{2}}{B_{2}} + \frac{C_{f}^{2} C_{g}^{2}}{B_{1}}}{\frac{1}{\eta} + \frac{1}{\nu}} + \frac{L_{g} C_{f}}{2} \left\| \mathbf{w}_{k+1} - \mathbf{w}_{k} \right\|_{2}^{2} + \frac{\sqrt{d_{1} C_{f} L_{g}}}{2} \left\| \mathbf{w}_{k} - \mathbf{w}^{*} \right\|_{2}^{2}$$
(24)

Proof. For this lemma we only highlight the difference to the proof of lemma 7 in ALEXR [5]. We define $\Delta_k := \frac{1}{B_1} \sum_{i \in \mathcal{B}_{*}^k} [\nabla g_i(\mathbf{w}_k; \tilde{\mathcal{B}}_{i,2}^k)]^{\top} y_{k+1}^{(i)} - \frac{1}{n} \sum_{i=1}^n [\nabla g_i(\mathbf{w}_k)]^{\top} \bar{y}_{k+1}^{(i)}$.

$$\frac{1}{n} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{*}), \overline{y}_{k+1}^{(i)} \right\rangle - \left\langle G_{k}, \mathbf{w}_{k+1} - \mathbf{w}_{*} \right\rangle \\
= \frac{1}{n} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{k}), \overline{y}_{k+1}^{(i)} \right\rangle + \frac{1}{n} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k}) - g_{i}(\mathbf{w}_{*}), \overline{y}_{k+1}^{(i)} \right\rangle \\
+ \left\langle \frac{1}{n} \sum_{i=1}^{n} \left[\nabla g_{i}(\mathbf{w}_{k}) \right]^{\top} \overline{y}_{k+1}^{(i)} + \Delta_{k}, \mathbf{w}_{*} - \mathbf{w}_{k+1} \right\rangle \\
+ \left\langle \frac{1}{n} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{k}), \overline{y}_{k+1}^{(i)} \right\rangle + \left\langle \frac{1}{n} \sum_{i=1}^{n} \left[\nabla g_{i}(\mathbf{w}_{k}) \right]^{\top} \overline{y}_{k+1}^{(i)}, \mathbf{w}_{k} - \mathbf{w}_{*} \right\rangle \\
+ \left\langle \frac{L_{g}}{2} \left\| \mathbf{w}_{k} - \mathbf{w}^{*} \right\|^{2} * \frac{1}{n} \sum_{i=1}^{n} \left\langle \overline{y}_{k+1}^{(i)}, \mathbf{1} \right\rangle + \left\langle \frac{1}{n} \sum_{i=1}^{n} \left[\nabla g_{i}(\mathbf{w}_{k}) \right]^{\top} \overline{y}_{k+1}^{(i)} + \Delta_{k}, \mathbf{w}_{*} - \mathbf{w}_{k+1} \right\rangle \\
= \frac{1}{n} \sum_{i=1}^{n} \left\langle \overline{y}_{k+1}^{(i)}, g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{k}) \right\rangle + \left\langle \frac{1}{n} \sum_{i=1}^{n} \left[\nabla g_{i}(\mathbf{w}_{k}) \right]^{\top} \overline{y}_{k+1}^{(i)}, \mathbf{w}_{k} - \mathbf{w}_{k+1} \right\rangle \\
+ \left\langle \Delta_{k}, \mathbf{w}_{*} - \mathbf{w}_{k+1} \right\rangle + \frac{\sqrt{d_{1}C_{f}L_{g}}}{2} \left\| \mathbf{w}_{k} - \mathbf{w}^{*} \right\|^{2} \tag{25}$$

Note that we have an extra term in blue in 25 comparing with lemma 7 in [5] due to the weakly convexity of g_i . And the remaining part is just the same so we omit it.

Lemma C.5 (Generalization of Lemma 8 in ALEXR to weakly-convex g_i). Suppose that assumptions 3.2(A3), 4.5 hold. Then, the following holds for Algorithm 2.

$$\frac{1}{n} \mathbb{E} \sum_{i=1}^{n} \left\langle g_{i}(\mathbf{w}_{k+1}) - g_{i}(\mathbf{w}_{*}), \, \bar{y}_{k+1}^{(i)} \right\rangle - \mathbb{E} \left\langle G_{k}, \, \mathbf{w}_{k+1} - \mathbf{w}_{*} \right\rangle \\
\leq \frac{C_{f}^{2} \sigma_{1}^{2}}{\frac{1}{B_{2}} + \frac{C_{f}^{2} C_{g}^{2}}{B_{1}} + 4C_{f}^{2} C_{g}^{2}}{\frac{1}{n} + \frac{1}{\nu}} + \frac{1}{\mu} \|\mathbf{w}_{k+1} - \mathbf{w}_{k}\|_{2}^{2} + \frac{\sqrt{d_{1}} C_{f} \rho_{g}}{2} \|\mathbf{w}_{k} - \mathbf{w}^{*}\|^{2} \tag{26}$$

Proof. Similar to lemma C.4, we have an extra term $\frac{\sqrt{d_1}C_f\rho_g}{2}\|\mathbf{w}_k-\mathbf{w}^*\|^2$ in the upper bound for weakly convex g_i .

27

⁶this inequality also holds with assumption 3.2(A3) so lemma C.5 can use the same technique to handle weakly convexity of g_i

C.2 Proof of Inner Loop

Proof of theorem C.1. Similar to theorem 1 in [5] but use lemma C.5 instead of lemma 8 in ALEXR, we have

$$\mathbb{E}[L(\mathbf{w}_{k+1}, y_{*}) - L(\mathbf{w}_{*}, \bar{y}_{k+1})] \\
\leq \frac{\frac{1}{\gamma} + (1 - \frac{B_{1}}{n})}{B_{1}} \mathbb{E}[U_{\bar{f}_{i}^{*}}(y_{*}, y_{k})] - \frac{\frac{1}{\gamma} + 1}{B_{1}} \mathbb{E}[U_{\bar{f}_{i}^{*}}(y_{*}, y_{k+1})] \\
+ \left(\frac{1}{2\eta} + \frac{\sqrt{d_{1}}C_{f}\rho_{g}}{2}\right) \mathbb{E} \|\mathbf{w}_{*} - \mathbf{w}_{k}\|_{2}^{2} - \left(\frac{1}{2\eta} + \frac{1}{2\nu}\right) \mathbb{E} \|\mathbf{w}_{*} - \mathbf{w}_{k+1}\|_{2}^{2} \\
- \left(\frac{1}{\gamma n} - \frac{\lambda_{2} + \lambda_{3}\theta}{\lambda n}\right) \mathbb{E} \left[U_{\bar{f}_{i}^{*}}(\bar{y}_{k+1}, y_{k})\right] - \left(\frac{1}{2\eta} - \frac{C_{g}^{2}}{2\lambda_{2}} - \frac{\frac{1}{\eta} + \frac{1}{\nu}}{4}\right) \mathbb{E} \|\mathbf{w}_{k+1} - \mathbf{w}_{k}\|_{2}^{2} + \frac{\theta C_{g}^{2}}{2\lambda_{3}} \mathbb{E} \|\mathbf{w}_{k} - \mathbf{w}_{k-1}\|_{2}^{2} \\
+ \mathbb{E}[\Gamma_{k+1} - \theta\Gamma_{k}] + \frac{2(1 + 2\theta)\sigma_{0}^{2}}{B\lambda(1 + \frac{1}{\gamma})} + \frac{\frac{C_{f}^{2}\sigma_{1}^{2}}{B_{2}} + \frac{C_{f}^{2}C_{g}^{2}}{B_{1}} + 4C_{f}^{2}C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}}. \tag{27}$$

Define $\Upsilon_k^{\mathbf{w}} \coloneqq \frac{1}{2}\mathbb{E} \|\mathbf{w}_* - \mathbf{w}_k\|_2^2$ and $\Upsilon_k^y = \frac{1}{S}\mathbb{E} U_{\bar{f}_i^*}(y_*, y_k)$. Note that $L(\mathbf{w}_{k+1}, y_*) - L(\mathbf{w}_*, \bar{y}_{k+1}) \ge 0$. Multiply both sides of (27) by θ^{-k} and do telescoping sum from k = 0 to $K_t - 1$. Add $\eta \theta^{-K_t} \Upsilon_{K_t}^{\mathbf{w}}$ to both sides.

$$\frac{\theta^{-K_{t}}\Upsilon_{K_{t}}^{\mathbf{w}}}{\eta} \leq \sum_{k=0}^{K_{t}-1} \theta^{-k} \left(\left(\left(\frac{1}{\eta} + \sqrt{d_{1}}C_{f}\rho_{g} \right) \Upsilon_{k}^{\mathbf{w}} + \left(\frac{1}{\gamma} + \left(1 - \frac{B_{1}}{n} \right) \right) \Upsilon_{k}^{y} - \theta \mathbb{E}\Gamma_{k} \right) \right) \\
- \left(\left(\frac{1}{\eta} + \frac{1}{\nu} \right) \Upsilon_{k+1}^{\mathbf{w}} + \left(\frac{1}{\gamma} + 1 \right) \Upsilon_{k+1}^{y} - \mathbb{E}\Gamma_{k+1} \right) \right) \\
+ \frac{\theta^{-K_{t}}\Upsilon_{K_{t}}^{\mathbf{w}}}{\eta} + \left(\frac{2(1 + 2\theta)\sigma_{0}^{2}}{\lambda B(1 + \frac{1}{\gamma})} + \frac{\frac{C_{f}^{2}\sigma_{1}^{2}}{B_{2}} + \frac{C_{f}^{2}C_{g}^{2}}{B_{1}} + 4C_{f}^{2}C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}} \right) \sum_{k=0}^{K_{t}-1} \theta^{-k} \\
- \sum_{k=0}^{K_{t}-1} \theta^{-k} \left(\frac{1}{\gamma n} - \frac{(\lambda_{2} + \lambda_{3}\theta)}{\lambda n} \right) \mathbb{E}[U_{\bar{f}_{i}^{*}}(\bar{y}_{k+1}, y_{k})] \\
- \sum_{k=0}^{K_{t}-1} \theta^{-k} \left(\frac{1}{2\eta} - \frac{\frac{1}{\eta} + \frac{1}{\nu}}{4} - \frac{C_{g}^{2}}{2\lambda_{2}} - \frac{C_{g}^{2}}{2\lambda_{3}} \right) \mathbb{E} \|\mathbf{w}_{k+1} - \mathbf{w}_{k}\|_{2}^{2} - \theta^{-K_{t}+1} \frac{C_{g}^{2}}{2\lambda_{3}} \mathbb{E} \|\mathbf{w}_{K_{t}} - \mathbf{w}_{K_{t}-1}\|_{2}^{2}$$
(28)

Let $\eta \geq \frac{1-\theta}{\theta/\nu - \sqrt{d_1}C_f\rho_g}$, $\theta \geq \frac{1}{2}$ and $1/\nu > 4\sqrt{d_1}C_f\rho_g$ such that $\theta \geq \frac{\frac{1}{\eta} + \sqrt{d_1}C_f\rho_g}{\frac{1}{\eta} + \frac{1}{\nu}}$ and $\frac{1}{\gamma} + 1 \leq \frac{B_1}{n(1-\theta)}$ such that $\theta \geq \frac{\frac{1}{\eta} + 1 - \frac{B_1}{\eta}}{\frac{1}{\gamma} + 1}$. Then,

$$\begin{split} &\sum_{k=0}^{K_t-1} \theta^{-k} \left(\left((\frac{1}{\eta} + \sqrt{d_1} C_f \rho_g) \Upsilon_k^{\mathbf{w}} + \left(\frac{1}{\gamma} + \left(1 - \frac{B_1}{n} \right) \right) \Upsilon_k^y - \theta \mathbb{E} \Gamma_k \right) - ((\frac{1}{\eta} + \frac{1}{\nu}) \Upsilon_{k+1}^{\mathbf{w}} + (\frac{1}{\gamma} + 1) \Upsilon_{k+1}^y - \mathbb{E} \Gamma_{k+1}) \right) \\ &\leq (\frac{1}{\eta} + \sqrt{d_1} C_f \rho_g) \Upsilon_0^{\mathbf{w}} + \left(\frac{1}{\gamma} + \left(1 - \frac{B_1}{n} \right) \right) \Upsilon_0^y - \theta \mathbb{E} \Gamma_0 - \theta^{-K_t+1} \left((\frac{1}{\eta} + \frac{1}{\nu}) \Upsilon_{K_t}^{\mathbf{w}} + (\frac{1}{\gamma} + 1) \Upsilon_{K_t}^y - \mathbb{E} \Gamma_{K_t} \right). \end{split}$$

By setting $\mathbf{w}_{-1} = \mathbf{w}_0$, we have $\Gamma_0 = 0$. Besides, we have $\Gamma_{K_t} \leq \frac{1}{n} \sum_{i=1}^n \|g_i(\mathbf{w}_{K_t}) - g_i(\mathbf{w}_{K_{t-1}})\| \left\|y_*^{(i)} - y_k^{(i)}\right\| \leq \frac{C_g}{n} \|\mathbf{w}_{K_t} - \mathbf{w}_{K_{t-1}}\| \|y_* - y_{K_t}\| \leq \frac{C_g^2}{2\lambda_3} \|\mathbf{w}_{K_t} - \mathbf{w}_{K_{t-1}}\|^2 + \frac{\lambda_3}{2\lambda n^2} U_{\bar{f}_i^*}(y_*, y_{K_t})$. Note that the first term in the RHS here can-

celled out with the last term of RHS in 28. Thus,

$$\frac{\theta^{-K_{t}}\Upsilon_{K_{t}}^{\mathbf{w}}}{\eta} \leq \left(\frac{1}{\eta} + \sqrt{d_{1}}C_{f}\rho_{g}\right)\Upsilon_{0}^{\mathbf{w}} + \left(\frac{1}{\gamma} + \left(1 - \frac{B_{1}}{n}\right)\right)\Upsilon_{0}^{y} \\
- \theta^{-K_{t}+1}\left(\left(\frac{1}{\eta} + \frac{1}{\nu}\right)\Upsilon_{K_{t}}^{\mathbf{w}} + \left(\frac{1}{\gamma} + 1 - \frac{\lambda_{3}S}{2\lambda n^{2}}\right)\Upsilon_{K_{t}}^{y} - \frac{1}{\eta\theta}\Upsilon_{K_{t}}^{\mathbf{w}}\right) \\
+ \left(\frac{2(1 + 2\theta)\sigma_{0}^{2}}{\lambda B(1 + \frac{1}{\gamma})} + \frac{\frac{C_{f}^{2}\sigma_{1}^{2}}{B_{2}} + \frac{C_{f}^{2}C_{g}^{2}}{B_{1}} + 4C_{f}^{2}C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}}\right) \sum_{k=0}^{K_{t}-1} \theta^{-k} \left(\frac{1}{\gamma n} - \frac{(\lambda_{2} + \lambda_{3}\theta)}{\lambda n}\right) \mathbb{E}[U_{\bar{f}_{i}^{*}}(\bar{y}_{k+1}, y_{k})] \\
- \sum_{k=0}^{K_{t}-1} \theta^{-k} \left(\frac{1}{2\eta} - \frac{\frac{1}{\eta} + \frac{1}{\nu}}{4} - \frac{C_{g}^{2}}{2\lambda_{2}} - \frac{C_{g}^{2}}{2\lambda_{3}}\right) \mathbb{E}\|\mathbf{w}_{k+1} - \mathbf{w}_{k}\|_{2}^{2}. \tag{29}$$

To make the \heartsuit terms in (29) be non-negative, we choose $\lambda_2 \simeq \lambda_3 \simeq \frac{C_g \sqrt{B_1 \lambda}}{\sqrt{n \sqrt{d_1} C_f \rho_g}}$ while ensuring that

$$\gamma \leq O\left(\frac{\lambda n^2}{\lambda_3 B_1} \wedge \frac{\lambda}{\lambda_2 + \lambda_3 \theta}\right) = O\left(\frac{\sqrt{n\sqrt{d_1}C_f \rho_g \lambda}}{C_g \sqrt{B_1}}\right), \quad \eta \leq O\left(\frac{\sqrt{B_1 \lambda}}{C_g \sqrt{n\sqrt{d_1}C_f \rho_g}} \wedge \frac{1}{\rho_g C_f}\right). \tag{30}$$

By selecting $\eta=\frac{1-\theta}{\theta/\nu-\sqrt{d_1}C_f\rho_g}$, $\frac{1}{\gamma}=\frac{B_1}{n(1-\theta)}-1$ and $\frac{1}{\nu}=4\sqrt{d_1}C_f\rho_g$ we have that $\frac{1}{\gamma}+\left(1-\frac{B_1}{n}\right)=\frac{B_1\theta}{n(1-\theta)}$.

$$\begin{split} \mu \Upsilon_{K_t}^{\mathbf{w}} &\leq \mu (1 + \eta \sqrt{d_1} C_f \rho_g) \theta^{K_t} \Upsilon_0^{\mathbf{w}} + \frac{\left(\frac{1}{\gamma} + \left(1 - \frac{B_1}{n}\right)\right) \mu (1 - \theta)}{\theta / \nu - \sqrt{d_1} C_f \rho_g} \theta^{K_t} \Upsilon_0^y \\ &+ \left(\frac{2(1 + 2\theta)\sigma_0^2}{\lambda B (1 + \frac{1}{\gamma})} + \frac{\frac{C_f^2 \sigma_1^2}{B_2} + \frac{C_f^2 C_g^2}{B_1} + 4C_f^2 C_g^2}{\frac{1}{\eta} + \frac{1}{\nu}}\right) \frac{(\theta^{-K_t} - 1)\theta^{K_t}}{\theta^{-1} - 1} \frac{(1 - \theta)\mu}{\theta / \nu - \sqrt{d_1} C_f \rho_g} \\ &\leq \mu (1 + \eta \sqrt{d_1} C_f \rho_g) \theta^{K_t} \Upsilon_0^{\mathbf{w}} + \frac{\frac{B_1}{n} \theta \mu}{\frac{1}{2} \theta / \nu} \theta^{K_t} \Upsilon_0^y + 2\mu \nu \left(\frac{2(1 + 2\theta)\sigma_0^2}{\lambda B_2 (1 + \frac{1}{\gamma})} + \frac{\frac{C_f^2 \sigma_1^2}{B_2} + \frac{C_f^2 C_g^2}{B_1} + 4C_f^2 C_g^2}{\frac{1}{\eta} + \frac{1}{\nu}}\right) \\ &\leq \mu (1 + \eta \sqrt{d_1} C_f \rho_g) \theta^{K_t} \Upsilon_0^{\mathbf{w}} + 2\theta^{K_t} \Upsilon_0^y + 2\left(\frac{2(1 + 2\theta)\sigma_0^2}{\lambda B_2 (\frac{1}{\gamma} + 1)} + \eta \left(\frac{C_f^2 \sigma_1^2}{B_2} + \frac{C_f^2 C_g^2}{B_1} + 4C_f^2 C_g^2\right)\right). \end{split}$$

We select

$$1 - \theta = O\left(\frac{1}{2} \wedge \frac{B_1}{n} \wedge \frac{1}{C_g} \sqrt{\frac{B_1 \lambda \sqrt{d_1} C_f \rho_g}{n}} \wedge \frac{\lambda B_2 B_1 \epsilon'}{\sigma_0^2 n} \wedge \frac{B_2 \sqrt{d_1} \rho_g \epsilon'}{C_f \sigma_1^2} \wedge \frac{B_1 \sqrt{d_1} \rho_g \epsilon'}{C_f C_g^2} \wedge \frac{\sqrt{d_1} \rho_g \epsilon'}{C_f C_g^2}\right)$$

to make (30) hold and

$$\frac{2(1+2\theta)\sigma_0^2}{\lambda B_2(\frac{1}{\gamma}+1)} + \eta \left(\frac{C_f^2 \sigma_1^2}{B_2} + \frac{C_f^2 C_g^2}{B_1} + 4C_f^2 C_g^2\right) \le \frac{2(1+2\theta)(1-\theta)\sigma_0^2 n}{\lambda B_2 B_1} + \frac{(1-\theta)\left(\frac{C_f^2 \sigma_1^2}{B_2} + \frac{C_f^2 C_g^2}{B_1} + 4C_f^2 C_g^2\right)}{\sqrt{d_1} C_f \rho_g} \le \epsilon'.$$

Besides, we show that $\Upsilon_0^{\mathbf{w}}$ can be bounded by constant.(boundedness of Υ_0^y is already guaranteed by assumption 4.9). Note that $\mathbf{w}_t^* = \operatorname{prox}_{\nu F_\lambda}(\mathbf{w}_{t,0})$ we have:

Since $L_f \coloneqq \frac{1}{\lambda} = O(\frac{1}{\epsilon})$, the number of inner loop iterations needed by Algorithm 2 to make $\mu \Upsilon_{K_t}^{\mathbf{w}} \le \epsilon'$ is

$$K_t = \tilde{O}\left(\frac{n}{B_1} + C_g\sqrt{\frac{nL_f}{B_1\sqrt{d_1}\rho_gC_f}} + \frac{nL_f\sigma_0^2}{B_2B_1\epsilon'} + \frac{C_f\sigma_1^2}{B_2\sqrt{d_1}\rho_g\epsilon'} + \frac{C_fC_g^2}{B_1\sqrt{d_1}\rho_g\epsilon'} + \frac{C_fC_g^2}{\sqrt{d_1}\rho_g\epsilon'}\right)$$

where $\tilde{O}(\cdot)$ hides the polylog $(C_f C_g \rho_g / \epsilon')$ factor and the green term is the dominant term so $K_t = \tilde{O}(\frac{n\sigma_0^2}{B_0B_1\epsilon\epsilon'})$.

C.3 Proof of Outer loop

Proof of Theorem 4.10. Let $\mathbf{w}_{-1} = \mathbf{w}_0$. By lemma C.2, $F_{\lambda,\nu}(\mathbf{w}_t)$ is $L_{F_{\lambda,\nu}}$ -smooth. Since $\beta \leq \frac{1}{2}$, we have

$$\|\mathbf{v}_{t+1} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\|^{2}$$

$$= \left\| (1 - \beta)(\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})) + (1 - \beta)(\nabla F_{\lambda,\nu}(\mathbf{w}_{t-1}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})) + \beta \left(\frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\right) \right\|^{2}$$

$$\leq (1 + \frac{\beta}{2})(1 - \beta)^{2} \left((1 + \frac{\beta}{2}) \|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} + (1 + \frac{2}{\beta}) \|\nabla F_{\lambda,\nu}(\mathbf{w}_{t-1}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\|^{2} \right)$$

$$+ (1 + \frac{2}{\beta})\beta^{2} \left\| \frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t}) \right\|^{2}$$

$$\leq (1 - \frac{\beta}{2}) \|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} + \frac{3L_{F_{\lambda,\nu}}^{2}}{\beta} \|\mathbf{w}_{t-1} - \mathbf{w}_{t}\|^{2} + 3\beta \left\| \frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t}) \right\|^{2}$$

$$\leq (1 - \frac{\beta}{2}) \|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} + \frac{3\alpha^{2}L_{F_{\lambda,\nu}}^{2}}{\beta} \|\mathbf{v}_{t}\|^{2} + 3\beta \left\| \frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t}) \right\|^{2}$$

$$\leq (1 - \frac{\beta}{2}) \|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} + \frac{3\alpha^{2}L_{F_{\lambda,\nu}}^{2}}{\beta} \|\mathbf{v}_{t}\|^{2} + 3\beta \left\| \frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t}) \right\|^{2}$$

$$\leq (31)$$

Since $\alpha \leq \frac{1}{2L_{F_{\lambda,\nu}}}$, by Lemma 2 in [46], we can obtain a result similar to (18) in Lemma B.1, that is,

$$F_{\lambda,\nu}(\mathbf{w}_{t+1}) \le F_{\lambda,\nu}(\mathbf{w}_t) + \frac{\alpha}{2} \left\| \mathbf{v}_{t+1} - \nabla F_{\lambda,\nu}(\mathbf{w}_t) \right\|^2 - \frac{\alpha}{2} \left\| \nabla F_{\lambda,\nu}(\mathbf{w}_t) \right\|^2 - \frac{\alpha}{4} \left\| \mathbf{v}_{t+1} \right\|^2$$
(32)

Multiplying both sides of (31) by $\frac{\alpha}{\beta}$ and adding them to both sides of (32), we have

$$\frac{\alpha}{2} \|\nabla F_{\lambda,\nu}(\mathbf{w}_{t})\|^{2}$$

$$\leq F_{\lambda,\nu}(\mathbf{w}_{t}) - F_{\lambda,\nu}(\mathbf{w}_{t+1}) + \left(\frac{\alpha}{\beta} - \frac{\alpha}{2}\right) \left(\|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} - \|\mathbf{v}_{t+1} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\|^{2}\right)$$

$$- \frac{\alpha}{4} \|\mathbf{v}_{t+1}\|^{2} + \frac{\alpha^{3} L_{F_{\lambda,\nu}}^{2}}{\beta^{2}} \|\mathbf{v}_{t}\|^{2} + 3\alpha \left\|\frac{1}{\nu}(\mathbf{w}_{t} - \hat{\mathbf{z}}_{t}) - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\right\|^{2}$$

$$\leq F_{\lambda,\nu}(\mathbf{w}_{t}) - F_{\lambda,\nu}(\mathbf{w}_{t+1}) + \left(\frac{\alpha}{\beta} - \frac{\alpha}{2}\right) \left(\|\mathbf{v}_{t} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t-1})\|^{2} - \|\mathbf{v}_{t+1} - \nabla F_{\lambda,\nu}(\mathbf{w}_{t})\|^{2}\right)$$

$$\frac{\alpha}{4} \left(\|\mathbf{v}_{t}\|^{2} - \|\mathbf{v}_{t+1}\|^{2}\right) + \frac{3\alpha}{\nu^{2}} \|\hat{\mathbf{z}}_{t} - \operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_{t})\|^{2},$$

where the second inequality is because $\nabla F_{\lambda,\nu}(\mathbf{w}_t) = \frac{1}{\nu}(\mathbf{w}_t - \operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_t))$ and $\alpha \leq \frac{\beta}{2L_{F_{\lambda,\nu}}}$ so $\frac{\alpha}{4} \geq \frac{\alpha^3 L_{F_{\lambda,\nu}}^2}{\beta^2}$.

By Theorem C.1, for $\epsilon' = \frac{\mu\nu^2\epsilon^2}{36}$, there exists $\theta \in (0,1)$ with $1-\theta = O(\epsilon^2)$ such that, by setting $\eta = \frac{1-\theta}{\theta\mu}$ and $\gamma = \frac{(1-\theta)n}{B_1}$, inner loop of Algorithm 2(i.e. ALEXR) guarantees $\mathbb{E}\left\|\hat{\mathbf{z}}_t - \mathrm{prox}_{\nu F_\lambda}(\mathbf{w}_t)\right\|_2^2 \leq \frac{\nu^2\epsilon^2}{18}$ for any t after $K_t = \tilde{O}\left(\frac{nL_f}{B_1B_2\epsilon^2} + \frac{L_f}{\epsilon^2}\right)$ iterations.

Recall that $\mathbf{v}_0 = 0$ and $\mathbf{w}_{-1} = \mathbf{w}_0$. Summing the inequality above over $t = 0, \dots T - 1$ and dividing both sides by $\frac{\alpha T}{2}$, we have:

$$\begin{split} & \mathbb{E}[\operatorname{dist}(0,\partial F_{\lambda}(\operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau})))^{2}] = \frac{1}{T}\sum_{t=0}^{T-1}\mathbb{E}\left\|\nabla F_{\lambda,\nu}(\mathbf{w}_{t})\right\|^{2} \\ \leq & \frac{2\mathbb{E}(F_{\lambda,\nu}(\mathbf{w}_{0}) - F_{\lambda,\nu}(\mathbf{w}_{T}))}{\alpha T} + \frac{\left(\frac{2}{\beta} - 1\right)\left\|\nabla F_{\lambda,\nu}(\mathbf{w}_{0})\right\|^{2}}{T} + \frac{6}{\nu^{2}T}\sum_{t=0}^{T-1}\mathbb{E}\left\|\hat{\mathbf{z}}_{t} - \operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_{t})\right\|^{2} \\ \leq & \frac{2\mathbb{E}(F_{\lambda,\nu}(\mathbf{w}_{0}) - F_{\lambda,\nu}(\mathbf{w}_{T}))}{\alpha T} + \frac{\left(\frac{2}{\beta} - 1\right)\left\|\nabla F_{\lambda,\nu}(\mathbf{w}_{0})\right\|^{2}}{T} + \frac{\epsilon^{2}}{3}, \end{split}$$

By setting $T=O(\epsilon^{-2})$, we'll have $\mathbb{E}[\operatorname{dist}(0,\partial F_{\lambda}(\operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau})))^{2}] \leq \epsilon^{2}$, meaning that \mathbf{w}_{τ} is a nearly ϵ -stationary point of (4) in expectation. Recall that $L_{f}=O(\frac{1}{\epsilon})$ so $K_{t}=O(\epsilon^{-3})$. Hence, the total complexity for finding such a solution is $\sum_{t=0}^{T-1}K_{t}=\tilde{O}(\epsilon^{-5})$.

Proof of Corollary 4.11. When g_i is L_g -smooth, $F_{\lambda}(\cdot)$ is L_F -smooth where $L_F = O(\frac{1}{\lambda}) = O(\frac{1}{\epsilon})$ is defined in lemma A.1. By Theorem 4.10,

$$\mathbb{E}\|\nabla F_{\lambda}(\mathrm{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}))\| = \frac{1}{\nu}\mathbb{E}\|\mathrm{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}) - \mathbf{w}_{\tau}\| \leq O(\epsilon)$$

According to Theorem C.1, for $\epsilon' = O(\epsilon^4)$, there exists $\theta \in (0,1)$ with $1-\theta = O(\epsilon^4)$ such that, by setting $\eta = \frac{1-\theta}{\theta} L_f$ and $\gamma = \frac{(1-\theta)n}{B_1}$ in inner loop of Algorithm 2(i.e. ALEXR), $\hat{\mathbf{w}}_{\tau} = \text{ALEXR}(\mathbf{w}_{\tau}, K)$ satisfies $\frac{\mu}{2} \mathbb{E} \left\| \hat{\mathbf{w}}_{\tau} - \text{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}) \right\|^2 \leq \epsilon^4$ and thus

$$\mathbb{E} \left\| \hat{\mathbf{w}}_{\tau} - \operatorname{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}) \right\| \leq O(\epsilon^{2})$$

after $K = \tilde{O}\left(\frac{nL_f}{B_1B_2\epsilon^4} + \frac{L_f}{\epsilon^4}\right)$ iterations. Then we have

$$\mathbb{E}\|\nabla F_{\lambda}(\hat{\mathbf{w}}_{\tau})\| \leq \mathbb{E}\|\nabla F_{\lambda}(\mathrm{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}))\| + L_{F}\mathbb{E}\left\|\mathrm{prox}_{\nu F_{\lambda}}(\mathbf{w}_{\tau}) - \hat{\mathbf{w}}_{\tau}\right\| \leq O(\epsilon),$$

which means $\hat{\mathbf{w}}_{\tau}$ is an ϵ -stationary solution of (3). By Proposition 4.4, $\hat{\mathbf{w}}_{\tau}$ is a nearly ϵ -stationary solution of (1). By Theorem 4.10, \mathbf{w}_{τ} is found within complexity $O(\epsilon^{-5})$ and $\hat{\mathbf{w}}_{\tau} = \text{ALEXR}(\mathbf{w}_{\tau}, K)$ has complexity $K = \tilde{O}\left(\frac{nL_f}{B_1B_2\epsilon^4} + \frac{L_f}{\epsilon^4}\right)$. The total complexity for computing $\hat{\mathbf{w}}_{\tau}$ is still $\tilde{O}\left(\frac{n}{B_1B_2\epsilon^5} + \frac{1}{\epsilon^5}\right)$.

D Convergence Analysis of SONEX and ALEXR2 applying to Constraint Optimization Problems

In this section, we present the complexity analysis of Algorithm 2 when it is applied to (8) to solve the constrained optimization problem 7.

D.1 Proof of Proposition 5.2 and its variant

Proof of Proposition 5.2. By the definition of $f_{\lambda}(\cdot)$, we have

$$\nabla f_{\lambda}(\cdot) = \frac{1}{\lambda} \min\{[\cdot]_{+}, \lambda \rho\}.$$

Suppose \mathbf{w} is a nearly ϵ -stationary point of (9), which means there exists $\hat{\mathbf{w}}$ such that $\|\mathbf{w} - \hat{\mathbf{w}}\| \le \epsilon$ and $\operatorname{dist}(0, \partial \Phi_{\lambda}(\hat{\mathbf{w}})) \le \epsilon$. This means there exist $\mathbf{h}_i(\hat{\mathbf{w}}) \in \partial g_i(\hat{\mathbf{w}})$ and

$$\nu_i = \frac{1}{m} \nabla f_{\lambda}(g_i(\hat{\mathbf{w}})) = \frac{1}{\lambda m} \min\{[g_i(\hat{\mathbf{w}})]_+, \lambda \rho\}$$

for $i = 0, 1, \dots, m$ such that

$$\|\mathbf{h}_0(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| \le \epsilon \tag{33}$$

where $\mathbf{J}(\hat{\mathbf{w}}) = [\mathbf{h}_1(\hat{\mathbf{w}}), \dots, \mathbf{h}_m(\hat{\mathbf{w}})] \in \mathbb{R}^{d \times m}$ and $\boldsymbol{\nu} = (\nu_1, \dots, \nu_m)^{\top} \in \mathbb{R}^m$.

Suppose $\max_{i=1,...,m} g_i(\hat{\mathbf{w}}) > \epsilon$. Then there exists k such that $[g_k(\hat{\mathbf{w}})]_+ > \epsilon$. Recall that $\lambda = \frac{\epsilon}{\rho}$. We have

$$\nu_k = \frac{1}{\lambda m} \min\{[g_k(\hat{\mathbf{w}})]_+, \lambda \rho\} = \frac{1}{\lambda m} \epsilon = \frac{\rho}{m}.$$

By Assumption 5.1, we have

$$\|\mathbf{h}_0(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| \ge \|\mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| - \|\mathbf{h}_0(\hat{\mathbf{w}})\| \ge \sigma_{min}(\mathbf{J}(\hat{\mathbf{w}})) \|\boldsymbol{\nu}\| - \|\mathbf{h}_0(\hat{\mathbf{w}})\| \ge \frac{\delta\rho}{m} - C_g > 1,$$

which contradicts with (33). Therefore, we must have

$$\max_{i=1,\dots,m} g_i(\hat{\mathbf{w}}) \le \epsilon. \tag{34}$$

Finally, when $\max_{i=1,\ldots,m} g_i(\hat{\mathbf{w}}) \leq \epsilon$, we have for $\forall i=1,2,\cdots,m$

$$|g_i(\hat{\mathbf{w}})\nu_i| \le \sum_{i=1}^m |g_i(\hat{\mathbf{w}})\nu_i| = \sum_{i=1}^m [g_i(\hat{\mathbf{w}})]_+ \frac{\min\{[g_i(\hat{\mathbf{w}})]_+, \lambda\rho\}}{\lambda m} \le \frac{\rho}{m} \sum_{i=1}^m [g_i(\hat{\mathbf{w}})]_+ \le O(\epsilon). \quad (35)$$

With (33), (34) and (35), $\hat{\mathbf{w}}$ is an ϵ -KKT point of (7) so $\hat{\mathbf{w}}$ is a nearly ϵ -KKT point of (7).

Proposition 5.2 is given when \mathbf{w} is a deterministic nearly ϵ -stationary solution to (9). However, the solution \mathbf{w} found by our algorithms is only a nearly ϵ -stationary solution to (9) in expectation. This means there exists $\hat{\mathbf{w}}$ such that $\mathbb{E} \|\mathbf{w} - \hat{\mathbf{w}}\| \le \epsilon$ and $\mathbb{E} \mathrm{dist}(0, \partial \Phi_{\lambda}(\hat{\mathbf{w}})) \le \epsilon$. In this case, we cannot prove the four inequalities in the conclusion of Proposition 5.2 hold deterministically. Instead, we can only show that the first two inequality hold in expectation while the last two hold in high probabilities. We present this variant of Proposition 5.2 below with its proof.

Proposition D.1. Suppose Assumptions 3.2(A3) and 5.1 hold. If $\rho > \frac{m(C_g+1)}{\delta}$ and $\lambda = \frac{\epsilon}{\rho}$, a nearly ϵ -stationary solution \mathbf{w} to (9) in expectation is also a nearly $O(\epsilon)$ -KKT solution to the original problem (7) in the sense that there exist $\hat{\mathbf{w}}$ and $\nu_i \geq 0$ for $i=1,\ldots,m$ such that $\mathbb{E}\|\mathbf{w}-\hat{\mathbf{w}}\| \leq \epsilon$ and $\mathbb{E} \text{dist}(0,\partial g_0(\hat{\mathbf{w}})+\sum_{i=1}^m \partial g_i(\hat{\mathbf{w}})\nu_i)) \leq O(\epsilon)$ and it holds with probability $1-O(\epsilon)$ that $\max_{i=1,\ldots,m} g_i(\hat{\mathbf{w}}) \leq O(\epsilon)$ and $|g_i(\hat{\mathbf{w}})\nu_i| \leq O(\epsilon)$, $\forall i=1,2,\cdots,m$.

Proof. By the definition of $f_{\lambda}(\cdot)$, we have

$$\nabla f_{\lambda}(\cdot) = \frac{1}{\lambda} = \min\{[\cdot]_{+}, \lambda \rho\}.$$

Suppose \mathbf{w} is a nearly ϵ -stationary point of (9) in expectation, which means there exists $\hat{\mathbf{w}}$ such that $\mathbb{E} \|\mathbf{w} - \hat{\mathbf{w}}\| \le \epsilon$ and $\mathbb{E} \text{dist}(0, \partial \Phi_{\lambda}(\hat{\mathbf{w}})) \le \epsilon$. This means there exist $\mathbf{h}_i(\hat{\mathbf{w}}) \in \partial g_i(\hat{\mathbf{w}})$ and

$$\nu_i = \frac{1}{m} \nabla f_{\lambda}(g_i(\hat{\mathbf{w}})) = \frac{1}{\lambda m} \min\{[g_i(\hat{\mathbf{w}})]_+, \lambda \rho\}$$

for $i = 0, 1, \ldots, m$ such that

$$\mathbb{E} \|\mathbf{h}_0(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| < \epsilon$$

where $\mathbf{J}(\hat{\mathbf{w}}) = [\mathbf{h}_1(\hat{\mathbf{w}}), \dots, \mathbf{h}_m(\hat{\mathbf{w}})] \in \mathbb{R}^{d \times m}$ and $\boldsymbol{\nu} = (\nu_1, \dots, \nu_m)^{\top} \in \mathbb{R}^m$.

Suppose $\max_{i=1,\dots,m} g_i(\hat{\mathbf{w}}) > \epsilon$. Then there exists k such that $[g_k(\hat{\mathbf{w}})]_+ > \epsilon$. Recall that $\lambda = \frac{\epsilon}{\rho}$. We have

$$\nu_k = \frac{1}{\lambda m} \min\{[g_k(\hat{\mathbf{w}})]_+, \lambda \rho\} = \frac{1}{\lambda m} \epsilon = \frac{\rho}{m}.$$

By Assumption 5.1, we have

$$\|\mathbf{h}_{0}(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| \ge \|\mathbf{J}(\hat{\mathbf{w}})\boldsymbol{\nu}\| - \|\nabla g_{0}(\hat{\mathbf{w}})\| \ge \sigma_{min}(\mathbf{J}(\hat{\mathbf{w}}))\|\boldsymbol{\nu}\| - \|\mathbf{h}_{0}(\hat{\mathbf{w}})\| \ge \frac{\delta\rho}{m} - C_{g} > 1$$

Therefore,

$$\begin{split} \epsilon &\geq \mathbb{E} \left\| \mathbf{h}_{0}(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}}) \boldsymbol{\nu} \right\| \\ &= \mathbb{E} \left[\left\| \mathbf{h}_{0}(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}}) \boldsymbol{\nu} \right\| \Big| \max_{i=1,\dots,m} g_{i}(\hat{\mathbf{w}}) > \epsilon \right) \right] \operatorname{Prob}(\max_{i=1,\dots,m} g_{i}(\hat{\mathbf{w}}) > \epsilon) \\ &+ \mathbb{E} \left[\left\| \mathbf{h}_{0}(\hat{\mathbf{w}}) + \mathbf{J}(\hat{\mathbf{w}}) \boldsymbol{\nu} \right\| \Big| \max_{i=1,\dots,m} g_{i}(\hat{\mathbf{w}}) \leq \epsilon \right) \right] \operatorname{Prob}(\max_{i=1,\dots,m} g_{i}(\hat{\mathbf{w}}) \leq \epsilon) \\ &\geq \operatorname{Prob}(\max_{i=1,\dots,m} g_{i}(\hat{\mathbf{w}}) > \epsilon) \left(\frac{\delta \rho}{m} - C_{g} \right) \end{split}$$

As a result,

$$\operatorname{Prob}(\max_{i=1,\dots,m} g_i(\hat{\mathbf{w}}) > \epsilon) \le \epsilon \left(\frac{\delta \rho}{m} - C_g\right)^{-1} = O(\epsilon). \tag{36}$$

Finally, when $\max_{i=1,\dots,m} g_i(\hat{\mathbf{w}}) \leq \epsilon$, we have for $\forall i=1,2,\cdots,m$

$$|g_i(\hat{\mathbf{w}})\nu_i| \le \sum_{i=1}^m |g_i(\hat{\mathbf{w}})\nu_i| = \sum_{i=1}^m [g_i(\hat{\mathbf{w}})]_+ \frac{\min\{[g_i(\hat{\mathbf{w}})]_+, \lambda\rho\}}{\lambda m} \le \frac{\rho}{m} \sum_{i=1}^m [g_i(\hat{\mathbf{w}})]_+ \le O(\epsilon).$$

It then follows from (36) that for $\forall i = 1, 2, \dots, m$

Prob
$$(|g_i(\hat{\mathbf{w}})\nu_i| \ge O(\epsilon)) \le O(\epsilon)$$
.

D.2 Sketch Proof of Theorem 5.3

Since the proof is almost the same as that of Theorem 4.10, we only highlight the difference in the proof.

We first consider the case where $\{g_i\}_{i=0}^m$ are weakly convex. We will slightly modify Algorithm 2 to solve the following problem

$$\min_{\mathbf{w} \in \mathbb{R}^d} \left\{ \Phi_{\lambda,\nu}(\mathbf{w}) := \min_{\mathbf{z} \in \mathbb{R}^d} \left\{ \Phi_{\lambda}(\mathbf{z}) + \frac{1}{2\nu} \|\mathbf{z} - \mathbf{w}\|^2 \right\} \right\}$$
(37)

$$= \min_{\mathbf{w} \in \mathbb{R}^d} \left\{ \min_{\mathbf{z} \in \mathbb{R}^d} \max_{y_i \in [0, \rho]} \left\{ g_0(\mathbf{z}) + \frac{1}{m} \sum_{i=1}^m y_i g_i(\mathbf{z}) - \frac{\lambda}{2} \mathbf{y}_i^2 + \frac{1}{2\nu} \left\| \mathbf{z} - \mathbf{w} \right\|^2 \right\} \right\}, \tag{38}$$

where $\Phi_{\lambda,\nu}(\mathbf{w})$ is defined in (9). According to Proposition D.2 in [6], the second (compositional) term in $\Phi_{\lambda}(\mathbf{w})$ is $(\rho\rho_g)$ -weakly convex. Hence, $\Phi_{\lambda}(\mathbf{w})$ is $\rho_{\Phi_{\lambda}}$ -weakly convex with $\rho_{\Phi_{\lambda}} = \rho_g + \rho\rho_g$. If $\nu < \rho_{\Phi_{\lambda}}^{-1}$, $\Phi_{\lambda,\nu}(\mathbf{w})$ is $(\frac{1}{\nu} - \rho_{\Phi_{\lambda}})$ -strongly convex. As done in Section (4.2), in the tth iteration of Algorithm 2, we apply ALEXR to solve the inner min-max problem in (38) with $\mathbf{w} = \mathbf{w}_t$, namely,

$$\min_{\mathbf{z}} \max_{\mathbf{y} \in [0,\rho]^m} \left\{ L_t(\mathbf{z}, \mathbf{y}) := g_0(\mathbf{z}) + \frac{1}{m} \sum_{i=1}^m \left(\mathbf{y}_i g_i(\mathbf{w}) - \frac{\lambda}{2} y_i^2 \right) + \frac{1}{2\nu} \|\mathbf{z} - \mathbf{w}_t\|_2^2 \right\}.$$

we only need to replace $G_{t,k}$ in Algorithm 2 to

$$G_{t,k} = G_{t,k}^{0} + G_{t,k}^{1} = \partial g_{0}(\mathbf{z}_{t,k}, \mathcal{B}_{0,2}^{t,k}) + \frac{1}{|\mathcal{B}_{1}^{t,k}|} \sum_{i \in \mathcal{B}_{1}^{t,k}} [\partial g_{i}(\mathbf{z}_{t,k}, \tilde{\mathcal{B}}_{i,2}^{t,k})]^{\top} \mathbf{y}_{t,k+1}^{(i)},$$
(39)

where $\mathcal{B}^{t,k}_{0,2}$ is a batch of data sampled from the distribution of ξ_0 .

We then consider the case where $\{g_i\}_{i=0}^m$ are smooth. In this case, we can directly apply Algorithm 1 to (9), which is different from (3) only in the extra term g_0 . To handle this difference, we only need to replace G_t in Algorithm 1 to

$$G_{t} = G_{t}^{0} + G_{t}^{1} = \nabla g_{0}(\mathbf{w}_{t}, \mathcal{B}_{0,2}^{t}) + \frac{1}{|\mathcal{B}_{1}^{t}|} \sum_{i \in \mathcal{B}_{1}^{t}} \nabla f_{i,\lambda}(u_{t,i}) \nabla g_{i}(\mathbf{w}_{t}, \mathcal{B}_{i,2}^{t})$$
(40)

where $\mathcal{B}_{0,2}^t$ is a batch of data sampled from the distribution of ξ_0 . With these changes, we can prove Theorem 5.3 as follows.

Sketch Proof of Theorem 5.3. When $\{g_i\}_{i=0}^m$ are weakly convex:

For simplicity of notation, we drop the dependence on t in the notation of all variables and parameters in the proof below. First, we need to modify Lemma 5 in [5]. We have

$$L_{t}(\mathbf{z}_{k+1}, \mathbf{y}) - L_{t}(\mathbf{z}, \bar{\mathbf{y}}_{k+1})$$

$$\leq \frac{\gamma^{-1}}{m} U_{\bar{f}_{i}^{*}}(\mathbf{y}, \mathbf{y}_{k}) - \frac{\gamma^{-1} + 1}{m} U_{\bar{f}_{i}^{*}}(\mathbf{y}, \bar{\mathbf{y}}_{k+1}) - \frac{\gamma^{-1}}{m} U_{\bar{f}_{i}^{*}}(\bar{\mathbf{y}}_{k+1}, \mathbf{y}_{k})$$

$$+ \frac{1}{m} \sum_{i=1}^{m} \left\langle g_{i}(\mathbf{z}_{k+1}) - \tilde{g}_{k}^{(i)}, y_{i} - \bar{\mathbf{y}}_{k+1}^{(i)} \right\rangle + \frac{1}{2\eta} \|\mathbf{z} - \mathbf{z}_{k}\|_{2}^{2}$$

$$- \left(\frac{1}{2\eta} + \frac{1}{4\nu} \right) \|\mathbf{z} - \mathbf{z}_{k+1}\|_{2}^{2} - \frac{1}{2\eta} \|\mathbf{z}_{k+1} - \mathbf{z}_{k}\|_{2}^{2}$$

$$+ \frac{1}{m} \sum_{i=1}^{m} \left\langle g_{i}(\mathbf{z}_{k+1}) - g_{i}(\mathbf{z}), \bar{\mathbf{y}}_{k+1}^{(i)} \right\rangle - \left\langle G_{k}^{1}, \mathbf{z}_{k+1} - \mathbf{z} \right\rangle$$

$$+ g_{0}(\mathbf{z}_{k+1}) - g_{0}(\mathbf{z}) - \left\langle G_{k}^{0}, \mathbf{z}_{k+1} - \mathbf{z} \right\rangle,$$

$$(41)$$

where the blue terms above are included due to the modifications to Algorithm 2. Next, we modify Lemma 8 in [5] to handle the additional terms above. Denote that $\Delta_k^0 := g_0'(\mathbf{z}_k) - G_k^0$. Note that

$$-\left\langle G_k^0, \, \mathbf{z}_{k+1} - \mathbf{z} \right\rangle = -\left\langle g_0'(\mathbf{z}_k), \, \mathbf{z}_{k+1} - \mathbf{z} \right\rangle + \left\langle \Delta_k^0, \, \mathbf{z}_{k+1} - \mathbf{z} \right\rangle,$$

where $g_0'(\mathbf{z}_k) \in \partial g(\mathbf{z}_k)$ is a subgradient at \mathbf{z}_k . The term $\langle \Delta_k^0, \mathbf{z}_{k+1} - \mathbf{z} \rangle$ can be handled in the same way as in (24) in [5]. Due to the weak convexity of g_0 , we have

$$g_0(\mathbf{z}) - g_0(\mathbf{z}_k) \ge \langle g_0'(\mathbf{z}_k), \mathbf{z} - \mathbf{z}_k \rangle - \frac{\rho_g}{2} \|\mathbf{z} - \mathbf{z}_k\|_2^2$$

Then, the remaining blue terms in (41) can be upper bounded as

$$\begin{split} &g_{0}(\mathbf{z}_{k+1}) - g_{0}(\mathbf{z}) - \langle g_{0}'(\mathbf{z}_{k}), \, \mathbf{z}_{k+1} - \mathbf{z} \rangle \\ &= g_{0}(\mathbf{z}_{k+1}) - g_{0}(\mathbf{z}_{k}) + g_{0}(\mathbf{z}_{k}) - g_{0}(\mathbf{z}) - \langle g_{0}'(\mathbf{z}_{k}), \mathbf{z}_{k+1} - \mathbf{z} \rangle \\ &\leq C_{g} \|\mathbf{z}_{k+1} - \mathbf{z}_{k}\|_{2} + \langle g_{0}'(\mathbf{z}_{k}), \mathbf{z}_{k} - \mathbf{z} \rangle - \langle g_{0}'(\mathbf{z}_{k}), \mathbf{z}_{k+1} - \mathbf{z} \rangle + \frac{\rho_{g}}{2} \|\mathbf{z} - \mathbf{z}_{k}\|_{2}^{2} \\ &\leq \frac{4C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}} + \frac{\left(\frac{1}{\eta} + \frac{1}{\nu}\right) \|\mathbf{z}_{k+1} - \mathbf{z}_{k}\|_{2}^{2}}{16} + \langle g_{0}'(\mathbf{z}_{k}), \mathbf{z}_{k} - \mathbf{z}_{k+1} \rangle + \frac{\rho_{g}}{2} \|\mathbf{z} - \mathbf{z}_{k}\|_{2}^{2} \\ &\leq \frac{8C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}} + \frac{\left(\frac{1}{\eta} + \frac{1}{\nu}\right) \|\mathbf{z}_{k+1} - \mathbf{z}_{k}\|_{2}^{2}}{8} + \frac{\rho_{g}}{2} \|\mathbf{z} - \mathbf{z}_{k}\|_{2}^{2}. \end{split}$$

Let $(\mathbf{z}_*, \mathbf{y}_*)$ be the saddle point of $L_t(\mathbf{z}, \mathbf{y})$. Then, we can get

$$\mathbb{E}[L_{t}(\mathbf{z}_{k+1}, \mathbf{y}_{*}) - L_{t}(\mathbf{z}_{*}, \bar{\mathbf{y}}_{k+1})] \\
\leq \frac{\gamma^{-1} + \left(1 - \frac{B_{1}}{m}\right)}{B_{1}} \mathbb{E}[U_{\bar{f}_{i}^{*}}(\mathbf{y}_{*}, \mathbf{y}_{k})] - \frac{\gamma^{-1} + 1}{B_{1}} \mathbb{E}[U_{\bar{f}_{i}^{*}}(\mathbf{y}_{*}, \mathbf{y}_{k+1})] \\
+ \left(\frac{1}{2\eta} + \frac{\sqrt{d_{1}}C_{f}\rho_{g}}{2} + \frac{\rho_{g}}{2}\right) \mathbb{E} \|\mathbf{z}_{*} - \mathbf{z}_{k}\|_{2}^{2} - \left(\frac{1}{2\eta} + \frac{1}{4\nu}\right) \mathbb{E} \|\mathbf{z}_{*} - \mathbf{z}_{k+1}\|_{2}^{2} \\
- \left(\frac{1}{m\gamma} - \frac{\lambda_{2} + \lambda_{3}\theta}{\lambda m}\right) \mathbb{E} \left[U_{\bar{f}_{i}^{*}}(\bar{\mathbf{y}}_{k+1}, \mathbf{y}_{k})\right] - \left(\frac{1}{2\eta} - \frac{C_{g}^{2}}{2\lambda_{2}} - \frac{3}{8}\left(\frac{1}{\eta} + \frac{1}{\nu}\right)\right) \mathbb{E} \|\mathbf{z}_{k+1} - \mathbf{z}_{k}\|_{2}^{2} \\
+ \frac{\theta C_{g}^{2}}{2\lambda_{3}} \mathbb{E} \|\mathbf{z}_{k} - \mathbf{z}_{k-1}\|_{2}^{2} + \mathbb{E}[\Gamma_{k+1} - \theta\Gamma_{k}] + \frac{2(1 + 2\theta)\sigma_{0}^{2}}{B_{2}\lambda(1 + \gamma^{-1})} + \frac{\frac{C_{f}^{2}C_{g}^{2}}{B_{1}} + \frac{C_{f}^{2}C_{g}^{2}}{B_{1}} + 8C_{f}^{2}C_{g}^{2} + 8C_{g}^{2}}{\frac{1}{\eta} + \frac{1}{\nu}}. \tag{44}$$

Following the similar proof as in Theorem C.1, the modified inner loop of ALEXR2 described in Section 5 can guarantee $\frac{\mu}{2}\mathbb{E}\left\|\hat{\mathbf{z}}_t - \mathrm{prox}_{\nu F_\lambda}(\mathbf{w}_t)\right\|_2^2 \leq \epsilon'$ for any t after $K_t = \tilde{O}(\frac{m}{\lambda B_2 B_1 \epsilon'})$. Because

 $\lambda = \frac{\epsilon}{\rho}$ and $\rho > \frac{C_g}{\delta}$ as specified in Proposition D.1, we have $K_t = \tilde{O}(\frac{C_g^2 m}{B_2 B_1 \delta^2 \epsilon \epsilon'})$. For outer loop, we directly leverage the result from Appendix C, i.e.

$$T = O(\max\{\rho_g + \rho\rho_g, \rho C_g\}\epsilon^{-2}) = O(\max\{\rho_g + \frac{C_g\rho_g}{\delta}, \frac{C_g^2}{\delta}\}\epsilon^{-2})$$

By setting $\epsilon' = \frac{\epsilon^2}{\rho_g + \rho \rho_g} = \frac{\epsilon^2}{\rho_g + \frac{C_g \rho_g}{\epsilon}}$, the total iteration will be

$$\tilde{\mathcal{O}}\left(\frac{m\rho_g^2}{B_2B_1\delta^4\epsilon^5}\right)$$

When $\{g_i\}_{i=0}^m$ are smooth:

The proof follows almost the same procedure as the proof of Theorem 4.7. In particular, since \mathbf{v}_t and G_t will contain the additional stochastic gradient from g_0 , we need to replace (14) by

$$V_{t+1} := \left\| \mathbf{v}_{t+1} - \nabla \Phi_{\lambda}(\mathbf{w}_t) \right\|^2 \tag{45}$$

while U_{t+1} , \mathbf{u}_t and $\mathbf{g}(\mathbf{w}_t)$ are still defined as in (15), (16) and (17). Like Lemma B.1, it holds that

$$\Phi_{\lambda}(\mathbf{w}_{t+1}) \le \Phi_{\lambda}(\mathbf{w}_{t}) + \frac{\eta}{2} V_{t+1} - \frac{\eta}{2} \|\nabla \Phi_{\lambda}(\mathbf{w}_{t})\|^{2} - \frac{\eta}{4} \|\mathbf{v}_{t+1}\|^{2},$$
(46)

if $\eta \leq \frac{1}{2L_F}$. By a proof similar to Lemma 3 in [2], we can still show (19) for V_{t+1} in the new definition in (45). Moreover, (20) and (21) still hold as they are not related to g_0 . Finally, we can still take the weighted summation of both sides (46), (19), (20), and (21) as in the proof of Theorem 4.7 to show that

$$\frac{1}{T} \sum_{t=0}^{T-1} \|\nabla \Phi_{\lambda}(\mathbf{w}_{t})\|^{2} \leq \frac{1}{T} O\left(\frac{m}{B_{1} \sqrt{B_{2}} \epsilon^{3}} + \frac{1}{\min\{B_{1}, B_{2}\} \epsilon^{2}} + \frac{m}{B_{1} B_{2} \epsilon^{3}}\right).$$

This means Algorithm 1 finds a nearly ϵ -stationary point of $\Phi_{\lambda}(\mathbf{w})$ in expectation in $T = O(\frac{m}{B_1\sqrt{B_2}\epsilon^5})$ iterations. Then the conclusion follows from Proposition D.1 and the fact that $C_f = O(\frac{1}{\delta})$.

E SONEX with Adaptive Learning Rates

In this section we will show that our algorithms can also be easily extended to adaptive learning rate while still retaining the same complexity under an additional assumption introduced later. We consider an adaptive step size update such as adam:

$$\mathbf{w}_{t+1} = \mathbf{w}_t - \tilde{\eta} \circ \mathbf{v}_{t+1}, \tilde{\eta} = \frac{\eta}{\sqrt{\mathbf{s}_t} + \varepsilon}, \mathbf{s}_{t+1} = (1 - \beta')\mathbf{s}_t + \beta'G_t \circ G_t$$

where G_t is the overall gradient estimator mentioned in algorithm 1 and \circ denotes Hadamard(elementwise) product. The following assumption has been justified for the adaptive step size by [48].

Assumption E.1. We assume that the adaptive learning rates $\tilde{\eta}$ are (element-wise)bounded, i.e.

$$\eta c_l \le \tilde{\eta}_i \le \eta c_u$$

for any element $\tilde{\eta}_i$ of $\tilde{\eta}$.

Below we provide lemmas which are straightforward extensions of lemma B.1 \sim B.4 to adaptive learning setting. The proof is similar except that $\|\mathbf{w}_{t+1} - \mathbf{w}_t\|^2 \le \|\tilde{\eta} \circ \mathbf{v}_{t+1}\|^2 \le \|\eta^2 c_u^2 \mathbf{v}_{t+1}\|^2 = \eta^2 c_u^2 \|\mathbf{v}_{t+1}\|^2$.

Lemma E.2. (Lemma 3 in [48]) Under assumption E.1, for $\mathbf{w}_{t+1} = \mathbf{w}_t - \tilde{\eta} \circ \mathbf{v}_{t+1}$, with $\eta c_l \leq \tilde{\eta} \leq \eta c_u$ and $\tilde{\eta} L_F \leq \frac{c_l}{2c_u^2}$, we have:

$$F(\mathbf{w}_{t+1}) \le F(\mathbf{w}_t) + \frac{\eta c_u}{2} V_{t+1} - \frac{\eta c_l}{2} \|\nabla F(\mathbf{w}_t)\|^2 - \frac{\eta c_l}{4} \|\mathbf{v}_{t+1}\|^2$$
(47)

where $V_{t+1} = \|\mathbf{v}_{t+1} - \nabla F(\mathbf{w}_t)\|^2$

Lemma E.3. If $\beta \leq \frac{2}{7}$, the gradient variance $V_{t+1} := \|\mathbf{v}_{t+1} - \nabla F(\mathbf{w}_t)\|^2$ can be bounded as

$$\mathbb{E}\left[V_{t+2}\right] \leq (1-\beta)\mathbb{E}\left[V_{t+1}\right] + \frac{2L_F^2\eta^2c_u^2\mathbb{E}\left[\|\mathbf{v}_{t+1}\|^2\right]}{\beta} + \frac{3L_f^2C_1^2}{n}\mathbb{E}\left[\sum_{\mathbf{z}_i \in \mathcal{B}_1^{t+1}} \|u_{t+2,i} - u_{t+1,i}\|^2\right] + \frac{2\beta^2C_f^2(\zeta^2 + C_g^2)}{\min\{B_1, B_2\}} + 5\beta L_f^2C_1^2\mathbb{E}\left[U_{t+2}\right],$$
(48)

where
$$U_{t+1} = \frac{1}{n} \|\mathbf{u}_{t+1} - \mathbf{g}(\mathbf{w}_t)\|^2$$
, $\mathbf{u}_t = [u_{t,1}, \dots, u_{t,n}]^\top$, $\mathbf{g}(\mathbf{w}_t) = [g_1(\mathbf{w}_t), \dots, g_n(\mathbf{w}_t)]^\top$.

Lemma E.4. Suppose $\gamma' = \frac{n-B_1}{B_1(1-\gamma)} + (1-\gamma)$ and $\gamma \leq \frac{1}{2}$ and the MSVR update is used in Algorithm 1. It holds that

$$\mathbb{E}\left[U_{t+2}\right] \le \left(1 - \frac{\gamma B_1}{n}\right) \mathbb{E}\left[U_{t+1}\right] + \frac{8nC_g^2 \eta^2 c_u^2}{B_1} \mathbb{E}\left\|\mathbf{v}_{t+1}\right\|^2 + \frac{2B_1 \gamma^2 \sigma_1^2}{nB_2}.$$
 (49)

Lemma E.5. Under the same assumptions as Lemma B.3, it holds that

$$\mathbb{E}[\|\mathbf{u}_{t+2} - \mathbf{u}_{t+1}\|^2] \le \frac{4B_1 \gamma^2}{n} \mathbb{E}[U_{t+1}] + \frac{9n^2 C_g^2 \eta^2 c_u^2}{B_1} \mathbb{E}\|\mathbf{v}_{t+1}\|^2 + \frac{2B_1 \gamma^2 \sigma_0^2}{B_2}.$$
 (50)

We then give the following theorem with similar proof technique.

Theorem E.6. Under Assumption 3.2 (A1), 4.5 and 4.6, by setting $\lambda = \Theta(\epsilon), \beta = \Theta(\frac{c_l \min\{B_1, B_2\}\epsilon^2}{c_u}), \gamma = \Theta(\frac{c_l B_2 \epsilon^4}{c_u}), \eta = \Theta(\frac{c_l^{1.5} B_1 \sqrt{B_2}}{c_u^{2.5} n} \epsilon^3)$, SONEX with $\gamma' = 1 - \gamma + \frac{n - B_1}{B_1(1 - \gamma)}$ converges to an approximate ϵ -stationary solution of (1) within $T = O(\frac{c_u^{2.5} n}{c_r^{2.5} B_1 \sqrt{B_2}} \epsilon^{-5})$ iterations.

Combining the above theorem with Theorem 4.4, we obtain the following guarantee:

Corollary E.7. Under Assumption 3.2(A1), 4.3, 4.5 and 4.6, with the same setting as in Theorem E.6, SONEX converges to a nearly ϵ -stationary solution of (1) within $T = O(\frac{c_u^{2.5}n}{c_r^{2.5}B_1\sqrt{B_2}}\epsilon^{-5})$ iterations.

Proof of theorem E.6 and corollary E.7. Since the proof is almost the same as the theorem 4.7 and corollary 4.8, we only highlight the difference.

Let P > 0 be a positive constant to be decided later. Taking the weighted summation of both sides (47), (48), (49), and (50) as specified in the following formula

$$\frac{1}{\eta c_u} \times (47) + \frac{1}{\beta} \times (48) + P \times (49) + \frac{3L_f^2 C_1^2}{n\beta} \times (50),$$

we have

$$\begin{split} &\frac{1}{\eta c_{u}} \mathbb{E} F_{\lambda}(\mathbf{w}_{t+1}) + \frac{1}{\beta} \mathbb{E} V_{t+2} + \left(P - 5L_{f}^{2}C_{1}^{2}\right) \mathbb{E} U_{t+2} \\ \leq &\frac{1}{\eta c_{u}} \mathbb{E} F_{\lambda}(\mathbf{w}_{t}) + \left(\frac{1}{\beta} - \frac{1}{2}\right) \mathbb{E} V_{t+1} + \left(P\left(1 - \frac{\gamma B_{1}}{n}\right) + \frac{12L_{f}^{2}C_{1}^{2}\gamma^{2}B_{1}}{\beta n^{2}}\right) \mathbb{E} U_{t+1} \\ &- \left(\frac{c_{l}}{4c_{u}} - \frac{2L_{F}^{2}\eta^{2}c_{u}^{2}}{\beta^{2}} - \frac{8nC_{g}^{2}P\eta^{2}c_{u}^{2}}{B_{1}} - \frac{27nL_{f}^{2}C_{1}^{2}C_{g}^{2}\eta^{2}c_{u}^{2}}{B_{1}\beta}\right) \mathbb{E} \left\|\mathbf{v}_{t+1}\right\|^{2} \\ &+ \frac{2\beta C_{f}^{2}(C_{g}^{2} + \sigma_{1}^{2})}{\min\{B_{1}, B_{2}\}} + \frac{2B_{1}\gamma^{2}\sigma_{1}^{2}P}{nB_{2}} + \frac{6L_{f}^{2}C_{1}^{2}\gamma^{2}\sigma_{0}^{2}B_{1}}{\beta nB_{2}} - \frac{c_{l}}{2c_{u}} \mathbb{E} \left\|\nabla F_{\lambda}(\mathbf{w}_{t})\right\|^{2} \end{split}$$

We choose η such that

$$\frac{c_l}{4c_u} - \frac{2L_F^2 \eta^2 c_u^2}{\beta^2} - \frac{8nC_g^2 P \eta^2 c_u^2}{B_1} - \frac{27nL_f^2 C_1^2 C_g^2 \eta^2 c_u^2}{B_1 \beta} \ge 0.$$
 (51)

We follow proof of 4.7 to set

$$P = \frac{n}{\gamma B_1} \left(5 L_f^2 C_1^2 + \frac{12 L_f^2 C_1^2 \gamma^2 B_1}{\beta n^2} \right).$$

which satisfies the following linear equation

$$P\left(1 - \frac{\gamma B_1}{n}\right) + \frac{12L_f^2 C_1^2 \gamma^2 B_1}{\beta n^2} = P - 5L_f^2 C_1^2,$$

Combining the results above gives

$$\begin{split} &\frac{1}{\eta c_{u}} \mathbb{E} F_{\lambda}(\mathbf{w}_{t+1}) + \frac{1}{\beta} \mathbb{E} V_{t+2} + \left(P - 5L_{f}^{2}C_{1}^{2}\right) \mathbb{E} U_{t+2} \\ \leq &\frac{1}{\eta c_{u}} \mathbb{E} F_{\lambda}(\mathbf{w}_{t}) + \frac{1}{\beta} \mathbb{E} V_{t+1} + \left(P - 5L_{f}^{2}C_{1}^{2}\right) \mathbb{E} U_{t+1} \\ &+ \frac{2\beta C_{f}^{2}(C_{g}^{2} + \sigma_{1}^{2})}{\min\{B_{1}, B_{2}\}} + \frac{2B_{1}\gamma^{2}\sigma_{1}^{2}P}{nB_{2}} + \frac{6L_{f}^{2}C_{1}^{2}\gamma^{2}\sigma_{0}^{2}B_{1}}{\beta nB_{2}} - \frac{c_{l}}{2c_{u}} \mathbb{E} \left\|\nabla F_{\lambda}(\mathbf{w}_{t})\right\|^{2}. \end{split}$$

Summing both sides of the inequality above over $t = 0, 1, \dots, T-1$ and organizing term lead to

$$\frac{1}{T} \sum_{t=0}^{T-1} \|\nabla F_{\lambda}(\mathbf{w}_{t})\|^{2} \leq \frac{2c_{u}}{c_{l}T} \left(\frac{1}{\eta c_{u}} (F_{\lambda}(\mathbf{w}_{0}) - F_{\lambda}(\mathbf{w}_{T})) + \frac{1}{\beta} \mathbb{E}V_{1} + \left(P - 5L_{f}^{2}C_{1}^{2} \right) \mathbb{E}U_{1} \right) \\
+ \frac{c_{u}}{c_{l}} \left(\frac{4\beta C_{f}^{2} (C_{g}^{2} + \sigma_{1}^{2})}{\min\{B_{1}, B_{2}\}} + \frac{4B_{1}\gamma^{2}\sigma_{1}^{2}P}{nB_{2}} + \frac{12L_{f}^{2}C_{1}^{2}\gamma^{2}\sigma_{0}^{2}B_{1}}{\beta nB_{2}} \right)$$
(52)

Note that $L_F=O(\frac{1}{\epsilon}),\,L_f=O(\frac{1}{\epsilon})$ and $P=O(\frac{n}{B_1\gamma\epsilon^2}+\frac{\gamma}{n\beta\epsilon^2})$. We require that all the terms in the right-hand side to be in the order of $O(\epsilon^2)$. To ensure $\frac{4c_u\beta C_f^2(C_g^2+\sigma_1^2)}{c_l\min\{B_1,B_2\}}=O(\epsilon^2)$, we must have $\beta=O(\frac{c_l\min\{B_1,B_2\}\epsilon^2}{c_u})$. To ensure $\frac{4c_uB_1\gamma^2\sigma_1^2P}{c_lnB_2}=\frac{c_u}{c_l}O(\frac{\gamma}{B_2\epsilon^2}+\frac{B_1\gamma^3}{n^2B_2\beta\epsilon^2})=O(\epsilon^2)$, we must have $\gamma=O(\frac{c_lB_2\epsilon^4}{c_u})$. With these orders of β and γ , we also have $P=O(\frac{c_un}{c_lB_1B_2\epsilon^6})$ and $\frac{12c_uL_f^2C_1^2\gamma^2\sigma_0^2B_1}{c_l\beta nB_2}=O(\epsilon^4)$. Therefore the last three terms in (52) are $O(\epsilon^2)$.

To satisfy (51), we will need to set

$$\eta = O\left(\sqrt{\frac{c_l}{c_u^3}} \min\left\{\frac{\beta}{L_F}, \sqrt{\frac{B_1}{nP}}, \frac{1}{L_f}\sqrt{\frac{B_1\beta}{n}}\right\}\right)
= O\left(\sqrt{\frac{c_l}{c_u^3}} \min\left\{\frac{c_l}{c_u} \min\{B_1, B_2\}\epsilon^3, \frac{B_1\sqrt{B_2}}{n}\epsilon^3, \sqrt{\frac{c_lB_1\min\{B_1, B_2\}}{c_un}}\epsilon^2\right\}\right)
= O\left(\frac{c_l^{1.5}B_1\sqrt{B_2}}{c_u^{2.5}n}\epsilon^3\right).$$

We also set the sizes of $\mathcal{B}_{i,2}$ to be $O(\frac{1}{\epsilon^3})$ so that $\mathbb{E}U_1 = O(\epsilon^3)$. Also, it is easy to see that $F_{\lambda}(\mathbf{w}_0) - F_{\lambda}(\mathbf{w}_T) = O(1)$ and $\mathbb{E}V_1 = O(1)$. Then the first term in (52) becomes

$$\frac{1}{T}O\left(\frac{1}{\eta c_{l}} + \frac{1}{\beta} + P\epsilon^{3}\right) = \frac{1}{T}O\left(\frac{c_{u}^{2.5}n}{c_{l}^{2.5}B_{1}\sqrt{B_{2}}\epsilon^{3}} + \frac{c_{u}}{c_{l}\min\{B_{1}, B_{2}\}\epsilon^{2}} + \frac{c_{u}n}{c_{l}B_{1}B_{2}\epsilon^{3}}\right)$$

which will become $O(\epsilon^2)$ when $T = O(\frac{c_u^{2.5}n}{c_l^{2.5}B_1\sqrt{B_2}\epsilon^5})$.

Then the remaining part is just the same as the proof of corollary 4.8.

F More Experiment Details

F.1 GDRO with CVaR divergence

Motivation of GDRO Modern machine learning models are typically trained under the empirical risk minimization (ERM) framework, which treats all samples in the dataset equally. Although ERM

often achieves strong average performance on test sets drawn from distributions similar to the training data, it can perform poorly on rare or underrepresented subpopulations, i.e., it lacks robustness to distributional shifts. The motivation of GDRO is to address the spurious correlation between features and labels, and distributional shift. GDRO partitions the dataset into groups representing different distributions and applies a robust optimization scheme that assigns more weight on the worst groups. Specifically, GDRO considers the following objective:

$$\min_{\theta} \max_{\mathbf{p} \in \Omega} \frac{1}{n} \sum_{g=1}^{n} p_g \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{D}_g} l(\theta; (\mathbf{x}, y))$$

where $\Omega \subseteq \Delta$ is the set of distribution under consideration and Δ is the simplex. We consider here a popular choice of $\Omega = \{\mathbf{p} \in \Delta : p_i \leq \frac{1}{k}, \forall i \in [n]\}$, which, by solving the inner-level maximization problem, leads to an equivalent reformulation 10 corresponding to the so-called CVaR divergence [5, 49]. Intuitively GDRO with CVaR divergence aims to minimize averaged loss of the top-k worst groups.

CAMELYON17-WILDS [34] is part of the WILDS benchmark suite and consists of histopathology whole-slide images from five medical centers, with the goal of detecting metastatic tissue in lymph node biopsies. Following the WILDS setup, we frame this as a binary classification task on image patches, where the primary challenge lies in distribution shift across hospitals (domains). We construct group with attributes 'hospital' and 'slide' which generates 30 groups.

Amazon-WILDS [50] is a text classification dataset derived from Amazon product reviews, where the goal is to predict binary sentiment (positive or negative) based on TF-IDF features of review text. The data spans multiple product categories. We construct group with the attribute 'user' which generates 1252 groups.

CelebA [35] is a large-scale facial attribute dataset containing over 200,000 celebrity images annotated with 40 binary attributes. We select 4 attributes 'Attractive', 'Mouth_Slightly_Open', 'Male' and 'Blonde_Hair' and construct 16 groups, where 'Blonde_Hair' also serves as the target attribute for us to do classification.

Hyperparameter tuning. We tune the same hyperparameters of different methods from the same candidates as follows for fair comparison. For the three tasks we train the models for 10, 4, 15 epochs with batch size and the number of groups within a mini batch of 256(8), 32(8), 64(4), respectively. We set $\alpha = 0.15$ for all the three dataset. We tune learning rate in {1e-5, 2e-5, 5e-5, 1e-4, 2e-4, 5e-4, 1e-3, 2e-3, 5e-3}, λ in {1, 0.1, 0.01}, γ and β in {0.1, 0.2, 0.5, 0.8} and γ' in {0.01, 0.02, 0.05, 0.1, 0.2}. We set weight decay to be 0.01, 0.01, 0.02 for the three tasks, respectively. We use step decay (decay by 0.3x for every 3 epochs), linear decay with 1st epoch warmup, step decay (decay by 0.2x for every 3 epochs) for learning rate for the three tasks, respectively.

F.2 AUC Maximization with ROC Fairness Constraints

In this part we perform experiment on learning a model with ROC fairness constraint [51] following the same experiment setting as [8]. Suppose the data are divided into two demographic groups $\mathcal{D}_p = \{(\mathbf{a}_i^p, b_i^p)\}_{i=1}^{n_p} \text{ and } \mathcal{D}_u = \{(\mathbf{a}_i^u, b_i^u)\}_{i=1}^{n_u}, \text{ where a denotes the input data and } b \in \{1, -1\}$ denotes the class label. A ROC fairness is to ensure the ROC curves for classification of the two groups are the same.

Since the ROC curve is constructed with all possible thresholds, we follow [51] by using a set of thresholds $\Gamma = \{\tau_1, \cdots, \tau_m\}$ to define the ROC fairness. For each threshold τ , we impose a constraint that the false positive rate (FPR) and true positive rate (TPR) of the two groups are close, formulated as follow:

$$h_{\tau}^{+}(\mathbf{w}) := \left| \frac{1}{n_{p}^{+}} \sum_{i=1}^{n_{p}} \mathbf{I}\{b_{i}^{p} = 1\} \sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{p}) - \tau) - \frac{1}{n_{u}^{+}} \sum_{i=1}^{n_{u}} \mathbf{I}\{b_{i}^{u} = 1\} \sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{u}) - \tau) \right| - \kappa \le 0$$

$$h_{\tau}^{-}(\mathbf{w}) := \left| \frac{1}{n_{p}^{-}} \sum_{i=1}^{n_{p}} \mathbf{I} \{ b_{i}^{p} = -1 \} \sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{p}) - \tau) - \frac{1}{n_{u}^{-}} \sum_{i=1}^{n_{u}} \mathbf{I} \{ b_{i}^{u} = -1 \} \sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{u}) - \tau) \right| - \kappa \leq 0,$$




Figure 4: Training curves of 5 constraint values in zero-one loss of different methods for continual learning with non-forgetting constraints when targeting the foggy class. Top: squared-hinge penalty method with different ρ ; Bottom: smoothed hinge penalty method with different ρ .

where $s_{\mathbf{w}}(\cdot)$ denotes a parameterized model, $\sigma(z)$ is the sigmoid function, and $\kappa > 0$ is a tolerance parameter. We use the pairwise AUC loss as the objective function:

$$F(\mathbf{w}) = \frac{-1}{|\mathcal{D}_{+}||\mathcal{D}_{-}|} \sum_{\mathbf{x}_{i} \in \mathcal{D}_{+}} \sum_{\mathbf{x}_{j} \in \mathcal{D}_{-}} \sigma(s(\mathbf{w}, \mathbf{x}_{i}) - s(\mathbf{w}, \mathbf{x}_{j})),$$

where \mathcal{D}_+ and \mathcal{D}_- denote the set of positive and negative examples regardless their groups, respectively. We follow [8] to recast original constraint optimization as the following hinge-penalized objective:

$$\min_{\mathbf{w}} F(\mathbf{w}) + \frac{1}{2|\Gamma|} \sum_{\tau \in \Gamma} \beta [h_{\tau}^{+}(\mathbf{w})]_{+} + \beta [h_{\tau}^{-}(\mathbf{w})]_{+}$$
(53)

It is notable that the penalty terms in the above problem can be formulated as $[h_{\tau}^{+}(\mathbf{w})]_{+} = f(g(\mathbf{w};\tau))$, where $f(g) = [|g_{1} - g_{2}| - \kappa]_{+}, \ g_{1}(\mathbf{w};\tau) = \frac{1}{n_{p}^{+}} \sum_{i=1}^{n_{p}} \mathbf{I}\{b_{i}^{p} = 1\}\sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{p}) - \tau), \ g_{2}(\mathbf{w};\tau) = \frac{1}{n_{u}^{+}} \sum_{i=1}^{n_{u}} \mathbf{I}\{b_{i}^{u} = 1\}\sigma(s_{\mathbf{w}}(\mathbf{a}_{i}^{u}) - \tau)$. As a result, f is convex, $g(\mathbf{w})$ is smooth. As a result, Algorithm 2 is applicable.

Hyperparameter tuning. For ALEXR2, we tune K_t in $\{5,10\}$, tune α (i.e. learning rate for outer loop) and η (i.e. learning rate for inner loop) in $\{1\text{e-3}, 1\text{e-2}, 0.1, 1\}$, and tune λ (i.e. smoothing coefficient for outer function) and ν (i.e. smoothing coefficient for overall objective) in $\{2\text{e-2}, 2\text{e-3}, 2\text{e-4}\}$ and $\{0.1, 0.01\}$, respectively. We fix the hyperparameters in MSVR update for both ALEXR2 7 and SONX, i.e. $\hat{\gamma}=0.8$ and $\hat{\gamma}'=\hat{\gamma}\theta=0.1$. For the two baseline algorithms, we tune the initial learning rate in $\{0.1, 1\text{e-2}, 1\text{e-3}, 1\text{e-4}\}$. We decay learning rate(outer Ir for ALEXR2) at 50% and 75% epochs by a factor of 10. We tuned $\rho=\{4,6,8,10,20,40\}$ for ALEXR2 and SONX and $\rho=\{10,40,80,100,200,400,800,1000\}$ for SOX. We also compare a double-loop method (ICPPAC) [10, Algorithm 4], where we tune their η in $\{0.1,0.01\}$, τ in $\{1,10,100\}$, μ in $\{1\text{e-2},1\text{e-3},1\text{e-4}\}$, and fix θ_t to 0.1.

F.3 Continual learning with non-forgetting constraints

We follow [7] and consider training a CLIP model [52] with global contrastive loss (GCL) [17] as the objective and a so-called model developmental safety (MDS) constraints on protected tasks, given by:

$$\begin{aligned} & \min_{\mathbf{w}} F(\mathbf{w}, \mathcal{D}) := \frac{1}{n} \sum_{(\mathbf{x}_i, \mathbf{t}_i) \in \mathcal{D}} L_{\text{GCL}}(\mathbf{w}; (\mathbf{x}_i, \mathbf{t}_i), (\mathcal{T}_i^-, \mathcal{I}_i^-)), \\ & \text{s.t. } h_k := \mathcal{L}_k(\mathbf{w}, D_k) - \mathcal{L}_k(\mathbf{w}_{\text{old}}, D_k) \leq 0, k = 1, \cdots, m. \end{aligned}$$

⁷As discuss in section 4.2, update of \mathbf{y} in inner loop is equivalent to MSVR update, i.e. $y_{i,t} = \nabla f_{i,\lambda}(u_{i,t}), u_{i,t+1} = (1-\hat{\gamma})u_{i,t} + \hat{\gamma}g_i(\mathbf{z}_{t,k}; \mathcal{B}_{i,2}^{t,k}) + \hat{\gamma}\theta(g_i(\mathbf{z}_{t,k}; \mathcal{B}_{i,2}^{t,k}) - g_i(\mathbf{z}_{t-1,k}; \mathcal{B}_{i,2}^{t,k}))$




Figure 5: Training curves of 4 constraint values in zero-one loss of different methods for continual learning with non-forgetting constraints when targeting the overcast class. Top: squared-hinge penalty method with different ρ ; Bottom: smoothed hinge penalty method with different ρ .

where $L_{GCL}(\mathbf{w}; (\mathbf{x}_i, \mathbf{t}_i), (\mathcal{T}_i^-, \mathcal{I}_i^-))$ is the two-way GCL for each image-text pair $(\mathbf{x}_i, \mathbf{t}_i)$,

$$L_{GCL}(\mathbf{w}; \mathbf{x}_i, t_i, \mathcal{T}_i^-, \mathcal{I}_i^-) := -\tau \log \frac{\exp(E_I(\mathbf{w}, \mathbf{x}_i)^\top E_T(t_i)/\tau)}{\sum_{t_j \in \mathcal{T}_i^-} \exp(E_I(\mathbf{x}_i)^\top E_T(t_j)/\tau)}$$
$$-\tau \log \frac{\exp(E_T(t_i)^\top E_I(\mathbf{x}_i)/\tau)}{\sum_{\mathbf{x}_j \in \mathcal{I}_i^-} \exp(E_T(t_j)^\top E_I(\mathbf{x}_i)/\tau)},$$

where $E_I(\mathbf{x})$ and $E_T(\mathbf{t})$ denote the normalized feature representation of an image \mathbf{x} and a text t, generated by visual encoder and text encoder of CLIP model, respectively. \mathcal{T}_i^- denotes the set of all texts to be contrasted with respect to (w.r.t) \mathbf{x}_i (including itself) and \mathcal{I}_i^- denotes the set of all images to be contrasted w.r.t t_i (including itself). Here, the data \mathcal{D} is a target dataset. $\mathcal{L}_k(\mathbf{w},\mathcal{D}_k) = \frac{1}{n_k} \sum_{(\mathbf{x}_i,y_i) \sim \mathcal{D}_k} \ell_k(\mathbf{w},\mathbf{x}_i,y_i)$ is the loss of the model \mathbf{w} on the k-th protected task, where ℓ_k is a logistic loss for the k-th classification task.

Hyperparameter tuning. The training data for the target tasks are sampled from the training set of BDD100K and LAION400M [53], while the data used to construct the MDS constraints are sampled exclusively from BDD100K. For all the method we use the same learning rate of 1e-6 and the same weight decay 0.1. For each experiment we run for 60 epochs * 400 iterations per epoch and the batch size is 256. We tune ρ in {0.01, 0.1, 1, 5, 10, 20, 40, 80} for SONEX+smooth Hinge and in {1, 20, 50, 100, 200, 400, 800} for SOX+squared hinge; we set $\gamma_1 = \gamma_2 = 0.8, \gamma_1' = \gamma_2' = 0.1, \tau = 0.05$ where γ_1, γ_1' is the hyperparameter in MSVR update for objective term (i.e. L_{GCL} ⁸) while γ_2, γ_2' is the hyperparameter in MSVR update for the penalty terms.

F.4 Ablation Study

We conduct a series of ablation studies to examine the effects of key hyperparameters and design choices.

Results for varying β We investigate the impact of different β on model performance. We evaluate $\beta \in \{0.1, 0.3, 0.5, 0.7, 1.0\}$ and conduct experiments for SONEX on Amazon dataset for GDRO task and ALEXR2 on Adult dataset for AUC maximization with ROC fairness constraints. The results are shown in table 3 which indicates the importance of setting $\beta \leq 1.0$.

Results for varying θ We analyze the influence of different θ of ALEXR2 on model performance, testing $\theta \in \{\frac{1}{16}, \frac{1}{8}, \frac{1}{4}, \frac{1}{2}, 1\}$ and conduct experiments for ALEXR2 on Adult dataset. As shown in Table 4, the final AUC of ALEXR2 remains relatively stable across this range, suggesting that it is not highly sensitive to θ .

⁸note that GCL is also a type of FCCO problem, we can also apply MSVR update to track the inner functions

Benefit of Adam-type update Finally we compare Adam-type update with momentum-type update. We conduct experiments for SONEX on Amazon dataset and ALEXR2 on Adult dataset and the results are summarized in table 5. They clearly demonstrate the advantage of using Adam-type updates, which consistently yield superior performance.

Table 3: Final loss and AUC of SONEX and ALEXR2 on Amazon dataset and Adult dataset respectively, with varying β

SONEX w/ varying β	0.1	0.3	0.5	1.0
Final loss	0.5657	0.5563	0.5768	0.578
	•			
ALEXR2 w/ varying β	0.1	0.3	0.5	1.0
Final AUC	0.8975	0.8976	0.8973	0.8969

Table 4: Final AUC of ALEXR2 with varying θ on Adult dataset

varying θ	0.1	0.3	0.5	1.0
Final AUC	0.8975	0.8976	0.8973	0.8969

Table 5: Comparison between Adam-type update and Momentum-type update for SONEX and ALEXR2 on Amazon dataset and Adult dataset, respectively

Algorithm	Adam-type	Momentum-type
SONEX(Final Loss)	0.5657	0.9666
ALEXR2(Final AUC)	0.8975	0.861

F.5 Verification of Assumption 4.3

We verify the assumption 4.3 by computing minimum eigenvalue λ_{min} of $\nabla \mathbf{g}(\mathbf{w}) \nabla \mathbf{g}(\mathbf{w})^{\top}$ in group DRO experiment. We compute λ_{min} for models trained on Camelyon17 and CelebA from the last epoch and report it in table 6. Our experiment results demonstrate that the minimum eigenvalue of $\nabla \mathbf{g}(\mathbf{w}) \nabla \mathbf{g}(\mathbf{w})^{\top}$ remains positive after training process finishes.

Besides, assumption 5.1 has been verified empirically in Appendix A.1 of [7].

F.6 Other Details of Experiments

Computing Resource and Running time: The experiments of AUC Maximization with ROC Fairness Constraints and the experiments of group DRO of Camelyon17 dataset and CelebA dataset in our paper is run on an A30 24G GPU, among which the first experiment takes less than 10 minutes for each run while for the second one, Camelyon17 takes about 4 hours and CelebA takes about 5 hours, for each run. The Amazon dataset of group DRO experiment is run on one A100 40GB GPUs and takes about 12 hours each run. The experiment of continual learning with non-forgetting constraints is run on two A100 40GB GPUs and takes about 12 hours each run.

Data Split: We perform data split of CelebA dataset ourselves: within each group, samples are divided into training, validation, and test sets in an 8:1:1 ratio. For all the other datasets mentioned in our paper we use default split.

Other Experiment results We show the training curves of individual constraint values for our two experiments about constraint optimization, as shown in Figure 4, 5 (continual learning with non-forgetting constraints) and Figure 6, 7 (AUC maximization with ROC fairness constraints). Note that we use other weather conditions except foggy to construct non-forgetting constraints since there is no foggy data in BDD100k for defining such a constraint.

Table 6: Minimum eigen values λ_{min} of $\nabla \mathbf{g}(\mathbf{w}) \nabla \mathbf{g}(\mathbf{w})^{\top}$ at different solution.

Camelyon17	seed 1	seed 2	seed 3
λ_{min}	0.0206	0.0349	0.0261

CelebA	seed 10	seed 20
λ_{min}	0.9000	0.2713




Figure 6: Training curves of 14 constraint values of different methods on adult dataset for AUC maximization with ROC fairness constraints. Top row: SOX with squared-hinge penalty method; Middle: SONX with Hinge penalty method; Bottom: ALEXR2 with smoothed hinge penalty method.




Figure 7: Training curves of 14 constraint values of different methods on COMPAS dataset for AUC maximization with ROC fairness constraints. Top row: SOX with squared-hinge penalty method; Middle: SONX with Hinge penalty method; Bottom: ALEXR2 with smoothed hinge penalty method.