
Identifiability of total effects from abstractions of time series causal graphs

Charles K. Assaad^{1,2} Emilie Devijver³ Eric Gaussier³ Gregor Gössler⁴ Anouar Meynaoui⁵

¹Sorbonne Université, INSERM, Institut Pierre Louis d’Epidémiologie et de Santé Publique, F75012, Paris, France

²EasyVista, F38000, Grenoble, France

³Univ Grenoble Alpes, CNRS, Grenoble INP, LIG, F38000, Grenoble, France

⁴Univ. Grenoble Alpes, INRIA, CNRS, Grenoble INP, LIG, F38000, Grenoble, France

⁵Université of Rennes 2, F35000, Rennes, France

Abstract

We study the problem of identifiability of the total effect of an intervention from observational time series in the situation, common in practice, where one only has access to abstractions of the true causal graph. We consider here two abstractions: the extended summary causal graph, which conflates all lagged causal relations but distinguishes between lagged and instantaneous relations, and the summary causal graph which does not give any indication about the lag between causal relations. We show that the total effect is always identifiable in extended summary causal graphs and provide sufficient conditions for identifiability in summary causal graphs. We furthermore provide adjustment sets allowing to estimate the total effect whenever it is identifiable.

1 INTRODUCTION

Over the last century and across numerous disciplines, experimentation has emerged as a potent methodology for estimating without bias the total effect of an intervention on a specific component of a given system [Neyman et al., 1990]. However, experimentation can be costly, unethical or even unfeasible. Both researchers and experts are thus interested in estimating the effect of an intervention directly from observational data. This can be done under some assumptions when relying on a complete causal graph [Pearl et al., 2000], and typically relies on two sequential steps: identifiability and estimation [Pearl, 2019]. The identifiability step involves distinguishing cases where a solution is possible and, when it exists, providing an estimand - an expression enabling the estimation of intervention effects from observational data. The subsequent step involves the actual estimation of this estimand from the available data.


The identifiability step received much attention for non-

temporal causal graphs [Pearl, 1993, 1995, Spirtes et al., 2000, Pearl et al., 2000, Shpitser and Pearl, 2008]. For abstraction of causal graphs, Perkovic [2020] derived necessary and sufficient conditions for identifying total effects in maximally oriented partially directed acyclic graphs and Anand et al. [2023] provided necessary and sufficient conditions when dealing with a directed acyclic graphs, where each vertex represent a cluster of variables and where relationships between clusters of variables are specified, but relationships between the variables within a cluster are left unspecified.


For temporal causal graph, Blondel et al. [2016] developed the do-calculus for the full-time causal graphs (FTCGs, Figure 1.1a). However, in dynamic systems, experts have difficulties in building full time causal graphs, while they can usually build an abstraction of those graphs such as an extended summary causal graph (ESCG, as in Figure 1.1b) where all lagged causal relations are conflated but lagged and instantaneous relations are clearly distinguished or such as a summary causal graph (SCG, as in Figure 1.1c) where all temporal information is omitted. Assuming no instantaneous relations, Eichler and Didelez [2007] demonstrated that the total effect is identifiable from an ESCG or an SCG, and Assaad et al. [2023] established identifiability in the presence of instantaneous relations for acyclic SCGs. Ferreira and Assaad [2024] addressed the identifiability problem for general SCGs, including cycles and instantaneous relations for the direct effect; however, the identifiability of total effects in this context remains unexplored.

Our main contributions consist in demonstrating, under causal sufficiency, that the total effect is always identifiable when working with an extended summary causal graph and providing sufficient conditions for identifying the total effect when working with a summary causal graph. The main difficulty lies in the fact that these abstractions may represent different full-time causal graphs with potentially different skeletons and orientations.


The remainder of the paper is structured as follows: Section



(a) Three FTCGs, \mathcal{G}_1^f , \mathcal{G}_2^f and \mathcal{G}_3^f .



(b) Three ESCGs, \mathcal{G}_1^e , \mathcal{G}_2^e and \mathcal{G}_3^e , resp. derived from \mathcal{G}_1^f , \mathcal{G}_2^f and \mathcal{G}_3^f .



(c) The SCG \mathcal{G}^s , derived from any FTCG in (a) and any ESCG in (b).

Figure 1.1: Illustration: (a) three FTCGs, (b) three ESCGs derived from them, (c) the SCG which can be derived from any FTCG in (a) and any ESCG in (b). Consider $f(y_t | do(x_{t-1}))$, red vertex: the variable we intervene on, blue vertex: the response we are considering. Bold edges correspond to directed paths from X_{t-1} to Y_t , and gray vertices correspond to nodes with different status depending on the FTCG (see Definition 6).

2 introduces the main notions, Section 3 presents the problem setup, identifiability conditions in ESCGs and SCGs are respectively presented in Sections 4 and 5. Section 6 discusses real applications for our theoretical results, and Section 7 concludes the paper. Omitted proofs can be found in the Supplementary Material.

2 PRELIMINARIES

Graph notions For a *directed acyclic graph* \mathcal{G} , a *path* from X to Y in \mathcal{G} is a sequence of distinct vertices $\langle X, \dots, Y \rangle$ in which every pair of successive vertices is adjacent. A *directed path* from X to Y is a path from X to Y in which all edges are directed towards Y in \mathcal{G} , that is $X \rightarrow \dots \rightarrow Y$. A *backdoor path* between X and Y is a path between X and Y with an arrowhead into X in \mathcal{G} . If $X \rightarrow Y$, then X is a *parent* of Y . If there is a directed path from X to Y , then X is an *ancestor* of Y , and Y is a *descendant* of X . A vertex counts as its own descendant and as its own ancestor. The sets of parents, ancestors and descendants of X in \mathcal{G} are denoted by $\text{Par}(X, \mathcal{G})$, $\text{Anc}(X, \mathcal{G})$ and $\text{Desc}(X, \mathcal{G})$ respectively. If a path π contains $X \rightarrow Z \leftarrow Y$ as a subpath, then Z is a *collider* on π . A vertex Z is a

definite non-collider on a path π if the edge $X \leftarrow Z$, or the edge $Z \rightarrow Y$ is on π . A vertex is of *definite status* on a path if it is a collider, a definite non-collider or an endpoint on the path. A path π is of *definite status* if every vertex on π is of definite status. A path π from X to Y of definite status is *active* given a vertex set \mathcal{Z} , with $X, Y \notin \mathcal{Z}$ if every definite non-collider on π is not in \mathcal{Z} , and every collider on π has a descendant in \mathcal{Z} . Otherwise, \mathcal{Z} *blocks* π . By a slight abuse of notation, we denote $\mathcal{G} \setminus \{Y\}$ as the subgraph of \mathcal{G} when removing the vertex Y and its corresponding edges. Lastly, the *skeleton* of a graph corresponds to all vertices and edges of the graph without considering edge orientations.

For a *directed graph* \mathcal{G} , a directed path from X to Y and a directed path from Y to X form a *directed cycle* in \mathcal{G} . A self-loop on X also forms a directed cycle. We denote by $\text{Cycles}(X, \mathcal{G})$ the set of all directed cycles containing X in \mathcal{G} , and by $\text{Cycles}^>(X, \mathcal{G})$ the subset of $\text{Cycles}(X, \mathcal{G})$ with at least two different vertices (i.e., excluding self-loops). In addition, all notions introduced before for directed acyclic graphs hold for *directed graphs*, with potential cycles. To avoid any ambiguity we would like to make some clarifications. For a *directed graph* \mathcal{G} , a *backdoor path* between X and Y is a path between X and Y which starts by either $X \leftarrow$ or $X \rightleftharpoons$. A path is blocked by an empty set if there exists a vertex Z such that $\rightarrow Z \leftarrow$ is on the path. Note that the above does not hold for $\rightleftharpoons Z \leftarrow$, $\rightarrow Z \rightleftharpoons$ or $\rightleftharpoons Z \rightleftharpoons$. For clarity, whenever a path is blocked by an empty set in a directed graph we will say that it is σ -blocked¹. Note that $X \rightleftharpoons Y$ and $X \leftarrow Y$ are the only σ -active backdoor paths of size 2 in \mathcal{G} .

If each vertex in a directed acyclic graph corresponds to an observed variable then, given an ordered pair of vertices (X, Y) in \mathcal{G} , a set of vertices \mathcal{Z} satisfies the *standard backdoor criterion* relative to (X, Y) if no vertex in \mathcal{Z} is a descendant of X , and \mathcal{Z} blocks every backdoor path between X and Y .

Causal graphs in time series Consider \mathcal{V} a set of p observational time series and $\mathcal{V}^f = \{\mathcal{V}_{t-\ell} | \ell \in \mathbb{Z}\}$ the set of temporal instances of \mathcal{V} where $\mathcal{V}_{t-\ell}$ correspond to the variables of the time series at time $t-\ell$. We suppose that the time series are generated from an *unknown* dynamic structural causal model (DSCM, Rubenstein et al. [2018]), an extension of structural causal models (SCM, Pearl et al. [2000]) to time series. This DSCM defines a full-time causal graph (FTCG, see below) which we call the *true* FTCG [Runge et al., 2019, Runge, 2021, Assaad et al., 2022a] and a joint distribution P over its vertices which we call the *true* probability distribution.

¹The notion of σ -blocked path by a set \mathcal{Z} is a generalization of the notion of blocked path by a set \mathcal{Z} (which was introduced for directed acyclic graphs) to directed graphs [Forre and Mooij, 2017]. These two notions becomes equivalent when $\mathcal{Z} = \emptyset$. In this paper, we will use the notion of σ -blocked only when $\mathcal{Z} = \emptyset$.

Definition 1 (Full-time causal graph (FTCG), Figure 1.1a). Let \mathcal{V} be a set of p observational time series and $\mathcal{V}^f = \{\mathcal{V}_{t-\ell}^e | \ell \in \mathbb{Z}\}$. The full-time causal graph (FTCG) $\mathcal{G}^f = (\mathcal{V}^f, \mathcal{E}^f)$ representing a given DSCM is defined by: $X_{t-\gamma} \rightarrow Y_t \in \mathcal{E}^f$ if and only if X directly causes Y at time t with a time lag of $\gamma > 0$ if $X = Y$ and with a time lag of $\gamma \geq 0$ for $X \neq Y$.

As common in causality studies on time series, we consider in the remainder acyclic FTCGs with potential instantaneous causal relations. Note that acyclicity is guaranteed for relations between variables at different time stamps. In addition, note that for any time series X , $\forall i > 0$, X_{t-i} can cause X_t ; for example, the stock price yesterday can affect the stock price today. We furthermore assume causal sufficiency:

Assumption 1 (Causal sufficiency). *There is no hidden common cause between any two observed variables.*

In practice, it is usually impossible to work with FTCGs and people have resorted to simpler causal graphs, exploiting the fact that causal relations between time series hold throughout time, as formalized in the following assumption which allows one to focus on a finite number of past slices, given by the maximum lag. We fix it to γ_{\max} in the remainder.

Assumption 2 (Consistency throughout time). *All the causal relationships in the the FTCTG \mathcal{G}^f remain constant in direction and magnitude throughout time².*

Experts are used to working with abstractions of causal graphs which summarize the information into a smaller graph that is interpretable, often with the omission of precise temporal information. We consider in this study two known causal abstractions for time series, namely *extended summary causal graphs* and *summary causal graphs*. An extended summary causal graph [Assaad et al., 2022c] distinguishes between past time slices, denoted as \mathcal{V}_{t-}^e , and present time slices, denoted as \mathcal{V}_t^e , thus enabling the differentiation between lagged and instantaneous causal relations.

Definition 2 (Extended summary causal graph (ESCG), Figure 1.1b). Let $\mathcal{G}^f = (\mathcal{V}^f, \mathcal{E}^f)$ be an FTCTG built from the set of time series \mathcal{V} satisfying Assumption 2 with maximal temporal lag γ_{\max} . The extended summary causal graph (ESCG) $\mathcal{G}^e = (\mathcal{V}^e, \mathcal{E}^e)$ associated to \mathcal{G}^f is given by $\mathcal{V}^e = (\mathcal{V}_{t-}^e, \mathcal{V}_t^e)$ and \mathcal{E}^e defined as follows:

- for any X in \mathcal{V} , we define two vertices, X_{t-} and X_t , respectively in \mathcal{V}_{t-}^e and \mathcal{V}_t^e ;

²In our context we consider a dynamic system with several univariate observational time series, thus the problem of finding a unique total effect would be ill-posed if Assumption 2 is not satisfied since violating the assumption would mean that the total effect would change over time.

- for all $X_t, Y_t \in \mathcal{V}_t^e$, $X_t \rightarrow Y_t \in \mathcal{E}^e$ if and only if $X_t \rightarrow Y_t \in \mathcal{E}^f$;
- for all $X, Y \in \mathcal{V}_{t-}^e$, $X_{t-} \rightarrow Y_t \in \mathcal{E}^e$ if and only if there exists at least one temporal lag $0 < \gamma \leq \gamma_{\max}$ such that $X_{t-\gamma} \rightarrow Y_t \in \mathcal{E}^f$.

In that case, we say that \mathcal{G}^e is derived from \mathcal{G}^f .

At a higher level of abstraction, a summary causal graph [Peters et al., 2013, Meng et al., 2020, Assaad et al., 2022a,b] represents causal relationships among time series, regardless of the time delay between the cause and its effect.

Definition 3 (Summary causal graph (SCG), Figure 1.1c). Let $\mathcal{G}^f = (\mathcal{V}^f, \mathcal{E}^f)$ be an FTCTG built from the set of time series \mathcal{V} satisfying Assumption 2 with maximal temporal lag γ_{\max} . The summary causal graph (SCG) $\mathcal{G}^s = (\mathcal{V}^s, \mathcal{E}^s)$ associated to \mathcal{G}^f is such that

- \mathcal{V}^s corresponds to the set of time series \mathcal{V} ,
- $X \rightarrow Y \in \mathcal{E}^s$ if and only if there exists at least one temporal lag $0 \leq \gamma \leq \gamma_{\max}$ such that $X_{t-\gamma} \rightarrow Y_t \in \mathcal{E}^f$.

In that case, we say that \mathcal{G}^s is derived from \mathcal{G}^f as well as from the ESCG derived from \mathcal{G}^f .

Since an FTCTG is assumed to be a directed acyclic graph, an ESCG is inherently a directed acyclic graph. In contrast, an SCG is a directed graph as it may include directed cycles and even self-loops. For example, the three FTCTGs in Figure 1.1a and the three ESCGs in Figure 1.1b are acyclic, while the SCG in Figure 1.1c has a cycle. We use the notation $X \rightleftarrows Y$ to indicate situations where there are time lags where X causes Y and other lags where Y causes X . Additionally, if an SCG is an abstraction of an ESCG, in cases where there is no instantaneous relation, ESCGs and SCGs convey the same information.

It is worth noting that if there is a single ESCG or SCG derived from a given FTCTG, different FTCTGs, with possibly different orientations and skeletons, can yield the same ESCG or SCG. For example, the SCG in Figure 1.1c can be derived from any FTCTG and any ESCG in Figures 1.1a and 1.1b, even though they may have different skeletons (for example, \mathcal{G}_1^f and \mathcal{G}_3^f or \mathcal{G}_1^e and \mathcal{G}_3^e) and different orientations (for example, \mathcal{G}_1^f and \mathcal{G}_2^f or \mathcal{G}_1^e and \mathcal{G}_2^e). Therefore, even if each vertex in an FTCTG is assumed to represent a single observed variable, a vertex in the past slice of an ESCG represent a set of variables while a vertex in the present time slice represents a single variable, and a vertex in the SCG corresponds to a time series. In the remainder, for a given ESCG or SCG \mathcal{G} , we call any FTCTG from which \mathcal{G} can be derived as a *candidate FTCTG* for \mathcal{G} . For example, in Figure 1.1, $\mathcal{G}_1^f, \mathcal{G}_2^f$ and \mathcal{G}_3^f are all candidate FTCTGs for \mathcal{G}^s . The set of all candidate FTCTGs for \mathcal{G} is denoted by $\mathcal{C}(\mathcal{G})$.

3 PROBLEM SETUP

We focus in this paper on the *total effect* [Pearl et al., 2000] of the *singleton* variable $X_{t-\gamma}$ on the *singleton* variable Y_t , written $P(Y_t = y_t | do(X_{t-\gamma} = x_{t-\gamma}))$ (as well as $P(y_t | do(x_{t-\gamma}))$ by a slight abuse of notation), when the only knowledge one has of the underlying DSCM consists in the ESCG or SCG derived from the unknown, true FTCCG. Y_t corresponds to the response and $do(X_{t-\gamma} = x_{t-\gamma})$ represents an intervention (as defined in Pearl et al. [2000] and Eichler and Didelez [2007, Assumption 2.3]) on the variable X at time $t - \gamma$, with $\gamma \geq 0$.

The above setting is very common in practice and entails that one neither knows the true FTCCG nor the true probability distribution. Furthermore, even if one has access to observed data, in practice such observations are finite, which prevents one from discovering the true FTCCG, and even from detecting it in the set of candidate FTCCGs, as no existing causal discovery method is guaranteed to yield the true FTCCG in the finite data setting [Ait-Bachir et al., 2023]. In the purely theoretical context of infinite data, discovering the true FTCCG is only possible with additional assumptions, beyond the scope of this study [Assaad et al., 2022b].

Each candidate FTCCG proposes a particular decomposition of the true joint probability distribution which is given by the standard recursive decomposition that characterizes Bayesian networks. Not all decompositions are however correct with respect to the true probability distribution P .

In general, a total effect $P(y_t | do(x_{t-\gamma}))$ is said to be identifiable from a graph if it can be uniquely computed with a do-free formula from the observed distribution [Pearl, 1995, Perkovic, 2020]. In our context, this means that the same do-free formula should hold in all candidate FTCCG so as to guarantee that it holds for the true one.

Definition 4 (Identifiability of total effects in ESCGs and SCGs). *In a given ESCG or SCG \mathcal{G} , $P(y_t | do(x_{t-\gamma}))$ is identifiable iff it can be rewritten with a do-free formula that is valid for any FTCCG in $\mathcal{C}(\mathcal{G})$.*

One way to rewrite $P(y_t | do(x_{t-\gamma}))$ with a do free-formula is by finding an adjustment set of variables for which:

$$P(y_t | do(x_{t-\gamma})) = \sum_{\mathbf{z}} P(y_t | x_{t-\gamma}, \mathbf{z}) P(\mathbf{z}). \quad (1)$$

Whenever a set of variables satisfy Equation (1), we call it a *valid adjustment set*. The standard backdoor criterion, introduced in Pearl [1995], allows one to obtain valid adjustment sets using the true FTCCG. We provide here another version of the backdoor criterion that allows us to find a valid adjustment set given all candidate FTCCGs without knowing which one is the true FTCCG.

Definition 5 (Backdoor criterion over all candidate FTCCGs). *Let $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ be an ESCG or SCG. A set of vertices \mathcal{Z}*

satisfies the backdoor criterion over all candidate FTCCGs relative to $(X_{t-\gamma}, Y_t)$ if

- (i) \mathcal{Z} blocks all backdoor paths between $X_{t-\gamma}$ and Y_t in any FTCCG in $\mathcal{C}(\mathcal{G})$,
- (ii) \mathcal{Z} does not contain any descendant of $X_{t-\gamma}$ in any FTCCG in $\mathcal{C}(\mathcal{G})$.

Note that when there is no backdoor path between $X_{t-\gamma}$ and Y_t in any FTCCG in $\mathcal{C}(\mathcal{G})$, $\mathcal{Z} = \emptyset$ satisfies the backdoor criterion over all candidate FTCCGs.

The backdoor criterion over all candidate FTCCGs is sound for the identification of the total effect $P(y_t | do(x_{t-\gamma}))$ in an ESCG or SCG, as stated in the following corollary that can be deduced from [Pearl, 1995, Theorem 1].

Corollary 1. *Let X and Y be distinct vertices in an ESCG or SCG \mathcal{G} of a DSCM with true (unknown) probability P . Under Assumptions 1 and 2 for \mathcal{G} , if there exists a set \mathcal{Z} satisfying the backdoor criterion over all possible FTCCGs relative to $(X_{t-\gamma}, Y_t)$, then the total effect of $X_{t-\gamma}$ on Y_t is identifiable in \mathcal{G} , and \mathcal{Z} is a valid adjustment set for the formulae given in Equation (1).*

However, enumerating all candidate FTCCGs is computationally expensive [Robinson, 1977], even when considering the constraints given by an ESCG or an SCG.

Formally, we address the following technical problem:

Problem 1. *Consider an ESCG or an SCG \mathcal{G} and the total effect $P(y_t | do(x_{t-\gamma}))$. We aim to find out conditions to identify $P(y_t | do(x_{t-\gamma}))$ when having access solely to an ESCG or an SCG without enumerating all candidate FTCCGs in $\mathcal{C}(\mathcal{G})$.*

Remarks

1. Our context is different from the one considered in Perkovic [2020] since the graphs we have to consider for a given ESCG or SCG, namely the candidate FTCCGs, may have different skeletons and may not all be compatible with the true underlying distribution. Furthermore, in ESCGs and SCGs, each vertex does not necessarily correspond to a single variable.
2. Our context is different from the one considered in Anand et al. [2023]. They consider cluster of variables, even for the response and the intervention variable, while we are interested in the total effect $P(y_t | do(x_{t-\gamma}))$ where the response variable and the intervention variable are singletons. Furthermore, we may have cycles in the SCGs, while they assume acyclic graphs.
3. The cycles that we consider in this work, namely in SCGs, do not hold the same conceptual meaning as the cycles considered in Bongers et al. [2021], as in our

case, cyclicity comes from the abstraction of an acyclic graph.

4 IDENTIFIABILITY IN ESCG

The total effect is always identifiable by adjustment in ESCGs, as stated in the following theorem.

Theorem 1. (*Identifiability in ESCG*) Consider an ESCG \mathcal{G}^e . Under Assumptions 1 and 2 for \mathcal{G}^e , the total effect $P(y_t|do(x_{t-\gamma}))$ is identifiable in \mathcal{G}^e for any $\gamma \geq 0$. Furthermore, the set

$$\mathcal{B}_\gamma = \{(Z_{t-\gamma-\ell})_{1 \leq \ell \leq \gamma_{\max}} | Z_{t-} \in \text{Par}(X_t, \mathcal{G}^e)\} \cup \{Z_{t-\gamma} | Z_t \in \text{Par}(X_t, \mathcal{G}^e)\},$$

is a valid adjustment set for $P(y_t|do(x_{t-\gamma}))$ for the formulae given in Equation (1).

If \mathcal{B}_γ is a valid adjustment set, it may still be very large. Additional adjustment sets, potentially smaller than \mathcal{B}_γ , can however be obtained in the densest candidate FTCTG, which is the candidate FTCTG which contains all potential edges and is thus maximal in the number of edges.

Proposition 1. Consider an ESCG \mathcal{G}^e and a maximal lag γ_{\max} and let $\gamma \geq 0$. Any adjustment set \mathcal{B}'_γ for the total effect $P(y_t|do(x_{t-\gamma}))$ that satisfies the standard backdoor criterion on the densest candidate FTCTG in $\mathcal{C}(\mathcal{G}^e)$ is a valid adjustment set for the total effect. In addition, \mathcal{B}_γ is a valid adjustment set with respect to the standard backdoor criterion on the densest candidate FTCTG.

Note however that smaller (in the number of variables) adjustment sets may exist in the true FTCTG when it is different from the densest candidate FTCTG.

5 IDENTIFIABILITY IN SCG

In this section, we start by presenting the main result of the paper which provides sufficient conditions for identifying the total effect only by using an SCG and providing an adjustment set that can be used whenever the sufficient conditions are satisfied. Then we provide another adjustment set that is more suitable in practice. Finally, we discuss several examples where the total is not identifiable using an SCG.

Note that we are only considering sufficient conditions because the backdoor criterion is not complete, meaning it does not provide all possible valid adjustment sets. Therefore, the backdoor criterion over all candidate FTCTGs is not necessarily complete.

5.1 MAIN RESULT: SUFFICIENT CONDITIONS FOR IDENTIFIABILITY

We provide sufficient conditions³ for the identifiability in SCG. Recall that $Cycles(X, \mathcal{G}^s)$ is the set of all directed cycles containing X in \mathcal{G}^s , and $Cycles^>(X, \mathcal{G}^s)$ is the subset where cycles contain at least 2 different vertices.

Theorem 2. (*Identifiability in SCG*) Consider an SCG $\mathcal{G}^s = (\mathcal{V}^s, \mathcal{E}^s)$ associated with a DSCM with true (unknown) probability distribution P . Under Assumptions 1 and 2, the total effect $P(y_t|do(x_{t-\gamma}))$, with $\gamma \geq 0$, is identifiable if $X \notin \text{Anc}(Y, \mathcal{G}^s)$ or $X \in \text{Anc}(Y, \mathcal{G}^s)$ and none of the following holds:

1. $\gamma \neq 0$ and $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) \neq \emptyset$, or
2. there exists a σ -active backdoor path

$$\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$$

from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq \text{Desc}(X, \mathcal{G}^s)$ and one of the following holds:

- (a) $n > 2$, i.e. $\langle V^2, \dots, V^{n-1} \rangle \neq \emptyset$, or
- (b) $n = 2$ and $\gamma \neq 1$, or
- (c) $n = 2$, $\gamma = 1$ and $Cycles(Y, \mathcal{G}^s \setminus \{X\}) \neq \emptyset$.

In the remainder, we prove the above theorem through Lemmas 5.1-5.3. To do so, for the total effect $P(y_t|do(x_{t-\gamma}))$, we consider the following set:

$$\mathcal{A}_\gamma = \{(Z_{t-\gamma-\ell})_{1 \leq \ell \leq \gamma_{\max}} | Z \in \text{Desc}(X; \mathcal{G}^s)\} \cup \{(Z_{t-\gamma-\ell})_{0 \leq \ell \leq \gamma_{\max}} | Z \in \mathcal{V}^s \setminus \text{Desc}(X, \mathcal{G}^s)\} \quad (2)$$

and we prove that it is a valid adjustment set when the total effect is identifiable. As one can note, it contains all possible parents of $X_{t-\gamma}$ in all candidate FTCTGs of \mathcal{G}^s . Thus, \mathcal{A}_γ blocks any backdoor path π between $X_{t-\gamma}$ and Y_t in any candidate FTCTG through the parent of $X_{t-\gamma}$ on that path.

We first introduce the notion of ambiguous vertices, represented in gray in every figure, that will be useful for the proofs of most of the lemmas.

Definition 6 (Ambiguous vertices). Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$, for $\gamma \geq 0$. A vertex $V_{t'}$ belonging to an active backdoor path for $(X_{t-\gamma}, Y_t)$ in a candidate FTCTG is ambiguous if there exists another candidate FTCTG in which $V_{t'}$ is a descendant of $X_{t-\gamma}$.

Ambiguous vertices are crucial for identifiability. In addition to ambiguous vertices, one can also define ambiguous paths, as follows.

³In Supplementary Material, we provide an equivalent version of Theorem 2 which might be easier to read to certain readers.




Figure 5.1: Three SCGs and a total effect which is identifiable. Each pair of red and blue vertices in the FTCGs represents the total effect we are interested in, and we precise the total effect and the lag considered in the caption. This illustrates Lemma 5.2 (Figure a-b) and Lemma 5.3 (Figure c).

Definition 7 (Ambiguous paths). Consider an SCG \mathcal{G}^s and a candidate FTCG \mathcal{G}^f . A path $\pi^f \in \mathcal{G}^f$ between $X_{t-\gamma}$ and Y_t , for $\gamma \geq 0$, is an ambiguous path if it does not contain any vertex at time $t - \gamma - \ell$ for $\ell \geq 1$. We note Π_γ^f the set of all ambiguous paths in \mathcal{G}^f .

When π is not an ambiguous path ($\pi \notin \Pi_\gamma^f$), then at least one vertex on π is in the past of $X_{t-\gamma}$ and thus cannot be ambiguous. One thus has the following property:

Property 1. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$, for $\gamma \geq 0$. Suppose π^f is a backdoor path between $X_{t-\gamma}$ and Y_t in a candidate FTCG \mathcal{G}^f . If $\pi^f \notin \Pi_\gamma^f$, then π^f is blocked by a subset of \mathcal{A}_γ containing at least one non-ambiguous vertex.

Example 5.1. For example, in Figure 5.5c, $\pi_1^f = \langle X_{t-1}, X_{t-2}, Y_{t-1}, Y_t \rangle$ is not an ambiguous path between X_{t-1} and Y_t since X_{t-2} precedes X_{t-1} in time. On the other hand, $\pi_2^f = \langle X_{t-1}, Y_{t-1}, Y_t \rangle$ is an ambiguous path between X_{t-1} and Y_t . The path π_1^f is blocked by X_{t-2} .

We now introduce the notion of compatible path that will allow us to relate backdoor paths in a given SCG and its candidate FTCGs.

Definition 8 (Compatible path). Consider an SCG \mathcal{G}^s , a candidate FTCG \mathcal{G}^f , and the total effect $P(y_t | do(x_{t-\gamma}))$, for $\gamma \geq 0$. We say that a path $\pi^f = \langle X_{t-\gamma}, W_{t_2}^2, \dots, W_{t_{m-1}}^{m-1}, Y_t \rangle$ in \mathcal{G}^f is compatible with a path $\pi^s = \langle X, V^2, \dots, V^{n-1}, Y \rangle$ in \mathcal{G}^s if for all $(W_{t_j}^j)_{2 \leq j \leq m-1}$: either $W^j \in \langle V^2, \dots, V^{n-1} \rangle$ or $\exists V \in \langle V^2, \dots, V^{n-1} \rangle$ such that $W^j \in Cycles(V, \mathcal{G}^s) \setminus Cycles(X, \mathcal{G}^s)$.

The following property relates backdoor paths in a given SCG and in any of its candidate FTCG.

Property 2. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$ for $\gamma \geq 0$. Then (i) \Rightarrow (ii), where:

- (i) $\gamma = 0$ or $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$,
- (ii) in any candidate FTCG \mathcal{G}^f , there exists no backdoor path $\pi^f \in \Pi_\gamma^f$ that is not compatible with any backdoor path in \mathcal{G}^s .

The two above properties allow one to prove the following lemmas which prove that each condition of Theorem 2 is sufficient. The first lemma is rather straightforward and concern the case where $X \notin Anc(Y, \mathcal{G}^s)$ for a given SCG \mathcal{G}^s .

Lemma 5.1. Consider an SCG \mathcal{G}^s , $\gamma \geq 0$ fixed and the total effect $P(y_t | do(x_{t-\gamma}))$. If $X \notin Anc(Y, \mathcal{G}^s)$ then $P(y_t | do(x_{t-\gamma}))$ is identifiable, and $P(y_t | do(x_{t-\gamma})) = P(y_t)$.

The following lemma excludes both Conditions 1 and 2 of Theorem 2 by considering the negation of Condition 1 (in (i)) and the situation in which there is no σ -active backdoor path from X to Y with $Z = \emptyset$.

Lemma 5.2. Consider an SCG \mathcal{G}^s , $\gamma \geq 0$ fixed and the total effect $P(y_t | do(x_{t-\gamma}))$. If $X \in Anc(Y, \mathcal{G}^s)$ and

- (i) either $\gamma = 0$ or $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and
- (ii) $\nexists \sigma$ -active backdoor path $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$,

then $P(y_t | do(x_{t-\gamma}))$ is identifiable by \mathcal{A}_γ .

This lemma is illustrated in Figure 5.1a - 5.1b.

When there is a σ -active backdoor path from X to Y with $Z = \emptyset$, the negation of Condition 2 of Theorem 2 is obtained with $n = 2$, $\gamma = 1$ and $Cycles(Y, \mathcal{G}^s \setminus \{X\}) = \emptyset$. The negation of Condition 1 of Theorem 2 is obtained in this setting with $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$. Note that, as before, having a σ -active backdoor path from X to Y with $Z = \emptyset$ and $n = 2$ is equivalent to $X \rightleftarrows Y$.

Lemma 5.3. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-1}))$ ($\gamma = 1$). If the only σ -active backdoor path from X to Y in \mathcal{G}^s with $Z = \emptyset$ is $X \rightleftarrows Y \in \mathcal{G}^s$ and

- (i) $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and
- (ii) $Cycles(Y, \mathcal{G}^s \setminus \{X\}) = \emptyset$,

then $P(y_t | do(x_{t-1}))$ is identifiable by \mathcal{A}_γ .

This lemma is illustrated in Figure 5.1c.




Figure 5.2: An example of an SCG \mathcal{G}_1^s in (a) satisfying Condition 1 in Theorem 2 and two candidate FTCGs in (b) and (c). Each pair of red and blue vertices in the FTCGs represents the total effect we are interested in. Gray vertices are ambiguous: they are on an active backdoor path in (b) and belong to a directed path in (c) (bold edges indicate direct paths from X_{t-1} to Y_t).

5.2 ADJUSTMENT SET

When the total effect is identifiable and when $X \in \text{Anc}(Y, \mathcal{G}^s)$, the set A_γ defined in Equation (2) is a valid adjustment set, but it has a large size, so we provide a smaller valid adjustment set, defined as follows:

$$\mathcal{A}'_\gamma = \{V_{t'} \in \mathcal{A}_\gamma \mid V \in \text{Anc}(X, \mathcal{G}^s) \cup \text{Anc}(Y, \mathcal{G}^s)\}.$$

Proposition 2. Consider an SCG \mathcal{G}^s and the total effect $P(y_t \mid \text{do}(x_{t-\gamma}))$, with $\gamma \geq 0$. Under conditions of identifiability provided by Theorem 2, the set \mathcal{A}'_γ is a valid adjustment set for the total effect.

5.3 NON IDENTIFIABLE EXAMPLES

In this section, we provide several examples of SCGs where the total effect cannot be identified by finding a valid adjustment set.

Example 5.2. Consider the SCG in Figure 5.2a and the two candidate FCTGs given in Figure 5.2b and 5.2c. Suppose we are interested in the total effect $P(y_t \mid \text{do}(x_{t-1}))$. In the first FCTG depicted in Figure 5.2b, the path $\langle X_{t-1}, Z_{t-1}, Y_t \rangle$ is an active back-door path. Since Z_{t-1} is the only vertex on this path that is not an endpoint, we need to adjust for it to eliminate the confounding bias induced by this path. However, in the second FCTG depicted in 5.2c, $\langle X_{t-1}, Z_{t-1}, Y_t \rangle$ forms a directed path. This implies that we should not adjust for Z_{t-1} to preserve the




Figure 5.3: An example of an SCG \mathcal{G}_2^s in (a) satisfying Condition 2a in Theorem 2 and two candidate FTCGs in (b) and (c). Each pair of red and blue vertices in the FTCGs represents the total effect we are interested in. Gray vertices are ambiguous: they are on an active backdoor path in (b) and belong to a directed path in (c) (bold edges indicate direct paths from X_{t-1} to Y_t).

influence of X_{t-1} on Y_t through the path passing by Z_{t-1} . Since we do not know which FTCG is the true one, then we cannot determine whether we should adjust for Z_{t-1} or not. Consequently, there is no valid adjustment set to identify the total effect $P(y_t \mid \text{do}(x_{t-1}))$.

Example 5.3. Consider the SCG in Figure 5.3a and the two candidate FTCGs in Figures 5.3b and 5.3c. Suppose we are interested in the total effect $P(y_t \mid \text{do}(x_{t-1}))$. The path $\langle X_{t-1}, Z_{t-1}, X_t, Y_t \rangle$ is an active back-door path in the first FTCG depicted in (b). Since Z_{t-1} is the only vertex on this path that is not an endpoint and that does not belong to a directed path in the same graph, we need to adjust for it to eliminate the confounding bias induced by this path. However, in the second FTCG depicted in 5.3c, $\langle X_{t-1}, Y_{t-1}, Z_{t-1}, X_t, Y_t \rangle$ forms a directed path. This implies that we should not adjust for Z_{t-1} to preserve the influence of X_{t-1} on Y_t through the path passing by Z_{t-1} . Since we do not know which FTCG is the true one, then we cannot determine whether we should adjust for Z_{t-1} or not. Consequently, there is no valid adjustment set to identify the total effect $P(y_t \mid \text{do}(x_{t-1}))$.

Example 5.4. Consider the SCG in Figure 5.4a and the two candidate FTCGs in Figures 5.4b and 5.4c. Suppose we are interested in the the total effect $P(y_t \mid \text{do}(x_{t-2}))$. The path $\langle X_{t-2}, Y_{t-2}, X_{t-1}, Y_t \rangle$ is an active back-door path in the first FTCG depicted in 5.4b. Since Y_{t-2} is the only vertex on this path that is not an endpoint and that does not belong to a directed path in the same graph, we need to adjust for it to eliminate the confounding bias induced




Figure 5.4: An example of an SCG \mathcal{G}_3^s in (a) satisfying Condition 2b in Theorem 2 with respect to $\Pr(y_t | do(x_{t-2}))$ and two candidate FT CGs in (b) and (c). Each pair of red and blue vertices in the FT CGs represents the total effect we are interested in. Gray vertices are ambiguous: they constitute a backdoor path in (b) and belong to a directed path in (c) (bold edges indicate direct paths from X_{t-2} to Y_t).

by this path. However, in the second FT CG depicted in 5.4c, $\langle X_{t-2}, Y_{t-2}, X_{t-1}, Y_t \rangle$ forms a directed path. This implies that we should not adjust for Y_{t-2} to preserve the influence of X_{t-2} on Y_t through the path passing by Y_{t-2} . Since we do not know which FT CG is the true one, then we cannot determine whether we should adjust for Y_{t-2} or not. Consequently, there is no valid adjustment set to identify the total effect $P(y_t | do(x_{t-2}))$.

Example 5.5. Consider the SCG in Figure 5.5a and the two candidate FT CGs in Figures 5.5b and 5.5c. Suppose we are interested in the total effect $P(y_t | do(x_{t-1}))$. The path $\langle X_{t-1}, Y_{t-1}, Y_t \rangle$ is an active back-door path in the first FT CG depicted in 5.5b. Since Y_{t-1} is the only vertex on this path that is not an endpoint, we need to adjust for it to eliminate the confounding bias induced by this path. However, in the second FT CG depicted in 5.5c, $\langle X_{t-1}, Y_{t-1}, Y_t \rangle$ forms a directed path. This implies that we should not adjust for Y_{t-1} to preserve the influence of X_{t-1} on Y_t through the path passing by Y_{t-1} . Since we do not know which FT CG is the true one, then we cannot determine whether we should adjust for Y_{t-1} or not. Consequently, there is no valid adjustment set to identify the total effect $P(y_t | do(x_{t-1}))$.

Notice that in Figure 5.5, removing the self-loop on Y makes the total effect identifiable. This is because the active backdoor path and the directed path discussed in Example 5.5 would no longer exist, leaving only directed paths or blocked (due to a collider) backdoor paths between X_{t-1} and Y_t .

6 DISCUSSION ON REAL-WORLD APPLICATIONS

Nephrology. Hypertension has long been considered as a risk factor for kidney function decline. At the same time, the kidney is known to have a major role in affecting blood pressure through sodium extraction and regulating electrolyte




Figure 5.5: An example of an SCG \mathcal{G}_4^s in (a) satisfying Condition 2c in Theorem 2 with respect to $\Pr(y_t | do(x_{t-1}))$ and two candidate FT CGs in (b) and (c). Each pair of red and blue vertices in the FT CGs represents the total effect we are interested in. Gray vertices are ambiguous: they constitute a backdoor path in (b) and belong to a directed path in (c) (bold edges indicate direct paths from X_{t-1} to Y_t).

balance [Yu et al., 2020]. This can be represented with the SCG in Figure 6.1a where the kidney function is represented by the creatinine level. Epidemiologists are interested to know if preventing kidney function decline can reduce the public health burden of hypertension and at the same time nephrologists are interested in knowing how much a treatment related to hypertension can improve the state of the kidney. Using Theorem 2 and assuming no hidden confounding, we can identify the total effect in each direction with a lag equal to 1 (if there are confounders that do not form additional cycles, the total effect remains identifiable if we measure them and take them into account in the SCG). We can collect data for estimation by conducting weekly blood tests on patients with kidney insufficiency, especially those whose hypertension and creatinine levels fluctuate.

Finance. It has been suggested that there exists a bidirectional causal relationship between the number of unique active wallets associated with bridge protocols and the mean transaction fees within the Ethereum network [Ante and Saggi, 2024]. Additionally, we consider that transaction fees causes itself over time, as depicted in the SCG shown in Figure 6.1b. In this scenario, the total effect of mean transaction fees on the number of unique active wallets is identifiable using Theorem 2 with a lag of 1. However, the same does not hold true for the opposite direction: the total effect of the number of unique active wallets on the mean transaction fees is not identifiable using Theorem 2.

System monitoring. Consider a subgraph of the SCG described in [Bystrova et al., 2024], representing the web activity in an IT system. Suppose that system experts observed a high number of queries at midnight for several weeks, likely due to a Distributed Denial of Service attack. Simultaneously, they noticed that CPU usage at midnight was very high, preventing the system from running some processes. Therefore, the system experts would like to determine (before intervening in the system) how much a reduction in

bandwidth in the network would reduce the global CPU usage. Theorem 2 shows that the total effect of Network input on CPU Global is identifiable for any lag. In addition, Theorem 2 implies that the total effect between all pairs of variables is identifiable since in the SCG there exists no cycles of size greater than 2. We can estimate those total effect using the data introduced in Bystrova et al. [2024].

Thermoregulation. Inspired by the experiment conducted in Peters et al. [2013], we consider maintaining a steady temperature in an apartment composed of four rooms: a living room, a kitchen, a bathroom, and an office. The living room is the only room containing a radiator, and all rooms are connected to each other through the living room. Additionally, all rooms contain a window except for the office. Temperature sensors were placed in the four rooms, plus one outside the apartment, and temperatures were recorded on an hourly basis. We consider the SCG presented in Figure 6.1d as the true one. Clearly, the outside temperature directly influences all rooms containing a window and the temperature in each room cannot cause the outside temperature. Since the living room contains a radiator, it can affect the temperatures in all other rooms. Additionally, since we may use fire in the kitchen for cooking, which can increase the temperature, we consider that the temperature in the kitchen can affect the temperature in the living room. Similarly, since we may use hot water in the bathroom, which can increase the temperature, we consider that the temperature in the bathroom can influence the temperature in the living room. All other vertices representing rooms in the graph are not connected to each other because they are not physically directly connected; they are all connected through the living room. Suppose we are specifically interested in estimating the total effect of the temperature in living room on the temperature in the office. Theorem 2 states that this total effect is identifiable for any lag since $Cycles(Living\ Room, \mathcal{G} \setminus \{Office\}) = \emptyset$ and there exists no σ -active backdoor path between Living Room and Office.

7 CONCLUSION

We studied in this paper the identification of total effects between singleton variables, under causal sufficiency, for both extended summary causal graphs and summary causal graphs. We showed that the total effect is always identifiable for extended summary causal graphs. The same does not hold for summary causal graphs for which we established graphical conditions which are sufficient, in any underlying probability distribution, for the identifiability of the total effect. In addition, in case of identifiability, we provided several valid adjustment sets for estimating the total effect in extended summary causal graphs, and two adjustment sets when considering summary causal graphs.

These results have significant implications, such as impact analysis in dynamic systems, particularly in scenarios where




Figure 6.1: Real summary causal graphs from (a) Nephrology, (b) Finance, (c) System Monitoring, and (d) Thermoregulation. Each pair of red and blue vertices represents the total effect(s) of interest where the red vertex represents the intervention and the blue vertex represents the response. According to Theorem 2, each of these total effects is either identifiable in general or identifiable under certain conditions on γ .

experts are unable to provide either a full temporal causal graph or an extended summary causal graph. They are also valuable in cases where the assumptions underlying causal discovery methods for inferring causal graphs with time lags are deemed overly restrictive. Furthermore, these results offer insights that can be useful in different disciplines such as Nephrology, Finance, System Monitoring, and Thermoregulation.

For future works, it would be valuable to establish necessary and sufficient conditions for the identifiability of total effects using SCGs, to extend this work to the case where the responses and interventions can be multivariate, and to the case where there are hidden confounding.

Acknowledgements

We thank Ali Aït-Bachir from EasyVista for discussions about the application of this work in system monitoring. We thank Benjamin Glemain and Nathanael Lapidus from IPLESP and Paolo Malvezzi from CHU Grenoble for discussions about the application of this work in Nephrology/Epidemiology. Finally, we thank Clément Yvernes from LIG for several discussions and five anonymous reviewers for their many insightful comments and suggestions. This work was partially supported by MIAI@Grenoble Alpes (ANR-19-P3IA-0003), by the CIPHOD project (ANR-23-CPJ1-0212-01), and by the CSPR R&D Booster Auvergne-Rhône-Alpes project.

References

- Tara V. Anand, Adele H. Ribeiro, Jin Tian, and Elias Bareinboim. Causal effect identification in cluster dags. *Proceedings of the AAAI Conference on Artificial Intelligence*, 37(10):12172–12179, Jun. 2023.
- Lennart Ante and Aman Saggiu. Time-varying bidirectional causal relationships between transaction fees and economic activity of subsystems utilizing the ethereum blockchain network. *Journal of Risk and Financial Management*, 17(1), 2024. ISSN 1911-8074. doi: 10.3390/jrfm17010019.
- Charles K. Assaad, Emilie Devijver, and Eric Gaussier. Entropy-based discovery of summary causal graphs in time series. *Entropy*, 24(8), 2022a. ISSN 1099-4300. doi: 10.3390/e24081156.
- Charles K. Assaad, Emilie Devijver, and Eric Gaussier. Survey and evaluation of causal discovery methods for time series. *J. Artif. Int. Res.*, 73, may 2022b. ISSN 1076-9757. doi: 10.1613/jair.1.13428.
- Charles K. Assaad, Emilie Devijver, and Eric Gaussier. Discovery of extended summary graphs in time series. In James Cussens and Kun Zhang, editors, *Proceedings of*

the Thirty-Eighth Conference on Uncertainty in Artificial Intelligence, volume 180 of *Proceedings of Machine Learning Research*, pages 96–106. PMLR, 01–05 Aug 2022c.

Charles K. Assaad, Imad Ez-Zejjari, and Lei Zan. Root cause identification for collective anomalies in time series given an acyclic summary causal graph with loops. In Francisco Ruiz, Jennifer Dy, and Jan-Willem van de Meent, editors, *Proceedings of The 26th International Conference on Artificial Intelligence and Statistics*, volume 206 of *Proceedings of Machine Learning Research*, pages 8395–8404. PMLR, 25–27 Apr 2023.

Ali Aït-Bachir, Charles K. Assaad, Christophe de Bignoncourt, Emilie Devijver, Simon Ferreira, Eric Gaussier, Hosein Mohanna, and Lei Zan. Case studies of causal discovery from it monitoring time series. The History and Development of Search Methods for Causal Structure Workshop at the 39th Conference on Uncertainty in Artificial Intelligence, 2023.

Gilles Blondel, Marta Arias, and Ricard Gavaldà. Identifiability and transportability in dynamic causal networks. *International Journal of Data Science and Analytics*, 3: 131–147, 2016.

Stephan Bongers, Patrick Forré, Jonas Peters, and Joris M. Mooij. Foundations of structural causal models with cycles and latent variables. *The Annals of Statistics*, 49(5):2885 – 2915, 2021. doi: 10.1214/21-AOS2064.

Daria Bystrova, Charles K. Assaad, Julyan Arbel, Emilie Devijver, Eric Gaussier, and Wilfried Thuiller. Causal discovery from time series with hybrids of constraint-based and noise-based algorithms. *Transactions on Machine Learning Research*, 2024.

Michael Eichler and Vanessa Didelez. Causal reasoning in graphical time series models. In *Proceedings of the Twenty-Third Conference on Uncertainty in Artificial Intelligence*, UAI’07, page 109–116, Arlington, Virginia, USA, 2007. AUAI Press. ISBN 0974903930.

Simon Ferreira and Charles K. Assaad. Identifiability of direct effects from summary causal graphs. *Proceedings of the AAAI Conference on Artificial Intelligence*, 38, 2024.

Patrick Forré and Joris M. Mooij. Markov properties for graphical models with cycles and latent variables, 2017.

Yuan Meng, Shenglin Zhang, Yongqian Sun, Ruru Zhang, Zhilong Hu, Yiyin Zhang, Chenyang Jia, Zhaogang Wang, and Dan Pei. Localizing failure root causes in a microservice through causality inference. In *2020 IEEE/ACM 28th International Symposium on Quality of Service (IWQoS)*, pages 1–10, 2020. doi: 10.1109/IWQoS49365.2020.9213058.

Jerzy Neyman, D. M. Dabrowska, and T. P. Speed. On the Application of Probability Theory to Agricultural Experiments. Essay on Principles. Section 9. *Statistical Science*, 5(4):465 – 472, 1990. doi: 10.1214/ss/1177012031.

Judea Pearl. [Bayesian Analysis in Expert Systems]: Comment: Graphical Models, Causality and Intervention. *Statistical Science*, 8(3):266 – 269, 1993. doi: 10.1214/ss/1177010894.

Judea Pearl. Causal diagrams for empirical research. *Biometrika*, 82(4):669–688, 1995.

Judea Pearl. The seven tools of causal inference, with reflections on machine learning. *Commun. ACM*, 62(3):54–60, feb 2019. ISSN 0001-0782. doi: 10.1145/3241036.

Judea Pearl et al. Models, reasoning and inference. *Cambridge, UK: CambridgeUniversityPress*, 19:2, 2000.

Emilija Perkovic. Identifying causal effects in maximally oriented partially directed acyclic graphs. In Jonas Peters and David Sonntag, editors, *Proceedings of the 36th Conference on Uncertainty in Artificial Intelligence (UAI)*, volume 124 of *Proceedings of Machine Learning Research*, pages 530–539. PMLR, 03–06 Aug 2020.

Jonas Peters, D. Janzing, and B. Schölkopf. Causal inference on time series using restricted structural equation models. In *Advances in Neural Information Processing Systems 26*, pages 154–162, 2013.

R. W. Robinson. Counting unlabeled acyclic digraphs. In Charles H. C. Little, editor, *Combinatorial Mathematics V*, pages 28–43, Berlin, Heidelberg, 1977. Springer Berlin Heidelberg. ISBN 978-3-540-37020-8.

P. K. Rubenstein, S. Bongers, B. Schölkopf, and J. M. Mooij. From deterministic ODEs to dynamic structural causal models. In *Proceedings of the 34th Conference on Uncertainty in Artificial Intelligence (UAI)*, pages 114–123, August 2018.

Jakob Runge. Necessary and sufficient graphical conditions for optimal adjustment sets in causal graphical models with hidden variables. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 15762–15773. Curran Associates, Inc., 2021.

Jakob Runge, Peer Nowack, Marlene Kretschmer, Seth Flaxman, and Dino Sejdinovic. Detecting and quantifying causal associations in large nonlinear time series datasets. *Science Advances*, 5(11):eaau4996, 2019. doi: 10.1126/sciadv.aau4996.

Ilya Shpitser and Judea Pearl. Complete identification methods for the causal hierarchy. *Journal of Machine Learning Research*, 9:1941–1979, 2008.

Peter Spirtes, Clark N Glymour, Richard Scheines, and David Heckerman. *Causation, prediction, and search*. MIT press, 2000.

Zhi Yu, Josef Coresh, Guanghao Qi, Morgan Grams, Eric Boerwinkle, Harold Snieder, Alexander Teumer, Cristian Pattaro, Anna Köttgen, Nilanjan Chatterjee, and Adrienne Tin. A bidirectional mendelian randomization study supports causal effects of kidney function on blood pressure. *Kidney International*, 98(3):708–716, 2020. ISSN 0085-2538. doi: <https://doi.org/10.1016/j.kint.2020.04.044>.

A SUPPLEMENTARY MATERIAL

A.1 PROOFS OF SECTION 4

Theorem 1. (*Identifiability in ESCG*) Consider an ESCG \mathcal{G}^e . Under Assumptions 1 and 2 for \mathcal{G}^e , the total effect $P(y_t | do(x_{t-\gamma}))$ is identifiable in \mathcal{G}^e for any $\gamma \geq 0$. Furthermore, the set

$$\mathcal{B}_\gamma = \{(Z_{t-\gamma-\ell})_{1 \leq \ell \leq \gamma_{\max}} | Z_{t-} \in \text{Par}(X_t, \mathcal{G}^e)\} \cup \{Z_{t-\gamma} | Z_t \in \text{Par}(X_t, \mathcal{G}^e)\},$$

is a valid adjustment set for $P(y_t | do(x_{t-\gamma}))$ for the formulae given in Equation (1).

Proof. If $X \notin \text{Anc}(Y, \mathcal{G}^e)$, then in every candidate FTCCG \mathcal{G}^f , $X_{t-\gamma} \notin \text{Anc}(Y_t, \mathcal{G}^f)$. Thus, $P(y_t | do(x_{t-\gamma}))$ is always identifiable in \mathcal{G}^e , and $P(y_t | do(x_{t-\gamma})) = P(y_t)$.

Assume now that $X \in \text{Anc}(Y, \mathcal{G}^e)$. Let γ_{\max} be the maximal lag, and \mathcal{G}^f be a candidate FTCCG. We prove that

$$\mathcal{B}_\gamma = \{(Z_{t-\gamma-\ell})_{1 \leq \ell \leq \gamma_{\max}} | Z_{t-} \in \text{Par}(X_t, \mathcal{G}^e)\} \cup \{Z_{t-\gamma} | Z_t \in \text{Par}(X_t, \mathcal{G}^e)\}$$

is an adjustment set for $P(y_t | do(x_{t-\gamma}))$ in \mathcal{G}^f .

First, we have to prove that $\text{Par}(X_{t-\gamma}, \mathcal{G}^f) \subseteq \mathcal{B}_\gamma$. Let $Z_{t-\gamma-\ell} \in \text{Par}(X_{t-\gamma}, \mathcal{G}^f)$. If $\ell = 0$, then Z_t causes X_t in \mathcal{G}^e by consistency throughout time, which means that $Z_{t-\gamma} \in \mathcal{B}_\gamma$. If $\ell > 0$, then Z_{t-} causes X_t in \mathcal{G}^e , that is $Z_{t-\ell-\gamma} \in \mathcal{B}_\gamma$. This shows that the set \mathcal{B}_γ blocks all backdoor paths relatively to $P(y_t | do(x_{t-\gamma}))$.

Then, we have to prove \mathcal{B}_γ does not contain any descendant of $X_{t-\gamma}$ in \mathcal{G}^f . If this is true, there exists $W_{t-\gamma} \in \mathcal{B}_\gamma \cap \text{Desc}(X_{t-\gamma}, \mathcal{G}^f)$, at time slice $t - \gamma$ because it is a parent and a descendant of $X_{t-\gamma}$. By consistency throughout time, $W_t \in \text{Desc}(X_t, \mathcal{G}^f)$. However, by definition of \mathcal{B}_γ , $X_t \in \text{Desc}(W_t, \mathcal{G}^f)$, which contradicts the acyclicity assumption of \mathcal{G}^e . It means that neither it blocks any directed path between $X_{t-\gamma}$ and Y_t , nor it contains any descendant of Y_t . \square

Proposition 1. Consider an ESCG \mathcal{G}^e and a maximal lag γ_{\max} and let $\gamma \geq 0$. Any adjustment set \mathcal{B}'_γ for the total effect $P(y_t | do(x_{t-\gamma}))$ that satisfies the standard backdoor criterion on the densest candidate FTCTG in $\mathcal{C}(\mathcal{G}^e)$ is a valid adjustment set for the total effect. In addition, \mathcal{B}_γ is a valid adjustment set with respect to the standard backdoor criterion on the densest candidate FTCTG.

Proof. Let \mathcal{G}_d^f be the densest candidate FTCTG, and \mathcal{B}'_γ be an adjustment set in \mathcal{G}_d^f . Let \mathcal{G}^f be another candidate FTCTG. By definition of \mathcal{G}_d^f , any back-door path in \mathcal{G}^f is also in \mathcal{G}_d^f (the last graph contains all possible edges). Then, \mathcal{B}'_γ blocks all back-door paths in \mathcal{G}^f . Moreover, since no vertex in \mathcal{B}'_γ is a descendant of $X_{t-\gamma}$ in \mathcal{G}_d^f , the same holds for \mathcal{G}^f . Thus, \mathcal{B}'_γ is also an adjustment set in \mathcal{G}^f . \square

A.2 PROOFS OF SECTION 5

Property 1. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$, for $\gamma \geq 0$. Suppose π^f is a backdoor path between $X_{t-\gamma}$ and Y_t in a candidate FTCTG \mathcal{G}^f . If $\pi^f \notin \Pi_\gamma^f$, then π^f is blocked by a subset of \mathcal{A}_γ containing at least one non-ambiguous vertex.

Proof. Suppose π^f is path between $X_{t-\gamma}$ and Y_t for $\gamma \geq 0$. If $\pi^f \notin \Pi_\gamma^f$ and then π^f contains at least one vertex $Z_{t-\gamma-\ell}$ for $\ell \geq 1$. $Z_{t-\gamma-\ell}$ is temporally prior to $X_{t-\gamma}$ which means π^f that if π^f is a backdoor path then adjusting on $Z_{t-\gamma-\ell}$ and the parents of $Z_{t-\gamma-\ell}$ on the path will block the path. Furthermore, for the same reason, there cannot be a directed path from $X_{t-\gamma}$ to $Z_{t-\gamma-\ell}$ in any FTCTG. Finally, again for the same reason, $Z_{t-\gamma-\ell}$ and the parents of $Z_{t-\gamma-\ell}$ are in \mathcal{A}_γ . \square

Property 2. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$ for $\gamma \geq 0$. Then (i) \Rightarrow (ii), where:

- (i) $\gamma = 0$ or $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$,
- (ii) in any candidate FTCTG \mathcal{G}^f , there exists no backdoor path $\pi^f \in \Pi_\gamma^f$ that is not compatible with any backdoor path in \mathcal{G}^s .

Proof. Assume first $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$. Suppose $\exists \pi^f = X_{t-\gamma} \leftarrow W_{t-\gamma} \cdots \rightarrow Y_t \in \Pi_\gamma^f$ which is a backdoor path between $X_{t-\gamma}$ and Y_t that is not compatible with any back-door path $\pi^s = \langle V^1 = X, V^2, \dots, V^{n-1}, V^n = Y \rangle$ in \mathcal{G}^s .

If $n = 2$, then the path compatible with the cycle $\langle X, X \rangle$ is of the form $X_{t-\gamma} \rightarrow X_{t-\gamma+i} \rightarrow \cdots \rightarrow X_{t-\gamma+j} \rightarrow Y_t$: it means that π^f cannot be a back-door path.

If $n > 2$, $W_{t-\gamma}$ is such that $W \notin \{V^2, \dots, V^{n-1}\}$ and $\nexists V \in \{V^2, \dots, V^{n-1}\}$ such that $W \in Cycles(V, \mathcal{G}^s)$. If the path between $W_{t-\gamma}$ and Y_t does not pass by $X_{t-\gamma+\ell}$

with $\ell > 0$, then there exists a back-door path between X and Y passing by W in \mathcal{G}^s as π^f lies in a candidate FTCTG, which contradicts our assumption. So the path necessarily passes by $X_{t-\gamma+\ell}$. Thus there is a cycle C_x on X such that $size(C_x) > 2$, which leads again to a contradiction. Thus, there does not exist a back-door path $\pi^f \in \Pi_\gamma^f$ between $X_{t-\gamma}$ and Y_t that is not compatible with any back-door path in \mathcal{G}^s .

The case $\gamma = 0$ is treated in the same way, with the fact that the path considered cannot go back to X_t as this would create a cycle in the FTCTG. \square

Lemma 5.1. Consider an SCG \mathcal{G}^s , $\gamma \geq 0$ fixed and the total effect $P(y_t | do(x_{t-\gamma}))$. If $X \notin Anc(Y, \mathcal{G}^s)$ then $P(y_t | do(x_{t-\gamma}))$ is identifiable, and $P(y_t | do(x_{t-\gamma})) = P(y_t)$.

Proof. If $X \notin Anc(Y, \mathcal{G}^s)$, then in every candidate FTCTG \mathcal{G}^f , $X_{t-\gamma} \notin Anc(Y_t, \mathcal{G}^f)$. Thus, $P(y_t | do(x_{t-\gamma}))$ is always identifiable in \mathcal{G}^s , and $P(y_t | do(x_{t-\gamma})) = P(y_t)$. \square

Lemma 5.2. Consider an SCG \mathcal{G}^s , $\gamma \geq 0$ fixed and the total effect $P(y_t | do(x_{t-\gamma}))$. If $X \in Anc(Y, \mathcal{G}^s)$ and

- (i) either $\gamma = 0$ or $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and
- (ii) $\nexists \sigma$ -active backdoor path $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$,

then $P(y_t | do(x_{t-\gamma}))$ is identifiable by \mathcal{A}_γ .

Proof. We will prove that \mathcal{A}_γ is an adjustment set for $P(y_t | do(x_{t-\gamma}))$ in any candidate FTCTG under conditions (i) and (ii). Let \mathcal{G}^f be a candidate FTCTG, and Π_γ^f the set of ambiguous paths. By Property 1, any back-door path $\pi^f \notin \Pi_\gamma^f$ can be blocked by \mathcal{A}_γ . Furthermore, by definition, elements of \mathcal{A}_γ cannot be descendant of $X_{t-\gamma}$.

We now turn our attention to paths in Π_γ^f . Let $\pi^f \in \Pi_\gamma^f$ be a back-door path between $X_{t-\gamma}$ and Y_t . Since $\gamma = 0$ or $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ then by Property 2, all back-door paths in Π_γ^f are compatible with back-door paths in \mathcal{G}^s . Let $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ be a σ -active back-door path in \mathcal{G}^s compatible with π^f . By (ii), there exists $m \geq 1$ vertices such that $\{V^{i_1}, \dots, V^{i_m}\} \subseteq \langle V^2, \dots, V^{n-1} \rangle$ and $\{V^{i_1}, \dots, V^{i_m}\} \not\subseteq Desc(X, \mathcal{G}^s)$. Then, $\forall V_{t-\gamma}$ such that $V \in \{V^{i_1}, \dots, V^{i_m}\}$, $V_{t-\gamma} \notin Desc(X_{t-\gamma}, \mathcal{G}^f)$ and since $X \in Anc(Y, \mathcal{G}^s)$ then it must be the case that $V \notin Desc(Y, \mathcal{G}^s)$ and by consequence $V_{t-\gamma} \notin Desc(Y_t, \mathcal{G}^f)$. Thus, $V_{t-\gamma}$ cannot be an ambiguous vertex. Its parent in π^f furthermore blocks π^f , is not ambiguous (as otherwise $V_{t-\gamma}$ would be ambiguous) and is a member of \mathcal{A}_γ by definition of \mathcal{A}_γ . Thus \mathcal{A}_γ blocks all back-door paths between $X_{t-\gamma}$ and Y_t in any candidate FTCTG \mathcal{G}^f . Furthermore, no node in \mathcal{A}_γ can block a directed path between $X_{t-\gamma}$ and Y_t or is a descendant of Y_t as nodes in \mathcal{A}_γ are either defined before

$t - \gamma$ or are not descendant of $X_{t-\gamma}$, and thus of Y_t . This concludes the proof. \square

Lemma 5.3. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-1}))$ ($\gamma = 1$). If the only σ -active backdoor path from X to Y in \mathcal{G}^s with $\mathcal{Z} = \emptyset$ is $X \leftrightarrow Y \in \mathcal{G}^s$ and

- (i) $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and
- (ii) $Cycles(Y, \mathcal{G}^s \setminus \{X\}) = \emptyset$,

then $P(y_t | do(x_{t-1}))$ is identifiable by \mathcal{A}_γ .

Proof. We will prove that \mathcal{A}_1 is an adjustment set for $P(y_t | do(x_{t-1}))$ in any candidate FTCCG under conditions (i) and (ii). Let \mathcal{G}^f be a candidate FTCCG, and Π_1^f the set of ambiguous paths.

Since (ii) then by Property 2, all back-door paths in Π_1^f are compatible with back-door paths in \mathcal{G}^s . In addition, by Property 1, any path $\pi^f \notin \Pi_1^f$ can be blocked by \mathcal{A}_1 . Therefore, in the following, we focus on paths in Π_1^f compatible with back-door paths in \mathcal{G}^s .

Consider the σ -active back-door path $\pi^s = \langle X, Y \rangle$. As there cannot be a loop on Y by (i), the only path $\pi^f \in \Pi_1^f$ from X_{t-1} to Y_t compatible with π^s that pass by Y_{t-1} is $\pi^f = \langle X_{t-1}, Y_{t-1}, X_t, Y_t \rangle$. Then, under consistency throughout time, acyclicity and temporal priority, the only choices are $X_{t-1} \rightarrow Y_{t-1} \rightarrow X_t \rightarrow Y_t$ and $X_{t-1} \leftarrow Y_{t-1} \rightarrow X_t \leftarrow Y_t$. The first is a directed path, the second a back-door path already blocked due to the collider $Y_{t-1} \rightarrow X_t \leftarrow Y_t$. Thus, all potential back-door paths between X_{t-1} and Y_t in any candidate \mathcal{G}^f are blocked, and \mathcal{A}_1 does not activate them. \square

Theorem 2. (Identifiability in SCG) Consider an SCG $\mathcal{G}^s = (\mathcal{V}^s, \mathcal{E}^s)$ associated with a DSCM with true (unknown) probability distribution P . Under Assumptions 1 and 2, the total effect $P(y_t | do(x_{t-\gamma}))$, with $\gamma \geq 0$, is identifiable if $X \notin Anc(Y, \mathcal{G}^s)$ or $X \in Anc(Y, \mathcal{G}^s)$ and none of the following holds:

- 1. $\gamma \neq 0$ and $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) \neq \emptyset$, or
- 2. there exists a σ -active backdoor path

$$\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$$

from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$ and one of the following holds:

- (a) $n > 2$, i.e. $\langle V^2, \dots, V^{n-1} \rangle \neq \emptyset$, or
- (b) $n = 2$ and $\gamma \neq 1$, or
- (c) $n = 2$, $\gamma = 1$ and $Cycles(Y, \mathcal{G}^s \setminus \{X\}) \neq \emptyset$.

Proof. The proof of this theorem is given by Lemmas 5.1-5.3. \square

Proposition 2. Consider an SCG \mathcal{G}^s and the total effect $P(y_t | do(x_{t-\gamma}))$, with $\gamma \geq 0$. Under conditions of identifiability provided by Theorem 2, the set \mathcal{A}'_γ is a valid adjustment set for the total effect.

Proof. Let \mathcal{G}^f be a candidate FTCCG. Consider $V_{t'} \in \mathcal{A}_\gamma \setminus \mathcal{A}'_\gamma$: by definition of \mathcal{A}'_γ , it follows that $V_{t'} \notin Anc(X_{t-\gamma}, \mathcal{G}^f) \cup Anc(Y_t, \mathcal{G}^f)$. Therefore $V_{t'}$ does not lie on any back-door path between $X_{t-\gamma}$ and Y_t : $V_{t'}$ is not necessary in the adjustment set, confirming that \mathcal{A}'_γ is also an adjustment set. \square

A.3 ANOTHER VERSION OF THEOREM 2

Theorem 2 - Version 2. Consider an SCG $\mathcal{G}^s = (\mathcal{V}^s, \mathcal{E}^s)$. The total effect $P(y_t | x_{t-\gamma})$ with γ is identifiable from \mathcal{G}^s if $X \notin Anc(Y, \mathcal{G}^s)$ or $X \in Anc(Y, \mathcal{G}^s)$ and one of the following conditions holds:

- 1. $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and there exists no σ -active backdoor path $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$ or
- 2. $\gamma = 0$ and there exists no σ -active backdoor path $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$ or
- 3. $Cycles^>(X, \mathcal{G}^s \setminus \{Y\}) = \emptyset$ and there exists a σ -active backdoor path $\pi^s = \langle V^1 = X, \dots, V^n = Y \rangle$ from X to Y in \mathcal{G}^s such that $\langle V^2, \dots, V^{n-1} \rangle \subseteq Desc(X, \mathcal{G}^s)$, and $n = 2$, and $\gamma = 1$, and $Cycles(Y, \mathcal{G}^s \setminus \{X\}) = \emptyset$.