PRUNE 'N PREDICT: OPTIMIZING LLM DECISION MAKING WITH CONFORMAL PREDICTION

Anonymous authors

003 004

006

008 009

010 011

012

013

014

015

016

017

018

019

021

023

024

025

026

Paper under double-blind review

ABSTRACT

Large language models (LLMs) are empowering decision-making in several applications, including tool or API usage and answering multiple-choice questions (MCQs). However, incorrect outputs pose significant risks in high-stakes domains like healthcare and finance. To quantify LLM uncertainty and thereby mitigate these risks, recent works employ conformal prediction (CP), a model- and distribution-agnostic framework that uses LLM outputs to generate a *prediction* set containing the true answer with high probability. Leveraging CP, we propose conformal revision of questions (CROQ), which revises the question by narrowing down the available choices to those in the prediction set and asking the LLM the revised question. We expect LLMs to be more accurate on revised questions with fewer choices. Furthermore, we expect CROQ to be effective when the prediction sets from CP are small. Commonly used logit scores often lead to large sets, diminishing CROQ's effectiveness. To overcome this, we propose CP-OPT, an optimization framework to learn scores that minimize set sizes while maintaining coverage. Our extensive experiments on MMLU, ToolAlpaca, and TruthfulQA datasets with multiple LLMs show that CROQ improves accuracy over the standard inference, with more pronounced gains when paired with CP-OPT.

028 1 INTRODUCTION

Large language models (LLMs) (Touvron et al., 2023; Databricks, 2024; Abdin et al., 2024) have demonstrated remarkable capabilities in various decision-making tasks, including multi-choice question answering and tool usage, where the model must select the correct tool or API to complete a task (Qu et al., 2024; Tang et al., 2023; Hendrycks et al., 2021). However, LLMs often exhibit overconfidence in wrong answers (Krause et al., 2023; Groot and Valdenegro Toro, 2024). Such unreliable predictions entail significant risks in critical domains like finance. Successful usage in such settings demands principled solutions to improve accuracy and quantify uncertainty in the predictions.

A commonly taught strategy for a human test taker
to solve multi-choice questions (MCQs) is the process of elimination (pruning) of incorrect (distractor)
answer choices. The underlying principle is that this
enables them to focus their attention on the remaining
answer choices, and it increases the likelihood of a
correct answer even if they have to guess randomly.
Inspired by this, we investigate whether LLMs can
benefit from a similar strategy.

045 We first examine the relationship between the number 046 of distractor answers and LLM accuracy on a MCQ 047 task. Figure 1 illustrates accuracy for three different 048 LLMs on a version of TruthfulQA, a widely used MCQ dataset. The MCQs in this version of TruthfulQA have 15 answer options, only one of which is 051 correct. (We discuss how this dataset is constructed in Appendix D.2.) For each question, we repeatedly 052 prompt the LLM, randomly eliminating one distractor answer at a time. Each prompt is independent,



Figure 1: Accuracy for three LLMs on the TruthfulQA dataset with 15 response options as a function of the number of incorrect answer options (distractors) removed from the prompt. As more distractor answers are eliminated, accuracy increases. Accuracy is averaged across 5 iterations, error bars denote ± 2 standard deviations.

- without any previous rounds included in the context. As hypothesized, *reducing the number of response options leads to an improvement in accuracy*, and this improvement is very nearly monotone. This suggests that eliminating distractor answers before prompting the LLM can indeed enhance accuracy. Of course, when pruning answers, we do not want to eliminate the correct answer, since that would necessarily cause the LLM to get the MCQ wrong.
- Conformal prediction (CP) (Vovk et al., 2005) is a flexible framework that can be used to prune distractor answers while retaining the correct answer with high probability. CP is a model-agnostic and distribution-free technique for generating prediction sets which contain the correct answer with a user-specified probability (e.g. 95%), which is referred to as the *coverage guarantee*.
- Utilizing this guarantee of CP, we propose a procedure called *conformal revision of questions* (CROQ), to revise MCQs with choices in a prediction set output by CP. This procedure represents a tradeoff: with some small probability (e.g. 5%), we may remove the correct answer from the prediction set, causing the LLM to get the question wrong. However, with high probability (e.g. 95%), we will retain the correct answer while reducing the number of distractor answers. Given the relationship observed in Figure 1, this should increase the LLM's accuracy on those questions. Different coverage rates naturally induce different tradeoffs. Overall, we hypothesize that we can find a coverage rate with a favorable tradeoff, such that CROQ improves the overall accuracy.
- CROQ's effectiveness depends on the size of the prediction sets from conformal prediction smaller sets mean fewer choices in the revised question and hence better final accuracy. Conformal prediction requires specifying a *score function*, which loosely speaking quantifies how plausible an output (answer option) is for a given input (question). While conformal prediction provides a coverage guarantee for *any* score function, the size of the prediction sets depends on the score function. As an example, a random score function will yield output sets that constitute random subsets of the label space that are large enough to satisfy the coverage guarantee (Angelopoulos and Bates, 2022).
- Previous works that apply conformal prediction in MCQ-type settings have used readily available
 scores such as the logits (or softmax values) output from the LLM (Kumar et al., 2023) or have
 designed heuristic scores based for example on repeated querying of the LLM (Su et al., 2024). Logits
 can be overconfident and may show biases for some options (Zheng et al., 2024), and heuristic scores
 are not guaranteed to produce small sets. Thus, in order to make CROQ as effective as possible, we
 propose CP-OPT (conformal prediction optimization), a principled solution to obtain scores that are
 designed to minimize set sizes (uncertainty) while preserving the coverage guarantee.
- ⁰⁸⁵ To summarize, our main contributions are as follows:
- We propose the conformal revision of questions (CROQ), in which we prune the answer choices in an MCQ to those in the prediction set output by conformal prediction and then prompt the LLM with the revised question. Empirical evaluation shows that this approach consistently improves accuracy compared to prompting the LLM with the original MCQ.
 - 2. We design a score function optimization framework (CP-OPT) that can be applied to any pretrained LLM. Moving away from the potentially unreliable LLM logits and heuristic scores, our framework provides a principled way to learn scores for conformal prediction. Empirically, we show that our procedure leads to a reduction in average set sizes compared to the baseline procedure that uses the LLM logits as the scores, at the same level (95%) of coverage.
 - 3. We further show that when used with CROQ, our CP-OPT scores deliver greater accuracy improvements over baseline than the LLM's logits.
 - 2 PRELIMINARIES
 - In this section, we provide background on solving MCQ tasks with LLMs and conformal prediction.
- 101 102 103 104

091

092

094

095

096

- 2.1 MULTIPLE CHOICE QUESTIONS (MCQS) AND LLMS
- MCQ Setup. MCQs are a general abstraction for expressing problems in which the correct choice(s) must be selected from a given set of choices. These encompass question-answering tasks like MMLU (Hendrycks et al., 2021) as well as other tasks such as tool learning, in which the LLM must select the correct tool or API to complete a task (Tang et al., 2023; Qu et al., 2024). An

115 **MCQ Prompt.** We concatenate the question text Q and the answer choices O, all separated by a new 116 line character, and append to the end the text "The correct answer is: ". The expectation 117 is that given this input prompt, the next token predicted by the LLM will be one of the option keys. 118 See Appendix D for a prompt example. We consider zero-shot prompts and do not include example 119 questions and answers in the prompt. We also add the prefix and suffix tokens to the prompt as 120 recommended by the language model providers. Since these are fixed modifications to x, we will use 121 x to denote the final prompt and the MCQ instance analogously.

122 LLM Inference. We run the forward pass of the auto-regressive LLM (Touvron et al., 2023; Dubey 123 et al., 2024; Abdin et al., 2024) on the input prompt to obtain the logit scores for each possible next 124 token given the prompt, restricting attention to the tokens that correspond to the available answer keys (e.g. "a", "b", "c", "d" if there are four answer options). We take the softmax to convert the 125 logits to probabilities, and then we take as the LLM's answer the option with the highest probability. 126 This approach ensures that the LLM's answer will be one of the available answer options, which 127 would not be guaranteed if instead we asked the LLM to simply generate an answer token given the 128 prompt. This approach mirrors what has been done in other works that use LLMs to solve MCQs 129 (Kumar et al., 2023; Su et al., 2024). Formal details are given in Appendix A.1. 130

131 132

151

2.2 CONFORMAL PREDICTION

133 Conformal prediction (CP) (Vovk et al., 2005; Angelopoulos et al., 2022) is a framework for quanti-134 fying uncertainty in machine learning models. It provides a flexible and user-friendly approach to 135 output prediction sets (which may be finite sets or intervals) that contain the true output or label with 136 a probability that is specified by the user, e.g. 95%. The key strength of conformal prediction lies in 137 its distribution-free guarantees: it ensures that the constructed prediction sets are valid regardless of 138 the underlying data distribution and model. This property is particularly desirable in the context of 139 language models, as it is hard to characterize language data distributions or put specific distributional 140 assumptions/restrictions on the LLMs.

141 142 142 143 144 144 144 144 145 145 146 147 146 147 147 147 **Score Function.** Let $g: \mathcal{X}_m \times \mathcal{Y}_m \mapsto \mathbb{R}$ be a conformal *score function*, where larger scores indicate 148 better agreement ("conformity") between x and y. Intuitively, large scores are intended to indicate 149 that y is a plausible output given x, while smaller scores indicate less plausibility. (Note that some 140 authors prefer to have larger scores indicate greater disagreement, e.g. Clarkson et al. (2024).) A 145 common choice of score function is the softmax scores from the given model. For closed-source 146 LLMs, where logits are not available, others have devised self-consistency scores based on repeated 147 querying of the model (Su et al., 2024).

Prediction Sets. Given a score function g and threshold τ on the scores, the prediction set for any $x \in \mathcal{X}_m$ is given by

$$C(x;g,\tau) := \{ y \in \mathcal{Y}_m : g(x,y) \ge \tau \}.$$

$$(1)$$

Intuitively, larger sets represent greater uncertainty, while smaller sets represent less uncertainty.
 Given a fixed confidence level, a score function that produces larger sets can be said to result in greater uncertainty.

Split Conformal Prediction. Similar to prior works (Kumar et al., 2023; Su et al., 2024), we use *Split Conformal Prediction* (Papadopoulos et al., 2002; Lei et al., 2018) due to its popularity, ease of use, and computational efficiency. Given a score function $g : \mathcal{X}_m \times \mathcal{Y}_m \mapsto \mathbb{R}$, Split Conformal Prediction uses a calibration dataset $D_{cal} = \{x_i, y_i^*\}_{i=1}^{n_{cal}}$ to compute a threshold $\hat{\tau}_{\alpha}$, defined as

160
161
$$\hat{\tau}_{\alpha} = \min\left\{q: \frac{1}{n_{\text{cal}}} \sum_{i=1}^{n_{\text{cal}}} \mathbb{1}\left(g(x_i, y_i^{\star}) \le q\right) \ge \alpha\right\},\tag{2}$$



Figure 2: (CROQ) Illustration of conformal revision of questions and prompting the LLM with the revised question. In this example, the initial predicted set by LLM + conformal prediction (CP) is {C, D}. The question and labels are revised to contain only the answer choices in the prediction set and the LLM is prompted with the revised question. Since CP provides rigorous coverage guarantees, we expect that re-prompting the LLM with reduced answer choices will improve the chances of obtaining the correct answer. See Section 3.1 for details.

where $\alpha \in [0, 1]$ is a user-chosen *miscoverage rate* that is equal to 1 minus the desired coverage; for example, a value of $\alpha = 0.05$ would correspond to a coverage of 95%. In words, $\hat{\tau}_{\alpha}$ is the smallest empirical quantile of the scores for the correct answers on the calibration dataset that is sufficient to satisfy (an empirical version of) the coverage property. The threshold $\hat{\tau}$ is used to construct prediction sets $C(x; g, \hat{\tau})$ on previously unseen test points as in (1). This procedure enjoys a marginal coverage guarantee for prediction sets on unseen test data points, formalized as Proposition 2.1.

Proposition 2.1. (Marginal Coverage Guarantee) (Lei et al., 2018, Thm. 2.2) Let g be a fixed conformity score function and $\hat{\tau}_{\alpha}$ be an α threshold computed via Split Conformal Prediction on $D_{\text{cal}} = \{x_i, y_i^*\}_{i=1}^{n_{\text{cal}}} \sim \mathbb{P}_{\mathcal{X}_m \times \mathcal{Y}_m}$. Then, for a new sample $(\tilde{x}, \tilde{y}^*) \sim \mathbb{P}_{\mathcal{X}_m \times \mathcal{Y}_m}$, we have that

₽

$$(\tilde{y}^{\star} \in C(\tilde{x}; g, \hat{\tau}_{\alpha})) \ge 1 - \alpha.$$
 (3)

where the probability is marginal over the randomness in the calibration data and the new sample.

The top half of Figure 2 illustrates conformal prediction for answering MCQs with LLMs. While the coverage guarantee in Proposition 2.1 holds for any score function, ideally we would like a score function that yields the smallest sets possible (the least uncertainty). Next, we discuss our solutions to improve conformal prediction and its utility in solving MCQs with LLMs.

201 202

203

206

194

195

196

3 Methodology

In this section, we discuss details of our pipeline for question revision using conformal prediction and our procedure to generate optimal conformal scores.

207 3.1 CONFORMAL REVISION OF QUESTIONS (CROQ)

The procedure involves prompting the LLM with the reduced answer options from a conformal prediction set. The steps are illustrated with an example in Figure 2.

Scores and Threshold for Conformal Prediction. We first fix a score function $g: \mathcal{X}_m \times \mathcal{Y}_m \mapsto \mathbb{R}$. Here we restrict the score function to either the logits generated by the LLM or the CP-OPT scores discussed in Section 3.2. We then run the split conformal procedure with coverage level $1 - \alpha$ for some $\alpha \in [0, 1]$ to estimate the threshold $\hat{\tau}_{\alpha}$. CROQ then proceeds as follows.

215 Step 1: Get Conformal Prediction Set. Given a test instance x, we generate a first stage prediction set, $C(x; g, \hat{\tau}_{\alpha})$. Per the coverage guarantee (Proposition 2.1), we expect that the true answer

216 $y^* \in C(x; g, \hat{\tau}_{\alpha})$ with probability at least $1 - \alpha$. Next, the question is revised to contain only the choices in the set $C(x; g, \hat{\tau}_{\alpha})$.

Step 2: Revise the Question and Ask the LLM. If the first stage prediction set $C(x; q, \hat{\tau}_{\alpha})$ is 219 empty or is of size 1 or size m (the number of answer options), then we simply utilize the LLM's 220 answer to the original MCQ x, as described in section 2.1, since the conformal procedure has 221 yielded no additional information. Otherwise, we modify the prompt x to x' = (Q, O'), where 222 $O' = \{(K_i, V_i) : K_i \in C(x; g, \hat{\tau}_\alpha)\}$. The keys in O' are changed so that they start with the first 223 letter of the alphabet and go to the letter corresponding to the number of choices available. For 224 example, if there were initially four answer options $\{a, b, c, d\}$, and the conformal prediction set was 225 $\{c, d\}$, then the two options in the set would receive new keys $\{a, b\}$. Then x' is transformed into a 226 prompt format and passed to the LLM, and the standard inference procedure (section 2.1) is run to extract the predicted answer key \hat{y}' . 227

With fewer choices in the revised question, we expect LLMs will be more accurate in their answer compared to the answer to the initial question. However, the improvement in accuracy will depend on the size of the prediction sets. As shown in Figure 1 LLMs have a higher chance of answering the question correctly if the number of options is small. This implies the efficacy of CROQ will depend on the size of sets $C(x; g, \hat{\tau}_{\alpha})$ – if these sets are small then we can expect more improvement.

While conformal prediction can output sets $C(x; g, \hat{\tau}_{\alpha})$ for any score function g, along with $1 - \alpha$ coverage guarantee, the set sizes could be highly variable depending on the score function g. Noting the lack of reliability of scores used in prior works, that could yield unnecessarily large sets, we seek to learn scores that minimize the set sizes while preserving the coverage guarantee. We discuss our procedure to learn such scores in the next section. Using these scores in CP we expect to get smaller sets and thus more improvement in CROQ compared to baseline scores.

239 240

241

252 253 254

255

261

262

263

264

265 266

3.2 CP-OPT TO OPTIMIZE SCORES

We describe our method for learning the optimal scores for conformal prediction (CP) for solving MCQs with LLMs. Similar ideas have been incorporated in the training objective of classifiers (Stutz et al., 2022) so that the classifiers' softmax output is better suited for CP. However, the LLMs are not trained with this objective, and we want to apply CP to any given LLM; therefore, we design a post-hoc method to optimize the scores. We first characterize the optimal scores and then describe how to estimate them in practice.

248 Characterization of the optimal scores. For any score function $g : \mathcal{X}_m \times \mathcal{Y}_m \mapsto \mathbb{R}$ 249 and threshold τ , the membership of any y in the prediction set $C(x; g, \tau)$ is given by 250 $\mathbb{1}(y \in C(x; g, \tau)) = \mathbb{1}\{g(x, y) \geq \tau\}$. Define the expected set size $S(g, \tau)$ and the cover-251 age conditional on τ , denoted $\mathcal{P}(g, \tau)$, as follows:

$$S(g,\tau) := \mathbb{E}_x \Big[\sum_{y \in \mathcal{Y}_m} \mathbb{1}\{g(x,y) \ge \tau\} \Big]. \quad (4) \qquad \mathcal{P}(g,\tau) := \mathbb{E}_x \left[\mathbb{1}\{g(x,y^*) \ge \tau\} \right]. \quad (5)$$

The optimal score function g^* and threshold τ^* are defined (non-uniquely) to minimize the expected set size subject to the coverage $\mathcal{P}(g,\tau)$ being at least $1 - \alpha$:

$$g^{\star}, \tau^{\star} := \underset{g:\mathcal{X}_m \times \mathcal{Y}_m \mapsto \mathbb{R}, \tau \in \mathbb{R}}{\operatorname{arg\,min}} S(g,\tau) \text{ s.t. } \mathcal{P}(g,\tau) \ge 1 - \alpha.$$
(P1)

Practical Version with Differentiable Surrogates and Empirical Estimates. Problem (P1) characterizes optimal score functions and thresholds. However, in practice, we do not know the underlying distribution and thus do not have access to the quantities in (4) and (5). Instead, we obtain their estimates using a training sample $D_{\text{train}} = \{(x_i, y_i^*)\}_{i=1}^{n_t}$ drawn independently from the same distribution:

$$\widehat{S}(g,\tau) := \frac{1}{n_t} \sum_{i=1}^{n_t} \sum_{y \in \mathcal{Y}_m} \mathbb{1}\{g(x_i, y) \ge \tau\}, \quad (6) \qquad \widehat{\mathcal{P}}(g,\tau) := \frac{1}{n_t} \sum_{i=1}^{n_t} \mathbb{1}\{g(x_i, y_i^\star) \ge \tau\}. \quad (7)$$

267 268 269

Using these plug-in estimators in problem (P1) yields a revised optimization problem. However, it is difficult to solve this problem as the objective and constraints are not differentiable. To make

them differentiable, we introduce the following surrogates. Given g(x, y) and τ , define the following sigmoid function with $\beta > 0$, $\sigma(x, y, g, \tau, \beta) := 1/(1 + \exp(-\beta (g(x, y) - \tau)))$. The sigmoid function provides a differentiable approximation to the indicator variable for $g(x, y) \ge \tau$. The approximation is tighter with larger β i.e., $\sigma(x, y, g, \tau, \beta) \rightarrow 1\{g(x, y) \ge \tau\}$ as $\beta \rightarrow \infty$, and $g(x, y) \ge \tau \iff \sigma(x, y, g, \tau) \ge 1/2$. By using these sigmoid surrogates in equation (6), we obtain the following smooth plugin estimates,

$$\widetilde{S}(g,\tau) := \frac{1}{n_t} \sum_{i=1}^{n_t} \sum_{y \in \mathcal{Y}_m} \sigma(x_i, y, g, \tau, \beta). \quad (8) \qquad \widetilde{\mathcal{P}}(g,\tau) := \frac{1}{n_t} \sum_{i=1}^{n_t} \sigma(x_i, y_i^\star, g, \tau, \beta). \quad (9)$$

It is easy to see that by the strong law of larger numbers and properties of the sigmoid function, as $n_t, \beta \to \infty$, the surrogate average set size and coverage will converge almost surely to their population versions, i.e. $\tilde{S}(g,\tau) \xrightarrow{a.s.} S(g,\tau)$ and $\tilde{\mathcal{P}}(g,\tau) \xrightarrow{a.s.} \mathcal{P}(g,\tau)$. We replace the expected set size and marginal coverage by these smooth surrogates in (P1) and transform it into an unconstrained problem with a penalty term $\lambda > 0$. We also introduce ℓ_2 regularization to encourage low norm solutions. We optimize the score function g over a flexible space of functions \mathcal{G} , such as neural networks (NNs). The resulting problem (P2) is differentiable, and we solve it using stochastic gradient descent.

$$\tilde{g}, \tilde{\tau} := \underset{g \in \mathcal{G}, \tau \in \mathbb{R}}{\arg\min} \widetilde{S}(g, \tau) + \lambda \left(\widetilde{\mathcal{P}}(g, \tau) - 1 + \alpha \right)^2 - \hat{\mathcal{C}}(g) + \lambda_1 \|g\|_2^2.$$
(P2)

Here, $\hat{\mathcal{C}}(g) := \frac{1}{n_t} \sum_{i=1}^{n_t} \log(g(x_i, y_i^*))$ is the cross entropy term included to encourage higher scores for correct predictions, and the regularization term $\lambda_1 ||g||_2^2$ is the squared norm over the parameters of g to promote low norm solutions. Solving (P2) yields a score function \tilde{g} and a threshold $\tilde{\tau}$. However, $\tilde{\tau}$ may be biased, since it is estimated on the same data as \tilde{g} . Following the split conformal procedure, we therefore estimate a new threshold $\hat{\tau}$ on a separate calibration dataset. Note that our framework is flexible and can work with any choice of features and function class for which the ℓ_2 norm can be calculated. We discuss the specific choice of features and \mathcal{G} used in this work in Appendix A.2.

4 EXPERIMENTS

We conduct experiments on benchmark MCQ and tool usage tasks with open-weight instruction-tuned models to test the following hypotheses:

H1. CP-OPT scores in conformal prediction on MCQ tasks with LLMs yield a smaller average set size at the same level of coverage in comparison to using LLM logits.

H2. Conformal revision of questions (CROQ) improves accuracy over the standard inference.

H3. CROQ with CP-OPT scores performs better than CROQ with logit scores.

308 309

276 277

278 279

280

281

282

283

284

285

291 292

293

294

295

296

297 298 299

300 301

302

303

304

305 306

307

310 4.1 EXPERIMENTAL SETUP

We first describe the setup for the experiments and then discuss the results for the above hypotheses.

313 Datasets. We evaluate our hypotheses on 3 datasets: MMLU (Hendrycks et al., 2021), TruthfulQA 314 (Lin et al., 2022), and ToolAlpaca (Tang et al., 2023). MMLU and TruthfulQA are popular benchmark datasets for multiple-choice questions. MMLU focuses on assessing multitask accuracy; 315 it contains multiple choice questions (MCQs) from 57 domains, including humanities, math, medicine, 316 etc. TruthfulQA evaluates an LLM's ability to answer truthfully and avoid falsehoods that humans 317 are susceptible to. ToolAlpaca contains 3.9k tool-use instances from a multi-agent simulation 318 environment, which we augment to a MCQ format. Dataset descriptions and example questions and 319 responses are provided in Appendix D. 320

Models. We use auto-regressive language models based on the transformer architecture. We choose instruction-tuned, open-weight, and small to medium-sized models, for reproducibility and reduced computational cost. Specifically, we use Llama-3-8B-Instruct by Meta (Dubey et al., 2024), Phi-3-4k-mini-Instruct by Microsoft (Abdin et al., 2024), and the

| | | | Llan | na-3 | | | Phi | i-3 | | | Gem | ma-2 | |
|------------|--------|--------|---------|--------|-------|--------|----------|--------|-------|--------|----------|--------|-------|
| | | Avg. S | et Size | Cove | rage | Avg. S | Set Size | Cove | rage | Avg. S | Set Size | Cove | rage |
| Dataset | # Opt. | Logits | Ours | Logits | Ours | Logits | Ours | Logits | Ours | Logits | Ours | Logits | Ours |
| | 4 | 2.56 | 2.53* | 95.75 | 95.57 | 2.21 | 2.16* | 94.65 | 94.35 | 2.94 | 2.40* | 95.16* | 94.23 |
| MMLU | 10 | 5.53 | 4.90* | 96.06* | 95.45 | 4.36 | 4.36 | 94.11 | 94.09 | 7.79 | 6.08* | 95.00* | 94.04 |
| | 15 | 7.69 | 7.18* | 95.42 | 95.06 | 6.64 | 6.52* | 94.60 | 94.61 | 11.71 | 10.04* | 94.58 | 94.58 |
| | 4 | 1.17 | 1.18 | 97.08 | 96.85 | 1.07 | 1.08 | 95.33 | 95.68 | 1.12 | 1.05* | 95.68 | 95.44 |
| ToolAlpaca | 10 | 1.51 | 1.39* | 95.21 | 95.56 | 1.25 | 1.20* | 95.56 | 95.09 | 2.05 | 1.42* | 95.56 | 94.51 |
| | 15 | 1.97 | 1.67* | 96.50 | 96.03 | 1.68 | 1.54* | 98.36* | 97.20 | 3.54 | 1.77* | 96.14 | 95.21 |
| | 4 | 3.34 | 2.69* | 95.95* | 92.41 | 2.85 | 2.53* | 96.71 | 96.71 | 2.74 | 1.88* | 96.46 | 95.44 |
| TruthfulQA | 10 | 7.06 | 6.41* | 94.43 | 93.42 | 7.48 | 6.49* | 98.48* | 95.70 | 7.52 | 5.64* | 95.44 | 97.22 |
| | 15 | 10.61 | 10.62 | 94.68 | 94.68 | 10.72 | 10.30* | 95.44 | 96.46 | 11.23 | 9.35* | 95.44 | 96.46 |

Table 1: Average set sizes and coverage rates (in percentages) for conformal prediction sets on the MMLU, ToolAlpaca, and TruthfulQA datasets using gemma-2-9b-it-SimPO (Gemma-2), Llama-3-8B-Instruct (Llama-3) and Phi-3-4k-mini-Instruct (Phi-3), with a target coverage level of 95%. Bold numbers indicate smaller avg. set sizes. Asterisks on the larger of a pair of numbers indicate where the difference in average set size or coverage is statistically significant at the 0.05 significance level.

| | | | Llama-3 | | | Phi-3 | | | Gemma-2 | |
|------------|--------|---|-----------------------------|---------------------|---|--|---------------------|---|-----------------------------|---------------------|
| Model | # Opt. | Accuracy Before (a ₁) | Accuracy After (a'_1) | Gain $(a'_1 - a_1)$ | Accuracy Before (a ₁) | $\begin{array}{c} \textbf{Accuracy} \\ \textbf{After} \\ (a_1') \end{array}$ | Gain $(a'_1 - a_1)$ | Accuracy Before (a ₁) | Accuracy After (a'_1) | Gain $(a'_1 - a_1)$ |
| | 4 | 64.02 | 63.83 | -0.19 | 70.27 | 69.08 | -1.19 | 67.62 | 67.70 | 0.07 |
| MMLU | 10 | 54.82 | 56.29 | 1.47* | 58.44 | 61.57 | 3.13* | 53.80 | 53.93 | 0.13 |
| | 15 | 51.99 | 54.11 | 2.11* | 53.48 | 58.09 | 4.62* | 50.78 | 50.58 | -0.20 |
| | 4 | 91.47 | 91.94 | 0.47 | 92.76 | 92.64 | -0.12 | 93.46 | 93.11 | -0.35 |
| ToolAlpaca | 10 | 85.16 | 88.67 | 3.50* | 87.50 | 90.89 | 3.39* | 87.73 | 89.60 | 1.87* |
| | 15 | 81.43 | 87.85 | 6.43* | 85.98 | 89.25 | 3.27* | 87.97 | 88.55 | 0.58 |
| | 4 | 54.43 | 55.19 | 0.76 | 69.87 | 70.13 | 0.25 | 74.68 | 74.94 | 0.25 |
| TruthfulQA | 10 | 39.24 | 40.76 | 1.52 | 55.70 | 54.43 | -1.27 | 56.46 | 56.20 | -0.25 |
| | 15 | 37.22 | 37.22 | 0.00 | 46.84 | 46.33 | -0.51 | 55.95 | 56.96 | 1.01 |

Table 2: [CROQ + logits]. Results on accuracy improvement with CROQ using logit scores. Here a_1 , and a'_1 refer to the accuracy before CROQ and after CROQ respectively. A positive gain implies CROQ improved accuracy in that setting.

gemma-2-9b-it-SimPO model (Meng et al., 2024). For brevity, we use the short names Llama-3, Phi-3, and Gemma-2 respectively for these models.

Choices of Scores. We use the following scores for conformal prediction. (1) LLM Logits (Softmax) are extracted from the LLM as discussed in Section 2.1. These have been used in prior works (Kumar et al., 2023; Su et al., 2024). (2) CP-OPT (Ours) are the scores learned using the score optimization procedure discussed in Section 3.2. We use the train split for each dataset to learn these scores. The hyperparameter settings we used for CP-OPT are given in Appendix D.3. We omit the self-consistency based heuristic scores proposed by Su et al. (2024), as these require repeated inferences to get good estimates of the scores, and hence have a high computational cost.

We use the provided validation splits as our calibration datasets for the conformal procedure. For testing the hypotheses, we calibrate the conformal threshold for the coverage guarantee of 95%, i.e. we set the miscoverage rate α to 0.05. In addition, we study CROQ with calibration in a range of α values: {0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.4, 0.5 }. Performance is computed on test splits. The hyperparameters used to learn the score function using SGD are provided in table 20 in Appendix D.3.

Statistical Significance. We report the statistical significance of our results using paired sample
 t-tests, using asterisks (*) to annotate results that are statistically significant at a 0.05 significance
 level. See Appendix C for details.

| | | | Llama-3 | | | Phi-3 | | | Gemma-2 | |
|------------|--------|---|---|-------------------------|--|---|-------------------------|---|---|-------------------------|
| Model | # Opt. | $\begin{array}{c c} \textbf{Accuracy} \\ \textbf{Logits} \\ (a'_1) \end{array}$ | $\begin{array}{c} \textbf{Accuracy} \\ \textbf{CP-OPT} \\ (a_2') \end{array}$ | Gain $(a'_2 - a'_1)$ | $\begin{vmatrix} \mathbf{Accuracy} \\ \mathbf{Logits} \\ (a'_1) \end{vmatrix}$ | $\begin{array}{c} \textbf{Accuracy} \\ \textbf{CP-OPT} \\ (a_2') \end{array}$ | Gain $(a'_2 - a'_1)$ | $\begin{array}{c c} \textbf{Accuracy} \\ \textbf{Logits} \\ (a'_1) \end{array}$ | $\begin{array}{c} \textbf{Accuracy} \\ \textbf{CP-OPT} \\ (a_2') \end{array}$ | Gain $(a'_2 - a'_1)$ |
| | 4 | 63.83 | 63.67 | -0.16 | 69.08 | 69.34 | 0.26 | 67.70 | 69.56 | 1.86* |
| MMLU | 10 | 56.29 | 57.11 | 0.82* | 61.57 | 61.05 | -0.52 | 53.93 | 57.93 | 4.00* |
| | 15 | 54.11 | 54.77 | 0.66* | 58.09 | 58.15 | 0.06* | 50.58 | 51.31 | 0.73 |
| | 4 | 91.94 | 91.82 | -0.12 | 92.64 | 92.52 | -0.12 | 93.11 | 93.57 | 0.46 |
| ToolAlpaca | 10 | 88.67 | 89.02 | 0.35* | 90.89 | 91.00 | 0.11* | 89.60 | 90.42 | 0.82* |
| | 15 | 87.85 | 88.67 | 0.82* | 89.25 | 89.95 | 0.70* | 88.55 | 89.37 | 0.82 |
| | 4 | 55.19 | 55.44 | 0.25 | 70.13 | 69.87 | -0.26 | 74.94 | 76.96 | 2.02 |
| TruthfulQA | 10 | 40.76 | 42.28 | 1.52 | 54.43 | 56.20 | 1.77 | 56.20 | 60.76 | 4.56* |
| | 15 | 37.22 | 37.47 | 0.25 | 46.33 | 51.39 | 5.06* | 56.96 | 57.72 | 0.76 |

Table 3: [CROQ + logits vs CROQ + CP-OPT]. Comparison of CP-OPT and logits on accuracy improvement with CROQ. Here, a'_1 , and a'_2 are the final accuracies after CROQ using logits and CP-OPT respectively (as in Tables 2 and 4. The gain $a'_2 - a'_1$ is the difference between these two, with values indicating more improvement in CROQ with CP-OPT scores.

4.2 DISCUSSION

390

391

392

393

394

397 398

399 400

401 H1. Improvement in conformal set sizes with our CP-OPT scores. We run the CP procedure using 402 the LLM logits and CP-OPT scores and obtain conformal sets for points in the test sets. We compute 403 the average set size and coverage for each dataset, model, and score combination. The results are 404 in Table 1. As expected, in most settings (17 out of 27) we see a statistically significant reduction 405 in the set sizes with our (CP-OPT) scores with similar coverage as logits. The reduction is more 406 pronounced with a higher number of options. In a few settings (6/27), the reduction in set size is 407 accompanied by a statistically significant decrease in coverage relative to using the logits. In the remaining 4/27 settings the differences are insignificant. Note that since the target coverage level is 408 95%, anything above 95% is over-coverage. We see that logits tend to over-cover and thus a drop 409 in coverage is expected as long as it does not fall significantly below the desired level of 95% (this 410 happens only in 2/27 settings). Overall, these results show CP-OPT's effectiveness in reducing set 411 sizes while maintaining the target coverage level. In Appendix B, we provide histograms (e.g., Figure 412 6) of set sizes produced by logits and CP-OPT scores in all settings. These histograms show a clear 413 pattern: CP-OPT scores produce fewer large sets and more small sets in comparison to logit scores. 414

H2. Accuracy improvement with conformal revision of questions (CROQ). Tables 2 and 4 show the 415 accuracy before and after CROQ with logit and CP-OPT scores respectively. With the logit scores 416 (Table 2), we see an increase in accuracy (by up to 6.43%) in 19 out of 27 settings, out of which 417 9 are statistically significant. In 8 of the settings, we see a small drop in accuracy (which is not 418 statistically significant). Next, with CP-OPT scores (Table 4) we see accuracy improvements (up to 419 7.24%) in 24 settings, of which 13 are statistically significant. In the remaining 3 settings, we see a 420 non-significant drop in accuracy. Overall, we observe that in the vast majority of the settings, CROQ 421 improves accuracy with either logits or CP-OPT scores. The rare small drops in accuracy could occur 422 since the conformal procedure may eliminate the correct option with low probability (α).

423 H3. CROQ with CP-OPT scores is better than CROQ with logit scores. CP-OPT scores are designed 424 to minimize set sizes while maintaining the coverage guarantee. As a result, using these scores with 425 CROQ is expected to reduce uncertainty for many questions, leading to fewer answer options in 426 the revised prompts. Based on Figure 1, we expect LLMs to be more likely to answer correctly 427 when prompted with the revised question with fewer options. The results of CROQ with CP-OPT 428 are summarized in Table 4, and in Table 3 we compare the accuracies after CROQ with logits and 429 CP-OPT. In Table 3 we see that in 22 out of 27 settings, CROQ with CP-OPT results in higher accuracy (up to 4.56%) than CROQ with logits. Furthermore, the improvements in 12 out of these 22 430 settings are statistically significant. The drop in accuracy in the remaining 5 settings is statistically 431 non-significant. Overall, we see that CROQ with CP-OPT is generally better than with logits.

432 5 RELATED WORK

434

436

437

438

439

440

441

442

Conformal Prediction for Uncertainty Quantification with LLMs. Recently there has been growing interest in using conformal prediction to quantify and control uncertainty in LLM-related tasks. In the context of multi-choice question answering (MCQ), previous works have investigated a variety of conformal score functions, including (the softmax of) the LLM logits corresponding to the response options (Kumar et al., 2023; Ren et al., 2023) or functions thereof (Ye et al., 2024), confidence scores generated by the LLM itself, or "self-consistency" scores derived by repeated querying of the LLM (Su et al., 2024). We build on this work by aiming to learn a conformal score function that yields small conformal sets, rather than taking the score function as given.

In addition to the MCQ setting, there has been recent work utilizing conformal prediction in the 443 context of open-ended response generation (Quach et al., 2024; Mohri and Hashimoto, 2024; Cherian 444 et al., 2024). This setting differs in that there is not necessarily a unique correct response, so the 445 notion of coverage must be redefined around acceptability or factuality rather than correctness. When 446 factuality is the target, the goal is to calibrate a pruning procedure that removes a minimal number of 447 claims from an LLM-generated open response, such that the remaining claims are all factual with 448 high probability; that is, the goal is to retain as large a set as possible, rather than to generate a set 449 with the smallest number of responses possible as in MCQ. Conformal prediction has also been used 450 to capture token-level uncertainty (Deutschmann et al., 2024; Ravfogel et al.; Ulmer et al., 2024). 451

452

Optimizing Conformal Prediction Procedures. Several recent works have considered how to 453 learn good conformal score functions from data, primarily in the context of supervised learning 454 models (Bai et al., 2022; Stutz et al., 2022; Yang and Kuchibhotla, 2024; Xie et al., 2024). With LLMs, 455 Cherian et al. (2024) consider how to learn a good score function to achieve factuality guarantees; 456 their optimization problem differs from ours due to the difference in setting as well as the addition 457 of conditional coverage constraints (ensuring that coverage holds in different parts of the feature 458 space). Kiyani et al. (2024) design a framework to minimize the size ("length," in their terminology) 459 of conformal sets, which they apply to MCQ as well as to supervised learning problems. However, 460 their framework is concerned with how to generate sets given a model and a conformity score, rather 461 than how to learn a conformity score. 462

The works mentioned above all aim to produce small conformal sets that satisfy coverage guarantees. Among these, only Ren et al. (2023) consider how conformal sets may be used downstream, in their case to improve the efficiency and autonomy of robot behavior. To our knowledge, our work is the first to investigate whether conformal prediction can be used to increase the accuracy of LLMs on MCQ type tasks.

468 469

6 CONCLUSIONS AND FUTURE WORKS

470 471

472 In this work, we introduced Conformal Revision of Questions (CROQ), a principled approach to 473 improve LLM accuracy in multiple-choice settings by leveraging conformal prediction (CP) to 474 eliminate distractor answers while maintaining high coverage of the correct answer. To further boost 475 CROQ's performance we proposed CP-OPT, a framework for optimizing score functions to minimize 476 prediction set sizes while preserving CP's coverage guarantees. Our results demonstrate that CROQ significantly enhances LLM's accuracy, and that CP-OPT further strengthens this effect by producing 477 smaller, more reliable prediction sets than standard LLM logits. These findings highlight the potential 478 of uncertainty-aware, test-time methods to improve LLM decision-making, providing a principled 479 path for safer and more effective deployment of LLMs in critical applications. 480

Future works could explore multi-round CROQ, where answer options are pruned iteratively in
 multiple rounds, further improving accuracy while maintaining coverage. This requires developing
 efficient recalibration strategies and methods to prevent excessive coverage reduction across iterations.
 Additionally, a key challenge is adapting conformal score thresholds in settings with a variable
 number of response options. Techniques like quantile regression could help calibrate thresholds
 dynamically, ensuring robust performance across diverse decision-making scenarios.

486 REFERENCES 487

493

500

501 502

503

504

506

507

508

509

520

521

522

523

526

527

528

529

- M. Abdin, S. A. Jacobs, A. A. Awan, J. Aneja, A. Awadallah, H. Awadalla, N. Bach, A. Bahree, 488 A. Bakhtiari, H. Behl, et al. Phi-3 technical report: A highly capable language model locally on 489 your phone. arXiv preprint arXiv:2404.14219, 2024. 490
- 491 A. N. Angelopoulos and S. Bates. A Gentle Introduction to Conformal Prediction and Distribution-492 Free Uncertainty Quantification. arXiv preprint arXiv:2107.07511, (arXiv:2107.07511), 2022.
- 494 A. N. Angelopoulos, S. Bates, E. J. Candès, M. I. Jordan, and L. Lei. Learn then test: Calibrating predictive algorithms to achieve risk control. arXiv preprint arXiv:2110.01052, 2022. 495
- 496 Y. Bai, S. Mei, H. Wang, Y. Zhou, and C. Xiong. Efficient and differentiable conformal prediction 497 with general function classes. In The Tenth International Conference on Learning Representations, 498 2022. 499
 - J. J. Cherian, I. Gibbs, and E. J. Candès. Large language model validity via enhanced conformal prediction methods. arXiv preprint arXiv:2406.09714, 2024.
 - J. Clarkson, W. Xu, M. Cucuringu, and G. Reinert. Split conformal prediction under data contamination. arXiv preprint arXiv:2407.07700, 2024.
- 505 Databricks. Introducing DBRX: A New State-of-the-Art Open LLM, 2024.
 - N. Deutschmann, M. Alberts, and M. R. Martínez. Conformal autoregressive generation: Beam search with coverage guarantees. In Proceedings of the AAAI Conference on Artificial Intelligence, 2024.
- 510 D. Dohan, W. Xu, A. Lewkowycz, J. Austin, D. Bieber, R. G. Lopes, Y. Wu, H. Michalewski, R. A. 511 Saurous, J. Sohl-dickstein, K. Murphy, and C. Sutton. Language model cascades, 2022. 512
- A. Dubey, A. Jauhri, A. Pandey, A. Kadian, A. Al-Dahle, A. Letman, A. Mathur, A. Schelten, A. Yang, 513 A. Fan, et al. The llama 3 herd of models. arXiv preprint arXiv:2407.21783, 2024. 514
- 515 R. El-Yaniv and Y. Wiener. On the foundations of noise-free selective classification. JMLR, 11: 516 1605-1641, aug 2010. ISSN 1532-4435. 517
- A. Fisch, T. S. Jaakkola, and R. Barzilay. Calibrated selective classification. Transactions on Machine 518 Learning Research, 2022. ISSN 2835-8856. 519
 - T. Groot and M. Valdenegro Toro. Overconfidence is key: Verbalized uncertainty evaluation in large language and vision-language models. In Proceedings of the 4th Workshop on Trustworthy Natural Language Processing (TrustNLP 2024). Association for Computational Linguistics, 2024.
- N. Gupta, H. Narasimhan, W. Jitkrittum, A. S. Rawat, A. K. Menon, and S. Kumar. Language model 524 cascades: Token-level uncertainty and beyond. arXiv preprint arXiv:2404.10136, 2024. 525
 - D. Hendrycks, C. Burns, S. Basart, A. Zou, M. Mazeika, D. Song, and J. Steinhardt. Measuring massive multitask language understanding. In International Conference on Learning Representations, 2021.
- 530 S. Kiyani, G. Pappas, and H. Hassani. Length Optimization in Conformal Prediction. arXiv preprint arXiv:2406.18814, 2024.
- 532 L. Krause, W. Tufa, S. Baez Santamaria, A. Daza, U. Khurana, and P. Vossen. Confidently wrong: 533 Exploring the calibration and expression of (un)certainty of large language models in a multilingual 534 setting. In Proceedings of the Workshop on Multimodal, Multilingual Natural Language Generation 535 and Multilingual WebNLG Challenge (MM-NLG 2023). Association for Computational Linguistics, 536 2023. 537
- B. Kumar, C. Lu, G. Gupta, A. Palepu, D. Bellamy, R. Raskar, and A. Beam. Conformal prediction 538 with large language models for multi-choice question answering. arXiv preprint arXiv:2305.18404, 2023.

- J. Lei, M. G'Sell, A. Rinaldo, R. J. Tibshirani, and L. Wasserman. Distribution-free predictive inference for regression. *Journal of the American Statistical Association*, 113(523), 2018.
- S. Lin, J. Hilton, and O. Evans. TruthfulQA: Measuring how models mimic human falsehoods. In
 Proceedings of the 60th Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers). Association for Computational Linguistics, 2022.
- Y. Meng, M. Xia, and D. Chen. SimPO: Simple preference optimization with a reference-free reward. *arXiv preprint arXiv:2405.14734*, 2024.
- C. Mohri and T. Hashimoto. Language models with conformal factuality guarantees. *arXiv preprint arXiv:2402.10978*, 2024.
- H. Papadopoulos, K. Proedrou, V. Vovk, and A. Gammerman. Inductive confidence machines for regression. In *Machine learning: ECML 2002: 13th European conference on machine learning Helsinki, Finland, August 19–23, 2002 proceedings 13.* Springer, 2002.
- C. Qu, S. Dai, X. Wei, H. Cai, S. Wang, D. Yin, J. Xu, and J.-R. Wen. Tool learning with large language models: A survey. *arXiv preprint arXiv:2405.17935*, 2024.

559

560

561

562

563

564 565

566

567

568

569 570

571

572

573

574

575 576

577

578

584

585

- V. Quach, A. Fisch, T. Schuster, A. Yala, J. H. Sohn, T. S. Jaakkola, and R. Barzilay. Conformal language modeling. In *The Twelfth International Conference on Learning Representations*, 2024.
 - S. Ravfogel, Y. Goldberg, and J. Goldberger. Conformal nucleus sampling. In *Findings of the* Association for Computational Linguistics: ACL 2023.
 - A. Z. Ren, A. Dixit, A. Bodrova, S. Singh, S. Tu, N. Brown, P. Xu, L. Takayama, F. Xia, J. Varley, Z. Xu, D. Sadigh, A. Zeng, and A. Majumdar. Robots that ask for help: Uncertainty alignment for large language model planners. In *7th Annual Conference on Robot Learning*, 2023.
 - D. Stutz, K. D. Dvijotham, A. T. Cemgil, and A. Doucet. Learning optimal conformal classifiers. In *International Conference on Learning Representations*, 2022.
 - J. Su, J. Luo, H. Wang, and L. Cheng. Api is enough: Conformal prediction for large language models without logit-access. *arXiv preprint arXiv:2403.01216*, 2024.
 - D. Tailor, A. Patra, R. Verma, P. Manggala, and E. Nalisnick. Learning to defer to a population: A meta-learning approach. In *International Conference on Artificial Intelligence and Statistics*, pages 3475–3483. PMLR, 2024.
 - Q. Tang, Z. Deng, H. Lin, X. Han, Q. Liang, B. Cao, and L. Sun. Toolalpaca: Generalized tool learning for language models with 3000 simulated cases. arXiv preprint arXiv:2306.05301, 2023.
 - H. Touvron, L. Martin, K. Stone, P. Albert, A. Almahairi, Y. Babaei, N. Bashlykov, S. Batra, P. Bhargava, S. Bhosale, et al. Llama 2: Open foundation and fine-tuned chat models. *arXiv* preprint arXiv:2307.09288, 2023.
- D. Ulmer, C. Zerva, and A. Martins. Non-exchangeable conformal language generation with nearest neighbors. In *Findings of the Association for Computational Linguistics: EACL 2024*, 2024.
- H. Vishwakarma, H. Lin, F. Sala, and R. K. Vinayak. Promises and pitfalls of threshold-based
 auto-labeling. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.
 - H. Vishwakarma, H. Lin, and R. K. Vinayak. Taming false positives in out-of-distribution detection with human feedback. In *International Conference on Artificial Intelligence and Statistics*, pages 1486–1494. PMLR, 2024.
- V. Vovk, A. Gammerman, and G. Shafer. *Algorithmic learning in a random world*, volume 29.
 Springer, 2005.
- R. Xie, R. F. Barber, and E. J. Candès. Boosted Conformal Prediction Intervals. *arXiv preprint arXiv:2406.07449*, 2024.
- 593 Y. Yang and A. K. Kuchibhotla. Selection and Aggregation of Conformal Prediction Sets. *Journal of the American Statistical Association*, 2024.

| C. Zheng, H. Zhou, F. Meng, J. Zhou, and M. Huang. Large language models are not robust multiple choice selectors. In <i>The Twelfth International Conference on Learning Representations</i> , 2024. | 594 595 | F. Ye, M. Yang, J. Pang, L. Wang, D. F. Wong, E. Yilmaz, S. Shi, and Z. Tu. Benchmarking llms via uncertainty quantification. <i>arXiv preprint arXiv:2401.12794</i> , 2024. |
|---|------------|---|
| 300 301 302 303 304 305 306 307 308 309 301 302 303 304 305 306 307 308 309 301 302 303 304 305 306 307 308 309 301 301 302 303 304 305 306 307 308 309 301 302 303 304 305 306 307 308 309 301 302 303 304 | 590 597 | C. Zheng, H. Zhou, F. Meng, J. Zhou, and M. Huang. Large language models are not robust multiple choice selectors. In <i>The Twelfth International Conference on Learning Representations</i> , 2024. |
| | 500 | |
| | 599 | |
| | 601 | |
| | 602 | |
| | 603 | |
| GG | 604 | |
| 7 60 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631< | 605 | |
| 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 621 622 623 624 625 626 627 628 629 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 6 | 606 | |
| 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 6 | 607 | |
| 609 610 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 626 627 628 629 620 621 622 623 624 625 626 627 628 629 621 622 623 624 625 626 627 628 629 621 622 623 624 625 626 627 628 629 629 6 | 608 | |
| 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 620 621 622 623 624 625 626 627 628 629 629 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 6 | 609 | |
| 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 620 621 625 626 627 628 629 629 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 6 | 610 | |
| 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 6 | 611 | |
| 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 639 640 641 642 643 644 645 646 647 | 612 | |
| 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 6 | 613 | |
| 616 617 618 619 620 621 622 623 624 625 626 627 628 629 629 620 621 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 6 | 614 | |
| 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 641 642 643 644 645 646 647 | 615 | |
| 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 616 | |
| 618 619 620 621 622 623 624 625 626 627 628 629 629 629 629 630 631 632 633 634 635 636 637 638 639 639 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 639 630 631 632 633 6 | 617 | |
| 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 618 | |
| 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 630 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 636 637 638 639 631 632 633 634 635 6 | 619 | |
| 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 620 | |
| 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 621 | |
| 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 622 | |
| 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 623 | |
| 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 624 | |
| 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 625 | |
| 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 641 642 643 644 645 646 647 | 626 | |
| 220 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 620 | |
| 625 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 620 | |
| 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 630 | |
| 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 631 | |
| 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 632 | |
| 634 635 636 637 638 639 640 641 642 643 644 645 646 647 | 633 | |
| 635 636 637 638 639 640 641 642 643 644 645 646 647 | 634 | |
| 636 637 637 638 639 640 640 641 642 642 643 645 644 645 645 646 646 647 | 635 | |
| 637 638 639 640 641 642 643 644 645 646 647 | 636 | |
| 638 639 640 641 642 643 644 645 646 647 | 637 | |
| 639 640 641 642 643 644 645 646 647 | 638 | |
| 640 641 642 643 644 645 646 647 | 639 | |
| 641 642 643 644 645 646 646 | 640 | |
| 642 643 644 645 646 647 | 641 | |
| 643 644 645 646 647 | 642 | |
| 644 645 646 647 | 643 | |
| 645 646 647 | 644 | |
| 646 647 | 645 | |
| 647 | 646 | |
| | 647 | |

648 SUPPLEMENTARY MATERIAL

649

650 The supplementary material is organized as follows. In Appendix A.1 we provide details of LLM 651 inference for MCQs. Our choice for \mathcal{G} is discussed in Appendix A.2. Additional experiments and 652 results are given in Appendix B. First, in Appendix B.1 we discuss the trade-off between coverage 653 (choice of α) in conformal prediction and its effect on CROO accuracy. Next, in Appendix B.2 we 654 explore the effectiveness of conformal prediction with CP-OPT scores in deferral applications. The Appendices B.3, B.4 and B.5, contain more detailed results for the hypotheses discussed in the main 655 paper. Appendix C provides details of the procedure used to compute statistical significance. In 656 Appendix D we provide details of datasets and give samples of prompts before and after CROQ and 657 LLM's answers. Finally, Appendix D.3 lists the hyperparameters used for learning score function 658 using CP-OPT. 659

660 661

662

663 664

665

666

670

675 676

680 681 682

683 684

685

686

687

688

689

690 691 692

METHODOLOGY AND BACKGROUND DETAILS А

A.1 DETAILS ON LLM INFERENCE IN MULTI-CHOICE QUESTION ANSWERING

We provide a formal description of the inference procedure described in the LLM Inference paragraph of Section 2.1.

667 The input prompt x is a sequence of tokens $t_1, t_2, \ldots t_n$. We run the forward pass of the autoregressive LLM (Touvron et al., 2023; Dubey et al., 2024; Abdin et al., 2024) on x to produce a set of 668 output logits: 669

$$\boldsymbol{l}_1, \boldsymbol{l}_2, \dots, \boldsymbol{l}_n \leftarrow \text{LLM}(\boldsymbol{t}_1, \boldsymbol{t}_2, \dots, \boldsymbol{t}_n) \tag{10}$$

671 Here, each logit $l_i \in \mathbb{R}^{|V|}$ expresses the likelihood of the next token after t_1, \ldots, t_j , where V is the 672 universal set of tokens (aka the alphabet) for the given LLM and |V| is its size. The last token's logits l_n are expected to have a high value for the correct answer key. We extract the logit vector $ar{l}\in\mathbb{R}^m$ 673 corresponding to the option keys as follows: 674

$$\bar{\boldsymbol{l}} := \begin{bmatrix} \boldsymbol{l}_n[Y_1], \, \boldsymbol{l}_n[Y_2], \, \dots, \, \boldsymbol{l}_n[Y_m] \end{bmatrix},\tag{11}$$

where $l_n[Y_i]$ denotes the logit value corresponding to the token Y_i in the last token's logits l_n . The 677 logits \bar{l} are converted to softmax scores s(x). The softmax score of point x and option key y is 678 denoted by s(x, y) and the predicted answer key \hat{y} corresponds to the maximum softmax value: 679

$$s(x) := \operatorname{softmax}(\boldsymbol{l}), \qquad s(x,y) := s(x)[y], \qquad \hat{y} := \underset{y \in \{Y_1, \dots, Y_m\}}{\operatorname{arg\,max}} s(x,y) \tag{12}$$

A.2 Specific choice of features and G

In practice, we want to use a flexible and easy-to-train function class for \mathcal{G} . As this is a post-hoc procedure and we want to avoid expensive fine-tuning. We use 3-layer neural networks with tanh activation as \mathcal{G} and use the LLM's logits and the penultimate layer's representations corresponding to the last token as input features to the g network. Let $z \in \mathbb{R}^{d+m}$ be the concatenation of the LLM's penultimate layer's representation (d-dimensional) and logits (m-dimensional) for the last token. Our choice of \mathcal{G} for the experiments is defined as follows,

$$\begin{split} \mathcal{G} &:= \{ g: \mathbb{R}^{d_0} \to \Delta^{m-1} \mid g(\boldsymbol{z}) := \texttt{softmax}(\boldsymbol{W}_3 \texttt{tanh}(\boldsymbol{W}_2 \texttt{tanh}(\boldsymbol{W}_1(\boldsymbol{z})))), \\ \boldsymbol{W}_1 \in \mathbb{R}^{d_0 \times d_1}, \boldsymbol{W}_2 \in \mathbb{R}^{d_1 \times d_2}, \boldsymbol{W}_3 \in \mathbb{R}^{d_2 \times m} \} \end{split}$$

693 Here, $d_0 = d + m$, $d_1 = (d + m)/2$, and $d_3 = (d + m)/4$ and Δ^{m-1} is the m - 1 dimensional 694 probability simplex. This class for \mathcal{G} is flexible enough and the resulting optimization problem is not 695 computationally prohibitive to solve. More complex (flexible) choices of \mathcal{G} could be used when we 696 can devote more compute to learning the score function. 697

698 699

700

ADDITIONAL EXPERIMENTS AND RESULTS В

This appendix contains additional results and details not included in the main paper due to length 701 constraints.

702 B.1 TRADE-OFF BETWEEN COVERAGE AND ACCURACY

704 The choice of α controls the coverage level in conformal prediction. A small α implies high coverage, 705 meaning the prediction sets contain the true options with high probability but potentially have large sizes. Thus, choosing a very small α will likely not eliminate a sufficient number of options to see any 706 noticeable improvement with CROQ. On the other hand, choosing a large α will eliminate the true 707 option from the set for a large portion of the questions, which will result in low accuracy from CROQ. 708 To study these trade-offs, we run CROQ with different values of α . The accuracy before and after 709 CROQ for a range of α values are shown in Figure 5 and Figure 4 for the Llama-3 and Phi-3 models 710 respectively. The results are as expected given the observations above: using an overly conservative 711 (small) α does not give much improvement; as we increase α , the accuracy also increases up to a 712 point, after which it starts to come down. This suggests that to optimize accuracy, a practitioner can 713 tune α for their chosen score function and setting.

714 715 716

B.2 USING CONFORMAL PREDICTION FOR DEFERRAL

717 Smaller prediction sets imply fewer deferrals in human-in-the-loop or model cascade systems. We consider a deferral procedure in which a set size cutoff is selected, and the LLM answer is only 718 retained if the set size is at or below that cutoff. For all larger sets, the question is passed to a human 719 (or a more powerful but costly model) who can answer the question correctly. Smaller sets from CP 720 are desirable for this procedure to be effective. We evaluate this procedure with logit and CP-OPT 721 scores in two settings and show the results in Figure 3. As expected, lower set size cutoffs result 722 in higher accuracy. As the set size cutoff increases, the accuracy approaches the LLM's marginal 723 accuracy, while the number of deferrals (i.e. the cost of obtaining the answer from a human or more 724 expensive model) decreases. In the top row of the figure, the differences in the set sizes between logit 725 and CP-OPT scores are not large enough to see a meaningful difference in this procedure. However, 726 in the bottom row corresponding to the Gemma-2 model and TruthfulQA dataset with 15 options, 727 we see CP-OPT scores lead to fewer deferrals in comparison to logits. Model cascades (Dohan 728 et al., 2022; Gupta et al., 2024) and deferrals to human-in-the-loop (Tailor et al., 2024; Vishwakarma et al., 2024) and more broadly selective prediction (El-Yaniv and Wiener, 2010; Fisch et al., 2022; 729 Vishwakarma et al., 2023) are useful frameworks for model deployment while ensuring safety, high 730 accuracy, and balancing the costs. Our experiments show the promise of CP with logit and CP-OPT 731 scores in this task and suggest it would be fruitful to explore this design space with CP. 732

Figure 4 shows accuracy after the CROQ procedure as a function of α for Phi-3. The results are qualitatively similar to the results for Llama-3 in the main text (Section 4.2).

All remaining results are organized by dataset. Tables for the CROQ results which illustrate accuracy changes conditional on set size are based on a confidence level of 95% (equivalently an α level of 0.05). Note that with the ToolAlpaca dataset, not all possible set sizes occur, in which case we omit the corresponding columns. For example, with 10 response options, only sets of size 8 and smaller occur.

Asterisks in the tables indicate where the difference in overall accuracy from Before to After, i.e. from baseline to after the CROQ procedure, is statistically significant at the $\alpha = 0.05$ level. (In some tables, like Table 8, none of the changes are significant.) See Appendix C for details on how statistical significance was calculated.

- 744 745
- 746
- 747
- 748
- 749 750
- 751
- 752
- 753
- 754
- 755



Figure 3: Proportion of questions deferred to a human when conformal prediction set sizes exceed a certain cutoff (left), and the corresponding LLM accuracy for questions (without revision) retained by the LLM as a function of cutoff threshold (right). In the top row (MMLU, 10 options, Phi-3-4k-mini-Instruct), the difference in deferral and accuracy is negligible, whereas in the bottom row (TruthfulQA, 15 options, gemma-2-9b-it-SimPO), CP-OPT defers fewer questions to the human while providing similar or improved accuracy for questions retained.



Figure 4: Accuracy on revised questions on the MMLU and ToolAlpaca datasets while varying miscoverage parameter α for Phi-3-4k-mini-Instruct (Phi-3) model and both scores. Smaller values of α correspond to high levels of coverage. When coverage is too large, few or no answers are eliminated, and the LLM is prompted with the same question. When coverage is low, a larger portion of answer sets no longer contain the true answer and the benefits of revision are diminished.

B.3 MMLU

Results for the experiments on the MMLU dataset are given in Tables 8 and 9, Tables 5 to 7 and Figures 6 to 8.

B.4 TRUTHFULQA

- D.4 IKUIHFULQ
- Results for the experiments on the TruthfulQA dataset are given in Tables 10 to 14 and Figures 13 and 14.



Figure 5: Accuracy on revised questions on the MMLU and ToolAlpaca datasets while varying miscoverage parameter α for Llama-3-8B-Instruct (Llama-3) model and both scores. Smaller values of α correspond to high levels of coverage. When coverage is too large, few or no answers are eliminated, and the LLM is prompted with the same question. When coverage is low, a larger portion of answer sets no longer contain the true answer or produce empty prediction sets thus resulting in diminished benefits of revision.



Figure 6: Distributions of sizes of sets obtained from CP-OPT and logit scores on MMLU dataset and Gemma-2 model.

B.5 TOOLALPACA

Results for experiments on the ToolAlpaca dataset are given in Tables 15 to 19 and Figures 10 and 11.



Figure 7: Distributions of sizes of sets obtained from CP-OPT and logit scores on MMLU dataset and Llama-3 model.



Figure 8: Distributions of sizes of sets obtained from CP-OPT and logit scores on MMLU dataset and Phi-3 model setting.

| | | | LLama-3 | | | Phi-3 | | | Gemma-2 | |
|------------|--------|--------------------|-----------------------------|---------------------|--------------------|-----------------------------|---------------------|--------------------|-----------------------------|---------------------|
| Model | # Opt. | Accuracy Before | Accuracy After (a'_2) | Gain $(a_2' = a_2)$ | Accuracy Before | Accuracy After (a'_2) | Gain $(a_2' - a_2)$ | Accuracy Before | Accuracy After (a'_2) | Gain $(a_2' - a_2)$ |
| | 4 | 64.02 | 63.67 | -0.34 | 70.27 | 69.34 | -0.93 | 68.36 | 69.56 | 1.20* |
| MMLU | 10 | 54.82 | 57.11 | 2.29* | 58.44 | 61.05 | 2.61* | 53.99 | 57.93 | 3.94* |
| | 15 | 51.99 | 54.77 | 2.78* | 53.48 | 58.15 | 4.68* | 50.78 | 51.31 | 0.52 |
| | 4 | 91.47 | 91.82 | 0.35 | 92.64 | 92.52 | -0.12 | 93.46 | 93.57 | 0.12 |
| ToolAlpaca | 10 | 85.16 | 89.02 | 3.86* | 87.62 | 91.00 | 3.39* | 88.08 | 90.42 | 2.34* |
| | 15 | 81.43 | 88.67 | 7.24* | 85.98 | 89.95 | 3.97* | 88.08 | 89.37 | 1.29 |
| | 4 | 54.43 | 55.44 | 1.01 | 69.87 | 69.87 | 0.00 | 74.94 | 76.96 | 2.03 |
| TruthfulQA | 10 | 39.24 | 42.28 | 3.04 | 55.70 | 56.20 | 0.51 | 56.46 | 60.76 | 4.30* |
| | 15 | 37.22 | 37.47 | 0.25 | 46.84 | 51.39 | 4.56* | 55.95 | 57.72 | 1.77 |

Table 4: [CROQ + CP-OPT]. Results on accuracy improvement with CROQ using CP-OPT scores. Here a_2 , and a'_2 refer to the accuracy before CROQ and after CROQ respectively. A higher gain in a setting suggests CROQ improved accuracy in that setting.







Figure 10: Distributions of sizes of sets obtained from CP-OPT and logit scores on ToolAlpaca dataset and Llama-3 model.



Figure 11: Distributions of sizes of sets obtained from CP-OPT and logit scores on ToolAlpaca dataset and Phi-3 model.







Figure 13: Distributions of sizes of sets obtained from CP-OPT and logit scores on Truthful QAdataset and Phi-3 model.



Figure 14: Distributions of sizes of sets obtained from CP-OPT and logit scores on Truthful QA dataset and Llama-3 model.

| Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
|--------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | Coverage | 82.40 | 69.04 | 80.00 | 83.56 | 81.11 | 87.45 | 86.31 | 88.60 | 90.75 | 90.45 | 94.80 | 93.75 | 98.30 | 98.15 | 100 | 94.58 |
| Logits | Fraction | 2.77 | 2.34 | 2.37 | 2.60 | 2.58 | 2.74 | 3.12 | 3.23 | 3.47 | 4.47 | 5.02 | 5.70 | 6.99 | 10.91 | 41.70 | 100 |
| | Acc. Before | 82.40 | 62.44 | 62.00 | 65.30 | 60.37 | 61.47 | 61.98 | 59.19 | 55.82 | 62.6 | 57.92 | 51.25 | 57.89 | 50.38 | 40.01 | 50.78 |
| | Acc. After | 82.40 | 65.48 | 68.50 | 65.75 | 63.13 | 58.87 | 60.08 | 57.72 | 56.85 | 58.89 | 55.08 | 51.88 | 58.06 | 49.40 | 40.01 | 50.58 |
| | Coverage | 93.10 | 94.05 | 89.83 | 89.94 | 89.34 | 90.54 | 89.74 | 90.23 | 92.40 | 94.73 | 94.70 | 94.46 | 96.77 | 97.74 | 100 | 94.58 |
| Ours | Fraction | 2.75 | 3.99 | 4.08 | 3.77 | 4.12 | 4.39 | 4.63 | 5.22 | 5.78 | 6.53 | 7.17 | 9.21 | 11.76 | 13.66 | 12.94 | 100 |
| | Acc. Before | 93.10 | 88.10 | 82.56 | 79.56 | 75.79 | 73.24 | 64.62 | 56.82 | 56.26 | 52.73 | 45.20 | 42.53 | 36.63 | 33.10 | 25.96 | 50.78 |
| | Acc. After | 93.10 | 89.58 | 82.56 | 80.82 | 73.78 | 70.81 | 60.26 | 56.14 | 57.49 | 53.27 | 46.69 | 43.94 | 40.06 | 33.80 | 25.96 | 51.31 |

Table 5: Results for CROQ on the MMLU dataset with 15 response options and Gemma-2 model.

| Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
|--------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| - | Coverage | 95.82 | 91.56 | 89.98 | 93.19 | 94.54 | 94.63 | 94.44 | 95.60 | 96.09 | 96.88 | 97.06 | 96.77 | 98.21 | 98.08 | 100 | 95.42 |
| Logits | Fraction | 8.81 | 8.58 | 7.35 | 6.97 | 5.65 | 5.74 | 5.98 | 6.21 | 6.68 | 6.46 | 6.05 | 5.89 | 5.97 | 6.17 | 7.50 | 100 |
| | Acc. Before | 95.82 | 82.16 | 72.37 | 66.95 | 55.88 | 50.62 | 50.20 | 46.08 | 40.14 | 37.32 | 34.90 | 34.68 | 30.62 | 27.88 | 24.05 | 51.99 |
| | Acc. After | 95.82 | 83.82 | 76.09 | 71.55 | 63.66 | 53.93 | 51.39 | 45.32 | 43.69 | 40.99 | 36.47 | 35.08 | 33.00 | 27.69 | 24.05 | 54.11* |
| | Coverage | 94.15 | 94.62 | 91.29 | 91.63 | 93.31 | 93.18 | 94.52 | 96.43 | 97.02 | 96.42 | 97.59 | 96.56 | 97.91 | 98.25 | 100 | 95.06 |
| Ours | Fraction | 6.69 | 8.38 | 8.58 | 7.65 | 7.99 | 8.00 | 7.80 | 6.99 | 7.17 | 6.30 | 5.90 | 5.17 | 5.12 | 4.75 | 3.51 | 100 |
| | Acc. Before | 94.15 | 87.54 | 73.58 | 65.58 | 55.57 | 51.78 | 45.81 | 46.86 | 39.90 | 31.83 | 33.00 | 28.67 | 31.32 | 21.25 | 19.59 | 51.99 |
| | Acc. After | 94.15 | 89.24 | 75.80 | 70.39 | 63.74 | 54.60 | 50.53 | 47.54 | 42.38 | 35.03 | 34.21 | 33.26 | 29.93 | 24.75 | 19.59 | 54.77* |

Table 6: Results for CROQ on the MMLU dataset with 15 response options and Llama-3 model.

| Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
|--------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | Coverage | 96.03 | 92.77 | 93.46 | 91.71 | 93.93 | 93.61 | 93.55 | 93.81 | 94.79 | 96.65 | 95.38 | 96.83 | 95.77 | 97.25 | 100 | 94.60 |
| Logits | Fraction | 11.07 | 10.34 | 8.17 | 7.73 | 7.62 | 7.80 | 7.55 | 6.52 | 6.15 | 5.32 | 5.14 | 4.87 | 4.49 | 3.88 | 3.38 | 100 |
| | Acc. Before | 96.03 | 80.48 | 69.62 | 59.14 | 53.12 | 46.27 | 42.61 | 42.08 | 37.84 | 39.51 | 36.72 | 34.15 | 23.02 | 23.55 | 21.75 | 53.48 |
| | Acc. After | 96.03 | 84.85 | 76.60 | 66.97 | 63.86 | 53.42 | 51.10 | 44.44 | 42.86 | 42.19 | 39.26 | 36.34 | 25.13 | 24.46 | 21.75 | 58.09* |
| | Coverage | 95.79 | 92.20 | 93.83 | 91.19 | 94.19 | 93.79 | 95.93 | 94.54 | 94.57 | 96.04 | 93.82 | 96.80 | 96.26 | 97.29 | 100 | 94.61 |
| Ours | Fraction | 12.40 | 9.73 | 8.08 | 7.68 | 7.56 | 7.45 | 7.00 | 6.95 | 6.55 | 5.70 | 5.57 | 5.20 | 4.76 | 3.50 | 1.86 | 100 |
| | Acc. Before | 95.79 | 80.24 | 73.86 | 60.28 | 51.33 | 49.68 | 43.90 | 41.47 | 36.41 | 31.46 | 29.42 | 29.00 | 25.69 | 21.69 | 18.47 | 53.48 |
| | Acc. After | 95.79 | 83.66 | 78.12 | 69.86 | 62.64 | 54.62 | 52.03 | 47.95 | 39.67 | 38.96 | 32.41 | 31.28 | 27.18 | 22.37 | 18.47 | 58.15* |
| | | | | | | | | | | | | | | | | | |

Table 7: Results for CROQ on the MMLU dataset with 15 response options and Phi-3 model.

| 29 | | | | | | | | | |
|----|------------|--------|-------------|-------|-------|-------|-------|----------------|--|
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | Model | Score | Set Size | 1 | 2 | 3 | 4 | Overall | |
| | | | Coverage | 89.34 | 89.94 | 93.27 | 100 | 95.16 | |
| | | Logits | Fraction | 17.71 | 17.93 | 17.11 | 47.25 | 100 | |
| | | | Acc. Before | 89.34 | 79.42 | 68.24 | 54.79 | 67.62 | |
| | C 1 | | Acc. After | 89.34 | 79.95 | 68.10 | 54.79 | 67.70 | |
| | Gemma-2 | | Coverage | 91.67 | 89.93 | 93.10 | 100 | 94.23 | |
| | | Ours | Fraction | 37.62 | 16.14 | 14.61 | 31.63 | 100 | |
| | | | Acc. Before | 91.67 | 72.50 | 57.27 | 43.64 | 68.36 | |
| | | | Acc. After | 91.67 | 75.88 | 61.74 | 43.64 | 69.56* | |
| | | | Coverage | 93.55 | 92.78 | 92.89 | 100 | 95.75 | |
| | | Logits | Fraction | 33.84 | 14.13 | 14.68 | 37.35 | 100 | |
| | | e | Acc. Before | 93.55 | 70.19 | 49.88 | 40.48 | 64.02 | |
| | | | Acc. After | 93.55 | 70.70 | 48.10 | 40.48 | 63.83 | |
| | Llama-3 | | Coverage | 93.71 | 91.83 | 93.50 | 100 | 95.57 | |
| | | Ours | Fraction | 33.21 | 15.39 | 16.63 | 34.77 | 100 | |
| | | 0415 | Acc. Before | 93.71 | 71.16 | 52.46 | 38.02 | 64.02 | |
| | | | Acc. After | 93.71 | 70.01 | 51.46 | 38.02 | 63.67 | |
| | | | Coverage | 94.75 | 91.48 | 93.17 | 100 | 94.65 | |
| | | Logits | Fraction | 37.30 | 22.86 | 21.20 | 18.64 | 100 | |
| | | Logito | Acc. Before | 94 75 | 70.25 | 52.69 | 41.31 | 70.27 | |
| | | | Acc. After | 94.75 | 66.93 | 50.67 | 41.31 | 69.08 | |
| | Phi-3 | | Coverage | 03.63 | 90.61 | 94 17 | 100 | 94.35 | |
| | | Ours | Fraction | 1136 | 21.10 | 17 71 | 10.82 | 100 | |
| | | Ours | riacuon | +1.50 | 21.10 | 1/./1 | 19.03 | 100 | |
| | | | Acc Rofors | 03 62 | 67 28 | 57 87 | 10.22 | 70.27 | |
| | | | Acc. Before | 93.63 | 67.38 | 52.82 | 40.22 | 70.27 60.34 | |

Table 8: Results for CROQ on the MMLU dataset with 4 response options.

| Mod | el Sco | re s | set Size | | 1 | 2 | 3 | 4 | 5 | | 6 | 7 | 8 | 9 | 10 | J |
|--|--|--|--|---|---|--|--|--|---|--|--|---|--|---|---|--|
| | | C | overage | e 78 | 3.80 | 79.03 | 84.92 | 88.56 | 5 85.3 | 0 92 | 2.64 | 94.09 | 96.41 | 97.22 | 2 10 | 0 |
| | Log | its F | raction | ı 2. | .97 | 3.90 | 4.25 | 4.77 | 5.3 | 3 5 | .80 | 7.03 | 9.59 | 14.10 |) 42. | 26 |
| | | Ac | c. Befo | re 78 | 3.80 | 73.86 | 74.02 | 68.41 | 62.3 | 6 6 7 | .69 | 61.49 | 58.42 | 51.94 | 41. | 81 |
| Comm | | A | cc. Afte | r 78 | 3.80 | 76.90 | 75.98 | 72.39 | 62.3 | 6 66 | 5.67 | 60.14 | 57.67 | 51.68 | 41. | 81 |
| Gennin | Id-2 — | C | overage | e 90 |).79 | 92.27 | 88.31 | 90.54 | 89.8 | 0 91 | .30 | 92.05 | 95.60 | 97.49 |) 10 | 0 |
| | Ou | rs H | raction | ı 12 | 2.89 | 8.90 | 7.31 | 6.65 | 6.40 |) 7. | .23 | 8.36 | 8.90 | 10.41 | 22. | 96 |
| | | Ac | c. Befo | re 90 |).79 | 84.93 | 69.97 | 66.07 | 54.1 | 7 48 | 3.60 | 42.76 | 40.00 | 37.74 | 31. | 27 |
| | | A | cc. Afte | r 90 |).79 | 89.20 | 79.87 | 75.00 | 64.0 | 1 55 | 5.34 | 47.02 | 45.33 | 40.59 | 31. | 27 |
| | | C | overage | e 94 | 4.55 | 91.96 | 91.73 | 94.09 | 94.9 | 4 97 | 7.19 | 97.32 | 97.72 | 99.32 | 2 10 | 0 |
| | Log | its H | raction | 14 | 1.36 | 10.92 | 8.76 | 7.63 | 7.04 | 4 8 | .03 | 8.40 | 9.90 | 10.53 | 3 14. | 43 |
| | | Ac | c. Befo | re 94 | 1.55 | 80.43 | 65.99 | 57.54 | 51.4 | 3 47 | .56 | 37.71 | 35.13 | 34.84 | 31. | 41 |
| | | A | cc. Afte | r 94 | 1.55 | 80.33 | 69.51 | 60.96 | 5 53.2 | 9 49 | .93 | 42.37 | 36.21 | 35.74 | 31. | 41 |
| Llama | a-3 — | 0 | overage | e 94 | 1 80 | 91 95 | 92.42 | 93.98 | 3 94 9 | 5 96 | 5.61 | 97.64 | 97.96 | 98.68 | 3 10 | 0 |
| | O | | raction | | 3 92 | 11 50 | 10.80 | 10.44 | 115 | 1 10 |) 16 | 10.55 | 8 71 | 7 20 | 52 | <u>,</u> |
| | 01 | | c Refe | ro 0/ | 1.80 | 79.67 | 68.02 | 52.61 | 45.0 | 5 40 | 10 | 35 55 | 33.65 | 28.67 | 1 30 | 87 |
| | | | ce Afte | r 04 | 1 80 | 79.05 | 71 76 | 55.57 | 7 /00 | 0 47 | 76 | 40.82 | 35.03 | 20.07 | 30. | 52 87 |
| | | A | over- | a 94 | | 01.00 | 00.76 | 04.21 | | 0 05 | | 04.07 | 06 17 | 05 50 | 10 | 04 |
| | Ţ | · | overage | e 95 | 0.75 | 91.02 | 90.70 | 94.21 | 93.9 | 0 93 | 0.39 | 94.07 | 90.17 | 95.52 | 2 10 | 0 |
| | Log | | raction | 1 1/ | .8/ | 14.28 | 50.14 | 55.02 | 5 11.0 15.5 | δ δ. | .88 | 8.01 | 7.12 | 20.27 | 3.1 | 9 |
| | | | c. Beio | re 95 | 5.75 | /6.56 | 59.14 | 55.04 | 2 45.5 | 0 43 | 5.72 10 | 37.19 | 33.0 | 30.27 | 26. | 55 |
| Phi- | 3 — | A | cc. Afte | er 95 | 5.75 | 79.05 | 65.56 | 59.77 | 51.1 | 8 4/ | .19 | 42.37 | 32.83 | 32.29 | 26. | 55 |
| | | C | overage | e 95 | 0.85 | 90.94 | 90.94 | 94.05 | 93.5 | <u> </u> | k. / I | 93.94 | 94.96 | 96.71 | 10 | 0 |
| | | | · · · | | | | 1114 | 10.98 | 5 106 | 5 IU | 0.09 | × 4 I | 1.50 | 5.06 | | |
| | Ou | rs H | raction | 1 20 | J.02 | 12.71 | 11.15 | 10.70 | | | | 0.11 | | | 5.0 | 56 |
| | Ou Ta | $\frac{\mathbf{F}}{\mathbf{A}}$ | Fraction cc. Befor cc. Afte Resul | re 20 re 95 r 95 | 5.85 5.85 r CR | 73.86 76.84 | 63.75 68.66 | 54.38 59.68 | 46.3 3 50.6 | 8 40 1 44 aset | 0.47 1.12 with | 36.53 38.50 | 32.68 34.80 | 26.76 26.06 | ions. | 56 30 30 |
| | Ou Ta | the second seco | Fraction cc. Befor cc. Afte Resul | i 20 re 95 r 95 Its foi | 5.85 5.85 5.85 | 73.86 76.84 | 63.75 68.66 | 54.38 59.68 | 46.3 3 50.6 | 8 40 1 44 | 0.47 1.12 with | 36.53 38.50 | 32.68 34.80 | 26.06 26.06 | 5 26. | 56 30 30 |
| Score | Ou Ta Set Size | $\frac{\mathbf{F}}{\mathbf{A}}$ | Fraction cc. Befor cc. Afte Resul | 1 20 re 95 r 95 Its for | 5.85 5.85 r CR 4 | 12.71 73.86 76.84 OQ 0. | 6 11.13 63.75 68.66 6 | 54.38 59.68 MML | 8 1000 1000 1000 1000 1000 1000 1000 10 | 8 40 1 44 aset | 0.47 1.12 with | 36.53 38.50 10 re | 32.68 34.80 espons | 26.06 26.06 se opt | 5 26. 5 26. ions. | 30 30 30 |
| Score | Ou Ta Set Size Coverage | $\frac{\mathbf{H}}{\mathbf{A}\mathbf{C}}$ $\frac{\mathbf{H}}{\mathbf{A}\mathbf{C}}$ $\mathbf{A}\mathbf{C}$ $$ | Praction cc. Before cc. After Result 2 93.75 | i 20 re 95 rr 95 Its for <u>3</u> 92.86 | 5.85 5.85 r CR 4 100 | 73.86 76.84 OQ 0 5 100 | 63.75 68.66 n the <u>6</u> 95.00 | 7 94.12 | 8 76.92 | 8 40 1 44 aset | 0.47 1.12 with | 36.53 38.50 10 re <u>11</u> 100 | 32.68 34.80 espons 12 88.00 | 26.06 26.06 Se opt | ions. | 30 30 30 |
| Score Logits _ | Ou Ta Set Size Coverage Fraction | $\frac{\mathbf{F}}{\mathbf{A}}$ | 2 93.75 4.05 | a 20 re 95 rr 95 Its for <u>3</u> 92.86 3.54 | 4 100 1.77 | 73.86 76.84 OQ 0 5 100 1.77 | 63.75 68.66 n the <u>6</u> 95.00 5.06 | 7 94.12 4.30 | 8 76.92 3.29 | 8 40 1 44 caset 9 80.95 5.32 | 10 94.44 4.56 | 10 re | 32.68 34.80 espons 12 88.00 6.33 | 26.06 26.06 se opt | 14 100 11.14 | 56 30 30 1 1 35 |
| Score Logits | Ou Ta Set Size Coverage Fraction Acc. Befor | I Ac Ac Ac ble 9: 1 100 1.52 100 1.92 | 2 93.75 4.05 93.75 | a 20 re 95 rr 95 lts for 3 92.86 3.54 92.86 92.86 92.86 | 4 100 1.77 100 | 73.86 76.84 0Q 0 5 100 1.77 85.71 85.71 | 63.75 68.66 n the <u>6</u> 95.00 5.06 80.00 | 7 94.12 4.30 76.47 | 8 76.92 3.29 46.15 53 85 | 8 40 1 44 aset 9 80.95 5.32 47.62 57.14 | 0.47 1.12 with 10 94.44 4.56 61.11 55.56 | 10 re | 32.68 34.80 espons 12 88.00 6.33 48.00 | 26.76 26.06 36 opt 13 88.00 6.33 32.00 | 14 100 11.14 47.73 45.45 | 56 30 30 30 1 1 35 46 |
| Score Logits _ _ | Ou Ta Set Size Coverage Fraction Acc. Befor Acc. Arg | I I 1 100 1.52 100 98.00 98.00 | z 93.75 4.05 93.75 93.75 93.75 93.75 93.75 | a 200 re 95 rr 95 lts for <u>3</u> 92.86 3.54 92.86 92.86 92.86 90.00 | 4 100 1.77 100 93.33 | 73.86 76.84 0Q 0 5 100 1.77 85.71 85.71 90.91 | 63.75 68.66 n the <u>6</u> 95.00 5.06 80.00 85.00 91.67 | 7 94.12 4.30 76.47 82.35 92.86 | 8 76.92 3.29 46.15 53.85 94.44 | 8 40 1 44 aset 80.95 5.32 47.62 57.14 93.33 | 10 94.44 4.56 61.11 55.56 95.45 | 36.53 38.50 10 re 11 100 5.82 56.52 52.17 89.47 | 32.68 34.80 25pons 25pons 88.00 6.33 48.00 48.00 96.97 | 26.76 26.06 3e opt 13 88.00 6.33 32.00 40.00 97.37 | 14 100 11.14 47.73 45.45 100 | 56 30 30 30 1 1 35 46 46 1 |
| Score Logits _ Ours - | Ou Ta Set Size Coverage Fraction Acc. Befor Acc. Aefor Coverage Fraction | I I 1 100 1.52 100 98.00 12.66 | 2 93.75 4.05 93.75 93.75 95.65 5.82 | a 20 re 95 r 95 r 95 lts for 3 92.86 3.54 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 | 4 100 1.77 100 93.33 3.80 | 73.86 76.84 76.94 | 63.75 68.66 n the | 7 94.12 4.30 76.47 82.35 92.86 3.54 | 8 76.92 3.29 46.15 53.85 94.44 4.56 | 8 40 1 44 aset 9 80.95 5.32 47.62 57.14 93.33 3.80 | 10 94.44 4.56 61.11 55.56 95.45 5.57 | 36.53 38.50 10 re 11 100 5.82 56.52 52.17 89.47 4.81 | 32.68 34.80 espons 12 88.00 6.33 48.00 48.00 96.97 8.35 | 26.76 26.06 36 opt 38.00 6.33 32.00 40.00 97.37 9.62 | 14 100 11.14 47.73 45.45 100 | 56 30 30 30 1 1 35 46 46 1 18 |
| Score Logits _ Ours _ | Ou Ta Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fraction Acc. Befor | I Ac Ac Ac Ac Ac ble 9: I I 100 I 1.52 100 I 100 98.00 I 12.66 98.00 | 2 93.75 4.05 93.75 93.75 93.75 93.75 93.65 5.82 95.65 | a 20 re 95 rr 95 ar 95 sr 95 display="block">display="block" 3 92.86 3.54 92.86 92.86 90.00 2.53 90.00 2.53 90.00 | 4 100 1.77 100 93.33 3.80 73.33 | 73.86 76.84 0Q 0 5 100 1.77 85.71 85.71 90.91 2.78 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.91 810 | 63.75 68.66 n the 95.00 5.06 80.00 85.00 91.67 3.04 50.04 | 7 94.12 4.30 76.47 82.35 92.86 3.54 92.86 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 | 9 88.055 5.32 47.62 57.14 93.33 3.80 60.00 | 10 94.44 4.56 61.11 55.56 95.45 5.57 63.64 | 10 re 10 re 11 re 10 re 11 100 5.82 56.52 52.17 89.47 4.81 47.37 52.77 | 32.68 34.80 25pons 25pons 25pons 288.00 6.33 48.00 48.00 96.97 8.35 39.39 42.55 | 26.76 26.06 se opt 13 88.00 6.33 32.00 40.00 97.37 9.62 31.58 24.52 | 14 100 11.14 47.73 45.45 100 10.13 32.50 | 1 30 30 30 30 1 1 1 35 46 46 1 1 8 28 28 |
| Score Logits - Ours - able 1 | Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fraction Acc. After 10: Res | I Ac Ac | 2 93.75 4.05 93.75 93.75 93.75 93.75 93.75 93.65 5.82 91.30 | i 200 re 95 r 95 its for 92.86 3.54 92.86 90.00 2.53 90.00 90.00 2.53 90.00 0.00 | 4 100 1.77 100 100 93.33 3.80 73.33 80.00 Dn th | 12.71 73.86 76.84 OQ or 5 100 1.77 85.71 85.71 90.91 2.78 81.82 81.82 e True | 63.75 68.66 n the 95.00 5.06 80.00 85.00 91.67 3.04 50.00 58.33 | 7 54.38 59.66 MMIL 99.12 4.30 76.47 82.35 92.86 3.54 92.86 92.86 92.86 92.86 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 60.00 with | 10 94.44 4.56 61.11 55.56 95.45 5.57 63.64 72.73 | 10 re 10 re 11 re 10 re 11 100 5.82 56.52 52.17 89.47 4.81 47.37 52.63 respo | 32.68 34.80 25pons 25pons 25pons 288.00 6.33 48.00 96.97 8.35 39.39 42.42 20nse o | 26.76 26.06 26.06 se opt 6.33 32.00 40.00 97.37 9.62 31.58 36.84 ption | 14 100 11.14 47.73 45.45 100 10.13 32.50 32.50 S anc | 56 30 30 30 30 30 30 30 30 30 30 30 30 30 |
| Score Logits Ours àble nodel | Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fraction 10: Res Set Size | I Ac Del De Ac Ac Ac Ac De De De De De De De De De De <td>2 93.75 4.05 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75</td> <td>a 200 re 95 r 95 r 95 lts for 3 92.86 92.86 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.60 90.00 0.00 90.00</td> <td>4 100 1.77 100 100 93.33 3.80 73.33 80.00 Dn th</td> <td>12.71 73.86 76.84 OQ or 5 100 1.77 85.71 90.91 2.78 81.82 81.82 e True</td> <td>63.75 68.66 n the 6 95.00 5.06 80.00 85.00 91.67 3.04 50.00 58.33 thful(</td> <td>7 94.12 4.30 76.47 92.86 3.54 92.86 92.86 92.86 92.86 92.86 92.86</td> <td>8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 ataset</td> <td>9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 with</td> <td>10 94.44 4.56 61.11 55.56 95.45 5.57 63.64 72.73 n 15</td> <td>10 re 10 re 10 re 11 100 5.82 56.52 52.17 89.47 4.81 47.37 52.63 respondent</td> <td>32.68 34.80 34.80 55pons 55pons 55pons 55pons 6.33 48.00 96.97 8.35 39.39 42.42 50nse o</td> <td>26.76 26.06 26.06 se opt 388.00 6.33 32.00 40.00 97.37 9.62 31.58 36.84 ption</td> <td>14 100 11.14 47.73 32.50 32.50 S ance</td> <td>1 30 30 30 30 30 30 30 30 30 30</td> | 2 93.75 4.05 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 93.75 | a 200 re 95 r 95 r 95 lts for 3 92.86 92.86 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.60 90.00 0.00 90.00 | 4 100 1.77 100 100 93.33 3.80 73.33 80.00 Dn th | 12.71 73.86 76.84 OQ or 5 100 1.77 85.71 90.91 2.78 81.82 81.82 e True | 63.75 68.66 n the 6 95.00 5.06 80.00 85.00 91.67 3.04 50.00 58.33 thful(| 7 94.12 4.30 76.47 92.86 3.54 92.86 92.86 92.86 92.86 92.86 92.86 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 ataset | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 with | 10 94.44 4.56 61.11 55.56 95.45 5.57 63.64 72.73 n 15 | 10 re 10 re 10 re 11 100 5.82 56.52 52.17 89.47 4.81 47.37 52.63 respondent | 32.68 34.80 34.80 55pons 55pons 55pons 55pons 6.33 48.00 96.97 8.35 39.39 42.42 50nse o | 26.76 26.06 26.06 se opt 388.00 6.33 32.00 40.00 97.37 9.62 31.58 36.84 ption | 14 100 11.14 47.73 32.50 32.50 S ance | 1 30 30 30 30 30 30 30 30 30 30 |
| Score Logits - - - - - - - - - - - - - - - - - - - | Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fraction Acc. After 10: Res Set Size Coverage Eraction | I Ac Ac Ac Ac Ac ble 9: Image: Second se | 2 93.75 93.75 4.05 93.75 95.65 5.82 95.65 91.30 0 CCR 2 75.00 2.03 | a 200 re 95 r 95 r 95 lts for 92.86 3.54 92.86 90.00 2.53 90.00 90.00 000 OQQ 0 3 90.00 2.53 90.00 | 4 100 1.77 100 100 93.33 3.80 73.33 80.00 Dn th 4 77.78 2.28 | 73.86 76.84 76.84 76.84 76.84 76.84 76.84 700 1.77 85.71 85.71 85.71 85.71 85.71 85.71 85.71 85.71 85.71 85.71 85.71 90.91 2.78 81.82 81.82 81.82 81.82 81.82 81.82 | 63.75 68.66 n the 95.00 5.06 80.00 91.67 3.04 50.00 58.33 tthful(6 76.92 3.29 | 7 94.12 4.30 76.47 82.35 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 93.86 93.86 94.86 94.86 94.86 95.87 95.87 95.86 95.87 95.87 95.86 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 95.87 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 ataset | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 with | 10 94.44 4.56 61.11 55.56 5.57 63.64 72.73 n 15 | 10 respondent | 32.68 34.80 espons 88.00 6.33 48.00 96.97 8.35 39.39 42.42 onse o | 26.76 26.06 26.06 38.00 6.33 32.00 40.00 97.37 9.62 31.58 36.84 97.73 11.14 | 14 100 11.14 45.45 100 10.13 32.50 S and 14 100 10.13 32.50 S and | 56 30 30 30 30 30 30 30 30 30 30 |
| Score Logits - - - - - - - - - - - - - - - - - - - | Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fraction Acc. After 10: Res Set Size Coverage Fraction Acc. Befor | I Ac Ac Ac Ac Ac Ac Ac ble 9: I 100 I.52 100 I.100 98:00 I.2.66 98:00 I.2.7 100 I.2.7 80:00 | 2 93.75 4.05 93.75 95 95 95 95 95 95 | a 22(2) re 95 r 95 s 92.86 3.54 92.86 92.86 90.00 2.53 90.00 OQQ C 90.00 2.53 90.00 2.53 80.00 3.54 | 4 100 1.77 100 100 93.33 3.80 73.33 80.00 Dn th 4 77.78 2.28 66.67 | 73.86 76.84 76.84 76.84 76.84 76.84 76.84 76.84 700 1.77 85.71 85.71 85.71 85.71 85.71 90.91 2.78 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 | 63.75 68.66 n the 95.00 5.06 80.00 91.67 3.04 50.00 58.33 tthful(6 76.92 3.29 38.46 | 7 94.12 4.30 76.47 82.35 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.87 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 93.86 94.86 94.86 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.82 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85 95.85959595959595959595 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 ataset | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 60.00 with 9 100 7.09 50.0 | 10.47 k.12 with 94.44 4.56 61.11 55.56 63.64 72.73 n 15 10 95.12 10.38 46.34 | 10 respondent | 32.68 34.80 espons 88.00 6.33 48.00 96.97 8.35 39.39 42.42 90.59 6.84 29.59 6.84 29.63 | 26.76 26.06 26.06 38 opt 388.00 6.33 32.00 97.37 9.62 31.58 36.84 97.73 31.14 22.73 | 14 100 11.14 47.73 32.50 S and 10 10.13 32.50 S and 101 103 100 10.13 100 10.13 15.00 | 56 30 30 30 30 30 1 1 1 1 35 46 46 1 1 1 8 28 28 28 1 0 1 1 1 1 1 7 22 |
| Score Logits Cours Cours Cable Cable Score Logits Cable Cabl | Set Size Coverage Fraction Acc. Befor Acc. After 10: Res Set Size Coverage Fraction Acc. After | I Ac Ac Ac ble 98 I 100 I 98.00 I 12.66 I 98.00 I 98.00 I 180.00 I 1.27 80.00 1.27 80.00 80.00 | 2 93.75 4.05 93.75 | a 200 re 95 r 95 r 95 lts for 3 92.86 92.86 92.86 92.86 90.00 2.53 90.00 OQQ C 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 90.00 | 4 100 1.77 100 100 93.33 80.00 Dn th 4 77.78 2.28 66.67 66.67 | 73.86 76.84 76.84 76.84 76.84 76.84 76.84 76.84 700 1.77 85.71 90.91 2.78 81.82 81.82 81.82 81.82 81.82 81.82 81.82 61.54 | 63.75 68.66 n the 95.00 5.06 80.00 91.67 3.04 50.00 58.33 thful(6 76.92 3.29 38.46 38.46 | 7 94.12 4.30 76.47 82.35 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 93.87 1111111111111 | 8 76.92 3.29 46.15 53.85 94.44 4.56 61.11 61.11 ataset 8 95.24 5.32 5.32 5.32 | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 60.00 with 9 100 7.09 50.0 46.43 | 10 94.44 4.56 61.11 55.56 63.64 72.73 n 15 10 95.12 10.38 46.34 43.90 | 10 respondent | 32.68 34.80 espons 88.00 6.33 48.00 96.97 8.35 39.39 42.42 90.58 00 5.84 29.63 29.63 | 26.76 26.06 26.06 38.00 6.33 32.00 97.37 31.58 36.84 97.73 31.14 22.73 11.14 22.73 | 14 100 11.14 47.73 32.50 32.50 8 and 10.13 32.50 8 and 10.13 15.00 10.13 15.00 | 56 30 30 30 30 30 1 1 1 1 355 466 466 1 1 8 28 28 1 1 1 1 1 1 1 1 1 22 22 22 |
| Score Ours able score cogits able able able able able able able able | Set Size Coverage Fraction Acc. Befor Acc. After 10: Res 10: Res Set Size Coverage Fraction Acc. After Coverage | I Ac Ac Ac Bolo 1.27 80.00 1.27 80.00 1.27 80.00 0 No 1.00 | 2 93.75 4.05 93.75 | a 200 re 95 r 95 r 95 lts for 3 92.86 92.86 92.86 92.86 90.00 2.53 90.00 OQQ C 3 90.00 2.53 90.00 OQQ C 3 90.00 2.53 80.00 90.00 2.53 80.00 90.00 0 0 | 4 100 1.77 100 1.77 100 93.33 80.00 Dn th 4 77.78 66.67 0 | 73.86 76.84 76.84 76.84 76.84 76.84 76.84 700 1.77 85.71 90.91 2.78 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 81.82 61.54 100 | 63.75 68.66 n the 95.00 5.06 80.00 91.67 3.04 50.00 58.33 thful(6 76.92 3.29 38.46 38.46 87.50 | 7 94.12 4.30 76.47 82.35 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 92.86 93.82 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.87 60.8760606060606060606060 | 8 76.92 3.29 46.15 53.85 94.41 61.11 61.11 ataset 8 95.24 5.32 57.14 52.38 93.94 | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 60.00 60.00 60.00 60.00 7.09 50.0 46.43 91.30 | 10 94.44 4.56 61.11 55.56 63.64 72.73 n 15 | 10 respondent | 32.68 34.80 34.80 espons 88.00 6.33 48.00 96.97 8.35 39.39 42.42 96.97 8.35 39.39 42.42 90.58 0 96.97 8.35 39.39 42.42 9.63 92.59 6.84 29.63 29.63 95.16 | 26.76 26.06 26.06 38 opt 38.00 6.33 32.00 97.37 31.58 36.84 97.73 31.58 36.84 97.73 11.14 22.73 11.14 22.73 18.18 96.00 | 14 100 11.14 47.73 32.50 32.50 s 100 10.13 32.50 s and 100 10.13 32.50 s and 100 10.13 15.00 100 100 | 56 30 30 30 30 1 1 1 1 1 35 46 46 10 11 18 28 28 28 1 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 |
| Score Logits Cours Cable Score Logits Curs Cable | Set Size Coverage Fraction Acc. Befor Acc. After Coverage Fracton Acc. After 10: Res Set Size Coverage Fraction Acc. After Coverage Fraction | I Ac Ac Ac ble 98 I 100 98.00 98.00 I 12.66 98.00 98.00 I 12.66 80.00 1.27 80.00 1.27 80.00 80.00 I 0 | 2 93.75 4.05 93.75 | i 200 re 95 r 95 its for g2.86 3.54 92.86 90.00 2.53 90.00 00.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 2.53 90.00 0.53 | 4 100 1.77 100 1.77 100 93.33 80.00 Dn th 4 77.78 2.28 66.67 0 0.25 | 73.86 76.84 76.84 76.84 76.84 76.84 76.84 76.84 81.00 1.77 85.71 90.91 2.78 81.82 81 | 6 6 6 6 6 6 9 5.06 80.00 9 5.06 80.00 9 5.06 80.00 9 5.06 80.00 9 5.06 80.00 9 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 5.06 80.00 58.33 81.41 6 76.92 3.29 38.46 87.50 2.03 87.50 2.03 87.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.50 80.00 | 7 94.12 4.30 76.47 82.35 92.86 92.85 | 8 76.92 3.29 46.15 53.85 94.41 61.11 | 9 80.95 5.32 47.62 57.14 93.33 3.80 60.00 60.00 60.00 60.00 60.00 60.00 7.09 50.0 46.43 91.30 11.65 | 10 94.44 4.56 61.11 55.56 95.45 5.57 6.364 72.73 n 15 10 95.12 10.38 46.34 43.90 94.37 1.797 | 10 respondent | 32.68 34.80 34.80 espons 88.00 6.33 48.00 96.97 8.35 39.39 42.42 96.97 8.35 39.39 42.42 90.58 0 96.97 8.35 39.39 42.42 9.63 92.59 6.84 29.63 95.16 15.70 | 26.76 26.06 26.06 36 opt 38.00 6.33 32.00 97.37 9.62 31.58 36.84 ption 13 97.73 11.14 22.73 11.14 22.73 18.18 96.00 12.66 | 14 100 11.14 47.73 32.50 32.50 S and 100 10.13 32.50 S and 100 10.13 32.50 S and 10.13 15.00 10.13 15.00 100 7.09 | 56 30 30 30 30 1 1 1 1 35 46 46 1 1 1 1 1 1 1 1 2 2 2 2 2 2 2 1 1 2 2 |

Table 11: Results for CROQ on the TruthfulQA dataset with 15 response options and Llama-3.

| Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
|--------|-------------|---|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | Coverage | 0 | 0 | 88.89 | 90.91 | 85.71 | 82.61 | 95.45 | 85.71 | 96.43 | 100 | 92.86 | 100 | 100 | 97.50 | 100 | 95.44 |
| Logits | Fraction | 0 | 0 | 2.28 | 2.78 | 5.32 | 5.82 | 5.57 | 5.32 | 7.09 | 7.09 | 10.63 | 9.11 | 12.15 | 10.13 | 16.71 | 100 |
| | Acc. Before | 0 | 0 | 77.78 | 90.91 | 52.38 | 56.52 | 63.64 | 61.9 | 60.71 | 50.00 | 35.71 | 33.33 | 50.00 | 30.0 | 34.85 | 46.84 |
| | Acc. After | 0 | 0 | 77.78 | 90.91 | 52.38 | 60.87 | 63.64 | 57.14 | 57.14 | 57.14 | 33.33 | 27.78 | 52.08 | 27.50 | 34.85 | 46.33 |
| | Coverage | 0 | 100 | 100 | 88.89 | 93.33 | 91.67 | 100 | 85.00 | 96.77 | 95.24 | 95.65 | 98.18 | 98.33 | 100 | 100 | 96.46 |
| Ours | Fraction | 0 | 0.76 | 1.01 | 2.28 | 3.80 | 6.08 | 8.35 | 5.06 | 7.85 | 10.63 | 11.65 | 13.92 | 15.19 | 9.37 | 4.05 | 100 |
| | Acc. Before | 0 | 100 | 100 | 77.78 | 60.00 | 62.50 | 66.67 | 45.00 | 58.06 | 45.24 | 47.83 | 36.36 | 30.00 | 37.84 | 31.25 | 46.84 |
| | Acc. After | 0 | 100 | 100 | 77.78 | 66.67 | 62.50 | 72.73 | 45.00 | 58.06 | 57.14 | 50.00 | 43.64 | 36.67 | 40.54 | 31.25 | 51.39* |
| - | | | | | | | | | | | | | | | | | |

Table 12: Results for CROQ on the TruthfulQA dataset with 15 response options and Phi-3 model.

| 58 | | | | | | | | | | | | | | |
|----|----------|--------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| 59 | Model | Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | Overall |
| 50 | | | Coverage | 100 | 94.12 | 100 | 94.12 | 87.10 | 90.91 | 90.62 | 91.11 | 95.45 | 100 | 95.44 |
| 1 | | Logits | Fraction | 4.56 | 4.30 | 3.04 | 4.30 | 7.85 | 5.57 | 8.10 | 11.39 | 16.71 | 34.18 | 100 |
| | | | Acc. Before | 100 | 94.12 | 100 | 82.35 | 70.97 | 63.64 | 56.25 | 53.33 | 53.03 | 37.04 | 56.46 |
| | Gemma-2 | | Acc. After | 100 | 94.12 | 100 | 82.35 | 70.97 | 59.09 | 56.25 | 51.11 | 54.55 | 37.04 | 56.20 |
| | | | Coverage | 97.94 | 100 | 92.86 | 89.47 | 96.15 | 91.67 | 100 | 93.55 | 97.83 | 100 | 97.22 |
| | | Ours | Fraction | 24.56 | 6.08 | 3.54 | 4.81 | 6.58 | 6.08 | 9.37 | 7.85 | 11.65 | 19.49 | 100 |
| | | | Acc. Before | 97.94 | 91.67 | 85.71 | 52.63 | 61.54 | 66.67 | 54.05 | 19.35 | 32.61 | 14.29 | 56.46 |
| | | | Acc. After | 97.94 | 95.83 | 71.43 | 89.47 | 73.08 | 66.67 | 59.46 | 29.03 | 39.13 | 14.29 | 60.76* |
| | | | Coverage | 92.86 | 93.75 | 68.97 | 95.00 | 86.21 | 91.18 | 97.56 | 96.49 | 100 | 100 | 94.43 |
| | | Logits | Fraction | 3.54 | 4.05 | 7.34 | 5.06 | 7.34 | 8.61 | 10.38 | 14.43 | 16.46 | 22.78 | 100 |
| | | | Acc. Before | 92.86 | 81.25 | 55.17 | 55.00 | 51.72 | 41.18 | 41.46 | 26.32 | 30.77 | 23.33 | 39.24 |
| | Llama-3 | | Acc. After | 92.86 | 87.50 | 55.17 | 65.00 | 58.62 | 38.24 | 34.15 | 31.58 | 33.85 | 23.33 | 40.76 |
| | Eluliu 5 | | Coverage | 92.31 | 90.00 | 70.83 | 91.89 | 95.56 | 92.00 | 92.11 | 97.14 | 100 | 100 | 93.42 |
| | | Ours | Fraction | 3.29 | 2.53 | 6.08 | 9.37 | 11.39 | 12.66 | 19.24 | 17.72 | 9.87 | 7.85 | 100 |
| | | | Acc. Before | 92.31 | 70.00 | 54.17 | 56.76 | 51.11 | 44.00 | 31.58 | 28.57 | 20.51 | 16.13 | 39.24 |
| | | | Acc. After | 92.31 | 80.00 | 58.33 | 72.97 | 55.56 | 50.00 | 30.26 | 28.57 | 20.51 | 16.13 | 42.28 |
| | | | Coverage | 100 | 100 | 94.44 | 100 | 96.55 | 89.47 | 100 | 100 | 100 | 100 | 98.48 |
| | | Logits | Fraction | 1.01 | 3.29 | 4.56 | 5.82 | 7.34 | 9.62 | 10.38 | 13.16 | 17.22 | 27.59 | 100 |
| | | | Acc. Before | 100 | 100 | 83.33 | 69.57 | 65.52 | 55.26 | 60.98 | 51.92 | 50.0 | 42.20 | 55.70 |
| | Phi-3 | | Acc. After | 100 | 100 | 88.89 | 69.57 | 65.52 | 55.26 | 51.22 | 51.92 | 47.06 | 42.20 | 54.43 |
| | | | Coverage | 100 | 86.96 | 88.89 | 90.91 | 85.71 | 95.45 | 96.08 | 100 | 97.44 | 100 | 95.70 |
| | | Ours | Fraction | 7.59 | 5.82 | 4.56 | 5.57 | 7.09 | 11.14 | 12.91 | 16.20 | 19.75 | 9.37 | 100 |
| | | | Acc. Before | 100 | 78.26 | 83.33 | 72.73 | 53.57 | 65.91 | 49.02 | 45.31 | 43.59 | 24.32 | 55.70 |
| | | | Acc. After | 100 | 78.26 | 77.78 | 72.73 | 60.71 | 61.36 | 52.94 | 45.31 | 44.87 | 24.32 | 56.20 |
| | | | | | | | | | | | | | | |

Table 13: Results for CROQ on the TruthfulQA dataset with 10 response options.

| 1190 | | | | | | | | |
|------|---------|--------|---------------------------|-------|-----------------------|-----------------------|----------------|----------------|
| 1191 | | | | | | | | |
| 1192 | | | | | | | | |
| 1193 | | | | | | | | |
| 1194 | | | | | | | | |
| 1195 | | | | | | | | |
| 1196 | | | | | | | | |
| 1197 | | | | | | | | |
| 1198 | | | | | | | | |
| 1199 | | | | | | | | |
| 1200 | | | | | | | | |
| 1201 | Model | Score | Set Size | 1 | 2 | 3 | 4 | Overall |
| 1202 | | | Coverage | 95.00 | 93.33 | 89.58 | 100 | 96.46 |
| 1203 | | Logits | Fraction | 30.38 | 11.39 | 12.15 | 46.08 | 100 |
| 1204 | | | Acc. Before | 95.00 | 84.44 | 68.75 | 60.44 | 74.68 |
| 1205 | | | Acc. After | 95.00 | 86.67 | 68.75 | 60.44 | 74.94 |
| 1206 | Gemma-2 | | Coverage | 97.00 | 90.48 | 87.04 | 100 | 95.44 |
| 1207 | | Ours | Fraction | 58.99 | 10.63 | 13.67 | 16.71 | 100 |
| 208 | | Ours | Acc. Before | 97.00 | 59.52 | 44 44 | 31.82 | 74 94 |
| 1209 | | | Acc. After | 97.00 | 66.67 | 53.70 | 31.82 | 76.96 |
| 1210 | | - | Coverage | 91.30 | 85 71 | 86 79 | 100 | 95.95 |
| 1211 | | Logits | Fraction | 11.50 | 8.86 | 13.42 | 66.08 | 100 |
| 212 | | Logits | Acc. Boforo | | 74.20 | 67.02 | 42.53 | 54.43 |
| 213 | | | Acc. After | 01.30 | 82.86 | 67.92 | 42.53 | 55 10 |
| 1214 | Llama-3 | | Covorago | 00.72 | 82.00 | 07.92 | 100 | 02.41 |
| 1215 | | 0 | Erection | 90.72 | 17.22 | 09.09 | 25.70 | 100 |
| 1216 | | Ours | Fraction | 24.50 | 17.22 | 42.55 | 35.70 | 54.42 |
| 217 | | | Acc. Before | 90.72 | 60.29 | 42.70 | 34.04 | 54.43 |
| 218 | | - | Acc. After | 90.72 | 63.24 | 44.94 | 34.04 | 55.44 |
| 1219 | | | Coverage | 98.65 | 90.54 | 94.05 | 100 | 96.71 |
| 1220 | | Logits | Fraction | 18.73 | 18.73 | 21.27 | 41.27 | 100 |
| 1221 | | | Acc. Before | 98.65 | 83.78 | 65.48 | 52.76 | 69.87 |
| 1222 | Phi-3 | | Acc. After | 98.65 | 81.08 | 69.05 | 52.76 | 70.13 |
| 1223 | | | Coverage | 96.75 | 95.31 | 92.86 | 100 | 96.71 |
| 1224 | | Ours | Fraction | 31.14 | 16.20 | 21.27 | 31.39 | 100 |
| 1005 | | | | | | | | |
| 1225 | | | Acc. Before | 96.75 | 82.81 | 58.33 | 44.35 | 69.87 |
| 1225 | | | Acc. Before Acc. After | 96.75 | 82.81 81.25 | 58.33 59.52 | 44.35 44.35 | 69.87 69.87 |

Table 14: Results for CROQ on the TruthfulQA dataset with 4 response options.

| Model | Score | Set Size | 1 | 2 | 3 | 4 | Overall |
|---------|--------|-------------|--------|-------|-------|-------|---------|
| | | Coverage | 95.71 | 95.71 | 92.86 | 100 | 95.68 |
| | Logits | Fraction | 89.84 | 8.18 | 1.64 | 0.35 | 100 |
| | Ū. | Acc. Before | 95.71 | 74.29 | 78.57 | 33.33 | 93.46 |
| | | Acc. After | 95.71 | 71.43 | 71.43 | 33.33 | 93.11 |
| Gemma-2 | | Coverage | 95.45 | 95.00 | 100 | 0 | 95.44 |
| | Ours | Fraction | 94.98 | 4 67 | 0.35 | 0 | 100 |
| | ouis | Acc. Before | 95.45 | 57 50 | 33 33 | 0 | 93.46 |
| | | Acc After | 95.15 | 57.50 | 66 67 | 0 | 93.57 |
| | | Coverage | 06.81 | 08.30 | 100 | 0 | 97.08 |
| | T : 4- | Enaction | 0.01 | 14.40 | 1.40 | 0 | 100 |
| | Logits | Fraction | 04.11 | 14.49 | 1.40 | 0 | 01.47 |
| | | Acc. Before | 96.81 | 62.90 | 66.67 | 0 | 91.47 |
| Llama-3 | | Acc. After | 96.81 | 66.13 | 66.67 | 0 | 91.94 |
| | | Coverage | 96.66 | 97.60 | 100 | 100 | 96.85 |
| | Ours | Fraction | 84.00 | 14.60 | 1.29 | 0.12 | 100 |
| | | Acc. Before | 96.66 | 64.00 | 63.64 | 100 | 91.47 |
| | | Acc. After | 96.66 | 68.80 | 36.36 | 100 | 91.82 |
| | | Coverage | 95.47 | 93.44 | 100 | 0 | 95.33 |
| | Logits | Fraction | 92.76 | 7.13 | 0.12 | 0 | 100 |
| | | Acc. Before | 95.47 | 59.02 | 0 | 0 | 92.76 |
| DL: 2 | | Acc. After | 95.47 | 55.74 | 100 | 0 | 92.64 |
| Phi-3 | | Coverage | 95.81 | 94.03 | 100 | 0 | 95.68 |
| | Ours | Fraction | 91.94 | 7.83 | 0.23 | 0 | 100 |
| | | Acc. Before | 95.81 | 56.72 | 50.00 | 0 | 92.64 |
| | | Acc. After | 95.81 | 55.22 | 50.00 | 0 | 92.52 |
| | | | 1 2000 | 50.22 | 20100 | • | |

Table 15: Results for CROQ on the ToolAlpaca dataset with 4 response options.

Table 17: Results for CROQ on the ToolAlpaca dataset with 15 response options and Gemma-2.

| Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
|--------|-------------|-------|-------|-------|-------|-------|-------|-------|------|---|----|----|----|----|----|----|---------|
| | Coverage | 95.73 | 96.98 | 96.21 | 100 | 100 | 80.00 | 100 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 96.50 |
| Logits | Fraction | 41.00 | 34.81 | 15.42 | 5.26 | 2.57 | 0.58 | 0.23 | 0.12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| | Acc. Before | 95.73 | 81.54 | 59.85 | 57.78 | 50.00 | 40.00 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 81.43 |
| | Acc. After | 95.73 | 86.91 | 75.76 | 84.44 | 68.18 | 60.00 | 50.00 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 87.85* |
| | Coverage | 96.10 | 95.00 | 97.80 | 100 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 96.03 |
| Ours | Fraction | 50.93 | 35.05 | 10.63 | 3.04 | 0.35 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| | Acc. Before | 96.10 | 72.33 | 57.14 | 30.77 | 33.33 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 81.43 |
| | Acc. After | 96.10 | 82.67 | 80.22 | 65.38 | 66.67 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 88.67* |

Table 18: Results for CROQ on the ToolAlpaca dataset with 15 response options and Llama-3 model.

| 350 | | | | | | | | | | | | | | | | | | |
|-----|--------|-------------|-------|-------|-------|-------|-------|------|------|---|---|----|----|----|----|----|----|---------|
| 351 | | | | | | | | | | | | | | | | | | |
| 352 | | | | | | | | | | | | | | | | | | |
| 353 | | | | | | | | | | | | | | | | | | |
| 354 | | | | | | | | | | | | | | | | | | |
| 355 | | | | | | | | | | | | | | | | | | |
| 356 | | | | | | | | | | | | | | | | | | |
| 357 | | | | | | | | | | | | | | | | | | |
| 358 | | | | | | | | | | | | | | | | | | |
| 359 | | | | | | | | | | | | | | | | | | |
| 360 | | | | | | | | | | | | | | | | | | |
| 861 | | | | | | | | | | | | | | | | | | |
| 362 | | | | | | | | | | | | | | | | | | |
| 363 | | | | | | | | | | | | | | | | | | |
| 364 | | | | | | | | | | | | | | | | | | |
| 365 | | | | | | | | | | | | | | | | | | |
| 366 | | | | | | | | | | | | | | | | | | |
| 867 | | | | | | | | | | | | | | | | | | |
| 368 | | | | | | | | | | | | | | | | | | |
| 369 | | | | | | | | | | | | | | | | | | |
| 370 | | | | | | | | | | | | | | | | | | |
| 371 | | | | | | | | | | | | | | | | | | |
| 372 | Score | Set Size | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | Overall |
| 373 | | Coverage | 97.93 | 98.67 | 98.89 | 100 | 100 | 100 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 98.36 |
| 374 | Logits | Fraction | 50.70 | 35.16 | 10.51 | 2.69 | 0.70 | 0.12 | 0.12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| 375 | | Acc. Before | 97.93 | 79.73 | 62.22 | 52.17 | 50.00 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 85.98 |
| 376 | | Acc. After | 97.93 | 86.71 | 66.67 | 56.52 | 66.67 | 0 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 89.25* |
| 377 | | Coverage | 97.76 | 96.13 | 98.46 | 93.33 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 97.20 |
| 378 | Ours | Fraction | 57.36 | 33.18 | 7.59 | 1.75 | 0.12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| 379 | | Acc. Before | 97.76 | 72.89 | 64.62 | 46.67 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 85.98 |
| 380 | | Acc. After | 97.76 | 82.75 | 69.23 | 60.00 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 89.95* |
| | | | | | | | | | | | | | | | | | | |

| Table 19: Results for CROQ on the ToolAlpaca data | aset with 15 response options and Phi-3 model. |
|---|--|
|---|--|

¹⁴⁰⁴ C CALCULATION OF STATISTICAL SIGNIFICANCE

1405

1422

All our statistical significance results are based on paired sample t-tests at level $\alpha = 0.05$ of the null hypothesis that the difference under consideration is 0. The relevant differences are the differences in set sizes or coverage values using logits vs. our CP-OPT scores (Table 1), and the differences in accuracy before and after applying the CROQ procedure (all other tables except for Table 20). This is equivalent to constructing 95% confidence intervals for the differences and marking results as significant whenever the corresponding confidence intervals exclude 0. We used paired rather than unpaired tests to account for the fact that each pair of values was measured on the same test set item.

Note that paired t-tests, like paired z-tests, assume that sample means are approximately normally distributed, which holds in our setting due to the central limit theorem and the relatively large sizes of the test sets. (The central limit theorem is often invoked to justify approximate normality when sample sizes are larger than 30.) At our sample sizes, t-tests are almost identical to z-tests, but they are very slightly more conservative.

For the CROQ results, hypothesis tests were conducted to compare overall accuracy before and after the CROQ procedure. Tests were not conducted to compare accuracy conditional on each possible set size, since many set sizes have small associated samples which results in little power to detect differences.

1423 D EXAMPLE QUESTIONS AND PROMPTS

1425 D.1 MMLU 1426

1427 Dataset Description

1428 **MMLU** (Hendrycks et al., 2021) is a popular benchmark dataset for multiple choice questions 1429 (MCQs) from 57 domains including humanities, math, medicine, etc. In the standard version, each 1430 question has 4 options, we create two augmented versions with 10 and 15 options for each question 1431 by adding options from other questions on the same topic. We ensure there is no duplication in 1432 options. The standard dataset has very little training points, so we randomly draw 30%, and 10% of 1433 the points from the test split and include them in the training set and validation set respectively. Note, 1434 that we remove these points from the test set. The resulting splits have 4.5k, 2.9k, and 8.4k points in 1435 the train, validation, and test splits.

- 1436 Dataset Examples
- The following is an example of an MCQ prompt in the CP-OPT format.
- 1439 Llama 3 Prompt:

| 441 | This question refers to the following information. |
|-----|---|
| 442 | In order to make the title of this discourse generally intelligible, I have translated the term |
| 443 | "Protoplasm," which is the scientific name of the substance of which I am about to speak, by |
| 444 | the words "the physical basis of life." I suppose that, to many, the idea that there is such a |
| 445 | thing as a physical basis, or matter, of life may be novel-so widely spread is the conception |
| 446 | of life as something which works through matter Thus the matter of life, so far as we |
| 447 | know it (and we have no right to speculate on any other), breaks up, in consequence of that |
| 448 | continual death which is the condition of its manifesting vitality, into carbonic acid, water, |
| 449 | and nitrogenous compounds, which certainly possess no properties but those of ordinary |
| 450 | matter. |
| 451 | Thomas Henry Huyley "The Physical Basis of Life" 1868 From the passage one may infer |
| 452 | that Huxley argued that "life" was |
| 453 | |
| 454 | A. essentially a philosophical notion |
| 455 | |
| 456 | B. a force that works through matter |
| 457 | |
| | |

| | C. merely a property of a certain kind of matter |
|----------|--|
| | D. a supernatural phenomenon |
| | the correct answer is |
| | |
| i | 3 Prompt: |
| | < user > This question refers to the following information. In order to make the title of this discourse generally intelligible, I have translated the "Protoplasm," which is the scientific name of the substance of which I am about to spea |
| | the words "the physical basis of life." I suppose that, to many, the idea that there is su thing as a physical basis, or matter, of life may be novel-so widely spread is the concep of life as something which works through matter Thus the matter of life, so far a know it (and we have no right to speculate on any other), breaks up, in consequence of |
| | continual death which is the condition of its manifesting vitality, into carbonic acid, v and nitrogenous compounds, which certainly possess no properties but those of ord matter. |
| | Thomas Henry Huxley, "The Physical Basis of Life," 1868 From the passage, one may that Huxley argued that "life" was |
| | A. essentially a philosophical notion |
| | B. a force that works through matter |
| | C. merely a property of a certain kind of matter |
| | D. a supernatural phenomenon |
| | < end > < assistant > the correct answer is |
| | male of the CDOO mineline on the MMI II detect where the correct ensurer is only a |
| (a 01 | npt revision. |
| | Initial Promot. |
| | The best explanation for drug addiction, according to Shapiro, appeals to |
| | A. one's individual mindset and social setting. |
| | C. one's genetic profile, which explains why some people have "addictive personalities |
| | D. specific psychological disorders such as obsessive-compulsive disorder. the correct answer is |
| | Output: |
| | Prediction: B. the pharmacological effects of drug use (e.g., withdrawal). Prediction Set: {A, B} |
| | Revised Prompt: |
| | The best explanation for drug addiction, according to Shapiro, appeals to |
| | |
| | |

| - 1 11 | the correct answer is |
|--|---|
| | Output: Prediction: A. one's individual mindset and social setting. |
| ſ | Initial Prompt: |
| | Answering multiple-choice questions is often easier than answering fill-in or completion questions, because multiple choice questions |
| | A. provide more retrieval cues |
| | B. enhance retention of information C. check memorization rather than critical thinking D are definitional rather than concentual |
| | the correct answer is |
| | Output: Prediction: C. check memorization rather than critical thinking |
| | Prediction Set: {A, C} |
| | Revised Prompt: |
| | Answering multiple-choice questions is often easier than answering fill-in or completion questions, because multiple choice questions |
| | A provide more retrieval cues |
| | B. check memorization rather than critical thinking |
| | the correct answer is |
| | Output: |
| U | Prediction: A. provide more retrieval cues |
| D | Prediction: A. provide more retrieval cuesD.2 TRUTHFULQA |
| | Prediction: A. provide more retrieval cues D.2 TRUTHFULQA Pataset Details |
| U D T L th T is the option | Prediction: A. provide more retrieval cues D.2 TRUTHFULQA Pataset Details the TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LLD the dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "Meargets", and resample questions using additional correct responses from "MC 2 Targets", so that ngle truthful answer is included in the response set. The dataset was split randomly by question that there was no overlap between splits. After resampling using the "MC 2 Targets", the train spontains 1,745 questions, the calibration split contains 695 questions, and the test split contains 2 uestions. Since we consider questions with a fixed number of response options, answers were eit pownselected, or additional responses from different, random questions from the same split wisserted. |
| U D T L th T sith c q q d in D | Prediction: A. provide more retrieval cues D.2 TRUTHFULQA Pataset Details the TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LL the dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "M argets", and resample questions using additional correct responses from "MC 2 Targets", so the ngle truthful answer is included in the response set. The dataset was split randomly by question that there was no overlap between splits. After resampling using the "MC 2 Targets", the train s contains 1,745 questions, the calibration split contains 695 questions, and the test split contains 2 uestions. Since we consider questions with a fixed number of response options, answers were eithownselected, or additional responses from different, random questions from the same split wasterted. |
| | Prediction: A. provide more retrieval cues 0.2 TRUTHFULQA Pataset Details the TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LL the dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "M argets", and resample questions using additional correct responses from "MC 2 Targets", so th ngle truthful answer is included in the response set. The dataset was split randomly by question there was no overlap between splits. After resampling using the "MC 2 Targets", the trains so pontains 1,745 questions, the calibration split contains 695 questions, and the test split contains 3 uestions. Since we consider questions with a fixed number of response options, answers were eit ownselected, or additional responses from different, random questions from the same split was retacted. Pataset Examples Initial Prompt: |
| | Prediction: A. provide more retrieval cues P.2 TRUTHFULQA Pataset Details he TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LL he dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "Margets", and resample questions using additional correct responses from "MC 2 Targets", so th ngle truthful answer is included in the response set. The dataset was split randomly by question nat there was no overlap between splits. After resampling using the "MC 2 Targets", the train s pontains 1,745 questions, the calibration split contains 695 questions, and the test split contains 1 vestions. Since we consider questions with a fixed number of response options, answers were eit ownselected, or additional responses from different, random questions from the same split westered. vestaset Examples Initial Prompt: In France they speak French. In England they speak English. In Wales they speak Welsh. In Australia they speak what? |
| C D T L th T si th co qu du irr | Prediction: A. provide more retrieval cues P.2 TRUTHFULQA Pataset Details he TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LL ne dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "M argets", and resample questions using additional correct responses from "MC 2 Targets", so the ngle truthful answer is included in the response set. The dataset was split randomly by question that there was no overlap between splits. After resampling using the "MC 2 Targets", the train sontains 1,745 questions, the calibration split contains 695 questions, and the test split contains 3 uestions. Since we consider questions with a fixed number of response options, answers were eit ownselected, or additional responses from different, random questions from the same split wasterted. Pataset Examples Initial Prompt: In France they speak French. In England they speak English. In Wales they speak Welsh. In Australia they speak Australian. B. Australia |
| | Prediction: A. provide more retrieval cues P.2 TRUTHFULQA Pataset Details The TruthfulQA dataset (Lin et al., 2022) contains 817 questions designed to evaluate truthfulness LM responses. Although the dataset is primarily use to evaluate open responses generated by LL the dataset is also prepared in an MCQA format. We perform evaluation of MCQA on the "M argets", and resample questions using additional correct responses from "MC 2 Targets", so th ngle truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the truthful answer is included in the response set. The dataset was split randomly by question the spontains 1,745 questions, the calibration split contains 695 questions, and the test split contains is uestions. Since we consider questions with a fixed number of response options, answers were eit ownselected, or additional responses from different, random questions from the same split westered. The truthful Prompt: In France they speak French. In England they speak English. In Wales they speak Welsh. In Australia they speak Australian. B. Australia. C. Australians speak Australian. C. Australians speak Australian. |

| _ | |
|---|---|
| | the correct answer is |
| | |
| | Output: |
| | Prediction: B. Australian. |
| | Prediction Set: {B, C, D} |
| | Derived Descents |
| | Kevised Prompt: In France they speak Franch In England they speak English In Wales they speak Waleh In |
| | Australia they speak ritericii. In England they speak English. In wates they speak weish. In |
| | Australia tiley speak what? |
| | A Australian |
| | B. Australians speak Australian. |
| | C. Australians speak English. |
| | the correct answer is |
| | |
| | Output: |
| | Prediction: C. Australians speak English. |
| | |
| | |
| | Initial Prompt: |
| | What are the richest countries in South America by GDP per capita? |
| | |
| | A. Venezuela, Chile, Uruguay and Argentina. |
| | B. The richest countries in South America by GDP per capita are Chile, Uruguay, Venezuela |
| | and Argentina. |
| | D. The richest countries in South America by GDP per capita are Chile and Uruguay |
| | the correct answer is |
| | |
| | Output: |
| | Prediction: B. The richest countries in South America by GDP per capita are Chile, Uruguay, |
| | Venezuela and Argentina. |
| | Prediction Set: {B, D} |
| | |
| | Revised Prompt: |
| | What are the richest countries in South America by GDP per capita? |
| | A The sightest countries in South America by ODD |
| | A. The richest countries in South America by GDP per capita are Chile, Uruguay, Venezuela |
| | and Argenuna. B. The richest countries in South America by CDP per conite are Chile and Uryguey |
| | the correct answer is |
| | |
| | Output: |
| | Prediction: B. The richest countries in South America by GDP per capita are Chile and |
| | Uruguay. |
| | |
| | |
| - | 3 TOOLALPACA |
| | taset Details |
| | |
| (| DAlpaca (Tang et al., 2023) contains 3.9k tool-use instances from a multi-agent simulation |
| , | ironment. The dataset was reformulated from a general purpose tool-selection task to an MC |
| | c. The LLW is prompted with an instruction and an API description and must select the corre |
| | ction based on the function name and a brief description. |
| | |

We filter out APIs that had an error in generating documentation, instances where a ground truth label
 was missing, and instances that required multiple, sequential function calls. After filtering, 2,703
 MCQ examples remain. The train split contains 856 synthetic examples, the calibration split contains

1620
1621 774 synthetic validation examples, and the test split contains 1040 real and synthetic API examples.
Splits are created to ensure no overlap in APIs occur. We follow a similar resampling procedure as used for TruthfulQA, so that the number of response options is fixed. Arguments are stripped from the provided function call so that the MCQ task was focuses towards tool selection, a critical task in the more general tool usage problem.

Dataset Examples

Initial Prompt:

Given the API Bugsnax, and the following instruction, "I need more information on a character called "Chandlo." Can you tell me about his role in the game, his description, location, and any quests associated with him?" Which of the following functions should you call?

A. searchItems Search for items based on a keyword or partial name.

B. getCharacterInfo Retrieve detailed information about a specific character in the game.

C. searchCharacters Search for characters based on a keyword or partial name.

D. getItemInfo Retrieve detailed information about a specific item in the game. the correct answer is

Output:

Prediction: C. searchCharacters Search for characters based on a keyword or partial name. Prediction Set: $\{B, C\}$

Revised Prompt:

Given the API Bugsnax, and the following instruction, "I need more information on a character called "Chandlo." Can you tell me about his role in the game, his description, location, and any quests associated with him?" Which of the following functions should you call?

A. getCharacterInfo Retrieve detailed information about a specific character in the game.B. searchCharacters Search for characters based on a keyword or partial name.the correct answer is

Output:

Prediction: A. getCharacterInfo Retrieve detailed information about a specific character in the game.

| 1654 |
|------|
| 1655 |
| 1656 |
| 1657 |
| 1658 |
| 1659 |
| 1660 |
| 1661 |
| 1662 |

Initial Prompt:

Given the API Cataas, and the following instruction, "I'm feeling a bit down and could use a pick-me-up. Could you find me a random picture of a cat? Make sure it's a cute one!" Which of the following functions should you call?

- A. getRandomCat Get random cat
- B. tags Will return all tags
- C. findCatById Get cat by id
- D. findCatByTag Get random cat by tag
- the correct answer is

Output:

Prediction: D. findCatByTag Get random cat by tag Prediction Set: {A, D}

1671 Revised Prompt:

Given the API Cataas, and the following instruction, "I'm feeling a bit down and could use a pick-me-up. Could you find me a random picture of a cat? Make sure it's a cute one!" Which

of the following functions should you call?

A. getRandomCat Get random cat B. findCatByTag Get random cat by tag the correct answer is

Output:

Prediction: A. getRandomCat Get random cat

1684 E HYPERPARAMETER SETTINGS

| Model | Dataset | # Opt. | λ | lr | weight decay | batch size |
|---------|------------|--------|-----------|------|--------------|------------|
| | | 4 | 5.0 | 1e-5 | 1e-7 | 128 |
| | MMLU | 10 | 0.1 | 1e-5 | 1e-9 | 128 |
| | | 15 | 1.0 | 1e-5 | 1e-9 | 256 |
| | | 4 | 0.5 | 1e-4 | 1e-6 | 128 |
| Gemma-2 | ToolAlpaca | 10 | 5.0 | 1e-4 | 1e-6 | 128 |
| | | 15 | 5.0 | 1e-4 | 1e-6 | 256 |
| | | 4 | 0.1 | 1e-4 | 1e-8 | 128 |
| | TruthfulQA | 10 | 0.1 | 1e-4 | 1e-7 | 128 |
| | | 15 | 5.0 | 1e-4 | 1e-6 | 128 |
| | | 4 | 1.0 | 5e-6 | 1e-9 | 128 |
| | MMLU | 10 | 0.5 | 1e-5 | 1e-8 | 128 |
| | | 15 | 0.5 | 5e-6 | 1e-8 | 256 |
| | | 4 | 0.5 | 1e-5 | 1e-8 | 128 |
| Llama-3 | ToolAlpaca | 10 | 1.0 | 5e-6 | 1e-7 | 128 |
| | | 15 | 0.5 | 1e-5 | 1e-9 | 128 |
| | | 4 | 0.5 | 1e-5 | 1e-8 | 128 |
| | TruthfulQA | 10 | 0.5 | 1e-4 | 1e-9 | 128 |
| | | 15 | 0.5 | 1e-5 | 1e-8 | 128 |
| | | 4 | 0.5 | 5e-6 | 1e-7 | 128 |
| | MMLU | 10 | 1.0 | 1e-5 | 1e-9 | 128 |
| | | 15 | 2.0 | 5e-6 | 1e-7 | 128 |
| | | 4 | 2.0 | 1e-5 | 1e-8 | 128 |
| Phi-3 | ToolAlpaca | 10 | 0.1 | 1e-5 | 1e-9 | 128 |
| | | 15 | 5.0 | 1e-5 | 1e-8 | 128 |
| | | 4 | 0.5 | 1e-5 | 1e-8 | 128 |
| | TruthfulQA | 10 | 10.0 | 5e-5 | 1e-8 | 128 |
| | | 15 | 0.1 | 1e-4 | 1e-10 | 128 |

1721Table 20: Hyperparameter settings for our score function learning procedure CP-OPT in our experi-
ments. For all settings we use SGD with momentum 0.9, learning rate (lr) as in the table with learning
rate decay, number of epochs = 1000 and $\beta = 1.0$.